

Published in final edited form as:


J Org Chem. 2009 October 2; 74(19): 7383–7388. doi:10.1021/jo901218g.

Solid Phase Synthesis of Peptidyl Thioacids Employing a 9-Fluorenylmethyl Thioester-Based Linker in Conjunction with Boc Chemistry

 David Crich^{a,b,*} and Kasinath Sana^b
^aCentre Scientifique de Gif-sur-Yvette, Institut de Chimie des Substances Naturelles, avenue de la Terrasse, 91198 Gif-sur-Yvette, France

^bDepartment of Chemistry, Wayne State University, 5101 Cass Avenue, Detroit, MI 48202

Abstract



A method for the synthesis of peptidyl thioacids is described based on the use of the *N*-[9-(thiomethyl)-9*H*-fluoren-2-yl]succinamic acid and cross-linked aminomethyl polystyrene resin. The method employs standard Boc chemistry SPPS techniques in conjunction with 9-fluorenylmethoxycarbonyl protection of side chain alcohols and amines, and 9-fluorenylmethyl protection of side chains acids and thiols. Cleavage from the resin is accomplished with piperidine, which also serves to remove the side chain protection and avoids the HF conditions usually associated with the resin cleavage stage of Boc chemistry SPPS. The so-obtained thioacids are converted to simple thioesters in high yield by standard alkylation according to well-established methods.

Introduction

Thioacids [RC(=O)SH]¹ are a fascinating but underappreciated class of compounds with a unique reactivity profile² that arises in part from their p*K*_a and the consequent ability of the conjugate base to act in a highly selective manner as nucleophile in aqueous media at pH3-6. Not surprisingly therefore most applications of thioacids have been in the field of peptide chemistry where they have been employed in amide bond forming reactions either directly³ or indirectly through their facile conversion to thioesters, key intermediates in a variety of chemical and enzymic amide ligation processes.⁴

The instability of thioesters toward the typical conditions of 9-fluorenylmethoxycarbonyl (Fmoc) chemistry solid phase peptide synthesis (SPPS), particularly the treatment with organic bases employed in the cleavage of Fmoc groups, led to an initial reliance on *tert*-butyloxycarbonyl-based (Boc) chemistry for the synthesis of peptidyl thioesters,^{4o,4,5} but the HF conditions typically required for cleavage from the resin following Boc chemistry SPPS limit the use of this chemistry. Fmoc chemistry-based methods have subsequently been developed that replace piperidine in the Fmoc removal step by cocktails of 1-

dcrich@icsn.cnrs-gif.fr.

Supporting Information Available. Complete experimental details, copies of the ¹H and ¹³C-NMR spectra of compounds 2–5, 7–12, and 22–30, and HPLC traces and mass spectra of peptidyl thioacids 22–27.

methylpyrrolidine, hexamethylenimine and 1-hydroxybenzotriazole but it has been found that these methods are prone to racemization at the thioester position.⁶ The backbone amide linker (BAL) strategy enables Fmoc chemistry SPPS with subsequent incorporation of a C-terminal thioester prior to cleavage from the resin but requires careful control of reaction conditions to avoid epimerization on introduction of the thioester to the C-terminal end of the peptide chain.⁷ To circumvent these problems numerous methods have been developed according to which, after completion of the peptide synthesis by Boc or Fmoc methods, the linker to the resin is activated in such a way as to permit its displacement by a thiol or thiolate resulting in the liberation of the peptide in the form of the desired thioester or thioacid.⁸ A variant on the BAL strategy, that avoids the epimerization problem, carries a C-terminal trithioorthoester through the Fmoc chemistry SPPS sequence before converting it to the required thioester by controlled hydrolysis.⁹ More recently, a number of strategies have been developed in which thioesters are generated by O-S or N-S shifts of mercapto esters and mercapto amides following unmasking of a protected thiol group.¹⁰ Despite the considerable ingenuity that has been deployed in the development of the above methods, none combine the directness that obviously results from the use of a simple C-terminal thioester-based linker with a method for release from the resin that avoids the use of HF. Previously we introduced the 9-fluorenylmethyl thioesters from which thioacids are liberated by simple treatment with piperidine, that is, under the conditions usually employed for the cleavage of Fmoc groups in Fmoc chemistry SPPS.¹¹ We conceived that a linker based on the 9-fluorenylmethyl thioester would be compatible with the general conditions of Boc chemistry SPPS and that following peptide assembly treatment of the resin with piperidine would release the Boc-protected peptide into solution in the form of a C-terminal thioacid that could be readily transformed into a thioester by simple alkylation. Of essence, this method, whose reduction to practice we report here, employs conditions no more forcing than those encountered in standard Boc and Fmoc chemistry SPPS protocols and circumvents the terminal HF treatment that limits most Boc chemistry SPPS methods. We further conceived that the utility of such this method would be enhanced by the application of a side chain protection strategies involving either a third orthogonal system enabling retention of side chain protecting groups post cleavage,¹² or a system according to which all protecting groups would be removed concomitantly with cleavage of the thioacid from the resin, depending on the ultimate application envisaged for the peptidyl thioacid.

Results and Discussion


The mercapto functionalized linker, *N*-[9-(tritylthiomethyl)-9*H*-fluoren-2-yl]succinamic acid (**5**) was prepared from commercially available 9*H*-fluoren-2-amine as shown in Scheme 1. The synthesis began with the conversion of 9*H*-fluoren-2-amine to corresponding hydroxyl functional compound **1** following a literature procedure¹³ involving formylation followed by reduction. Tosylation of **1** under standard conditions gave the sulfonate **2**, from which the amine **3** was liberated with trifluoroacetic acid. Reaction of **3** with succinic anhydride provided the hemisuccinate **4** that was subjected to treatment with tritylmercaptan and Hunig's base to give the protected linker **5** in excellent yield (Scheme 1). Treatment of **5** with diisopropyl carbodiimide and *N*-hydroxybenzotriazole in DMF gave an activated intermediate that was allowed to react with 1% divinylbenzene cross linked aminomethylpolystyrene resin (0.41 mmol/g loading). Following washing with DMF the functionalized resin was exposed to a 50% solution of TFA in dichloromethane to yield the desired resin-bound 9-fluorenylmethylthiol derivative **6** (Scheme 1). The attachment of linker **5** to the aminomethylpolystyrene resin was also accomplished in a satisfactory manner with the *O*-benzotriazolyl tetramethyluronium hexafluorophosphate (HBTU) reagent¹⁴ with the aid of diisopropylethylamine as base.

The preparation of a series of suitably protected amino acids was then undertaken. Thus, *N*-*tert*-butoxycarbonyl *L*-serine and the analogous *L*-threonine and *L*-tyrosine derivatives were converted to their allyl esters with potassium carbonate and then to the 9-fluorenylmethyl

carbonates with Fmoc chloride and pyridine. A combination of palladium(II) acetate, triphenylphosphine and phenylsilane was the reagent of choice for the liberation the allyl esters required to complete this short sequence (Scheme 2).

Following a literature protocol¹⁵ treatment of powdered L-aspartic and L-glutamic acids with triethylborane in THF at reflux for 24 h, then with 9-fluorenylmethanol, dicyclohexylcarbodiimide and 4-dimethylaminopyridine, and finally with gaseous hydrogen chloride gave the mono esters **13** and **14**. These HCl salts were then converted to the *N*-Boc derivatives in the standard manner (Scheme 3).

The corresponding mono allyl esters **17** and **18** were accessed according to the literature method¹⁶, as were the 9-fluorenylmethyl thioether of Boc-L-cysteine **19**,¹⁵ the Fmoc-protected L-arginine derivative **20**,¹⁷ and Alloc-protected L-histidine **21**.¹⁸



With all building blocks in hand attention was turned to SPPS using standard Boc techniques with DIC/HOBt as the coupling agent and TFA to liberate the N-terminus of the growing chains from their Boc derivatives. A number of peptides were assembled in this manner as set out in Table 1 (Entries 1–5). As with the preparation of the resin-bound thiol **6** (Scheme 1), this methodology is not limited to carbodiimide chemistry but is perfectly adaptable to the other methods as evidenced by the application of the HBTU protocol (Table 1, entry 6). After completion of the on-resin procedure, treatment with a solution of piperidine in DMF or, to enable direct loading of the reaction mixture to the HPLC column, acetonitrile released the desired thioacids protected at the N-terminal ends in the form of the Boc derivatives,¹⁹ which were typically obtained with a high degree of purity as determined by ESI-TOF and HPLC methods. Similar treatment of individual beads was used to systematically monitor the individual reaction steps during the course of the peptide assembly sequence. Although mass spectrometry was the method of choice for monitoring these SPPS reactions, the Kaiser ninhydrin test also performed in a perfectly satisfactory manner for both the coupling and Boc removal steps. While, the thioacid **22** was a simple model tetrapeptidyl thioacid (Table 1, entry 1), the peptidyl thioacids **23**, **24**, **25** and **27** are all natural sequences. The sequences of thioacids **23** and **24** (Table 1, entries 2 and 3) were selected from the Glucagon-like peptide-1 (GLP-1)²⁰ and represent segments 7–16 and 17–26, respectively, of that peptidyl hormone. Peptide **25** (Table 1, entry 4) represents the 94–101 segment of Human Secretory Phospholipase A₂

(hsPLAA₂),²¹ and peptide **27** (Table 1, entry 5) is the 65–84 unit of Human Parathyroid Hormone (hPTH).²²

The ¹H and ¹³C-NMR spectra of the crude tetrapeptidyl thioacid **22** as obtained on simple release from the resin, acidification with HCl, and drying are presented in the supporting information (Figures 1 and 2) to illustrate the high degree of purity typically obtained by this method. In a similar vein the ESI-TOF mass spectrum of the peptidyl thioacid **27**, immediately after release from the resin and prior to purification by HPLC is presented in the supporting information as Figure 3.

Particular attention is drawn to entries 4 and 5 of Table 1 in which the C-terminal amino acid is asparagine and glutamine, respectively. The amino acid building blocks for these residues were employed without protection of the side chain amide functionality and it is noteworthy that cyclization of these amides onto the resin-bound thioester with premature peptide release in the form of an imide did not occur to any significant extent.²⁴ In general, the strategy of employing Fmoc protection for side chain amines and hydroxyl groups, coupled with the protection of side chain carboxylates and thiols ensures clean chemistry, while eliminating the need for extra deprotection steps pre- or post cleavage of the peptidyl thioacid from the resin. Nevertheless, should the retention of side chain protection be required on cleavage from the resin this may be conveniently accomplished through the use of building blocks whose side chains are covered with either the allyl or allyloxycarbonyl system¹² depending on the case (Table 1, entries 2 and 3).

By way of example two of the peptidyl thioacids obtained in this manner were converted to the corresponding *S*-benzyl thioesters by simple alkylation with benzyl bromide and *sym*-collidine in DMF (Scheme 4).^{5c}

Finally, as a demonstration of the broad scope of the chemistry of thioacids, a single decapeptidyl thioacid was subjected to reaction with a sulfonyl azide, under conditions described by the Williams and Liskamp groups²⁵ for much simpler substrates, resulting in the isolation of a C-terminal sulfonamide (Scheme 5).

Overall, we describe the successful implementation of a straightforward method for the SPPS of peptidyl thioacids using standard Boc chemistry with release from the resin using conditions typically used for Fmoc removal during the course of Fmoc chemistry SPPS. In conjunction with the protection of side chain amino and hydroxyl groups as Fmoc carbonates, and of side chain acids and thiols as 9-fluorenylmethyl esters and thioethers this chemistry provides a very convenient and mild means of access to peptidyl thioacids, and, by simple alkylation, of their *S*-esters.

Experimental

General

Unless otherwise stated ¹H and ¹³C NMR were recorded in CDCl₃ solution and optical rotations in CHCl₃ solutions. All organic extracts were dried over sodium sulfate, and concentrated under aspirator vacuum. Chromatographic purifications were carried out over silica gel. All peptide thioacid syntheses were carried out on a 0.1 mmol scale employing 1% DVB cross linked aminomethyl polystyrene resin (244 mg, resin loading 0.41 mmol/g) in a 10 mL manual synthesizer glass reaction vessel with a Teflon-lined screw cap. The peptide resin was shaken during the both *N*^α-*tert*-butoxycarbonyl deprotection and coupling steps. After each coupling step, formation of the desired peptide thioacid was confirmed by cleavage of a small amount (~ 5 mg) of resin using a 20% solution of piperidine in DMF for 20 min., followed by examination by ESI- TOF mass spectrometry. Isolated yields were determined based the

theoretical yield calculated for the use of 0.1 mmol of resin with a loading of 0.41 mmol/g. These yields take no account of the aliquots removed for monitoring and are therefore minimum yields.

[2-(*tert*-Butoxycarbonylamino)-9*H*-fluoren-9-yl]methyl 4-methylbenzenesulfonate (2)

To a stirred solution of [2-(*tert*-butoxycarbonylamino)-9*H*-fluoren-9-yl]methanol¹³ (1.8 g, 5.8 mmol) and 4-methylbenzenesulfonyl chloride (1.65 g, 8.7 mmol) in CHCl₃ (20 mL) was added pyridine (0.9 mL, 11.6 mmol) at 0 °C. The reaction mixture was stirred at room temperature for 6 h. Then the organic layer was washed with 1M HCl, water, brine, dried and concentrated. Chromatographic purification using 30% ethyl acetate in hexane afforded **2** (2.42 g, 90%). Yellowish syrup; ¹H NMR (500 MHz) δ 7.78-7.76 (d, *J* = 8.0 Hz, 2H), 7.66-7.61 (dd, *J* = 8.5, 12.8 Hz, 2H), 7.57 (s, 1H), 7.51-7.50 (d, *J* = 7.5 Hz, 1H), 7.38-7.35 (t, *J* = 7.0 Hz, 2H), 7.31-7.29 (d, *J* = 8 Hz, 2H), 7.25-7.22 (t, *J* = 7.5 Hz, 1H), 6.61 (s, 1H), 4.31-4.24 (m, 2H), 4.20-4.17 (t, *J* = 7.5 Hz, 1H), 2.43 (s, 3H), 1.56 (s, 9H); ¹³C NMR (125 MHz) δ 153.0, 145.1, 143.7, 142.5, 141.3, 138.0, 136.5, 133.0, 130.1, 128.3, 128.2, 126.7, 125.3, 120.7, 119.8, 118.8, 115.6, 80.9, 72.0, 46.9, 28.6, 21.8; ESI-HRMS Calcd for C₂₆H₂₇NO₅S [M + Na]⁺: 488.1508. Found: 488.1486.

(2-Amino-9*H*-fluoren-9-yl)methyl 4-methylbenzenesulfonate (3)

To a stirred solution of **2** (2.4 g, 5.2 mmol) in CH₂Cl₂ (16 mL), was added TFA (4 mL) dropwise at 0 °C. The reaction mixture was stirred at same temperature for 20 min before it was neutralized at 0 °C by saturated aqueous NaHCO₃. Then the organic layer was washed with water, and brine, and dried and concentrated. Chromatographic purification using 40% ethyl acetate in hexane afforded **3** (1.9 g, 100%). Light yellow syrup; ¹H NMR (400 MHz) δ 7.78-7.76 (d, *J* = 8.4 Hz, 2H), 7.57-7.55 (d, *J* = 7.2 Hz, 1H), 7.50-7.48 (d, *J* = 8.0 Hz, 1H), 7.44-7.42 (d, *J* = 7.2 Hz, 1H), 7.34-7.29 (m, 3H), 7.17-7.13 (t, *J* = 6.4 Hz, 1H), 6.86 (s, 1H), 6.71-6.68 (dd, *J* = 1.6, 8.4 Hz, 1H), 4.26-4.18 (m, 2H), 4.13-4.09 (t, *J* = 7.2 Hz, 1H), 3.67 (br s, 2H), 2.41 (s, 3H); ¹³C NMR (100 MHz) δ 146.4, 145.1, 144.7, 142.0, 141.7, 133.1, 132.3, 130.1, 128.2, 128.1, 125.7, 125.1, 121.1, 119.0, 115.2, 112.2, 72.5, 46.8, 21.9; ESI-HRMS Calcd for C₂₁H₁₉NO₃S [M + H]⁺: 366.1164. Found: 366.1152.

***N*-[9-(Tosylloxymethyl)-9*H*-fluoren-2-yl]succinamic acid (4)**

To a stirred solution of **3** (1.8 g, 4.9 mmol) in THF (10 mL) was added solid succinic anhydride (590 mg, 5.9 mmol) portion wise over a period of 10 min at room temperature. The reaction mixture was stirred at room temperature for 1 h. Then the reaction mixture was concentrated and subjected to chromatographic purification using 5% methanol in dichloromethane when it afforded **4** (2.1 g, 91%). White solid, crystallized from chloroform/hexane, mp: 165.8-166.2 °C. ¹H NMR (400 MHz) δ 7.65-7.63 (m, 3H), 7.55-7.51 (m, 3H), 7.37-7.7.36 (d, *J* = 7.2 Hz, 1H), 7.26-7.2 (m, 4H), 7.14-7.11 (t, *J* = 7.2 Hz, 1H), 4.20-4.13 (m, 2H), 4.07-4.04 (t, *J* = 7.2 Hz, 1H), 2.67-2.62 (m, 4H), 2.32 (s, 3H); ¹³C NMR (100 MHz) δ 175.6, 171.1, 145.3, 143.3, 142.4, 141.1, 137.7, 137.3, 132.5, 130.1, 128.2, 128.0, 126.8, 125.1, 120.5, 120.1, 120.0, 119.8, 116.8, 72.0, 46.8, 31.7, 29.4, 21.7; ESI-HRMS Calcd for C₂₅H₂₃NO₆S [M + Na]⁺: 488.1144. Found: 488.1120.

***N*-[9-(Tritylthiomethyl)-9*H*-fluoren-2-yl]succinamic acid (5)**

To a stirred solution of **4** (2.0 g, 4.3 mmol) and triphenylmethanethiol (1.5 g, 5.4 mmol) in DMF (15 mL) was added diisopropylethylamine (1.8 mL, 10.8 mmol) at 0 °C. The reaction mixture was stirred at room temperature for 15 h, after which the DMF was removed under high vacuum and the crude mixture was dissolved in EtOAc and washed with water, and brine, and dried and concentrated. Chromatographic purification of the residue using 4% methanol in dichloromethane afforded **5** (2.23 g, 91%). Light brown solid, crystallized from chloroform/

hexane, mp: 84–85 °C. ^1H NMR (500 MHz) δ 7.80 (br s, 1H), 7.57–7.53 (m, 4H), 7.43–7.41 (m, 6H), 7.31–7.24 (m, 8H), 7.21–7.17 (m, 4H), 3.57–3.54 (t, $J = 7.0$ Hz, 1H), 2.74–2.70 (m, 3H), 2.67–2.62 (m, 3H); ^{13}C NMR (125 MHz) δ 177.5, 170.5, 147.2, 146.1, 144.9, 140.5, 137.5, 136.8, 130.0, 128.2, 127.7, 127.0, 126.7, 124.9, 120.4, 119.9, 119.7, 116.9, 67.6, 47.2, 36.1, 32.0, 29.7; ESI-HRMS Calcd for $\text{C}_{37}\text{H}_{31}\text{NO}_3\text{S} [\text{M} + \text{Na}]^+$: 592.1922. Found: 592.1892.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments


We thank the NIH (GM62160) for support of this work.

References



1. a) Kato S, Kawahara Y, Kageyama H, Yamada R, Niyomura O, Murai T, Kanda T. *J. Am. Chem. Soc.* 1996;118:1262–1267. b) Hadad CM, Rablen PR, Wiberg KB. *J. Org. Chem.* 1998;63:8668–8681.
2. a) Bauer, W.; Kühlein, K. *Methoden der Organischen Chemie*, 4th Ed; Carbonsäure und Carbonsäure Derivate. Falbe, J., editor. Vol. 1. Stuttgart: Thieme; 1985. p. 832–890. b) Niyomura O, Kato S. *Top. Curr. Chem.* 2005;251:1–12. c) Kato, S.; Murai, T. *The Chemistry of Acid Derivatives*. Patai, S., editor. Vol. 2. Chichester: Wiley; 1992. p. 803–847. d) Scheithauer S, Mayer R. *Topics in Sulfur Chemistry* 1979;4:1–373.
3. a) Blake J. *Int. Pept. Prot. Res.* 1981;17:273–274. b) Yamashiro D, Blake JF. *Int. J. Pept. Prot. Chem.* 1981;18:383–392. c) Blake J, Yamashiro D, Ramasharma K, Li CH. *Int. J. Peptide Protein Res.* 1986;28:468–476. [PubMed: 3102393] d) Yamashiro D, Li CH. *Int. J. Peptide Protein Res.* 1988;31:322–334. [PubMed: 3372135] e) Mitin YV, Zapevalova NP. *Int. J. Pept. Prot. Chem.* 1990;35:352–356. f) Høeg-Jensen T, Olsen CE, Holm A. *J. Org. Chem.* 1994;59:1257–1263.
4. a) Dawson PE, Muir TW, Clark-Lewis I, Kent SBH. *Science* 1994;266:776–779. [PubMed: 7973629] b) Dawson PE, Kent SBH. *Ann. Rev. Biochem.* 2000;69:923–960. [PubMed: 10966479] c) Yeo DSY, Srinivasan R, Chen GYJ, Yao SQ. *Chem. Eur. J.* 2004;10:4664–4672. d) Macmillan D. *Angew. Chem. Int. Ed.* 2006;45:7668–7672. e) Hackenberger CPR, Schwarzer D. *Angew. Chem. Int. Ed.* 2008;47:10030–10074. f) Kent SBH. *Chem. Soc. Rev.* 2009;38:338–351. [PubMed: 19169452] g) Flavell RR, Muir TW. *Acc. Chem. Res.* 2009;42:107–116. [PubMed: 18939858] h) Mihara H, Maeda S, Kurosaki R, Ueno S, Sakamoto S, Niidome T, Hojo H, Aimoto S, Aoyagi H. *Chem. Lett.* 1995:397–398. i) Schnolzer M, Kent SBH. *Science* 1992;256:221–225. [PubMed: 1566069] j) Baca M, Kent SBH. *Proc. Natl. Acad. Sci. U. S. A.* 1993;90:11638–11642. [PubMed: 8265601] k) Williams MJ, Muir TW, Ginsberg MH, Kent SBH. *J. Am. Chem. Soc.* 1994;116:10797–10798. l) Dawson PE, Kent SBH. *J. Am. Chem. Soc.* 1993;115:7263–7266. m) Futaki S, Sogawa K, Maruyama J, Asahara T, Niwa M. *Tetrahedron Lett.* 1997;38:6237–6240. n) Camarero JA, Pavel J, Muir TW. *Angew. Chem., Int. Ed.* 1998;37:347–349. o) Zhang L, Tam JP. *J. Am. Chem. Soc.* 1997;119:2363–2370. p) Shao Y, Lu W, Kent SBH. *Tetrahedron Lett.* 1998;39:3911–3914. q) Tam JP, Lu Y-A, Liu C-F, Shao J. *Proc. Natl. Acad. Sci. U. S. A.* 1995;92:12485–12489. [PubMed: 8618926]
5. a) Canne LE, Walker SM, Kent SBH. *Tetrahedron Lett.* 1995;36:1217–1220. b) Canne LE, Ferre-D'Amare AR, Burley SK, Kent SBH. *J. Am. Chem. Soc.* 1995;117:2998–3007. c) Lu W, Qasim MA, Kent SBH. *J. Am. Chem. Soc.* 1996;118:8518–8523. d) Hojo H, Aimoto S. *Bull. Chem. Soc. Jpn.* 1991;64:111–117. e) Kawakami T, Kogure S, Aimoto S. *Bull. Chem. Soc. Jpn.* 1996;69:3331–3338. f) Hojo H, Kwon Y, Kakuta Y, Tsuda S, Tanaka I, Hikichi K, Aimoto S. *Bull. Chem. Soc. Jpn.* 1993;66:2700–2706. g) Zhang L, Tam JP. *J. Am. Chem. Soc.* 1999;121:3311–3320. h) Li Y, Yu Y, Giulianotti M, Houghten RA. *J. Comb. Chem.* 2008;10:613–616. [PubMed: 18710293]
6. a) Li X, Kawakami T, Aimoto S. *Tetrahedron Lett.* 1998;39:8669–8672. b) Hasegawa K, Sha YL, Bang JK, Kawakami T, Akaji K, Aimoto S. *Let. Pept. Sci.* 2002;8:277–284.
7. a) Jensen KJ, Alsina J, Songster MF, Vagner J, Albericio F, Barany G. *J. Am. Chem. Soc.* 1998;120:5441–5452. b) Alsina J, Yokum TS, Albericio F, Barany G. *J. Org. Chem.* 1999;64:8761–

8769. [PubMed: 11674777] c) Alsina J, Yokum TS, Albericio F, Barany G. *Tetrahedron Lett* 2000;41:7277–7280.
8. a) Schwabacher AW, Maynard TL. *Tetrahedron Lett* 1993;34:1269–1270. b) Ingenito R, Bianchi E, Fattori D, Pessi A. *J. Am. Chem. Soc* 1999;121:11369–11374. c) Shin Y, Winans KA, Backes BJ, Kent SBH, Ellman JA, Bertozzi CR. *J. Am. Chem. Soc* 1999;121:11684–11689. d) Sweing A, Hilvert D. *Angew. Chem., Int. Ed* 2001;40:3395–3396. e) Camarero JA, Hackel BJ, De Yoreo JJ, Mitchell AR. *J. Org. Chem* 2004;69:4145–4151. [PubMed: 15176841] f) Blanco-Canosa JB, Dawson PE. *Angew. Chem., Int. Ed* 2008;47:6851–6855. g) Yamamoto N, Tanabe Y, Okamoto R, Dawson PE, Kajihara Y. *J. Am. Chem. Soc* 2008;130:501–510. [PubMed: 18085777]
9. Brask J, Albericio F, Jensen KJ. *Org. Lett* 2003;5:2951–2953. [PubMed: 12889916]
10. a) Botti P, Villain M, Manganiello S, Gaertner H. *Org. Lett* 2004;6:4861–4864. [PubMed: 15606085] b) Warren JD, Miller JS, Keding SJ, Danishefsky SJ. *J. Am. Chem. Soc* 2004;126:6576–6578. [PubMed: 15161285] c) George EA, Novick RP, Muir TW. *J. Am. Chem. Soc* 2008;130:4914–4924. [PubMed: 18335939] d) Kawakami T, Aimoto S. *Tetrahedron* 2009;65:3871–3877.
11. Crich D, Sana K, Guo S. *Org. Lett* 2007;9:4423–4426. [PubMed: 17900128]
12. The use of allyl esters in alloxycarbamates as a protecting group system orthogonal with both Boc and Fmoc chemistries is widely established. a) Grieco P, Gitu PM, Hruby VJ. *J. Peptide. Res* 2001;57:250–256. [PubMed: 11298927] b) Kates SA, Daniels SB, Albericio F. *Anal. Biochem* 1993;212:303–310. [PubMed: 8214570] c) Kates SA, Solé NA, Johnson CR, Hudson D, Barany G, Albericio F. *Tetrahedron Lett* 1993;34:1549–1552. d) Bloomberg GB, Askin D, Gargaro AR, Tanner MAJ. *Tetrahedron Lett* 1993;34:4709–4712. Alternative possibilities include the nitrobenzenesulfonyl protecting group for amines. e) Kan T, Fukuyama T. *Chem. Commun* 2004:353–359. f) Halpin DR, Lee JA, Warren SJ, Harbury PB. *PLOS Biology* 2004;2:1031–1038.
13. Albericio F, Cruz M, Debethune L, Eritja R, Giralt E, Grandas A, Marchan V, Pastor JJ, Pedroso E, Rabanal F, Royo M. *Synth. Commun* 2001;31:225–232.
14. a) Dourtoglou V, Ziegler JC, Gross B. *Tetrahedron Lett* 1978;19:1269–1272. b) Knorr R, Trzeciak A, Bannwarth W, Gillessen D. *Tetrahedron Lett* 1989;30:1927–1930. c) Fields, GB.; Tian, Z.; Barany, G. *Synthetic Peptides: A User's Guide*. Grant, GA., editor. New York: Freeman; 1992. p. 77-183. d) Schnoelzer M, Alewood P, Jones A, Alewood D, Kent SBH. *Int. J. Peptide Protein Res* 1992;40:180–183. [PubMed: 1478777]
15. Albericio F, Nicolas E, Rizo J, Ruiz-Gayo M, Pedroso E, Giralt E. *Synthesis* 1990:119–122.
16. Webster KL, Maude AB, O'Donnell ME, Mehrotra AP, Gani D. *J. Chem. Soc., Perkin Trans* 2001;1:1673–1695.
17. Katzhendler, J.; Klauzner, Y.; Beylis, I.; Mizhiritskii, M.; Shpernat, Y.; Ashkenazi, B.; Fridland, D. *PCT Int. Appl.* 076744. 2005.
18. Dangles O, Guibe F, Balavoine G, Lavielle S, Marquet A. *J. Org. Chem* 1987;52:4984–4993.
19. The peptides were isolated in the form of their N-Boc derivatives simply owing to the use of Boc-protected building blocks. Use of Fmoc-protected building blocks in the final coupling step will necessarily yield peptides with the N-terminus unprotected.
20. Lee S-H, Lee S, Youn YS, Na DH, Chae SY, Byun Y, Lee KC. *Bioconjugate Chem* 2005;16:377–382.
21. Hackeng TM, Griffin JH, Dawson PE. *Proc. Natl. Acad. Sci. U. S. A* 1999;96:10068–10073. [PubMed: 10468563]
22. Fairwell T, Hospattankar AV, Ronan R, Brewer HB Jr, Chang JK, Shimizu M, Zitzner L, Arnaud CD. *Biochemistry* 1983;22:2691–2697. [PubMed: 6871156]
23. Fields GB, Fields CG. *J. Am. Chem. Soc* 1991;113:4202–4207 and references therein cited
24. This is evident simply from the yields of the isolated thioacids **25** and **27**, which require average minimum coupling yields of >93% and >97%, respectively, for each coupling deprotection cycle. For comparable reasons premature peptide cleavage by diketopiperazine formation at the level of deprotection of the dipeptide and the migration of side chain protecting groups to N-terminal amines on neutralization do not appear to be major concerns, at least for the examples provided.
25. a) Shangguan N, Katukojvala S, Greenberg R, Williams LJ. *J. Am. Chem. Soc* 2003;125:7754–7755. [PubMed: 12822965] b) Merckx R, Brouwer AR, Rijkers DTS, Liskamp RMJ. *Org. Lett* 2005;7:1125–1128. [PubMed: 15760155] c) Barlett KN, Kolakowski RV, Katukojvala S, Williams LJ. *Org. Lett*


2006;8:823–826. [PubMed: 16494450] d) Kolakowski RV, Shanguan N, Sauers RR, Williams LJ. *J. Am. Chem. Soc* 2006;128:5695–5702. [PubMed: 16637636] e) Merx R, Van Haren MJ, Rijkers DTS, Liskamp RMJ. *J. Org. Chem* 2007;72:4574–4577. [PubMed: 17497928]




Scheme 1.
Preparation of a functionalized resin.




7, 86%




9, 92%




11, 95%



8, 91%




10, 90%





12, 92%


Scheme 2.
Preparation of protected hydroxyl amino acids

**Scheme 3.**

Preparation of mono 9-fluorenylmethyl esters of aspartic and glutamic acid



Scheme 4.
Peptidyl thioester synthesis



Scheme 5.
Formation of a sulfonamide.

