



Published in final edited form as:

J Immunotoxicol. 2009 December ; 6(4): 285–292. doi:10.3109/15476910903258260.

Tetrabromobisphenol A has Immunosuppressive Effects on Human Natural Killer Cells

Esther Caroline Kibakaya, Krishna Stephen, and Margaret M. Whalen

Department of Chemistry, Tennessee State University, Nashville, TN 37209

Abstract

Human natural killer (NK) cells are lymphocytes that destroy tumor cells, virally infected cells, and antibody-coated cells. Tetrabromobisphenol A (TBBPA) is used both as a reactive and an additive flame retardant in a variety of materials and appears to contaminate the environment. TBBPA has been found in human blood samples and if it interferes with NK cell function, this could increase risk of tumor development and/or viral infection. This study examines the effects of exposure to various concentrations of TBBPA for 24 hr, 48 hr, and 6 d on the lytic function, tumor-target-binding function, and ATP levels of NK cells. These same parameters were also monitored in NK cells that were exposed to TBBPA for 1 hr followed by 24 hr, 48 hr, and 6 d in TBBPA-free media. A 24 hr exposure of NK cells to 5 μM TBBPA caused a greater than 95% decrease in NK lytic function, a 70% decrease in binding function, and a 34% decrease in ATP levels in NK cells. Exposure to 2.5 μM TBBPA for 24 hr decreased lytic function by 76%, binding function by 20%, and had no effect on ATP levels. Exposure of NK cells to 5 μM TBBPA for 1 hr followed by 24 hr in TBBPA-free media caused a progressive and persistent loss of lytic function (41%) while not affecting either binding ability or ATP levels. The results indicate that TBBPA exposures decrease the lytic function of human NK cells and that an initial brief (1 hr) exposure can cause a progressive loss of function. Additionally, the data also indicate that TBBPA-induced loss of NK lytic function can occur at concentration of TBBPA that do not affect target-binding ability and ATP levels of NK cells.

Keywords

ATP; NK cells; Tetrabromobisphenol A; lytic function

Introduction

Tetrabromobisphenol A (TBBPA) is used as a flame retardant in a wide variety of materials including plastics, paper, and textiles (IPCS/WHO, 1995; HSBD, 2001). TBBPA has been used as a replacement for the polybrominated diphenylethers (PBDEs), whose persistence in the environment and potential for negative effects on health have prompted concerns (Sjodin et al., 1999; Eriksson et al, 2001). Its main use is as a reactive flame retardant in epoxy resin circuit boards (IPCS/WHO, 1995; HSDB, 2001). However, it is also applied to carpeting and upholstery (Gain, 1997). TBBPA has additional uses as a plasticizer and as an intermediate in the production of other flame retardants (IUPCS/WHO, 1995; HSBD, 2001). In 1998, the annual U.S. production of TBBPA was greater than 1 million pounds with estimates ranging between 100-500 million pounds (U.S. EPA, 2000a; Environ Defense, 2001). Its production

Please address all correspondence to: Dr. Margaret M. Whalen, Department of Chemistry, Tennessee State University, 3500 John A. Merritt Blvd., Nashville, TN 37209 (T) 615-963-5247 (F) 615-963-5326 mwhalen@tnstate.edu.

Declaration of Interest The Authors report no conflicts of interest. The Authors are alone responsible for the content and writing of the paper.

and use have the potential to release TBBPA into the environment. It is a highly hydrophobic compound and would be expected to adsorb onto suspended solids and sediment in streams (U.S. EPA, 1999, 2002b). Its half-life in water is expected to be 48-84 days (U.S. EPA, 1999, 2000b). TBBPA is found in sewage sludge (Sellstrom and Jansson, 1995; de Boer et al., 2002; de Wit, 2002;) and appears to bioaccumulate (IPCS/WHO, 1995; HSBD 2001; Birnbaum and Staskal, 2004).

TBBPA has been found in human serum samples (Hagmar et al., 2000; Nagayama et al., 2001; Thomsen et al., 2002). Levels reported were 0.24-0.71 ng/g lipid in serum samples from Norway (approximately 1.8-5.3 pM, using a conversion factor of 400 mg lipid/100ml serum) (Thomsen et al., 2002) and averaged 4.5 ng/g lipid (approximately 33.8 pM) in a Japanese study (Nagayama et al., 2001). A toxicokinetic study following a single oral dose of 300 mg/kg TBBPA to adult humans showed a peak plasma concentration of 103 µM within 3 hr that declined, with an elimination half-life of 13 hr (Schauer et al., 2006). Human exposure to TBBPA is believed to occur by inhalation, dermal contact, and ingestion (IPCS/WHO, 1995; Peterman et al., 2000). It has been found in commercial drinking water that was stored in polycarbonate containers (Peterman et al., 2000) and in seafood (IPCS/WHO, 1995). There has been little study of the toxicological effects of TBBPA on humans. However there have been studies of its effects in rodents and rainbow trout. TBBPA was shown to increase the activity of glutathione reductase in rainbow trout (Ronisz et al., 2004). In mice it has been shown to cause decreases in serum proteins and red blood cells as well as increases in spleen weight (IPCS/WHO, 1995). Administration of TBBPA to newborn rats caused polycystic kidney lesions (Fukuda et al., 2004). Administration of TBBPA to Wistar rats showed effects on circulating thyroid hormone, T4, levels (Van der Ven et al., 2008). This same study showed that there were no effects of TBBPA exposure on the function of splenic natural killer (NK) cells in exposed rats or their offspring (Van der Ven et al., 2008). *In vitro* studies of TBBPA showed that it was able to compete with thyroid hormone, T4, for binding to human transthyretin (thyroid hormone transport protein) (Meerts et al., 2000).

Human NK cells are immune system cells (lymphocytes) that are capable of destroying tumor cells, virally infected cells, and antibody-coated cells. NK cells have the markers CD56 and/or CD16 on their cell surface and lack the T-lymphocyte receptor/CD3 complex (Lotzova, 1993). NK cells are a crucial immune defense against malignant neoplasms and viral infections (Lotzova, 1993; Vivier et al., 2004). Their role in protecting against viral infections is demonstrated by increased incidences of viral infection in individuals that lack NK cells (Fleisher et al., 1982; Biron et al., 1989). Unlike T-lymphocytes, NK cells lyse (destroy) appropriate target cells without prior sensitization making them a frontline immune defense. Thus, if a compound such as TBBPA were to alter NK-cell function, this could result in increased risk of viral infection and tumor formation. A recent study has shown that another flame retardant inhibits the function of human NK cells (Hinkson and Whalen, 2009). Suppression of human NK cell function by contaminants such as these may account in part for increases in cancer incidence that have been seen, especially those seen in lymphomas and leukemias (SEER, 2009). Thus, it is important to address capacity of TBBPA as well as other environmental contaminants to diminish the function of human NK cells.

In the current study TBBPA was examined for its potential to disrupt the lytic function of NK cells. Both acute (i.e., 1 hr) and chronic (as long as 6 d) exposures were examined. The brief (1 hr) exposures were followed by periods of time in TBBPA-free media prior to assessing NK function (1 hr exposures, followed by up to 6 d in TBBPA-free media). These types of studies were important in that lasting (or developing) negative effects of a brief exposure to TBBPA would suggest greater risk from exposure to the compound. Finally, the effects of TBBPA on the ability of NK cells to bind to their target cells and on the intracellular ATP levels in NK cells were studied. Binding of NK cells to targets is a necessary step in the lytic process and

interference with binding could lead to loss of lytic function (Lotzova et al., 1993). Additionally, NK lytic function is an ATP dependent process (Andzelm et al., 2007), and decreases in ATP could lead to loss of lytic function.

Materials and Methods

Isolation of NK cells

Peripheral blood from healthy adult (male and female) volunteer donors was used for this study. Buffy coats (source leukocytes) obtained from the American Red Cross (Portland, OR) or Key Biologics, LLC (Memphis, TN) were used to prepare NK cells. Highly-purified NK cells were obtained using a rosetting procedure. Buffy coats were mixed with 0.8 ml of RosetteSep human NK cell enrichment antibody cocktail (StemCell Technologies, Vancouver, British Columbia, Canada) per 45 ml of buffy coat. The mixture was incubated for 20 min at room temperature (~25°C). Following the incubation, 7-8 ml of the mixture was layered onto 4 ml of Ficoll-Hypaque (1.077 g/ml; MP Biomedicals, Irvine, CA) and centrifuged at $1200 \times g$ for 30-40 min. The cell layer was then collected and washed twice with phosphate-buffered saline (PBS; pH 7.2) and stored in complete media (RPMI-1640 supplemented with 10% heat-inactivated bovine calf serum [BCS], 2 mM L-glutamine, and 50 U penicillin G/50 µg streptomycin/ml) at 1 million cells/ml (Whalen et al., 2002).

Chemical preparation

TBBPA (purchased from Fisher Scientific, 97% pure) was dissolved in dimethyl sulfoxide (DMSO) (Sigma-Aldrich, St. Louis, MO) to yield a 100 mM stock solution. Desired concentrations of TBBPA were then prepared by dilution of the stock into complete media containing 0.5% gelatin in the place of 10% bovine serum. Gelatin was used in place of bovine serum to avoid binding of TBBPA to the serum's albumin, which could interfere with its delivery the cells. The final concentration of DMSO in any of the TBBPA exposures did not exceed 0.01%.

Cell Treatments

NK cells (at a concentration of 1.5×10^6 cells/ml) were exposed to TBBPA or control (DMSO-bearing medium) for 24 hr, 48 hr, or 6 d. Following exposures, the cells were assayed for tumor-destroying function, binding function, and/or ATP levels. Additionally, NK cells were exposed to TBBPA for 1 hr; following the 1-hr exposure period, the TBBPA-containing or control media was removed and replaced with fresh TBBPA-free media and the cells were incubated in compound-free media for 24 hr, 48 hr, or 6 d prior to being assayed for tumor-destroying function, binding function, and/or ATP levels. The concentration range examined was 0.05 - 10 µM TBBPA.

Cell Viability

Cell viability was determined by trypan blue exclusion. Cell numbers and viability were assessed at the beginning and end of each exposure period. Viability was determined at each concentration for each of the exposure periods. The viability of treated cells was then compared to that of control cells at each length of exposure (Whalen et al., 2003). Only those concentrations where viability was unaffected were used at a given length of exposure. Viability was 100% of that seen in the control NK cells at the concentrations used at a specific exposure time.

Cytotoxicity assay

The ability of NK cells to lyse tumor cells was measured using a ^{51}Cr release assay (Whalen et al., 1999). The target cell in all cytotoxicity assays was the NK-susceptible K562 (human

chronic myelogenous leukemia) cell line. Briefly, K562 cells were incubated with ^{51}Cr (Perkin-Elmer Life Sciences, Boston, MA) in 0.2 - 0.5 ml of unadulterated BCS (BCS containing no additions) for 1-1.5 hr at 37 °C in an air:CO₂ (19:1) incubator. Following this incubation, the target cells were washed twice with gelatin media. NK (effector) cells ($1.2 \times 10^5/100 \mu\text{l}$ for 12:1 ratio with target cells) were added to the wells of round-bottom micro-well plates. The effector cells were diluted to a 6:1 ($0.6 \times 10^5/100 \mu\text{l}$) and a 3:1 ratio ($0.3 \times 10^5/100 \mu\text{l}$); each ratio was tested in triplicate. Target cells were added ($1 \times 10^4/100 \mu\text{l}$) to each well of the micro-well plate; the plate was then centrifuged at $300 \times g$ for 3.5 min and incubated for 2 hr at 37 °C (air:CO₂, 19:1). After this incubation, a 0.1 ml aliquot of supernatant was collected from each well and each assessed for radioactivity (in 60 sec) in a Packard COBRA γ - radiation counter (Packard Instrument Co., Meriden, CT). Target lysis was then calculated as: $100 \times [\text{test cpm} - \text{spontaneous cpm}]/[\text{maximum cpm} - \text{spontaneous cpm}]$. Maximum release was produced by adding 100 μl of 10% Triton X-100 to wells containing only the labeled target cells; spontaneous release was assessed from wells containing cells plus medium only.

Binding Assay

The percentage of target cells with bound NK cells was determined at two effector:target ratios, i.e., 12:1 and 6:1. The NK cells were treated as described above for the cytotoxicity assay. Tumor cells were then incubated with an appropriate number of control or TBBPA exposed lymphocytes in round-bottomed micro-well plates (20,000 tumor cells/well). Each condition was tested in triplicate. The cells were incubated at 37° C (air:CO₂, 19:1) for 10 min and then placed on ice. Following the incubation period, the cells were gently re-suspended with a micropipette, placed in a hemocytometer and viewed under a light microscope. The number of target cells with two or more lymphocytes bound to their surface was counted as well as the total number of targets to determine the percentage of tumor cells with lymphocytes bound. The minimum number of targets counted per determination was 50-100 (Whalen et al., 1999).

ATP Assay

Following exposures of cells to the various concentrations of TBBPA; 250,000 NK cells were added to 300 μl of somatic cell ATP releasing reagent (Sigma-Aldrich). To measure ATP, 100 μl of lysed cell suspension was added to 100 μl of a Luciferin/Luciferase mixture (Sigma). The light emission was then measured using a Fluoroskan Ascent FL luminescence plate reader (Thermo Electron Corp., Vantaa, Finland). ATP levels were then determined by extrapolation from a standard curve (Dudimah et al., 2007b).

Statistical Analysis

The data from TBBPA treated cells at a given concentration were compared to control data by pair wise analysis using Student's t-test (Figures 1, 3, 5, and 6). Two-way ANOVA, as well as *post-hoc* linear regression, were used to evaluate the effect of length of exposure at a given concentration of TBBPA in Figures 2 and 4.

Results

Lytic Function and Binding Function of NK Cells Exposed to TBBPA for 24 hr, 48 hr, and 6 d

Exposures of NK cells to 2.5 and 5 μM TBBPA for 24 hr caused very significant reductions in NK cell lytic function of $75.8 \pm 6.6\%$ and $96.2 \pm 3.1\%$, respectively ($p < 0.0001$) (Figure 1A). The decrease in lytic function after a 24 hr exposure to 1 μM TBBPA was $38.6 \pm 10.7\%$ ($p < 0.001$), while that at 0.5 μM was $23.3 \pm 11.2\%$ ($p < 0.001$). The binding function of NK cells exposed to 2.5 and 5 μM TBBPA for 24 hr was reduced by $19.1 \pm 14.3\%$ and $69.9 \pm$

16.6%, respectively ($p < 0.0001$) (Figure 1A). NK cells exposed to 1 μM TBBPA for 24 hr showed no loss of binding function.

NK cells were exposed to concentrations of 0.25 - 2.5 μM TBBPA for 48 hr because higher concentrations caused decreased viability of the NK cells. Exposure at all of the concentrations tested except 0.25 μM TBBPA caused significant decrease in the lytic function of NK cells. The decreases were $31.0 \pm 9.8\%$ at 0.5 μM , $56.0 \pm 7.1\%$ at 1 μM , and $86.1 \pm 6.3\%$ at 2.5 μM ($p < 0.0001$) (Figure 1B). NK cells that were exposed to concentrations below 1 μM TBBPA showed no loss of binding function. A 48 hr exposure to 1 μM TBBPA caused a small loss of binding function of $12.2 \pm 11.0\%$ ($p < 0.05$). Exposure to 2.5 μM TBBPA for 48 hr caused a decrease in binding function of $51.2 \pm 18.7\%$ ($p < 0.0001$) (Figure 1B).

The 6-d exposures of NK cells to TBBPA were carried out at concentrations of 0.05 μM - 1 μM . There was a significant decrease in tumor-cell-lysing function with a 6 d exposure to all of the concentrations of TBBPA except the 0.05 μM (Figure 1C). TBBPA caused decreases in lytic function of $20.3 \pm 16.0\%$ at 0.1 μM , $33.2 \pm 19.1\%$ at 0.25 μM , $63.7 \pm 21.4\%$ at 0.5 μM , and $89.2 \pm 16.0\%$ at 1 μM ($p < 0.01$) (Figure 1C). NK cells exposed to 0.25 - 1 μM TBBPA for 6 d showed a significant decrease in binding function ($33.2 \pm 23.7\%$, $p < 0.01$) only with the 1 μM exposure (Figure 1C).

Figure 2 plots the lytic function of NK cells exposed to 0.5 and 1 μM TBBPA as a function of length of exposure (using the data from Figure 1). Only the 0.5 and 1 μM TBBPA concentrations are shown as these are the only concentrations that were tested at all three lengths of exposure. This is done to show that there appears to be an increased loss of lytic function with length of exposure at a given concentration of TBBPA, when comparing the 24 hr exposures to the 6 d exposures ($p < 0.01$).

Lytic Function and Binding Function of NK Cells Exposed to TBBPA for 1 hr, Followed by 0 hr, 24 hr, 48 hr, and 6 d in TBBPA-Free Medium

In order to assess whether a brief exposure of NK cells to TBBPA could cause lasting and/or ensuing affects on lytic or binding function, NK cells were exposed to 0.5 - 10 μM TBBPA for 1 hr which was removed and the NK cells were then suspended in TBBPA-free media for 0 hr, 24 hr, 48 hr, or 6 d before assaying for lytic function or binding function. NK cells exposed to 10 μM for 1 hr and then tested immediately for lytic function (0 hr in TBBPA-free media) exhibited a statistically-significant decreases in lytic function of $30.1 \pm 12.5\%$ ($p < 0.001$) lower concentrations caused no significant decreases in lytic function. When NK cells were exposed to 0.5 - 10 μM , TBBPA for 1 hr and then incubated in TBBPA-free media for 24 hr prior to assaying for lytic function, decreases in lytic function of $22.4 \pm 9.9\%$ (1 μM), $27.0 \pm 11.3\%$ (2.5 μM), $41.4 \pm 10.9\%$ (5 μM), and $53.9 \pm 15.0\%$ (10 μM) ($p < 0.001$) were seen (Figure 3A). The binding function of NK cells exposed to TBBPA for 1 hr, followed by 24 hr in TBBPA-free media prior to assaying for binding function was unaffected by any of the concentrations tested (Figure 3A).

NK cells exposed to these same concentrations of TBBPA and incubated for 48 hr in TBBPA-free media prior to assay, showed reductions in lytic function of $18.3 \pm 8.9\%$ (2.5 μM), $22.6 \pm 14.7\%$ (5 μM), and $37.4 \pm 11.7\%$ (10 μM) ($p < 0.05$) (Figure 3B). The binding function of NK cells exposed to TBBPA for 1 hr, followed by 48 hr in TBBPA-free media prior to assaying for binding function was unaffected by any of the concentrations tested (Figure 3B).

In NK cells exposed to these same concentrations of TBBPA for 1 hr, followed by a 6-d period in TBBPA-free media lytic function decreased $33.6 \pm 23.5\%$ (2.5 μM), $41.2 \pm 9.9\%$ (5 μM), and $64.3 \pm 6.7\%$ (10 μM) ($p < 0.01$) (Figure 3C). NK cells that were exposed to 10 μM TBBPA for 1 hr and then given 6 d in TBBPA-free media prior to assaying for binding function showed

a loss of binding function of $49.6 \pm 8.4\%$ ($p < 0.0001$) (Figure 3C). All other concentrations tested 6 d following 1 hr exposures to TBBPA showed no loss of binding function (Figure 3C)

Figure 4 plots the results of 1 hr exposures to 5 and 10 μM TBBPA followed by 0 hr, 24 hr, 48 hr, and 6 d in TBBPA-free media as a function of length of time in TBBPA-free media (such as was done in Figure 2). There again appears to be an increasing loss of lytic function with time after a 1 hr exposure ($p < .01$).

ATP Levels in NK Cells Exposed to TBBPA for 24 hr, 48 hr, and 6 d

NK cells exposed to TBBPA for 24 hr showed no significant decreases in ATP levels except at 5 μM where ATP was decreased by $33.5 \pm 12.6\%$ ($p < 0.0001$) (Figure 5A). Lytic function data from Figure 1 were placed on the plots of ATP changes (Figure 5) in order to facilitate comparison of changes in ATP levels with those of lytic function. Reduction in ATP levels of $16.8 \pm 9\%$ and $32.7 \pm 8.4\%$ were seen with 48 hr exposures to 1 and 2.5 μM HBCD, respectively ($p < 0.0001$) (Figure 5B). A 6-d exposure of NK cells to 0.25, 0.5, and 1 μM HBCD led to decreases in ATP levels of $38.4 \pm 15\%$, $55.6 \pm 29.6\%$, and $78.3 \pm 27.2\%$, respectively ($p < 0.001$) (Figure 5C).

Effects of Exposures to TBBPA for 1 hr Followed by, 24 hr, 48 hr, and 6 d in TBBPA-Free Media on ATP Levels in NK Cells

As with lytic function and binding function, it is possible that an initial exposure to TBBPA for 1 hr could cause changes in NK cell ATP levels that would manifest later, even though the compound has been removed. Thus, we examined if the effects of exposures to TBBPA intensified following removal of the compound. To address this possibility, NK cells were exposed to 0.5 - 10 μM TBBPA for 1 hr that was then removed and the NK cells were then suspended in TBBPA-free media for 24 hr, 48 hr, or 6 d before assaying for ATP levels. NK cells exposed to TBBPA for 1 hr and then incubated for 24 hr in TBBPA-free media prior to measuring ATP levels showed no significant decreases in ATP levels (Figure 6A). As was done above, lytic function data from Figure 3 were placed on the plots of ATP changes (Figure 6) in order to facilitate comparison of changes in ATP levels with those of lytic function. NK cells exposed to 2.5, 5 and 10 μM TBBPA for 1 hr and given a 48 hr period in TBBPA-free media before assay showed very small decreases in ATP levels ($p < 0.05$) (Figure 6B). A 1 hr exposure of NK cells to 1-10 μM TBBPA followed by 6 d in TBBPA-free media lead to decreases in ATP of about 30% at the 1 - 5 μM concentrations of TBBPA and of $43.6 \pm 39.6\%$ at 10 μM ($p < 0.01$) (Figure 6C).

Discussion

Chemically-induced diminution of human NK cell immune function could leave an individual vulnerable to malignant neoplasias and/or viral infection. Previous studies have indicated that humans with a defect that leaves their NK cells unable to lyse target cells have a predisposition to development of certain tumors (Ortaldo et al., 1992) and that individuals lacking NK cells had severe herpes virus infection (Biron et al., 1989) and Epstein-Bar infections (Fleisher, et al., 1982). Studies in mice have shown that a decrease in NK cell activity of 80% results in an increase in tumor formation upon challenge with melanoma cells (Wilson et al., 2001). Decreased NK cell function and increased tumor metastases have been demonstrated in mice exposed to cigarette smoke (Lu et al., 2007) and exposure of mice to the chemical urethane causes tumors and has been shown to suppress NK function (Luebke et al., 1986).

A large number of chemicals currently contaminate the environment and are found in human tissues (Thornton et al. 2002). The use of TBBPA as a flame retardant and plasticizer (HSBD, 2001; IPCS/WHO, 1995) has resulted in its being found in human blood samples (Hagmar et

al., 2000; Nagayama et al., 2001; Thomsen et al., 2002). While it appears to be eliminated to some extent from the human system (Schauer et al., 2006), its detection in human blood samples indicates that a certain fraction accumulates in the body. Levels of TBBPA in human serum of 4.5 ng/g lipid (33.8 pM, using 400 mg lipid per 100 ml serum) have been reported (Nagayama et al., 2001). We examined the effects of TBBPA exposures on the lytic function of human NK cells. Further, the ability of an initial exposure to TBBPA to cause lasting and/or ensuing disruption of lytic function was examined, as a lasting disruption of NK function by a brief encounter with TBBPA would indicate a greater level of potential harm from exposure to this compound. The mechanism of TBBPA-induced decreases in lytic function was also addressed by examining the effects of TBBPA on the ability of NK cells to bind to tumor target cells and on ATP levels in NK cells. NK cells must first bind to target cells before they can initiate the process by which the target is lysed (Lotzova, 1993; Vivier et al. 2004). ATP plays an important role in lysis of tumor cells by NK cells (Andzelm et al., 2007).

A 24 hr exposure to as little as 0.5 μ M TBBPA caused a significant loss of lytic function in NK cells. When the concentration was 5 μ M there was a nearly complete loss of lytic function at 24 hr. These decreases in lytic function were not accompanied by comparable losses in binding function. While exposure to 2.5 μ M TBBPA for 24 hr caused a loss of lytic function of about 76% the loss of binding function was only 19%. Thus, these studies indicate that the TBBPA-induced loss of lytic function cannot be accounted for simply by TBBPA causing reduced ability of the NK cells to bind to their tumor targets.

An important finding of the current study was that a brief exposure to TBBPA could cause alteration(s) of the NK cell such that the loss of lytic function persisted and/or developed after the compound was removed. An exposure to 10 μ M TBBPA for 1 hr caused a 30% loss of lytic function while lower concentrations had no effect on lytic function. When cells were tested 24 hr following the 1-hr exposures, there was a 54% loss of lytic function at 10 μ M TBBPA. A 1 hr exposure to 5 μ M TBBPA caused no loss of lytic function; however, 24 hr after that exposure, the NK cells had lost 41% of their lytic ability. There was no loss of ability of NK cells to bind to target cells 24 hr after a 1 hr exposure to any of the concentrations of TBBPA that were tested. Thus, as was seen with continuous exposures to TBBPA loss of lytic function was not accompanied by a concomitant loss of binding function.

Since a loss of binding function could not account for the losses of lytic function seen with TBBPA exposures, we went on to examine whether TBBPA caused decreases in the intracellular ATP levels of NK cells. As mentioned earlier, ATP is required for the release of cytotoxic proteins from the NK cells onto target cells (Andzelm et al., 2007). As with binding function we found that a decrease in ATP levels did not necessarily accompany a loss of lytic function in TBBPA-exposed NK cells. For instance a 24 hr exposure to 2.5 μ M TBBPA caused a nearly 80% loss of lytic function without having any effect on ATP levels. We did find that after a 6-d exposure to TBBPA the loss of lytic function was associated with a decrease in ATP levels.

The current study indicates that exposures to TBBPA can cause very significant losses of human peripheral blood NK cell lytic function. However, a recent study in Wistar rats found that *in vivo* exposure to TBBPA had no effect on the function of splenic natural killer (NK) cells in exposed rats or their offspring (Van der Ven et al., 2008). As mentioned, this study examined the function of NK cells from the spleen and did not examine peripheral blood NK cells. A previous study indicated that *in vivo* exposure of rats to the chemical carcinogens, diethylnitrosamine and 2-acetylaminofluorene, decreased the function of peripheral blood NK cells but had no effect on splenic NK cells (Leu et al., 1985). Thus, the difference between splenic and peripheral blood NK cells may be a potential explanation for the fact that TBBPA dramatically decreased human NK cell lytic function but did not decrease the NK function in

rats. Additionally, species differences in susceptibility to TBBPA as well as exposure levels may also account for the differing results between rats and humans.

Previous studies have shown that a variety of environmental contaminants cause decreases in NK cell function that persist once the compound is removed. These include another brominated flame retardant, hexabromocyclododecane (HBCD) (Hinkson and Whalen, 2009), several organotin compounds (tributyltin [TBT], dibutyltin [DBT], and triphenyltin [TPT]) (Whalen et al., 2002; Dudimah et al., 2007a,b; Whalen et al., 2003) an organochlorine compound, pentachlorophenol (PCP) and a carbamate compound, ziram (Taylor et al., 2005). As with TBBPA, neither a decrease in target binding nor a decrease in ATP levels could fully account for the losses of lytic function seen with many of these compounds (HBCD, TBT, DBT, PCP, ziram) (Whalen et al., 1999, 2002; Dudimah et al., 2007a,b; Taylor et al., 2005; Hinkson and Whalen, 2009; Taylor and Whalen, 2009). However, as seen with TBBPA a decrease in ATP appears to be associated with the loss of function seen after 6 d with some of these compounds (HBCD, TBT, DBT, and PCP) (Dudimah et al., 2007a,b; Nnodu and Whalen, 2008; Hinkson and Whalen, 2009).

While significant decreases in NK lytic function were seen in human NK cells exposed to TBBPA, the concentrations at which these losses of function were seen were high compared to what has been measured in human serum (Nagayama et al., 2001). However, because the effects of TBBPA exposure do not subside once the compound is removed, there may be potential for cumulative effects over time.

In summary, the data indicate that exposure of NK cells to TBBPA can cause very significant losses of NK lytic function. Further, loss of lytic function occurs after a brief (1 hr) exposure, indicating that the process(es) is/are stimulated during a brief exposure that lead to a sustained loss of lytic function. TBBPA has effects on the ability of NK cells to bind to their targets and also is able to decrease intracellular levels of ATP under certain conditions. However, decreases in binding or in ATP levels cannot explain the losses of lytic function seen under several conditions. The current study shows that TBBPA interferes with the essential immune function of NK cells and thus has the potential for increasing the risk of cancer development in exposed individuals.

Acknowledgments

This research was supported by Grant 2S06GM-08092-34 from the National Institutes of Health.


References

- Andzelm MM, Chen X, Krzewski K, Orange JS, Strominger JL. Myosin IIA is required for cytolytic granule exocytosis in human NK cells. *J Exp Med* 2007;204:2285–2291. [PubMed: 17875677]
- Birnbaum LS, Staskal DF. Brominated flame retardants: Cause for concern? *Environ Health Perspect* 2004;112:9–17. [PubMed: 14698924]
- Biron CA, Byron KS, Sullivan JL. Severe herpes virus in an adolescent without natural killer cells. *New Engl J Med* 1989;320:1731–1735. [PubMed: 2543925]
- de Boer, J.; Allchin, C.; Zegers, B.; Boon, JP.; Brandsma, SH.; Morris, S.; Kruijt, AW.; van der Hesseligen, JM.; Haftka, JJ. HBCD and TBBP-A in Sewage Sludge, Sediments and Biota, Including Inter-laboratory Study. CO33/02. Wageningen, The Netherlands: RIVO-Netherlands Institute for Fisheries Research; 2002. p. 40
- de Wit CA. An overview of brominated flame retardants in the environment. *Chemosphere* 2002;46:583–624. [PubMed: 11999784]
- Dudimah FD, Gibson C, Whalen MM. Effect of dibutyltin on ATP levels in human natural killer cells. *Environ Toxicol* 2007a;22:117–123. [PubMed: 17366562]


- Dudimah FD, Odman-Ghazi SO, Hatcher F, Whalen MM. Effect of tributyltin (TBT) on ATP levels in human natural killer (NK) cells: Relationship to TBT-induced decreases in NK function. *J Appl Toxicol* 2007b;27:86–94. [PubMed: 17149696]
- Environ Defense. 2,2',6,6'-Tetrabromo-4,4'-isopropylidenediphenol - Industrial uses. 2001. http://www.scorecard.org/chemical-profiles/uses.tcl?edf_substance_id=79%2d94%2d7
- Eriksson P, Jakobsson E, Fredriksson A. Brominated flame retardants: A novel class of developmental neurotoxicants in our environment. *Environ Health Perspect* 2001;109:903–908. [PubMed: 11673118]
- Fleisher G, Koven N, Kamiya H, Henle W. A non-X-linked syndrome with susceptibility to severe Epstein-Bar virus infections. *J Pediatr* 1982;100:727–730. [PubMed: 6279813]
- Fukuda N, Ito Y, Yamaguchi M, Mitumori K, Koizumi M, Hasegawa R, Kamata E, Ema M. Unexpected nephrotoxicity induced by tetrabromobisphenol A in newborn rats. *Toxicol Lett* 2004;150:145–155. [PubMed: 15093670]
- Gain B. Flame retardants' Albemarle boosts capacity. *Chem July 2;1997 :88*. Full text available from PROMT 97 369076
- Hagmar L, Jakobsson K, Thuresson K, Rylander L, Sjodin A, Bergman A. Computer technicians are occupationally exposed to polybrominated diphenyl ethers and tetrabromobisphenol A. *Organohalogen Comp* 2000;47:202–205.
- Hinkson NC, Whalen MM. Hexabromocyclododecane decreases the lytic function and ATP levels of human natural killer cells. *J Appl Toxicol*. 2009 In press
- HSDB (Hazardous Substances Data Bank). 2,2',6,6'-Tetrabromobisphenol A. Bethesda, MD: National Library of Medicine; 2001. <http://www.toxnet.nlm.nih.gov/cgi-bin/sis/search>
- IPC/WHO (International Program on Chemical Safety/World Health Organization). Environmental Health Criteria 172: Tetrabromobisphenol A and Derivatives. Geneva: World Health Organization; 1995.
- Kakimoto K, Akutsu K, Konishi Y, Tanaka Y. Time trend of hexabromocyclododecane in the breast milk of Japanese women. *Chemosphere* 2008;71:1110–1114. [PubMed: 18076970]
- Lotzova E. Definition and function of natural killer cells. *Nat Immunol* 1993;12:177–193.
- Leu RW, Norton TR, Herriott MJ, Ringer DP, Kearns RJ. Suppression of natural killer and lymphocyte functions associated with carcinogen-induced premalignant hyperplastic nodules in rat liver. *Cancer Res* 1985;45:3282–3287. [PubMed: 2860970]
- Lu LM, Zavitz CC, Chen B, Kianpour S, Wan Y, Stampfli MR. Cigarette smoke impairs NK cell-dependent tumor immune surveillance. *J Immunol* 2007;178:936–943. [PubMed: 17202355]
- Luebke RW, Riddle MM, Rogers RR, Rowe DG, Garner RJ, Smialowicz RJ. Immune function in adult C57BL/6J mice following exposure to urethane pre-or post natally. *J Immunopharmacol* 1986;8:243–257. [PubMed: 3722835]
- Meerts IA, van Zanden JJ, Luijckx EA, van Leeuwen-Bol I, Marsh G, Jakobsson E, Bergman A, Brouwer A. Potent competitive interactions of some brominated flame retardants and related compounds with human transthyretin *in vitro*. *Toxicol Sci* 2000;56:95–104. [PubMed: 10869457]
- Nagayama, J.; Takasuga, T.; Tsuji, H., editors. Contamination levels of brominated flame retardants, dioxins, and organochlorine compounds in the blood of Japanese adults; *Human Levels and Trends*, Part 4. 2001. p. 218–221. Located at http://www.kemi.se/aktuellt/BFR/bfr_de14.pdf
- Nnodu U, Whalen MM. Pentachlorophenol decreases ATP levels in human natural killer cells. *J Appl Toxicol* 2008;28:1016–1020. [PubMed: 18623605]
- Ortaldo JR, Glenn GM, Young HA, Frey JL. Natural killer (NK) cell lytic dysfunction and putative NK cell receptor expression abnormality in members of a family with chromosome 3p-linked von Hippel-Lindau disease. *J Natl Cancer Inst* 1992;84:1897–1903. [PubMed: 1460671]
- Peterman PH, Orazio CE, Gale RW. Detection of tetrabromobisphenol A and formation of brominated [¹³C]-bisphenol A's in commercial drinking water stored in reusable polycarbonate containers. *ACS Div Environ Chem Extended Abstr* 2000;40:431–433.
- Ronisiz D, Farnen Finne E, Karlsson H, Forlin L. Effects of the brominated flame retardants hexabromocyclododecane (HBCDD), and tetrabromobisphenol A (TBBPA), on hepatic enzymes and other biomarkers in juvenile rainbow trout and feral eelpout. *Aquat Toxicol* 2004;69:229–245. [PubMed: 15276329]

- Schauer UM, Volkel W, Dekant W. Toxicokinetics of tetrabromobisphenol A in humans and rats after oral administration. *Toxicol Sci* 2006;91:49–58. [PubMed: 16481339]
- SEER (Surveillance Epidemiology and End Results) National Cancer Institute, U.S National Institutes of Health. Apr. 2009 www.cancer.gov
- Sellstroem U, Jansson B. Analysis of tetrabromobisphenol A in a product and environmental samples. *Chemosphere* 1995;31:3085–3092.
- Sjödin A, Hagmar L, Klasson-Wehler E, Kronholm-Diab K, Jakobsson E, Bergman A. Flame retardant exposure: polybrominated diphenyl ethers in blood from Swedish workers. *Environ Health Perspect* 1999;107:643–648. [PubMed: 10417362]
- Taylor TR, Tucker T, Whalen MM. Persistent inhibition of human natural killer cell function by ziram and pentachlorophenol. *Environ Toxicol* 2005;20:418–424. [PubMed: 16007644]
- Taylor TR, Whalen MM. Effects of ziram on tumor-cell-binding capacity, cell-surface marker expression, and ATP levels of human natural killer cells. *Cell Biol Toxicol*. 2009In press
- Thomsen C, Lundanes E, Becher G. Brominated flame retardants in archived serum samples from Norway: A study on temporal trends and the role of age. *Environ Sci Technol* 2002;36:1414–1418. [PubMed: 11999045]
- Thornton, JW.; McCally, M.; Houlihan, J. Biomonitoring of industrial pollutants: Health and policy implications of the chemical body burden; Public Health Rep. 2002. p. 315–323. www.ewg.org/reports/bodyburden/findings.php
- U.S. EPA. 40 CFR Part 372: Persistent Bioaccumulation Toxic (PBT) Chemicals; Final Rule. 1999
- U.S. EPA. Non-confidential Production Volume Information Submitted by Companies for Chemicals Under the 1998 Inventory Update Rule (IUR). CAS no. 79947. Washington, DC: U.S. EPA, Office of Pollution Prevention and Toxics (OPPT); 2000a. Internet: <http://www.epa.gov/cgi-bin/iursrch3.cgi>
- U.S. EPA. Emergency Planning and Community Right-To-Know Act Section 313 Reporting Guidance for Rubber and Plastics Manufacturing. EPA 745-B-00-017. Washington, DC: Office of Environmental Information; 2000b. p. 2-16. Internet: <http://www.epa.gov/tri/00rubber.pdf>
- Vivier E, Nunes JA, Vely F. Natural killer cell signaling pathways. *Science* 2004;306:1517–1519. [PubMed: 15567854]
- Whalen MM, Loganathan BG, Kannan K. Immunotoxicity of environmentally-relevant concentrations of butyltins on human natural killer cells *in vitro*. *Environ Res* 1999;81:108–116. [PubMed: 10433842]
- Whalen MM, Green SA, Loganathan BG. Brief butyltin exposure induces irreversible inhibition of the cytotoxic function on human natural killer cells *in vitro*. *Environ Res* 2002;88:19–29. [PubMed: 11896664]
- Whalen MM, Wilson S, Gleghorn C, Loganathan BG. Brief exposure to triphenyltin produces irreversible inhibition of the cytotoxic function of human natural killer cells. *Environ Res* 2003;92:213–220. [PubMed: 12804518]
- Wilson SD, McCay JA, Butterworth LF, Munson AE, White KL Jr. Correlation of suppressed natural killer cell activity with altered host resistance models in B₆C₃F₁ mice. *Toxicol Appl Pharmacol* 2001;177:208–218. [PubMed: 11749120]

(A) 24 hr



(B) 48 hr



(C) 6 days




Figure 1. Effects of TBBPA exposures on the ability of NK cells to lyse or bind tumor cells
 NK cells were exposed to 0.05 - 5 μM TBBPA for (A) 24 hr, (B) 48 hr, or (C) 6 d, and the tested for lytic function (\blacklozenge) or binding function (\blacksquare). To combine results from separate experiments (using cells from different donors), levels of lysis or binding were normalized as the percentage of the lytic or binding function of the control cells (untreated cells from the same donor) in a given experiment. Results were pooled from three separate experiments using different donors (triplicate determinations for each experiment; $n = 9$; mean \pm SD). Significant decreases in **Lytic Function** as compared to control: (A) 0.5, 1, 2.5 and 5 μM ($p < 0.001$); (B) 0.5, 1, and 2.5 μM ($p < 0.0001$); and, (C) 0.1, 0.25, 0.5, and 1 μM at ($p < 0.01$). Significant decreases in **Binding Function** as compared to control: (A) 2.5 and 5 μM ($p < 0.0001$); (B) 1 and 2.5 μM ($p < 0.05$); and, (C) 1 μM ($p < 0.01$).






Figure 2. Time course of the effects of 0.5 μM (black bar) and 1 μM (gray bar) TBBPA on the ability of NK cells to lyse tumor cells

Results are the 0.5 and 1 μM data from Figure 1 plotted versus length of exposure.

(A) 24 hr in TBBPA-free media**(B) 48 hr in TBBPA-free media**


(C) 6 days in TBBPA-free media

Figure 3. Effects of a 1-hr exposure to TBBPA followed by various periods in TBBPA-free media on the ability of NK cells to lyse or bind tumor cells

NK cells were exposed to 0.5 - 10 μM TBBPA for 1 hr, and then tested for lytic function (\blacklozenge) or binding function (\blacksquare) after (A) 24 hr, (B) 48 hr, or (C) 6 d. Results were normalized as described in Figure 1. Significant decreases in **Lytic Function** as compared to control: (A) 1, 2.5, 5 and 10 μM ($p < 0.001$); (B) 2.5, 5, and 10 μM ($p < 0.05$); (C) 2.5, 5, and 10 μM at ($p < 0.01$). Significant decreases in **Binding Function** as compared to control: (A) none; (B) none; and, (C) 10 μM ($p < 0.0001$).






Figure 4. Time course of the effects of 1-hr exposures to 5 μM (black bar) and 10 μM (gray bar) TBBPA followed by a period in TBBPA-free media on the ability of NK cells to lyse tumor cells. Results are the 5 and 10 μM data from Figure 3 plotted versus length of time in TBBPA-free media.

(A) 24 hr



(B) 48 hr



(C) 6 days






Figure 5. Effects of TBBPA exposures on the ATP levels and lytic function of NK cells
 NK cells were exposed to 0.05 - 10 μM TBBPA for (A) 24 hr, (B) 48 hr, or (C) 6 d, and then tested for ATP levels (■) or lytic function (◆). Results were normalized as described in Figure 1. Significant decreases in **ATP levels** as compared to control: (A) 5 μM ($p < 0.0001$); (B) 1 and 2.5 μM ($p < 0.0001$); and, (C) 0.25, 0.5 and 1 μM ($p < 0.001$). Significant decreases in **Lytic Function** as compared to control: (A) 0.5, 1, 2.5 and 5 μM ($p < 0.001$); (B) 0.5, 1, and 2.5 μM ($p < 0.0001$); and, (C) 0.1, 0.25, 0.5, and 1 μM at ($p < 0.01$).

(A) 24 hr in TBBPA-free media



(B) 48 hr in TBBPA-free media



(C) 6 days in TBBPA-free media




Figure 6. Effects of TBBPA exposures on the ATP levels and lytic function of NK cells
 NK cells were exposed to 0.5 - 10 μM TBBPA for 1 hr, followed by periods of time in TBBPA-free media prior to assaying for ATP levels (■) or lytic function (◆). (A) 24 hr, (B) 48 hr, or (C) 6 d. Results were normalized as described in Figure 1. Significant decreases in **ATP Levels** as compared to control: (A) none; (B) 2.5, 5, and 10 μM ($p < 0.05$); and, (C) 1, 2.5, 5, and 10 μM ($p < 0.01$). Significant decreases in **Lytic Function** as compared to control: (A) 1, 2.5, 5 and 10 μM ($p < 0.001$); (B) 2.5, 5, and 10 μM ($p < 0.05$); (C) 2.5, 5, and, 10 μM at ($p < 0.01$).