

Evolutionary Dynamics of the Genomic Region Around the Blast Resistance Gene *Pi-ta* in AA Genome *Oryza* Species

Seonghee Lee,^{*,†,1} Stefano Costanzo,[†] Yulin Jia,^{†,2} Kenneth M. Olsen[‡] and Ana L. Caicedo[§]

^{*}Rice Research and Extension Center, University of Arkansas, Stuttgart, Arkansas 72160, [†]U. S. Department of Agriculture–Agricultural Research Service, Dale Bumpers National Rice Research Center, Stuttgart, Arkansas 72160, [‡]Department of Biology, Washington University, St. Louis, Missouri 63130 and [§]Department of Biology, University of Massachusetts, Amherst, Massachusetts 01003

Manuscript received August 6, 2009
Accepted for publication October 3, 2009

ABSTRACT

The race-specific resistance gene *Pi-ta* has been effectively used to control blast disease, one of the most destructive plant diseases worldwide. A single amino acid change at the 918 position of the *Pi-ta* protein was known to determine resistance specificity. To understand the evolutionary dynamics present, we examined sequences of the *Pi-ta* locus and its flanking regions in 159 accessions composed of seven AA genome *Oryza* species: *O. sativa*, *O. rufipogon*, *O. nivara*, *O. meridionalis*, *O. glaberrima*, *O. barthii*, and *O. glumaepatula*. A 3364-bp fragment encoding a predicted transposon was found in the proximity of the *Pi-ta* promoter region associated with the resistance phenotype. Haplotype network analysis with 33 newly identified *Pi-ta* haplotypes and 18 newly identified *Pi-ta* protein variants demonstrated the evolutionary relationships of *Pi-ta* haplotypes between *O. sativa* and *O. rufipogon*. In *O. rufipogon*, the recent directional selection was found in the *Pi-ta* region, while significant deviation from neutral evolution was not found in all *O. sativa* groups. Results of sequence variation in flanking regions around *Pi-ta* in *O. sativa* suggest that the size of the resistant *Pi-ta* introgressed block was at least 5.4 Mb in all elite resistant cultivars but not in the cultivars without *Pi-ta*. These findings demonstrate that the *Pi-ta* region with transposon and additional plant modifiers has evolved under an extensive selection pressure during crop breeding.

PLANT resistance (*R*) genes have evolved to fight against a wide range of pathogens in a race-specific manner where a particular *R* gene in a plant recognizes the corresponding avirulence (*AVR*) gene in a pathogen race (FLOR 1971). Thus far, a number of *R* genes have been identified and characterized from diverse plant species. Most characterized *R* genes to date encode putative proteins with nucleotide binding sites (NBS) and leucine-rich repeats (LRR) (HULBERT *et al.* 2001). Most *R* genes are highly polymorphic and diversified, which is consistent with the ability to interact with diverse random molecules encoded by diverse pathogen *AVR* genes (MEYERS *et al.* 2003; BAKKER *et al.* 2006; SHEN *et al.* 2006).

Blast disease, caused by the filamentous ascomycete *Magnaporthe oryzae* B.C. Couch [formerly *M. grisea* (T. T. Hebert) M. E. Barr] (ROSSMAN *et al.* 1990; COUCH and KOHN 2002), has been one of the major constraints to

stable crop production. Currently, *Oryza sativa* and *M. oryzae* have been an excellent model pathosystem for uncovering the molecular coevolution mechanisms of host–pathogen (VALENT *et al.* 1991; TALBOT 2003). At least 80 race-specific *R* genes that confer resistance to specific pathogen races have been described in rice germplasm (BALLINI *et al.* 2008). Eleven blast *R* genes (*Pi-ta*, *Pib*, *Pi2/Piz-t*, *Pi5*, *Pi9*, *Pi21*, *Pi36*, *Pi37*, *Pi-d2*, *Pikm*, and *Pit*) have been cloned, and most of them, except *Pi21* and *Pi-d2*, were also predicted to encode receptor proteins with NBS (CHEN *et al.* 2006; FUKUOKA *et al.* 2009; JIA *et al.* 2009b). In most cases, blast *R* genes are members of small gene families with a single family member required for resistance. *Pikm* and *Pi5* are exceptions that require two members of the same gene family for *Pikm*- and *Pi5*-mediated resistance, respectively (ASHIKAWA *et al.* 2008; LEE *et al.* 2009). Recently, a retrotransposon was predicted to be involved in the *Pit* resistance (HAYASHI and YOSHIDA 2009).

The evolutionary dynamics and mechanisms of resistance mediated by *Pi-ta* is one of the best-studied *R* genes. *Pi-ta* has been effectively deployed in the United States and around the globe for controlling blast disease (BRYAN *et al.* 2000; JIA *et al.* 2000; JIA 2003; JIA *et al.* 2004a,b; HUANG *et al.* 2008; JIA and MARTIN 2008; WANG *et al.* 2008; JIA *et al.* 2009a). *Pi-ta* encodes a predicted cytoplasmic protein with a centrally located NBS and

Supporting information is available online at <http://www.genetics.org/cgi/content/full/genetics.109.108266/DC1>.

Sequence data from this article have been deposited with the EMBL/GenBank Data Libraries under accessions nos. GQ918334–GQ918489 and GQ984160.

¹Present address: Samuel Roberts Noble Foundation, 2510 Sam Noble Pky., Ardmore, OK 73401.

²Corresponding author: USDA-ARS Dale Bumpers National Rice Research Center, Stuttgart, AR 72160. E-mail: yulin.jia@ars.usda.gov

a highly interrupted LRR domain (referred to as the LRD) at the carboxyl terminus that recognizes the corresponding avirulence gene *AVR-Pita*, triggering race-specific resistance. A single amino acid substitution, serine (Ser) to alanine (Ala) at the position of 918, in the LRD of the *Pi-ta* protein was demonstrated to determine the direct interaction with AVR-Pita and the resistance specificity to blast pathogen *M. oryzae* (BRYAN *et al.* 2000; JIA *et al.* 2000). The resistant *Pi-ta* allele (Ala-918) was found in *O. sativa* and its ancestor *O. rufipogon* (JIA *et al.* 2004b; HUANG *et al.* 2008). Surveys of *Pi-ta* nucleotide sequences with limited accessions of *Oryza* species have revealed that the degree of nucleotide diversity is higher at the intron of the *Pi-ta* gene (JIA *et al.* 2003; HUANG *et al.* 2008; WANG *et al.* 2008; YOSHIDA and MIYASHITA 2009). HUANG *et al.* (2008) further suggested that a selective sweep occurred recently at the *Pi-ta* gene in *O. rufipogon*, but the extent of selection around the *Pi-ta* genomic region has not been demonstrated in either *O. rufipogon* or *O. sativa*.

Knowledge of the historical introduction of the *Pi-ta* gene can help to understand the extent of selection at the *Pi-ta* locus. The landraces Tadukan and Tetep, containing *Pi-ta* and other blast *R* genes in chromosome 12, have been used as breeding parents for preventing blast disease worldwide. Tadukan was confirmed to be the *Pi-ta* donor for various Asian *japonica* cultivars (RYBKA *et al.* 1997) whereas Tetep was the *Pi-ta* donor for the U. S. cultivars (GRAVOIS *et al.* 1995; MOLDENHAUER *et al.* 1998; MCCLUNG *et al.* 1999; GIBBONS *et al.* 2006; MOLDENHAUER *et al.* 2007). Recently, the large introgressed chromosomal segments surrounding the *Pi-ta* locus were identified in backcross BC₅ progenies and elite rice cultivars (JIA 2009). This suggests that the broad spectrum of the *Pi-ta* resistance in the United States may include the effects of other loci in the *Pi-ta* region, inherited as a “superlocus.” Toward this end, *Ptb(t)*, a nuclear gene that is required for the *Pi-ta*-mediated resistance, was also mapped at the *Pi-ta* region (JIA and MARTIN 2008). Further determination of DNA sequences around the *Pi-ta* gene should help to determine the minimal genomic region that is essential for *Pi-ta*-mediated resistance.

The two cultivated rice species, *O. sativa* and *O. glaberrima*, belong to the AA genome of *Oryza* species. *O. rufipogon* and *O. nivara* are wild progenitors of the Asian rice *O. sativa*, whereas *O. barthii* is a wild progenitor of the African cultivated rice *O. glaberrima* (LINARES 2002; YAMANAKA *et al.* 2003; LONDO *et al.* 2006). The comparison of *R*-gene diversity between cultivated rice and its wild ancestors is important to understand the selection effects of crop domestication and breeding.

The objectives of this study were (1) to characterize distributions of the *Pi-ta* allele in *O. sativa* and to detect the potential presence/absence of polymorphism(s) associated with the resistance phenotype; (2) to examine the molecular evolution and patterns of selection in the *Pi-ta* gene in *O. sativa* and *O. rufipogon*; (3) to analyze

molecular diversity around the *Pi-ta* locus in AA genome *Oryza* species; and (4) to understand the pattern and extent of selection for *Pi-ta*-mediated resistance in *Oryza* species during crop domestication.

MATERIALS AND METHODS

Plant materials and DNA preparation: A total of 159 geographically diverse accessions of *O. sativa*, *O. rufipogon*, and five other closely related AA genome *Oryza* species were selected for this study. These included 43 Asian landraces, 18 U. S. domesticated cultivars, and 58 U. S. weedy rice strains in *O. sativa*; 28 geographically diverse accessions of *O. rufipogon*; 4 accessions of *O. glaberrima*; and 2 accessions each of *O. nivara*, *O. barthii*, *O. meridionalis*, and *O. glumaepatula* (Table S1). U. S. cultivars and weedy rice seeds were obtained from the USDA-ARS Dale Bumpers National Rice Research Center, and all Asian landrace accessions consisting of 15 *indica*, 7 *aus*, 3 *aromatic*, 12 *tropical japonica*, and 4 *temperate japonica* were obtained from Susan McCouch at Cornell University and the International Rice Research Institute. Plants were grown in greenhouses at Washington University and the University of Massachusetts. DNA extracted from 2- to 4-week-old seedlings was diluted to 2 ng/ μ l for further analysis.

Primer design and DNA sequencing: Primer pairs were designed using the Primer3 program (ROZEN and SKALETSKY 2000) to amplify overlapping fragments (~700 bp each) for *Pi-ta*, including 5' upstream, 3' downstream, and a coding region with an intron (Table S2). All primers were verified by BLAST against both 93-11 (*indica*) and Nipponbare (*japonica*) genome sequences. Primers were also designed to amplify 400- to 700-bp fragments of six flanking genes in the regions from 9.6 to 11.6 Mb on chromosome 12. The six flanking loci around the *Pi-ta* gene were LOC_OS12G16690 (9.6 Mb), LOC_OS12G17080 (9.8 Mb), and LOC_OS12G17830 (10.2 Mb) and LOC_OS12G18690 (10.8 Mb), LOC_OS12G19290 (11.2 Mb), and LOC_OS12G20260 (11.8 Mb) (<http://rice.plantbiology.msu.edu/>). For 11 resistant cultivars carrying *Pi-ta* (Tadukan, Tetep, Te Qing, Yashiro-mochi, Pi4, Reiho, IR64, Katy, Banks, Drew, and Madison), fragments from six additional flanking loci were sequenced: LOC_OS12G12370 (6.8 Mb), LOC_OS12G13570 (7.6 Mb), LOC_OS12G14330 (8.2 Mb), LOC_OS12G22360 (12.6 Mb), LOC_OS12G24020 (13.7 Mb), and LOC_OS12G25630 (14.8 Mb) (<http://rice.plantbiology.msu.edu/>) (Figure 1).

Sequence data analysis: All DNA sequences from *Pi-ta* and 12 flanking genes were aligned using Vector NTI 10 (Invitrogen) and MEGA 4 (TAMURA *et al.* 2007). The genomic sequence from Nipponbare, a *temperate japonica* cultivar, was included as the reference sequence (<http://rice.plantbiology.msu.edu/>). Additional sequences of the *Pi-ta* gene of 50 accessions of *O. rufipogon*, 3 accessions of *O. nivara*, 2 accessions of *O. meridionalis*, 6 accessions of *O. glaberrima*, and 6 accessions of *O. barthii* were obtained from the GenBank database (Table S1), yielding a total of 226 accessions. For the sequence analysis, accessions of *temperate japonica*, *tropical japonica*, and *aromatics* collectively formed the *japonica* subspecies, and *aus* and *indica* together formed the *indica* subspecies. Nucleotide polymorphisms at and around the *Pi-ta* region were analyzed using the software DnaSP 4.9 (ROZAS *et al.* 2003). The level of nucleotide diversity at silent sites (π_{silent}) and the population mutation parameter θ_w (Watterson estimator) of *Pi-ta* and the flanking gene fragments were estimated for each group of *O. sativa* and compared with that of other *Oryza* species. Average rates of nonsynonymous (K_a) and synonymous (K_s) substitutions were calculated to examine selections at the *Pi-ta*




FIGURE 1.—Patterns of DNA sequence variation at and around the *Pi-ta* locus in the AA genome of *Oryza* species. (A) Sliding-window analysis of the *Pi-ta* locus in 159 accessions (top). The gene structure of *Pi-ta* is shown at the bottom. (B) Sliding-window analysis at the *Pi-ta* coding region (top). The structure of the *Pi-ta* coding region is shown at the bottom. Values were assigned to the nucleotide at the midpoint of 5 bp for A and 25 bp for B, respectively. The parameter of difference per site (y-axis) is plotted against the nucleotide position (x-axis). Each line indicates synonymous (red) or nonsynonymous variation (blue). (C) Graphic presentation of the genomic region of the *Pi-ta* locus and 12 flanking loci. Sequenced fragments and physical locations on the chromosome are indicated, and the names of loci are represented as A–L. A: outer envelope protein (LOC_OS12G12370); B: Myb-like protein (LOC_OS12G13570); C: NBS–LRR disease resistance protein (LOC_OS12G14330); D: ubiquitin–protein ligase/zinc ion-binding protein (LOC_OS12G16690); E: pentatricopeptide repeat-containing protein (LOC_OS12G17080); F: unknown (LOC_OS12G17830); G: unknown (LOC_OS12G18690); H: serine/threonine-protein kinase (LOC_OS12G19290); I: unknown (LOC_OS12G20260); J: unknown (LOC_OS12G22360); K: senescence-associated protein DIN1 (LOC_OS12G24020); L: sulfite oxidase (LOC_OS12G25630).

coding region in all accessions of *O. sativa* and *O. rufipogon*. Joint analyses of interspecific comparisons using *O. barthii* as an outgroup species were used for estimating the ratio of K_a/K_s and for determining deviations from neutral evolution (AKASHI 1999). Sliding-window analysis was performed to examine nucleotide polymorphism across the *Pi-ta* gene in all *Oryza* species. Statistical tests of neutrality such as Tajima's D , Fu and Li's D^* and F^* , and Fay and Wu's normalized H were calculated to examine the selection present at and around *Pi-ta*. Extended haplotype homozygosity (EHH) (SABETI *et al.* 2002) was calculated to visualize the effect of selection on the alleles containing Ala-918 or Ser-918. A haplotype network was also constructed for comparisons of genealogical relationships among *Pi-ta* haplotypes using TCS 1.21 (CLEMMENT *et al.* 2000).

RESULTS

Nucleotide diversity at the *Pi-ta* region: High levels of nucleotide variation were observed in the intron,

5'-UTR, and 3'-UTR regions of *Pi-ta* in 159 accessions (Figure 1A). Insertions and deletions (indels) ranging from 10 to 540 bp in the noncoding regions were distinguished among the *Pi-ta* haplotypes. A 242-bp deletion in an intron of *Pi-ta* was found only in *O. glaberrima*, *O. barthii*, and *O. glumaepatula*. Within the coding region, levels of nucleotide and amino acid polymorphism were substantially higher in the first exon. Comparisons of amino acid mutations among partitions of the coding region showed that nonsynonymous were more common than synonymous changes in the NBS region (Figure 1B). Nucleotide diversity in *O. sativa* was lower than that in *O. rufipogon*. A total of 175 polymorphic sites, excluding indels, were found in the coding region, including an intron; of these polymorphic sites, 29 occurred in *O. sativa*, 121 in *O. rufipogon*, and 25 in other *Oryza* species. Average

TABLE 1
Molecular evolutionary parameters of the *Pi-ta* gene in *Oryza* species analyzed in this study

Species	Sample no.	Nucleotide	θ_w	π_{silent}	D	D^*	F^*	H_n
<i>O. sativa</i>	55	4250	0.00206	0.00287	1.36790	0.43956	0.92805	0.16426
<i>O. sativa indica</i>	23	4250	0.00235	0.00257	-0.02930	-0.18210	-0.15867	-0.29793
<i>O. sativa japonica</i>	32	4250	0.00143	0.00230	1.63577	1.25707	1.62162	0.41988
<i>O. sativa japonica</i> Asian cultivar	16	4250	0.00174	0.00244	0.78421	0.47668	0.64840	0.01420
<i>O. sativa japonica</i> U. S. cultivar	16	4250	0.00174	0.00226	1.41409	1.53348**	1.72883*	0.64640
<i>O. rufipogon</i>	91	3988	0.00888	0.00522	-2.14289*	-2.09795	-2.54113*	-3.65945
<i>O. nivara</i>	5	4003	0.01520	0.01322	-1.06420	-1.06420	-1.15583	-3.39370
<i>O. glaberrima</i>	10	4002	0.00966	0.01366	1.88503	1.03161	1.41069	0.19870
<i>O. barthii</i>	9	4002	0.01066	0.01336	1.21069	1.07971	1.24886	-1.63819

θ_w , Watterson's nucleotide diversity estimator (1975) based on silent site; π , Nei's nucleotide diversity (1987) based on silent site; D , Tajima's D statistics (1989) based on the differences between the number of segregating sites and the average number of nucleotide differences; D^* and F^* , the neutral test proposed by Fu and Li (1993); and H_n , normalized Fay and Wu's H test statistics. Statistical significance: ** $P < 0.02$ and * $P < 0.05$.

pairwise nucleotide diversity (π_{silent}) and silent Watterson's nucleotide diversity estimator (θ_w) over the *Pi-ta* gene was lowest in *O. sativa* ($\pi_{\text{silent}} = 0.00292$, $\theta_w = 0.00180$) compared to other *Oryza* species, including *O. rufipogon* ($\pi_{\text{silent}} = 0.00522$ – 0.01366 , $\theta_w = 0.00888$ – 0.01520) (Table 1). The levels of diversity in African cultivated rice *O. glaberrima* and its wild progenitor *O. barthii* were similar to *O. rufipogon* and *O. nivara* (Table 1). Analyses for *O. glumaepatula* and *O. meridionalis* were not included because of sample limitation.

A total of 53 *Pi-ta* haplotypes were identified (Table S3) from seven AA genome *Oryza* species, including the previously reported 20 haplotypes (HUANG *et al.* 2008; WANG *et al.* 2008; YOSHIDA and MIYASHITA 2009). Among them, 32 *Pi-ta* haplogroups were identified

from different *Oryza* species in the haplotype network, suggesting that the diversification of *Pi-ta* haplotypes occurred before the divergence of these *Oryza* species (Figure 2). Nineteen haplotypes were from *O. sativa* and 25 haplotypes were from *O. rufipogon* (Table S3). A total of 26 *Pi-ta* variants from PT1 to PT26 were identified on the basis of the amino acid sequence of the *Pi-ta* protein in *Oryza* species (Table 2); these include 8 *Pi-ta* variants previously identified (WANG *et al.* 2008). Five *Pi-ta* variants—PT1, PT2, PT3, PT4, and PT20—were the most prevalent type of the variants in *O. sativa* (Figure 2 and Table S1). PT1 containing the functional amino acid alanine at 918 was found only in accessions of *O. sativa* and *O. rufipogon*. PT22, PT23, PT24, PT25, and PT26, were the major types of *Pi-ta* variants found in




FIGURE 2.—A haplotype network based on nucleotide polymorphisms of the *Pi-ta* coding region of 226 accessions of seven AA genome *Oryza* species: *O. sativa*, *O. rufipogon*, *O. nivara*, *O. meridionalis*, *O. glaberrima*, *O. barthii*, and *O. glumaepatula*. Each group of haplotypes is shown as a solid circle, and seven major haplotypes are marked in larger circles. The *Pi-ta* variants are in parentheses. Each branch represents a single mutational step. Branches with small solid circles indicate that there is more than a single mutational step between haplotypes. A number next to a branch represents the length of the mutational steps. Different sizes of circles represent the different numbers of each haplotype.




FIGURE 3.—Comparison of the EHH of two core haplotypes (alanine-918 and serine-918) in the *Pi-ta* region in *O. sativa*. The core was defined by a single amino acid change at the position of 918 (serine: TCT or alanine: GCT) that determines the resistance specificity of *Pi-ta*. The starting EHH value for alanine-918 is 0.331 while the EHH value is 0.669 for serine-918.

determined if it was due to selection or population structure in both species because of sample limitation.

The level of synonymous divergence (K_s) exceeded that of nonsynonymous divergence (K_a) in all partitions of the coding region of the *Pi-ta* protein except the NBS region in *O. sativa* and *O. rufipogon*, indicating purifying selection against amino acid substitutions in most portions of the gene (Table 3). These findings were also confirmed in comparisons between synonymous nucleotide polymorphism (π_{syn}) and nonsynonymous nucleotide polymorphism (π_{non}) in *O. rufipogon* (Table 3). However, the $\pi_{syn}:\pi_{non}$ ratio was smaller than one ($\pi_{syn}:\pi_{non} < 1$) in the NBS in *O. sativa* due to the very low polymorphism present in the species. The NBS of the *Pi-ta* protein in both *O. sativa* and *O. rufipogon* showed a greater number of interspecies nonsynonymous-to-synonymous substitutions ($K_a/K_s > 1$), indicating that positive directional selection has favored amino acid substitutions in this domain (Table 3).

Nucleotide polymorphisms in genomic regions around *Pi-ta*: We sequenced all fragments of targeted flanking loci around *Pi-ta* except one locus encoding a NBS-LRR disease resistance protein (LOC_OS12G14330), the RPM-1 homolog located at 8.2 Mb. The presence and absence of the RPM-1 homolog was found in both *O. sativa* and *O. rufipogon* accessions. The absence of the RPM-1 homolog was found in two Asian cultivars, Yashiro-mochi (*japonica*) and Te Qing (*indica*), and in all U. S. weedy rice carrying resistant *Pi-ta* (Table S4).

Nucleotide data sets shown in Figure 1 were aligned for 433–659 bp of six loci in 2 Mb around *Pi-ta* in all 159 accessions. The estimated values of nucleotide diversity for these loci were 0–0.00391 in *O. sativa* and 0.0015–0.00508 in *O. rufipogon*. The levels of sequence variation in flanking loci around *Pi-ta* were similar to the levels in the *Pi-ta* locus found in both species (Table 4). The test of Tajima's *D* in the region around *Pi-ta* in *O. sativa* and *O. rufipogon* revealed that no significant pattern of

TABLE 3

Molecular variation and selection at the *Pi-ta* gene in *O. sativa* (*indica*, *japonica*, and weedy rice) and *O. rufipogon*

Gene segment	S	π_{syn}^a	π_{non}^a	π_{non}/π_{syn}	$K_s(JC)^b$	$K_a(JC)^b$	K_a/K_s^b
<i>O. sativa</i> ($n = 113$)							
Coding	12	0.0015	0.00098	0.654	0.00871	0.00560	0.643
5' coding to NBS	6	0.00305	0.00337	1.105	0.01513	0.01034	0.683
NBS	2	0.00009	0.00003	0.295	0.00004	0.00451	102.5
NBS to LRD	0	0	0	0	0.03457	0.00371	0.107
LRD	4	0.00196	0.00063	0.319	0.00774	0.00536	0.692
<i>O. rufipogon</i> ($n = 91$)							
Coding	62	0.00249	0.00166	0.668	0.00849	0.00530	0.625
5' coding to NBS	24	0.00491	0.00278	0.564	0.01211	0.00698	0.577
NBS	15	0.00101	0.00155	1.543	0.00052	0.00484	9.341
NBS to LRD	4	0.00401	0.00049	0.121	0.03384	0.00359	0.106
LRD	19	0.00174	0.00139	0.801	0.00694	0.00514	0.741

^a π_{syn} , nucleotide diversity at synonymous site; π_{non} , nucleotide diversity at nonsynonymous site.

^b Jukes-Cantor (JC) corrected synonymous differences per synonymous site (K_s) and nonsynonymous differences per nonsynonymous site (K_a) using intraspecific and interspecific comparisons using *O. barthii*.

TABLE 4
Molecular diversity of genomic regions around *Pi-ta* in *O. sativa* (*indica*, *japonica*, and weedy rice) and *O. rufipogon*

Physical location (Mb)	Nucleotide polymorphism (π)										Tajima's <i>D</i>							
	9.6	9.8	10.2	10.8	11.2	11.8	9.6	9.8	10.2	10.8	11.2	11.8	9.6	9.8	10.2	10.8	11.2	11.8
<i>O. sativa</i>	0.00178	0.00115	0.00108	0.0018	0.0015	0.00391	0.66938	0.69958	0.82595	0.92414	NA	0.91739	0.17149	0.82595	0.92414	NA	0.91739	11.8
<i>O. indica</i>	0.0019	0.00109	0.00204	0.00218	0.00213	0.00479	-0.96803	-1.51481	-0.10605	1.08052	NA	1.02022	-0.70826	1.08052	1.08052	NA	1.02022	11.2
<i>O. japonica</i>	0.00118	0.00148	0.00047	0.00253	0.00048	0.00416	0.97327	0.27501	0.44003	0.6426	NA	0.13462	1.05235	0.6426	0.6426	NA	0.13462	11.2
Weedy rice	0.001	0.00129	0.0004	0.00166	0.00168	0.00365	-0.93379	-0.45271	-0.51132	-0.42536	NA	1.13812	-0.44628	-0.42536	-0.42536	NA	1.13812	11.2
<i>O. rufipogon</i>	0.00508	0.0015	0.0057	0.00477	0.00301	0.0015	-0.15872	0.23998	0.16801	0.22133	-1.36029	-0.3066	-2.57275	0.22133	-1.36029	-1.36029	-0.3066	11.2

^aThe sequence of *Pi-ta* including flanking region (2 kb upstream and downstream of *Pi-ta*) and coding region with intron was used for nucleotide polymorphism and Tajima's *D*.



selection presents around the *Pi-ta* locus in *O. sativa*. However, a significant negative value of Tajima's *D* was detected around the *Pi-ta* locus in *O. rufipogon*, similar to the result found in the *Pi-ta* gene (Table 4).

Interestingly, a 3364-bp insertion located 1012 bp upstream of the start codon (ATG) was found only in all accessions carrying the resistance *Pi-ta* allele (Figure 4A). The presence of the insertion in resistant accessions was verified by Southern blot analysis using a probe derived from the 5' region of *Pi-ta* (Figure 4B). The inserted fragment was cloned and sequenced from the U. S. cultivar, Katy (GenBank accession no. GQ984160). Sequences of the 3364-bp fragment were predicted to encode a protein with 844 amino acids with domains commonly found in zinc fingers and transcription factors and with domains commonly found in hAT family dimerization (hATC) of a transposable element (Figure 4C). Using the rice sequence database of Nipponbare (*japonica*) and 93-11 (*indica*), a highly homologous sequence with the insertion was found on chromosome 2 of 93-11 while no homologous sequence was found in the Nipponbare. From a Southern blot using the probe in the insertion and PCR analysis with primers amplifying the flanking region of the insertion, the 3364-bp insertion was determined on chromosome 2 in susceptible *indica* cultivars; however, the insertion was on both chromosomes 2 and 12 in resistant *indica* cultivars or *japonica* cultivars possessing *indica*-derived resistant *Pi-ta* (data not shown).

After surveying in the 2-Mb region around *Pi-ta* in 118 accessions of *O. sativa*, no polymorphism was detected in all resistant *O. sativa* accessions. Six additional flanking gene fragments were sequenced in the 8-Mb region to identify polymorphisms in those accessions (4 Mb upstream and 4 Mb downstream of *Pi-ta*). The different sizes of the *Pi-ta* introgressed block in resistant cultivated rice were estimated by detecting the initial breaking point of recombination surrounding the *Pi-ta* locus. A range from 5 to 8 Mb of the *Pi-ta* introgression block (the average being 7 Mb) was identified in 11 resistant cultivars (JIA *et al.* 2004b; WANG *et al.* 2007). Among them, the smallest block (5.4 Mb) was identified in Yashiro-mochi and the largest *Pi-ta* introgression (>8 Mb) was found in the two Japanese cultivars Pi4 and Reiho whose *Pi-ta* region was derived from Tadukan. A 6.8-Mb portion of the *Pi-ta* region in Tetep was identified in the U. S. cultivars Katy, Drew, Banks, and Madison (Figure 5).

DISCUSSION

In this study, we analyzed DNA sequence polymorphisms in and around the genomic region of *Pi-ta* in 159 geographically diverse *Oryza* accessions composed of several *Oryza* species to gain insight into the origin and evolution of *Pi-ta*. We discovered that the extended genomic region (>5 Mb) surrounding resistant *Pi-ta*



C

```

HSSRNRYDYGAEKRRKRRLAQAQSGKALDKFFLRETPNANIEDDISDDMAEVDANIAESDDAVEENVVDGDIGHDLADEGRDLASEGNEENIADDAD [100]
                                     ZnF_TTF
DNVSFRPDMFDPRTWDGLDPRKIDILLQKQPKRDLSEIHGPRDNLSSRFLASSYTKVLSNGEKCDREWLVYSKELDKVFCCKLLRKLGLVRGQLANDGV [200]
NDWNHLANRLKEHEVSRHEVTNMSTUYELRLRMQKNQITDKVAQRELEKEREHURRVLRLILLIVKFLAEHNIAFRGNSKLYQDSNGNGLGLVEMLVEF [300]
DPVIKEHVDRITNDKIRDHYLGPSIQNELINLLAVAIAKSSIIAKIKEAKYFSVILDCTPDASHQEQMSLIIRYVDVTTCSIEESFLGFLDNDTSGQGLF [400]
DVLVEELNSLDLVDVANVRGQYDNGSNMRGKHQGVQKLLDINPRAFYSACGCHSLNLTLCDMAKSCRKATEFFGVQIRIYTTFFANSTRKWKILKDNLSG [500]
LTLKSLSSSTRWESRVDSVKAIRFQIPEIREALLQVAETDNDPLTVSEVNSLSLENELGGFEFLVAIIWYEILSSINVSKQLQSKDMVIDIAIESVQGLI [600]
SLFKKYRENGFSKALEAAKQIALEMDPIEFRTKRKIKRKRQFDEGTSIDASIDSQSGEESFRINYPVVDQAIAASLIRRFEQYQGYEKTFFGLFTSDRL [700]
RLDDDSLAAACENLEVALKSGEHKIDGKELSDDELGLIQIILKKSNGPLDILQLKERPFYPNATVAYRILLTIPVTVASAERSFSKLLKLLKSLRSTM [800]
TQERLNGLATIALEKDILEKINYEDIIEDFISRNRTRMMLFSTS* [845]
    
```





FIGURE 5.—Sizes of *Pi-ta* introgressions in *O. sativa* Asian and U. S. cultivated rice through breeding selection during domestication. The *Pi-ta* region of Tadukan, which is the major donor for *Pi-ta* in Asian cultivars, was used to compare the size of the introgression block with other *Pi-ta*-containing cultivars. The solid bar represents the identical sequence of *Pi-ta* introgressed into resistant cultivars. The shaded bar represents sequence polymorphisms unrelated to the *Pi-ta* introgression that resulted from recombination events at the genomic region of *Pi-ta*. Sequence polymorphisms are marked on the breakpoint of the *Pi-ta* introgression block. P/A indicates the presence and absence of polymorphism.

was consistently maintained in resistant accessions to *M. oryzae* containing *AVR-Pita*. Significantly, one of the largest linkage blocks of resistant *Pi-ta* was identified in backcrossing and elite rice cultivars (JIA 2009). The identification of a large linkage block around *Pi-ta* raised at least two possibilities. First, other blast *R* genes in the *Pi-ta* region also introgressed into diverse elite rice cultivars. Other *R* genes such as *Pi-ta²*, *Pi39*, and *Pi20(t)* (RYBKA *et al.* 1997; LIU *et al.* 2007; LI *et al.* 2008) were also mapped at the *Pi-ta* region, but it was unknown if these and/or other unknown *R* genes were clustered in the *Pi-ta* region that have been introgressed as a large linkage block. Second, other components for the *Pi-ta*-mediated resistance reside within the 5-Mb region to form a superlocus. *R*-gene-mediated resistance may involve additional *R* genes that may be physically linked to provide a complete resistance to a plant pathogen. In tomato, Prf, a NBS–LRR protein, was identified to be involved in the Pto-mediated resistance (MUCYNA *et al.* 2006). In rice, at least two NBS–LRR proteins at the *Pikm* and *Pi5* loci have been identified as providing complete resistance to blast (ASHIKAWA *et al.* 2008; LEE *et al.* 2009). At the *Pikm* locus, *Pikm1-TS* and *Pikm2-TS* within

2.5 kb are required for *Pikm*-mediated disease resistance (ASHIKAWA *et al.* 2008). Similarly, two NBS–LRR proteins within 50 kb, *Pi5-1* and *Pi5-2*, were required for complete resistance (LEE *et al.* 2009). At the *Pi-ta* locus, another gene *Ptr(t)* was found to be essential for *Pi-ta*-mediated resistance (JIA and MARTIN 2008). The possible artificial selection of the large *Pi-ta* genomic region has been reported for maintaining the broad spectrum of *Pi-ta*-mediated blast resistance (JIA 2009). Taken together with other studies, this study suggests that other components such as *Ptr(t)* or *R* genes for the *Pi-ta*-mediated resistance may occur within at least 5 Mb of the *Pi-ta* region.

Simple insertion/deletion or transposon may play an important role in *R*-gene evolution. It has been reported that 18.8% of total *R* genes in Arabidopsis and 22.2% in rice are under presence/absence polymorphism (MEYERS *et al.* 2003; SHEN *et al.* 2006). An example of transposon and *R*-gene activation was found in the *Pit* gene. The insertion of a long-terminal-repeat retrotransposon in the promoter of *Pit* was predicted to regulate *Pit* transcription and its function for resistance (HAYASHI and YOSHIDA 2009). In our study, we found a transposon

FIGURE 4.—Genomic organization around *indica* (resistant *Pi-ta*) and *japonica* (susceptible *Pi-ta*) cultivars. (A) Comparisons of genomic regions around the *Pi-ta* locus between Nipponbare and Katy. (B) An insertion in the proximate *Pi-ta* promoter region differentiates the size of hybridized bands between two susceptible cultivars (Nipponbare and M202) and six resistant cultivars (*Pi-ta*) (top). Schematic of the *Pi-ta* genomic region with indicated restriction enzymes (bottom). (C) The two domains (shaded)—zinc finger in transposases and transcription factors (ZnF_TTF) and hAT family dimerization (hATC)—were identified by searching the conserved domain of proteins from NCBI database.

in the proximity of the *Pi-ta* promoter in resistant cultivars carrying *Pi-ta*, which was absent in accessions without *Pi-ta*. This finding suggests that the transposon may activate the *Pi-ta*-mediated resistance. Further study may lead to a better understanding of any associations of the transposon with *Pi-ta*-mediated resistance.

The divergence of *indica* and *japonica* subgroups in *O. sativa* was predicted to be caused by two independent domestications from geographically divergent *O. rufipogon* populations (LONDO and SCHAAL 2007). The *Pi-ta* haplotypes of *indica* or *japonica* origin were identified in this study (Figure 2). Resistant *Pi-ta* was found only in *indica*, weedy rice, *japonica* cultivars carrying the *indica*-derived *Pi-ta* region and *O. rufipogon*, suggesting that resistant *Pi-ta* did not originate from *japonica*. The *Pi-ta* variants in H5 and H6 were found only in *japonica* accessions, while H2 and H3 were found only in *indica* (Figure 2), consistent with a previous study (LONDO and SCHAAL 2007). The *Pi-ta* variant containing Ala-918 (PT1) separates the resistant *Pi-ta* variant from other variants in both *O. sativa* and *O. rufipogon*. This suggests that PT1 existed before the divergence of the two subspecies *indica* and *japonica*. The recent divergence of resistant *Pi-ta* from susceptible *Pi-ta* has also been proposed from the previous studies (HUANG *et al.* 2008; YOSHIDA and MIYASHITA 2009). Most of the *Pi-ta* variants possess serine at the position of 918. There was no amino acid sequence polymorphism in the group with PT1; however, significant amino acid polymorphism was identified in groups containing Ser-918, consistent with previous reports (HUANG *et al.* 2008; WANG *et al.* 2008; YOSHIDA and MIYASHITA 2009). These findings further suggest that there was recently a strong selection constraint on the resistant *Pi-ta* protein (PT1), and such pressures were not observed on other *Pi-ta* protein variants.

An excess of amino acid substitutions over neutral expectations were observed in the NBS region in both *O. sativa* and *O. rufipogon*, indicating that positive directional selection favored amino acid substitutions in the domain. The NBS domain in diverse proteins with ATP or GTP binding activity is involved in activating the NBS-LRR protein in resistance. It has been documented that the Toll-interleukin 1 receptor region of the *L* class of flax rust *R* genes (ELLIS *et al.* 1999) and the N-terminal domain with the NBS region of tomato MI protein (HWANG *et al.* 2000) are key regulators of signal transduction of disease resistance. Our findings suggest that the highly diversified NBS region may be important for maintaining the integrity of the *Pi-ta* protein with the LRD domain. In the LRD of the *Pi-ta* protein, the level of synonymous diversity was found to exceed the level of nonsynonymous diversity, which is suggestive of possible purifying selection acting on this domain. The $K_a:K_s$ ratio for the LRD of *Pi-ta* ($K_a:K_s = 0.692-0.741$) is relatively low compared to that observed in other LRRs (ELLIS *et al.* 1999; MAURICIO *et al.* 2003; ROSE *et al.* 2004;

BAKKER *et al.* 2006; ORGIL *et al.* 2007). It is possible that conservation of LRD in the *Pi-ta* protein may be necessary for recognizing AVR-Pita for the signal transduction (JIA *et al.* 2000). High nucleotide diversity and a large number of AVR-Pita haplotypes were recently identified, suggesting that AVR-Pita is under diversifying selection (Y. DAI and Y. JIA, unpublished data). Diversified selection at NBS and purifying selection against amino acid variants in the conserved functional LRD region may have played a major role in shaping the molecular evolution of *Pi-ta*.

In conclusion, this study revealed that (1) a transposon may be a part of the evolution with resistant *Pi-ta*, (2) all components needed for the *Pi-ta*-mediated resistance may be embedded within 5 Mb, and (3) strong artificial selection has acted at and around resistant *Pi-ta* in the modern cultivated rice *O. sativa*, while such selection is absent in cultivars without resistant *Pi-ta*. These findings suggest that the evolution of *Pi-ta* is much more complicated than previously documented. Further studies will be necessary for a better understanding of the molecular mechanism of *Pi-ta*-mediated signal recognition and transduction pathway.

The authors thank Briana Gross (Washington University), Michael Reagon (University of Massachusetts), David Gealy, and Anna McClung for extensive support throughout the research. We are also grateful to all members of the Molecular Plant Pathology lab and other staff members at the U. S. Department of Agriculture-Agricultural Research Service Dale Bumpers National Rice Research Center for their technical assistance (<http://ars.usda.gov/spa/dbnrrc/mpp>). This work was supported by the National Science Foundation under grant no. 0638820.

LITERATURE CITED

- AKASHI, H., 1999 Inferring the fitness effects of DNA mutations from polymorphism and divergence data: statistical power to detect directional selection under stationary and free recombination. *Genetics* **151**: 221-238.
- ASHIKAWA, I., N. HAYASHI, H. YAMANE, H. KANAMORI, J. WU *et al.*, 2008 Two adjacent nucleotide-binding site-leucine-rich repeat class genes are required to confer *Pikm*-specific rice blast resistance. *Genetics* **180**: 2267-2276.
- BAKKER, E. G., C. TOOMAJIAN, M. KREITMAN and J. BERGELSON, 2006 A genome-wide survey of *R* gene polymorphisms in *Arabidopsis*. *Plant Cell* **18**: 1803-1818.
- BALLINI, E., J.-B. MOREL, G. DROC, A. PRICE, B. COUTOIS *et al.*, 2008 A genome-wide meta-analysis of rice blast resistance genes and quantitative trait loci provides new insights into partial and complete resistance. *Mol. Plant-Microbe Interact.* **21**: 859-868.
- BRYAN, G. T., K. S. WU, L. FARRALL, Y. JIA, H. P. HERSHEY *et al.*, 2000 A single amino-acid difference distinguishes resistant and susceptible alleles of the rice blast resistance gene *Pi-ta*. *Plant Cell* **12**: 2033-2045.
- CHEN, X. W., J. SHANG, D. CHEN, C. LEI, Y. ZOU *et al.*, 2006 A B-lectin receptor kinase gene conferring rice blast resistance. *Plant J.* **46**: 794-804.
- CLEMENT, M., D. POSADA and K. A. GRANDALL, 2000 TCS: a computer program to estimate gene genealogies. *Mol. Ecol.* **9**: 1657-1660.
- COUCH, B. C., and L. M. KOHN, 2002 A multilocus gene genealogy concordant with host preference indicates segregation of a new species, *Magnaporthe oryzae*, from *M. grisea*. *Mycologia* **94**: 683-693.
- ELLIS, J. G., G. J. LAWRENCE, J. E. LUCK and P. N. DODDS, 1999 Identification of regions in alleles of the flax rust resis-

- tance gene L that determine differences in gene-for-gene specificity. *Plant Cell* **11**: 495–506.
- FLOR, H. H., 1971 Current status of the gene-for-gene concept. *Annu. Rev. Phytopathol.* **9**: 275–296.
- FU, Y. X., and W. H. LI, 1993 Statistical tests of neutrality of mutations. *Genetics* **133**: 693–709.
- FUKUOKA, S., N. SAKA, H. KOGA, K. ONO, T. SHIMIZU *et al.*, 2009 Loss of function of a proline-containing protein confers durable disease resistance in rice. *Science* **325**: 998–1001.
- GIBBONS, J. W., K. A. K. MOLDENHAUER, K. GRAVOIS, F. N. LEE, J. L. BERNHARDT *et al.*, 2006 Registration of 'Cybonnet' rice. *Crop Sci.* **46**: 2317–2318.
- GRAVOIS, K. A., K. A. K. MOLDENHAUER, F. N. LEE, R. J. NORMAN, R. S. HELMS *et al.*, 1995 Registration of 'Kaybonnet' rice. *Crop Sci.* **35**: 586–587.
- HAYASHI, K., and H. YOSHIDA, 2009 Refunctionalization of the ancient rice blast disease resistance gene *Pit* by the recruitment of a retrotransposon as a promoter. *Plant J.* **57**: 413–425.
- HUANG, C., S. HWANG, Y. CHIANG and T. LIN, 2008 Molecular evolution of the *Pi-ta* gene resistant to rice blast in wild rice (*Oryza rufipogon*). *Genetics* **179**: 1527–1538.
- HULBERT, S. H., C. A. WEBB, S. M. SMITH and Q. SUN, 2001 Resistance gene complexes: evolution and utilization. *Annu. Rev. Phytopathol.* **39**: 285–312.
- HWANG, C. F., A. V. BHAKTA, G. M. TRUESDELL, W. M. PUDLO and V. M. WILLIAMSON, 2000 Evidence for a role of the N terminus and leucine-rich repeat region of the *Mi* gene product in regulation of localized cell death. *Plant Cell* **12**: 1319–1329.
- JIA, Y., 2003 Marker assisted selection for the control of rice blast disease. *Pesticide Outlook* **14**: 150–152.
- JIA, Y., 2009 Artificial introgression of a large chromosome fragment around the rice blast resistance gene *Pi-ta* in backcross progeny and several elite rice cultivars. *Heredity* **103**: 333–339.
- JIA, Y., and R. MARTIN, 2008 Identification of a new locus, *Pit(t)*, required for rice blast resistance gene *Pi-ta*-mediated resistance. *Mol. Plant-Microbe Interact.* **21**: 396–403.
- JIA, Y., S. A. McADAMS, G. T. BRYAN, H. P. HERSHEY and B. VALENT, 2000 Direct interaction of resistance gene and avirulence gene products confers rice blast resistance. *EMBO J.* **19**: 4004–4014.
- JIA, Y., G. T. BRYAN, L. FARRALL and B. VALENT, 2003 Natural variation at the *Pi-ta* rice blast resistance locus. *Phytopathology* **93**: 1452–1459.
- JIA, Y., M. REDUS, Z. WANG and J. N. RUTGER, 2004a Development of a SNLP marker from the *Pi-ta* blast resistance gene by tri-primer PCR. *Euphytica* **138**: 97–105.
- JIA, Y., Z. WANG, R. G. FJELLSTROM, K. A. K. MOLDENHAUER, M. A. AZAM *et al.*, 2004b Rice *Pi-ta* gene confers resistance to the major pathotypes of the rice blast fungus in the US. *Phytopathology* **94**: 296–301.
- JIA, Y., F. N. LEE and A. McCLUNG, 2009a Determination of resistance spectra of the *Pi-ta* and *Pi-k* genes to U.S. races of *Magnaporthe oryzae* causing rice blast in a recombinant inbred line population. *Plant Dis.* **93**: 639–644.
- JIA, Y., X. WANG, S. COSTANZO and S. LEE, 2009b Understanding the coevolution of rice blast resistance gene *Pi-ta* and *Magnaporthe oryzae* avirulence gene *AVR-Pita*, pp. 137–147 in *Advances in Genetics, Genomics and Control of Rice Blast Disease*, edited by G. L. WANG and B. VALENT. Springer Science, New York.
- LEE, S.-K., M.-Y. SONG, Y.-S. SEO, H.-K. KIM, S. KO *et al.*, 2009 Rice *Pi5*-mediated resistance to *Magnaporthe oryzae* requires the presence of two CC-NB-LRR genes. *Genetics* **181**: 1627–1638.
- LI, W., C. L. LEI, Z. J. CHENG, Y. L. JIA, D. Y. HUANG *et al.*, 2008 Identification of SSR markers for a broad-spectrum blast resistance gene *Pi20(t)* for marker-assisted breeding. *Mol. Breed.* **22**: 141–149.
- LINARES, O. F., 2002 African rice (*Oryza glaberrima*): history and future potential. *Proc. Natl. Acad. Sci. USA* **99**: 16360–16365.
- LIU, X. Q., Q. Z. YANG, F. LIN, L. X. HUA, C. T. WANG *et al.*, 2007 Identification and fine mapping of *Pi39(t)*, a major gene conferring the broad-spectrum resistance to *Magnaporthe oryzae*. *Mol. Genet. Genomics* **278**: 403–410.
- LONDO, J. P., and B. A. SCHAAL, 2007 Origins and population genetics of US weedy red rice in the USA. *Mol. Ecol.* **16**: 4523–4535.
- LONDO, J. P., Y. CHIANG, K. HUNG, T. CHIANG and B. A. SCHAAL, 2006 Phylogeography of Asian wild rice, *Oryza rufipogon*, reveals multiple independent domestications of cultivated rice, *Oryza sativa*. *Proc. Natl. Acad. Sci. USA* **103**: 9578–9583.
- MAURICIO, R., E. A. STAHL, T. KORVES, D. TIAN, M. KREITMAN *et al.*, 2003 Natural selection for polymorphism in the disease resistance gene *Rps2* of *Arabidopsis thaliana*. *Genetics* **163**: 735–746.
- McCLUNG, A. M., M. A. MARCHETTI, B. D. WEBB and C. N. BOLLICH, 1999 Registration of 'Madison' rice. *Crop Sci.* **39**: 1256.
- MEYERS, B. C., A. KOZIK, A. GRIEGO, H. KUANG and R. W. MICHELMORE, 2003 Genome-wide analysis of NBS-LRR-encoding genes in *Arabidopsis*. *Plant Cell* **15**: 809–834.
- MOLDENHAUER, K. A. K., K. A. GRAVOIS, F. N. LEE, R. J. NORMAN, J. L. BERNHARDT *et al.*, 1998 Registration of 'Drew' rice. *Crop Sci.* **38**: 896–897.
- MOLDENHAUER, K. A. K., F. N. LEE, J. W. GIBBONS, J. L. BERNHARDT, R. J. NORMAN *et al.*, 2007 Registration of 'Ahrent' rice. *Crop Sci.* **47**: 446–447.
- MUCYNA, T. S., A. CLEMENTEA, V. M. E. ANDRIOTISA, A. L. BALMUTHA, G. E. D. OLDROYDB *et al.*, 2006 The tomato NBARC-LRR protein Prf interacts with Pto kinase in vivo to regulate specific plant immunity. *Plant Cell* **18**: 2792–2806.
- ORGIL, U., H. ARAKIT, S. TANGCHAIBURANA, R. HERKEY and S. XIAO, 2007 Intraspecific genetic variations, fitness cost and benefit of *RPW8*, a disease resistance locus in *Arabidopsis thaliana*. *Genetics* **176**: 2317–2333.
- ROSE, L. E., P. D. BITTNER-EDDY, C. H. LANGLEY, E. B. HOLUB, R. W. MICHELMORE *et al.*, 2004 The maintenance of extreme amino acid diversity at the disease resistance gene, *RPPI3*, in *Arabidopsis thaliana*. *Genetics* **166**: 1517–1527.
- ROSSMAN, A. Y., R. J. HOWARD and B. VALENT, 1990 *Pyricularia oryzae*, the correct name for the rice blast fungus. *Mycologia* **82**: 509–512.
- ROZAS, J., J. C. SÁNCHEZ-DELBARRIO, X. MESSEGYER and R. ROZAS, 2003 DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* **19**: 2496–2497.
- ROZEN, S., and H. SKALETSKY, 2000 Primer3 on the WWW for general users and for biologist programmers. *Methods Mol. Biol.* **132**: 365–386.
- RYBKA, K., M. MIYAMOTO, I. ANDO, A. SAITO and S. KAWASAKI, 1997 High resolution mapping of indica-derived rice blast resistance genes II. *Pi-ta²* and *Pi-ta* and a consideration of their origin. *Mol. Plant-Microbe Interact.* **10**: 517–524.
- SABETI, P. C., D. E. REICH, J. M. HIGGINS, H. Z. P. LEVINE, D. J. RICHTER *et al.*, 2002 Detecting recent positive selection in the human genome from haplotype structure. *Nature* **419**: 832–837.
- SHEN, J., H. ARAKI, L. CHEN, J.-Q. CHEN and D. TIAN, 2006 Unique evolutionary mechanism in *R*-genes under the presence/absence polymorphism in *Arabidopsis thaliana*. *Genetics* **172**: 1243–1250.
- TALBOT, N. J., 2003 On the trail of a cereal killer: exploring the biology of *Magnaporthe grisea*. *Annu. Rev. Microbiol.* **57**: 177–202.
- TAMURA, K., J. DUDLEY, M. NEI and S. KUMAR, 2007 MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* **24**: 1596–1599.
- VALENT, B., L. FARRALL and F. G. CHUMLEY, 1991 *Magnaporthe grisea* genes for pathogenicity and virulence identified through a series of backcrosses. *Genetics* **127**: 87–101.
- WANG, X., Y. JIA, Q. Y. SHU and D. WU, 2008 Haplotype diversity at the *Pi-ta* locus in cultivated rice and its wild relatives. *Phytopathology* **98**: 1305–1311.
- WANG, Z., Y. JIA, J. N. RUTGER and Y. XIA, 2007 Rapid survey for presence of a blast resistance gene *Pi-ta* in rice cultivars using the dominant DNA markers derived from portions of the *Pi-ta* gene. *Plant Breed.* **126**: 36–42.
- YAMANAKA, S., I. NAKAMURA, H. NAKAI and Y. SATO, 2003 Dual origin of the cultivated rice based on molecular markers of newly collected annual and perennial strains of wild rice species, *Oryza nivara* and *O. rufipogon*. *Genet. Resour. Crop Evol.* **50**: 529–538.
- YOSHIDA, K., and N. T. MIYASHITA, 2009 DNA polymorphism in the blast disease resistance gene *Pita* of the wild rice *Oryza rufipogon* and its related species. *Genes Genet. Syst.* **84**: 121–136.

GENETICS

Supporting Information

<http://www.genetics.org/cgi/content/full/genetics.109.108266/DC1>

Evolutionary Dynamics of the Genomic Region Around the Blast Resistance Gene *Pi-ta* in AA Genome *Oryza* Species

Seonghee Lee, Stefano Costanzo, Yulin Jia, Kenneth M. Olsen and Ana L. Caicedo

Copyright © 2009 by the Genetics Society of America

DOI: 10.1534/genetics.109.108266

TABLE S1

Description of rice accessions of seven *Oryza* species used in the present study

Accession	Species	Sub population	Origin	V ^a
Rathuwee	<i>O. sativa</i>	<i>indica</i>	Sri Lanka	PT3
Khao Dawk Mali -105	<i>O. sativa</i>	<i>indica</i>	Thailand	PT3
LalAman	<i>O. sativa</i>	<i>indica</i>	India	PT8
Dholi Boro	<i>O. sativa</i>	<i>indica</i>	Bangladesh	PT1
Ai-chiao-hong	<i>O. sativa</i>	<i>indica</i>	China	PT2
Chau	<i>O. sativa</i>	<i>indica</i>	Vietnam	PT3
Chhote-Dhan	<i>O. sativa</i>	<i>indica</i>	Nepal	PT3
Popot-165	<i>O. sativa</i>	<i>indica</i>	Indonesia	PT1
Dee-Geo-Woo-Gen	<i>O. sativa</i>	<i>indica</i>	USA	PT2
C101A51	<i>O. sativa</i>	<i>indica</i>	The Philippines	PT2
C101PKT	<i>O. sativa</i>	<i>indica</i>	The Philippines	PT1
IR64	<i>O. sativa</i>	<i>indica</i>	The Philippines	PT1
Raminad Str. 3	<i>O. sativa</i>	<i>indica</i>		PT3
Tadukan	<i>O. sativa</i>	<i>indica</i>	The Philippines	PT1
Tetep	<i>O. sativa</i>	<i>indica</i>	Vietnam	PT1
Tc Qing	<i>O. sativa</i>	<i>indica</i>	China	PT1
Jhona-349	<i>O. sativa</i>	<i>Aus-indica</i>	India	PT2
Kasalath	<i>O. sativa</i>	<i>Aus-indica</i>	India	PT2
DV85	<i>O. sativa</i>	<i>Aus-indica</i>	Bangladesh	PT2
BJ-1	<i>O. sativa</i>	<i>Aus-indica</i>	India	PT2
Dhala-Shaitta	<i>O. sativa</i>	<i>Aus-indica</i>	Bangladesh	PT2
Bei Khe	<i>O. sativa</i>	<i>Aus-indica</i>	Cambodia	PT8
Aus 196	<i>O. sativa</i>	<i>Aus-indica</i>	Bangladesh	PT3
DA 13	<i>O. sativa</i>	Aromatic	Bangladesh	PT2
Dom Sofid	<i>O. sativa</i>	Aromatic	Iran	PT2
ARC-13829	<i>O. sativa</i>	Aromatic	India	PT3
Ta hung ku	<i>O. sativa</i>	<i>Temperate japonica</i>	China	PT22
Kamenoo	<i>O. sativa</i>	<i>Temperate japonica</i>	Japan	PT22
Nep-Hoa-Vang	<i>O. sativa</i>	<i>Temperate japonica</i>	Vietnam	PT22
SHOEMED	<i>O. sativa</i>	<i>Temperate japonica</i>	The Philippines	PT22
Khao Hawm	<i>O. sativa</i>	<i>Tropical japonica</i>	Thailand	PT22
Mirti	<i>O. sativa</i>	<i>Tropical japonica</i>	Bangladesh	PT22
KU115	<i>O. sativa</i>	<i>Tropical japonica</i>	Thailand	PT22
Cicik Beton	<i>O. sativa</i>	<i>Tropical japonica</i>	Indonesia	PT8
Gotak Gatik	<i>O. sativa</i>	<i>Tropical japonica</i>	Indonesia	PT1
Asse Y Pung	<i>O. sativa</i>	<i>Tropical japonica</i>	The Philippines	PT22
Kotobuki-mochi	<i>O. sativa</i>	<i>Tropical japonica</i>	Japan	PT22
Trembese	<i>O. sativa</i>	<i>Tropical japonica</i>	Indonesia	PT22
Pi4	<i>O. sativa</i>	<i>Tropical japonica</i>	Japan	PT1
Reiho	<i>O. sativa</i>	<i>Tropical japonica</i>	Japan	PT1
Yashiro-mochi	<i>O. sativa</i>	<i>Tropical japonica</i>	Japan	PT1

Fukunoshiki	<i>O. sativa</i>	<i>Tropical japonica</i>	Japan	PT22
Shimokita	<i>O. sativa</i>	<i>Tropical japonica</i>	Japan	PT3
Tsuyake	<i>O. sativa</i>	<i>Tropical japonica</i>	Japan	PT22
M202	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT8
Bengal	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT22
Blue Rose	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT8
Carolina Gold	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT8
CL121	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT22
CL161	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT22
Cypress	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT22
Delitus	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT22
Drew	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT1
Edith	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT8
Palmyra	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT22
Rexoro	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT22
Zenith	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT22
Lemont	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT22
Katy	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT1
Banks	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT1
Madison	<i>O. sativa</i>	<i>Tropical japonica</i>	USA	PT1
1004-01 ^b	<i>O. sativa</i>	Weedy rice (SH)	Dunklin Co., MO, USA	PT1
1025-01	<i>O. sativa</i>	Weedy rice (BHA)	Clay Co., AR, USA	PT8
1081-01	<i>O. sativa</i>	Weedy rice (BHA)	Perry Co., AR, USA	PT2
1091-01	<i>O. sativa</i>	Weedy rice (SH)	Poinsett Co., AR, USA	PT1
1096-01	<i>O. sativa</i>	Weedy rice (BHA)	Arkansas Co., AR, USA	PT2
1196-01	<i>O. sativa</i>	Weedy rice (SH)	Crittenden Co., AR, USA	PT1
1098-01	<i>O. sativa</i>	Weedy rice (SH)	Bollinger Co., AR, USA	PT1
1134-01	<i>O. sativa</i>	Weedy rice (SH)	Lee Co., AR, USA	PT1
1135-01	<i>O. sativa</i>	Weedy rice (SH)	Desha Co., AR, USA	PT1
1141-01	<i>O. sativa</i>	Weedy rice (SH)	Lawrence Co., AR, USA	PT1
1160-01	<i>O. sativa</i>	Weedy rice (SH)	Morehouse Co., MO, USA	PT1
1179-01	<i>O. sativa</i>	Weedy rice (SH)	Coahoma Co., MS, USA	PT1
1188-01	<i>O. sativa</i>	Weedy rice (BHA)	East Carrol Co., LA, USA	PT8
10A	<i>O. sativa</i>	Weedy rice (BHA)	Prairie Co., AR, USA	PT2
16B	<i>O. sativa</i>	Weedy rice (SH)	Prairie Co., AR, USA	PT1
18A	<i>O. sativa</i>	Weedy rice (BHA)	Arkansas Co., AR, USA	PT2
1995-15	<i>O. sativa</i>	Weedy rice (SH)	Shaw, MS, USA	PT1
1996-1	<i>O. sativa</i>	Weedy rice (MIX)	Arkansas Co., AR, USA	PT3
1996-5	<i>O. sativa</i>	Weedy rice (SH)	Mississippi, USA	PT1
1996-9	<i>O. sativa</i>	Weedy rice (BHA)	Mississippi, USA	PT2
LA3	<i>O. sativa</i>	Weedy rice (BHA)	Crowley, LA, USA	PT1
MS4	<i>O. sativa</i>	Weedy rice (MIX)	Mississippi, USA	PT2
StgB	<i>O. sativa</i>	Weedy rice (BHA)	Stuttgart, AR, USA	PT2
StgS	<i>O. sativa</i>	Weedy rice (BHA)	Stuttgart, AR, USA	PT1

TX4	<i>O. sativa</i>	Weedy rice (BHA)	Katy, TX, USA	PT8
1001-01	<i>O. sativa</i>	Weedy rice (SH)	Cross Co., AR, USA	PT1
1002-02	<i>O. sativa</i>	Weedy rice (SH)	Independence Co., AR, USA	PT1
1005-02	<i>O. sativa</i>	Weedy rice (BHA)	Faulkner Co., AR, USA	PT2
1042-01	<i>O. sativa</i>	Weedy rice (BHA)	Jefferson Co., AR, USA	PT2
1047-01	<i>O. sativa</i>	Weedy rice (SH)	Morehouse Co., LA, USA	PT1
1073-02	<i>O. sativa</i>	Weedy rice (SH)	Butler Co., MO, USA	PT1
1092-02	<i>O. sativa</i>	Weedy rice (BR)	Coahoma Co., MS, USA	PT1
1111-01	<i>O. sativa</i>	Weedy rice (BR)	Woodruff Co., AR, USA	PT1
1190-01	<i>O. sativa</i>	Weedy rice (SH)	East Carroll Co., LA, USA	PT1
1199-01	<i>O. sativa</i>	Weedy rice (SH)	Ripley Co., MO, USA	PT1
1300-02	<i>O. sativa</i>	Weedy rice (BR)	Dunklin Co., MO, USA	PT1
1344-02	<i>O. sativa</i>	Weedy rice (SH)	Stoddard Co., MO, USA	PT1
Prairie Co. Short - 8	<i>O. sativa</i>	Weedy rice (BHA)	Prairie Co., AR, USA	PT2
Prairie Co. Tall - 10	<i>O. sativa</i>	Weedy rice (BHA)	Prairie Co., AR, USA	PT2
Prairie Co. Tall - 11	<i>O. sativa</i>	Weedy rice (BHA)	Prairie Co., AR, USA	PT2
Prairie Co. Tall - 17	<i>O. sativa</i>	Weedy rice (BHA)	Prairie Co., AR, USA	PT2
1995-12	<i>O. sativa</i>	Weedy rice (SH)	Crowley, LA, USA	PT1
1995-13	<i>O. sativa</i>	Weedy rice (BHA)	Crowley, LA, USA	PT1
1995-14	<i>O. sativa</i>	Weedy rice (BHA)	Crowley, LA, USA	PT1
1996-8	<i>O. sativa</i>	Weedy rice (SH)	Mississippi, USA	PT1
2002-51	<i>O. sativa</i>	Weedy rice (MIX)	Arkansas, USA	PT3
2004-1-A	<i>O. sativa</i>	Weedy rice (MIX)	Amagon, AR, USA	PT22
1183-01	<i>O. sativa</i>	Weedy rice (BR)	Chicot Co., AR, USA	PT1
1166-02	<i>O. sativa</i>	Weedy rice (BHA)	Coahoma Co., MS, USA	PT2
1107-01	<i>O. sativa</i>	Weedy rice (BHA)	Cross Co., AR, USA	PT2
1210-02	<i>O. sativa</i>	Weedy rice (SH)	Dunklin Co., MO, USA	PT1
1214-02	<i>O. sativa</i>	Weedy rice (BHA)	East Carroll Co., LA, USA	PT8
1163-01	<i>O. sativa</i>	Weedy rice (SH)	Morehouse Co., LA, USA	PT1
1120-02	<i>O. sativa</i>	Weedy rice (BR)	St. Francis Co., AR, USA	PT1
1333-02	<i>O. sativa</i>	Weedy rice (SH)	Stoddard Co., MO, USA	PT1
1202-02	<i>O. sativa</i>	Weedy rice (BHA)	Jefferson Co., AR, USA	PT1
2002-2-pot 1	<i>O. sativa</i>	Weedy rice (BHA)	Jackson Co., AR, USA	PT2
2002-2-pot 21	<i>O. sativa</i>	Weedy rice (MIX)	Lawrence Co., AR, USA	PT22
IRGC81990	<i>O. rufipogon</i>	Wild rice	Myanmar	PT12
IRGC100588	<i>O. rufipogon</i>	Wild rice	Taiwan	PT23
IRGC100904	<i>O. rufipogon</i>	Wild rice	Thailand	PT3
IRGC104501	<i>O. rufipogon</i>	Wild rice	India	PT14
IRGC104599	<i>O. rufipogon</i>	Wild rice	Sri Lanka	PT17
IRGC104624	<i>O. rufipogon</i>	Wild rice	China	PT9
IRGC104714	<i>O. rufipogon</i>	Wild rice	Thailand	PT15
IRGC104833	<i>O. rufipogon</i>	Wild rice	Thailand	PT11
IRGC104871	<i>O. rufipogon</i>	Wild rice	Thailand	PT8
IRGC105388	<i>O. rufipogon</i>	Wild rice	Thailand	PT16

IRGC105491	<i>O. rufipogon</i>	Wild rice	Malaysia	PT2
IRGC105711	<i>O. rufipogon</i>	Wild rice	India	PT10
IRGC105720	<i>O. rufipogon</i>	Wild rice	Cambodia	PT16
IRGC105855	<i>O. rufipogon</i>	Wild rice	Thailand	PT21
IRGC105888	<i>O. rufipogon</i>	Wild rice	Bangladesh	PT5
IRGC106086	<i>O. rufipogon</i>	Wild rice	India	PT1
IRGC106103	<i>O. rufipogon</i>	Wild rice	India	PT3
IRGC106122	<i>O. rufipogon</i>	Wild rice	India	PT9
IRGC106134	<i>O. rufipogon</i>	Wild rice	India	PT3
IRGC106150	<i>O. rufipogon</i>	Wild rice	Laos	PT3
IRGC106163	<i>O. rufipogon</i>	Wild rice	Laos	PT3
IRGC106168	<i>O. rufipogon</i>	Wild rice	Vietnam	PT18
IRGC106169	<i>O. rufipogon</i>	Wild rice	Vietnam	PT11
IRGC106321	<i>O. rufipogon</i>	Wild rice	Cambodia	PT13
IRGC106346	<i>O. rufipogon</i>	Wild rice	Myanmar	PT12
IRGC106453	<i>O. rufipogon</i>	Wild rice	Indonesia	PT19
IRGC106518	<i>O. rufipogon</i>	Wild rice	Vietnam	PT3
IRGC106523	<i>O. rufipogon</i>	Wild rice	Papau New	PT3
IRGC86662	<i>O. nivara</i>	Wild rice	Thailand	PT8
IRGC103821	<i>O. nivara</i>	Wild rice	China	PT2
IRGC101226	<i>O. barthii</i>	Wild rice	Mali	PT25
IRGC104081	<i>O. barthii</i>	Wild rice	Nigeria	PT26
IRGC86779	<i>O. glaberrima</i>	African cultivated rice	Liberia	PT26
IRGC100983	<i>O. glaberrima</i>	African cultivated rice	Nigeria	PT26
IRGC104587	<i>O. glaberrima</i>	African cultivated rice	Burkina Faso	PT26
IRGC102410	<i>O. glaberrima</i>	African cultivated rice	Mali	PT26
IRGC105561	<i>O. glumaepatula</i>	Wild rice	Colombia	PT28
IRGC105670	<i>O. glumaepatula</i>	Wild rice	Brazil	PT27
IRGC93261	<i>O. meridionalis</i>	Wild rice	Indonesia	PT3
IRGC101148	<i>O. meridionalis</i>	Wild rice	Australia	PT7

The sequences obtained from GenBank^c

O. rufipogon: EU346961, EU346962, EU346963, EU346964, EU346965, EU346966, EU346968, EU346970, EU346972, EU346974, EU346976, EU346978, EU346979, EU346980, EU346981, EU346982, EU346984, EU346986, EU346987, EU346988, EU346990, EU346991, EU346992, EU346994, EU346995, EU346997, EU346998, EU346999, EU347000, EU347003, EU347004, EU347006, EU770209, EU770213, EU770214, AB364479, AB364480, AB364481, AB364482, AB364483, AB364484, AB364485, AB364486, AB364487, AB364488, AB364489, AB364239, AB364240, AB364241, AB364242,

O. nivara: EU770220

O. glaberrima: EU770215, EU770219

O. barthii: EU346955, EU346957, EU346959, EU346960, EU770218

O. meridionalis: AB364494, AB364497

^a The Pi-ta variants.

^b USDA-ARS DB NRRC weedy rice accession numbers. SH: Strawhull awnless, BHA: Blackhull awned, BR: Brownhull awned, MIX: crop-weed hybrids.

^c Additional sequences of the *Pi-ta* coding region with an intron downloaded from the GenBank database.

TABLE S2

A list of primers for PCR amplification and sequencing the *Pi-ta* locus and its flanking region in seven *Oryza* species, *O. sativa*, *O. rufipogon*, *O. nivara*, *O. meridionalis*, *O. glumaepatula*, *O. barthii*, and *O. glaberrima* used in the present study.

Primer pair	Forward	Reverse
Pita001 ^a	AACACGGGACAAGAAATAGG	CTTCCATTAATGCCCTCTCC
Pita002 ^a	GGCAGCCAAGAATTAACAACATAGGC	AACTTGAGCTGCCATGCATTCTCC
Pita003 ^a	TCGGTCCATGCAAAGATCGTAAGC	CAGGGACCCGCATGATGACACC
Pita004 ^a	GCGATCCATGCTGTCAAATCAGC	GCCTGCCAAGATGGTAGCTCTCC
Pita005 ^a	GACGACTTCCTCGACGAGCTAACG	CAAATGCGTCCGGATAGTTTCAAGG
Pita006 ^a	TCAGTCTTCGGATGTTTGGGAGGT	ACAAGGGAGGCCCTTACGACTATT
Pita007 ^a	CCACAATGGCATGTAACTTATAGCAG	CTCCAAATCATCTAGAGCCAAATAGCC
Pita008 ^a	GAGTATTTGCTTAGGAGTACGTGTCT	CTTGCCCAACAACCTCCACTGAA
Pita009 ^a	CTGCCGTGGCTTCTATCTTTACCT	GCGAACTGCTCCATTGTGTGG
Pita010 ^a	CATTGGATCATTCTCAGACG	AGTACTTGAGAGACCCGAACC
Pita011 ^a	CTTGACACTCTCAAAGGACTGG	CTCTACTCTGAAGACGTGAAGAGG
Pita012 ^a	CTGCCGTGGCTTCTATCTTTACCT	GCGAACTGCTCCATTGTGTGG
Pita013 ^a	TCCCATGGTACTAACTCATGTC	ATGTTGCAAGGGTAATCAGAAG
Pita014 ^a	GGAAGGGCACCTTGTTTAATGTAGC	CAGACTCATTTATTGCCGCTTTGC
Pita015 ^a	AGCTACTGCGTCCCTCGATTTCC	TGCATTGGCTACAATGCGTAGG
Pita22F/Pita22R ^a	AACACGGGACAAGAAATAGG	GCTTACGATCTTTGCATGGACCGA
YL62/YL92 ^a	GGGAGACAGCACCATCGGTG	GCTCGCGAACCTCCTTAGCC
YL63/Pita7R ^a	CAACGTTACTCTCAAGCGAG	CTGCTATAAGTTAACATGCCATTGTGG
YL69/XW14 ^a	GGATGTTTGGGAGGTTGATC	TGCCAAAGCTACAGGTTCAATT
YL74/YL86 ^a	ACCATGTTTGCAAAGTTGAT	CATACACTTGACTTGTCCGA
Pita21F/Pita26R ^a	GGTTCGGGTCTCTCAAGTACT	TGACTACGAGCTTGTGGGATTGCT
Pita24F/Pita23R ^a	CCACACAATGGAGCAGTTCGC	GCTACAAACCTCACCCGAGCA
Weedy2F/Weedy2R ^a	ATGCCAGATGCGATCTATGACCGT	ATGGTGGGATCCAAGCCTACACAA
XW3/XW4 ^a	CTTATGTTGTATGTTGTCCTTC	ACGAGATACCCCTTCCCCTA
WPita17F/WPita17R ^a	GGGATGGTGCATGCATTAAGT	TAGCGTTGGGCAGTGGATAGTTGA
WPita15F/WPita16R ^a	CATGCATCCATGCACACAAGT	TCCAGATGAGCGACAAGTGGTTGA
3-2 ^b	CGCAGCTATCTTTCCGATTTGG	CAATCTCCCGTCATTGTCTTGG
3-4 ^b	GCTCAAGGAGGCACAAAATGG	ATCATGTCACCACCAGGAAGG
3-7 ^b	AGAGTAGGGCAGGGCTAGG	GCATCATGTTGGATGTTGTGG
3-11 ^b	CGAGGAAGTCGCAAATGAGG	TCTGATAGTGTCTAGAGGTGGGAAGG
3-16 ^b	GAGACGGAGAGGGTGATGAGG	GGGTGCGGTTATTGTTGATTAGC
3-22 ^b	GGAAGACACAAATGCCATCATTCC	TGTATTGTCAGAAGGCCGAACAG
5-7 ^c	CTGACCAAAGGCTTGGATGG	TGCAAGAGAGATGAGCTGAACG
5-11 ^c	ACATTGACGAGCAGGCAAGG	AACAAGATCATCACCGTTGTGC
5-14 ^c	CTCCAGGGCGTCTACTTTGC	CTTCCCCAGATGTGAAACTAATGC
5-21 ^c	CTTCTCAGCCGCGTTTCC	TCTTCTCCTTCACCTCCTTCTTGC
5-22 ^c	TACACGCGCAAGAGATACAGG	TAGTGTATCGTCCCATGCTTGTGC
5-24 ^c	TTGGTAGGGATGCTGTTGAGG	CACAGGGTCTTCAGATGAATTGG

^a Primers for the *Pi-ta* locus.

^b Primers for 5' upstream flanking region of the *Pi-ta* locus.

^c Primers for 3' downstream flanking region of the *Pi-ta* locus.

TABLE S3

The *Pi-ta* haplotypes from the *Pi-ta* coding region with an intron in 159 accessions of seven *Oryza* species, *O. sativa*, *O. rufipogon*, *O. nivara*, *O. meridionalis*, *O. glumaepatula*, *O. barthii*, and *O. glaberrima*.

OS1	GTTCCGGTCTGACAGGGCCCGCCCGGATCCAGCCGGCC	-----GT-----	GATTCACACTACTATCGTC	-T-GAT-CCATCCACCT-GACCCGATTCAGACTTTTGAATC-C-G-G-CAATCAATCATGTATGA	---GT-C
OS2C.....	-----	-----	-----A.....
OS3	-----	-----	-----A..A.....
OS4C.....	-----A.....	-----A.....
OS5C.....	-----A.....	-----A.....
OS6	-----C.....	-----	-----
OS7C.....	-----A.....	-----A.....
OS8	G.....AC.....	-----C.....A.....A..A.....
OS9	G.....C.....T.....	-----C.....A.....A..A.....
OS10	G.....C.CG.....T.....	-----C.....A.....T.....A..A.....
OS11	G.....C.CG.....T.....	-----C.....A.T.....T.....A..A.....
OS12	G.....C.CG.....T.....	-----C.....A.T.....T.....A..A.....
OS13	G.....C.....T.....	-----C.....A.....A..A.....
OS14	G.....C.....	-----C.....A.....T.....A..A.....
OS15	G.....C.....T.....	-----C.....A.....A..A.....
OS16	G.....AC.....A.....	-----C.....A.....A..A.....
OS17	G.....C.C.....	-----C.....A.....A..A.....
OS18	G.....C.....T.....	-----C.....A.....A.....
OS19	G.....C.CG.....T.....	-----C.....A.T.....A..A.....
OR20	G.....C.....	-----C.....A.....A..A.....
OR21	GG.....C.G.....T.....	-----C.....A.T.....T.....A..A.....
OR22	G.....C.....T.....	-----C.....A.....A..A.....
OR23	G.....G.C.....T.....AA.....	-----C.....A.A.....A..TA.....
OR24	G.....CT.....	-----C.....A.A.....G.....T.....G.....A..A.....
OR25	G.....C.....	-----G.....C.....A.....T.....A..A.....
OR26	G.....C.....T.....	-----C.....A.....AAT..A..A.....
OR27	G.....AC.....	-----C.....A.....A..A.....
OR28	G.....G.C.....	-----C.....A.....C.....A..A.....-T
OR29C.....	-----	-----	-----A.....
OR30	G.....C.....T.CG.....	-----GC.....-T.....A.....T.....A..A.....
OR31	G.....G.C.....	-----C.....CA.....A..A.....-T
OR32	G.....C.....T.....	-----C.....A.T.....T.....A..A.....
OR33	G.....T.....C.....	-----C.....A.....T.....A..A.....-A..
OR34	G.....C.....	-----C.....C.....A.....T.....A..TA.....
OR35	G.....CT.....	-----C.....A.....G.....T.....AA.....A..A.....
OR36	G.....C.....	-----C.....A.T.G.....T.....A..A.....
OR37	G..A.....C.....T.....	-----C.....A.....G.....A..A.....
OR38	G.....T.G.C.....T.....AA.....GATTTATCCTA	-----C.....A.A.....A..TA.....
OR39	G.....C.....T.....	-----C.....A.T.....T.....A..A.....
OR40	G.G.....C.....	-----C.....A.....A..A.....
OR41	G.....C.....	-----C.....A.....T.....A..A.....
OR42	G.....T.G.C.....	-----C.....C.....A.....T.....A..TA.....
OR43	G.....C.....A.....	-----C.....C.....A.....T.....A..TA.....
OR44	G.....C.....	-----C.....A.....T..T.....A..A.....
ON45	G.....AC.....	-----C.....A.....A..A.....
ON46	G.....CA.C.....A.....G.....	-----C.....C.....A.....T.....A..TA.....
OB47	TG...C.TG...C...G...C...G...A...	-----	CGATTTTCARACT..AGTTGGTA..CTTGGTANT..CMTA...AT...AAC..ATTTA-A..TC..CA-GCCGTAA-ATTT..C..TTT..TTCA...AGAAA..GA.		
OB48	TG...C.TG...C...G...G...A...	-----	CGATTTTCARACT..AGTTGGTA..CTTGGTANT..CMTA...AT...AAC..ATTTA-A..TC..CA...GTAA-ATTT..C..TTT..TTCA...AGAAA..GA.		
OG49	TG...C.TG...C...G...G...A...	-----	CGATTTTCARACT..AGTTGGTA..CTTGGTANT..CMTA...AT...AAC..ATTTA-A..TC..CA...GTAA-ATTT..C..TTT..TTCA...AGAAA..GA.		
OG50	TG...C.TG...C...G...G...A...	-----	CGATTTTCARACT..AGTTGGTA..CTTGGTANT..CMTA...AT...AAC..ATTTA-A..TC..CA...GTAA-ATTT..C..TTT..TTCA...AGAAA..GA.		
OG51	TG...C.T...C...G...G...A...	-----	CGATTTTCARACT..AGTTGGTA..CTTGGTANT..CMTA...AT...AAC..ATTTA-A..TC..CA...GTAA-ATTT..C..TTT..TTCA...AGAAA..GA.		
OG52	G...TGA...C...GA...G...A...	-----	CGATTTTCARACT..AGTTGGT...CTTGGTANT..CMTAC...AT...AAC..ATTTA..A..TC..C...GTAA...TTT..CG..TTTTC...ATAAA..A-		
OG53	C..A..TC...C...C..A...C...A...	-----	CGATTTTCARACT..AGTTGGT...CTTGGTANT..CMTAC...AT...AAC..ATTTA-A..TC..C...GT-A...TTT..CG..TTTTC...ATAAA..A-		


```

TGTATGARTTGGCTATTTGGCTCTAGATGATTTGGAGCTAGTATGTGGCTATATCTATTAATCTTGCCTTT-----RTAGATTTGAGCTGRTCGGCCGTGTCCOCGTTGGGGGGGTCTGA
-----T.
-----T.
-----R. T.
-----T.
-----A. TG
-----C. T.
-----T.
-----T.
-----R. T.
-----R C. T.
-----T. T.
-----T. T.
-----R. TG
-----C. T.
-----C. T.
-----R. T.
-----G. T.
-----R. T.
-----T. T.
-----C. T.
-----T. T.
-----T. T.
-----T. GT.
-----T. T.
-----G. R. TG
-----R. T.
-----T. T.
-----C. T.
-----R. T.
-----R. R. T.
-----T. T.
-----G. T.
-----T. T.
-----T. T.
-----T. T.
-----T. T.
-----T. T.
-----T. TG
-----T. T.
-----R. RT.
-----C. T.
-----C. T.
-----T. T.
-----G. T.
-----G. T.
-----T. T.
-----T. T.
-----T. T.
-----T. TG
-----T. T.
-----T. A. C. TTT A. RT. AG. T. A. T.
-----C. T. A. TTT A. RT. AG. T. A. T.
-----T. A. TTT A. RT. AG. TCA. T.
-----T. A. TTT A. RT. AG. TCA. T.
-----T. A. TTT A. RT. AG. TCA. T.
-----T. A. TT A. A. AGA T. A. T.
-----CC- T. A. TT A. A. AGA T. A. T.

```


TABLE S4**The presence and absence of *R* gene polymorphism in the *Pi-ta* genomic region among *Oryza* species**

Accessions	Number of accessions	Presence / Absence ^a
<i>O. sativa</i> (containing resistance <i>Pi-ta</i>)	17	15 / 2 ^b
<i>O. sativa</i> weedy rice	50	2 / 48 ^c
<i>O. rufipogon</i>	30	17 / 13
<i>O. nivara</i> , <i>O. barthii</i> , <i>O. glaberrima</i> , <i>O. glumaepatula</i>	10	10 / 0

^a Presence and absence polymorphism of the NBS-LRR gene (LOC_OS12G14330) at 8.5 Mb

^b Absence of the gene in two rice cultivars Yashiro-mochi and Te Qing.

^c The NBS-LRR gene is absent in all red rice accessions (33) containing resistance *Pi-ta*.