

Review

Isomer specificity of conjugated linoleic acid (CLA): 9E,11E-CLA

Yunkyoung Lee[§]

Obesity and metabolism laboratory, Jean Mayer USDA HNRCA at Tufts University, Boston, MA 02111, USA

Received October 11, 2008, Revised November 3, 2008, Accepted November 21, 2008

Abstract

Conjugated linoleic acids (CLA) were identified in 1980's, since then it has been intensively studied due to its various beneficial health effects such as anti-inflammatory, anti-atherogenic, anti-carcinogenic and anti-diabetic/obesity effects. Isomer specificity of a number of CLA isomers, especially predominant isomer 9Z,11E- and 10E,12Z-CLA, is now recognized. However, the less prevalent CLA isomers have not been well characterized. Recently, studies have reported the distinctively different effects of 9E,11E-CLA in colon cancer cells, endothelial cells, and macrophage cells compared to the rest of CLA isomers. In this review, various effects of CLAs, especially anti-inflammatory and anti-atherogenic effects, will be discussed with focusing on the isomer-specific effects and potential mechanism of action of CLA. At last, recent studies about 9E,11E-CLA in *in vitro* and animal models will be discussed.

Key Words: Conjugated linoleic acid (CLA), 9E,11E-CLA, anti-inflammatory, isomer-specificity

Overview of conjugated linoleic acids

CLAs are fatty acids that are mainly found in foods derived from ruminant animals (Pariza & Ha, 1990). They are geometrical and positional isomers of its parent molecule, linoleic acid (*cis*-9,*cis*-12-18:2, n-6 (LA)). The *cis*-9, *trans*-11 (9Z,11E-octadecenoic acid, C18:2) isomer, also known as ruminic acid (RA), is generated via biohydrogenation of dietary linoleic acids by ruminant microflora and is the most abundant natural CLA isomer (over 75-80% of total CLA). It is not clear whether humans are able to produce RA from other fatty acids (Yurawecz, 1999); however, it is likely that the majority of RA found in the human body originates from the diet. Estimated CLA intake for humans is between 0.5 and 1 g/d (Ip *et al.*, 1994); however, more recent estimates suggest that CLA and RA intakes are much lower (Yurawecz, 1999). For example, the average intake is 430 mg/day for men and 350 mg/day for women in Germany, 212 mg/day for men and 151 mg/day for women in the USA, and 500-1,000 mg/day in Australia (Yurawecz, 1999).

Studied for its potential beneficial effects of CLA on animal health have focused on anti-inflammatory, anti-atherogenic, anti-carcinogenic, and anti-diabetic/anti-obesity effects; however, the beneficial effects of CLA in humans is inconclusive (Kelly, 2001; Pariza *et al.*, 2001; Rainer & Heiss, 2004; Wahle *et al.*, 2004; Whigham *et al.*, 2000). It is important to note that a majority of the CLA studies was performed with a chemically synthesized CLA mixture, although in recent years, some were done with relatively pure 9Z,11E-CLA or 10E,12Z-CLA isomers.

CLA supplements which predominantly contain two major CLA isomers are on the market; however, the safety of CLA is a concern in public health. To obtain beneficial amounts of CLA, one must ingest 10-200 times what can be achieved in diet (Gaullier *et al.*, 2002). Although animal studies on CLA showed that it is reasonably safe and well-tolerated in rats, dogs, and pigs (O'Hagan & Menzel, 2003; Pariza, 2004; Scimeca, 1998), whether the same is true for humans is not known. Gaullier *et al.* (2002) stated that CLA enriched with 9Z,11E- and 10E,12Z-CLA may be better for human consumption than one enriched with 8E,10Z-, 9Z,11E-, 10E,12Z-, and 11Z,13E-CLA. In spite of these conclusions, concerns on CLA safety remain due to its adverse effects which induce fatty liver, insulin resistance, and lipodystrophy in mice and increase C-reactive protein levels, lipid peroxidation, and reduction of milk fat in humans (Pariza, 2004); therefore, it is essential for long-term human studies to make solid conclusions on recommending a proper amount of CLA supplementation to the public. It is also important to consider differences in CLA isomers in terms of both benefits and potential toxicity.

Anti-inflammatory effects

Levels of fatty acids are known to have a great impact on immune function. Dietary fat intake directly influences the fatty acid profile of blood lipids and cells. The immune-modulating effects of saturated and polyunsaturated fatty acids have been

[§] Corresponding Author: Yunkyoung Lee, Tel. 1-617-556-3146, Fax. 1-617-556-3344, Email. Yunkyoung.lee@tufts.edu

reported (Calder & Grimble, 2002). However, the available knowledge about the effects of CLAs on immune function in animals or humans is limited. The inflammatory-modulating effects of CLA are mainly observed in *in vitro* and animal models, while CLA's anti-inflammatory effects are not conclusive in human studies. Chicks and rats fed 0.5% CLA for 4 wks gained weight even after LPS injection, whereas chicks and rats without CLA lost weight. This led to the idea that CLA inhibited the catabolic effects of endotoxin (Cook *et al.*, 1993). CLA fed mice had higher levels of IL-2 compared to control diet fed mice (Hayek *et al.*, 1999; Wong *et al.*, 1997), while TNF- α and IL-6 levels were decreased (Rahman *et al.*, 2007). However, Kelley *et al.* (2002) reported that both 9Z,11E- and 10E,12Z-CLA significantly and similarly increased TNF- α and IL-6 levels without altering prostaglandin levels in CLA fed mice.

In *in vitro* models, CLA was positively related to anti-inflammatory effects. For example, Yu *et al.* (2002) showed that CLA reduced TNF- α and IL-6 levels induced by IFN- γ , at least partially by a PPAR γ -dependent pathway. CLA induced IFN- γ and IL-2 levels in Juckat T cells (Luongo *et al.*, 2003). Moreover, IL-10 receptor and IL-10 levels were induced by CLA in mouse dendritic cells (Loscher *et al.*, 2005). In human epithelial cells, 9Z,11E-CLA attenuated cell growth and reduced IL-8 levels while 10E,12Z-CLA had no effect (Jaudszus *et al.*, 2005).

The results from human studies focusing on anti-inflammatory effects of CLA are inconclusive. Song *et al.* (2005) demonstrated that CLA supplementation could be beneficial in humans (3 g/d CLA for 12 wks) by enhancing IgA, IgM, and IL-10 but inhibiting IgE, TNF- α , and IL-1 β . In addition, CLA boosted the hepatitis B vaccination in healthy human subjects although it did not alter other aspects of immune function including NK cell activity, lymphocyte proliferation, and production of TNF- α , IL-1 β , IL-6, IL-2, IL-4, IFN- γ , and PGE₂ (Albers *et al.*, 2003). In contrast, other human studies showed no significant relationship between CLA and anti-inflammatory effect (Albers *et al.*, 2003; Kelley *et al.*, 2000; Nugent *et al.*, 2005; Ramakers *et al.*, 2005; Ritzenthaler *et al.*, 2005; Tricon *et al.*, 2004a). There were no changes on the levels of PGE₂, leukotriene B₄, IL-1 β , and TNF- α that were induced by LPS in young healthy women (Kelley *et al.*, 2000).

The regulation of immune functions by CLA does not seem to be affected by a single mechanism. CLA affects the production of eicosanoids either directly or indirectly, enhances PPAR γ activation, attenuates the NF- κ B pathway, or directly decreases pro-inflammatory cytokines to have beneficial effects on inflammation which ultimately influence metabolic syndromes including obesity, insulin resistance, and atherosclerosis (Zulet *et al.*, 2005). CLA decreased the production of eicosanoids such as PGE₂, PGF₂ α , and LTB₄ (Li & Watkins, 1998; Sugano *et al.*, 1998), although the reduction of the eicosanoid production was not consistent (Hayek *et al.*, 1999; Kelley *et al.*, 2000). The mechanism by which CLA reduces AA-derived eicosanoids can be explained as follows: (1) CLA displaces AA in phospholipids;

(2) CLA competes with LA for desaturation and elongation; (3) CLA or CLA metabolites may act as substrates or antagonists for enzymes involved in the prostaglandin production process; and (4) CLA-derived eicosanoids themselves may have anti-inflammatory properties. The PPAR γ subtype is highly expressed in adipose tissue and macrophages (Olefsky, 2001). Most importantly, PPAR γ plays a critical role in the regulation of inflammation (Cuzzocrea *et al.*, 2004). Various CLA isomers activate PPAR γ in RAW 264.7 cells and decrease pro-inflammatory cytokines induced by IFN- γ (Yu *et al.*, 2002). NF- κ B is a transcription factor that is involved in cytokine gene expression, cellular adhesion, cell cycle activation, apoptosis, and carcinogenesis. CLA negatively regulates inflammatory mediators via the NF- κ B pathway by inhibiting I κ B phosphorylation (Cheng *et al.*, 2004). Finally, CLA could exert direct anti-inflammatory properties by regulating the gene expression of inflammation mediators (Loscher *et al.*, 2005; Luongo *et al.*, 2003; Song *et al.*, 2005), perhaps due to effects on NF- κ B or PPAR γ .

In conclusion, the anti-inflammatory action of CLA brings potential therapies for diseases such as atherosclerosis, diabetes, cancers, rheumatism, inflammatory bowel disease and obesity. Meanwhile, the lack of understanding regarding CLAs mechanism (s) of action, safety issue surrounding CLA as a dietary supplement, and isomer-specific effects on regulating inflammatory mediators require further studies.

Anti-atherogenic effects

Atherosclerosis is a complex disease that is influenced in part by inflammation (reviewed in (Ross, 1999)). Interactions between lipoproteins, monocyte-derived macrophages, T cells, and the normal cellular elements of the arterial wall are important contributors for the development of atherosclerosis. CLA-fed rabbits had significantly lower low density lipoprotein (LDL) cholesterol and TG levels as well as a lower ratio of LDL/HDL (high density lipoprotein) and LDL/total cholesterol compared to control diet fed rabbits (Kritchevsky *et al.*, 2000; Lee *et al.*, 1994). In hamsters, CLA reduced total cholesterol, LDL, VLDL (very low density lipoprotein), and TG although there were no changes on HDL levels (Nicolosi *et al.*, 1997). Unlike the results in rabbits and hamsters, the results from mice were somewhat conflicting (Arbones-Mainar *et al.*, 2006; Munday *et al.*, 1999; Nestel *et al.*, 2006; Toomey *et al.*, 2006). A mixture of CLA (9Z, 11E-CLA:10E, 12Z-CLA (80:20 blend)) was tested in ApoE^{-/-} mice and showed a suppression of atherosclerotic lesions by decreasing pro-inflammatory genes (i.e., metalloproteinases (MMP)-9 and PECAM-1) and an increase in apoptotic genes (Toomey *et al.*, 2006). Isomer-specificity of regulation of the lipid profile by CLA isomers was shown in a few studies. Mitchell *et al.* (2005) showed that 10E,12Z-CLA increased HDL levels in 1% CLA fed hamsters. However, Wilson *et al.* (2006) reported that 9Z,11E-CLA decreased cholesterol whereas

10E,12Z-CLA increased cholesterol levels in 0.5% CLA isomer fed hamsters. This discrepancy may be due to the relative amount and composition of the CLA mix used in the studies.

Different types of cell lines related to atherosclerosis were used to study the anti-atherogenic effects of CLA in *in vitro* systems. Decreased levels of 6-keto-prostaglandin F(1 α) (6k-PGF(1 α)), a stable breakdown product of PGI₂, were observed in bovine aortic endothelial cells treated with CLA (Coen *et al.*, 2004). CLA increased CD36 (scavenger receptor) in THP-1, human macrophages as well as reduced total lipids, PGE₂, PGI₂, and cytokine-induced NF- κ B binding activity in human smooth muscle cells (Ringseis *et al.*, 2006; Weldon *et al.*, 2004). However, CLA failed to inhibit adhesion molecules which are important for the initiation step of atherosclerotic lesion formation, such as ICAM-1, VCAM-1, and E-selectin induced by TNF- α in human aortic EC (Schleser *et al.*, 2006). Although there were no changes in ICAM-1 and VCAM-1 levels by CLA, CLA inhibited the adhesion activity of THP-1 cells to human umbilical vein EC (HUVEC) (Sneddon *et al.*, 2006).

CLA's fat-lowering effects have been reported in human subjects, although they have not been consistent. Benito *et al.* (2001) showed no alteration of cholesterol, TG, HDL, and LDL cholesterol levels in human on a 3.9 g/d CLA supplementation for 63 days. Desroches *et al.* (2005) reported that healthy subjects on CLA had lower total cholesterol, total cholesterol/HDL ratio compared to the subjects on control diet. On the other hand, Mougios *et al.* (2001) and Gaullier *et al.* (2004) reported a detrimental HDL-lowering effect of CLA mix in humans. Tricon *et al.* (2004b) showed that 10E,12Z-CLA increased a ratio of cholesterol/HDL, LDL:HDL, and TG levels compared to 9Z,11E-CLA in healthy humans. In spite of isomer-specific effects of CLA on lipid profile, the same group reported a similar effect of those two CLA isomers on immune cell function as inhibiting mitogen-induced T cell activation (Tricon *et al.*, 2004a). The CLA isomer-specificity varies depending on the results observed; therefore, studies on pure CLA isomers are needed to elucidate these questions.

Some proposed mechanisms for the anti-atherogenic and lipid-lowering effects of CLA include roles for PPARs, sterol regulatory element-binding proteins (SREBPs) and stearoyl-CoA desaturase (SCD) (reviewed in (Bhattacharya *et al.*, 2006)). Further studies using each CLA isomer and long-term clinical studies are needed to establish their potential anti-atherogenic effects and mechanism (s) of action.

9E,11E-CLA: its effects and mechanism

The isomer-specific effects of 9Z,11E-CLA and 10E,12Z-CLA are now well-recognized (reviewed in (Evans *et al.*, 2002)). However, the less prevalent CLA isomers have not been intensively studied. Recently, more studies have reported the effects of 9E,11E-CLA in colon cancer cells, endothelial cells, and macrophage

cells (Beppu *et al.*, 2006; Lai *et al.*, 2005; Yasui *et al.*, 2007). Lai *et al.* (2005) showed that among the CLA isomers, 9E,11E-CLA attenuated proliferation of bovine endothelial cells the most by enhancing caspase-3 activity. In the following year, Beppu *et al.* (2006) reported that 9E,11E-CLA showed the strongest apoptosis among 9Z,11Z-, 9Z,11E-, and 10E,12Z-CLA isomer on colon cancer cells, Caco-2 cells. The same researchers further reported the 9E,11E-CLA's chemopreventive effect in rats fed 0.1% or 1% 9E,11E-CLA for 4 wks (Yasui *et al.*, 2007). Ecker *et al.* (2007) showed that 9E,11E-CLA, but not 9Z,11E- and 10E,12Z-CLA, induces SREBP target genes such as ABCG-1 via a SREBP-1c-dependent mechanism in primary human monocyte-derived macrophages. Comparison of five different CLA isomers showed that 9E,11E-CLA indeed regulates gene expression distinctively different compared to the rest of CLA isomers in a mouse macrophage cell line, RAW 264.7 (Lee *et al.*, 2008). Thus, isomer specificity of 9E,11E-CLA has been recognized in various cell types.

Although the portion of 9E,11E-CLA in American beef fat and commercial CLA mixtures ranged from 1.5-3.7% of the total CLA isomers (Kramer *et al.*, 1998; Yurawecz, 1999), 9E,11E- and 9Z,11E-CLA were the main CLA isomers in serum isolated from women on 2.1 g of CLA supplementation for 45 days which represented 41% of the CLA total (Petridou *et al.*, 2003). These isomers were also the two major CLA isomers in platelet lipids, 17% and 33% of the total respectively (Al-Madaney *et al.*, 2003). Based on the basal plasma CLA levels in healthy men as 7 μ M, the 9E,11E-CLA concentration is estimated to be in the 1-3 μ M range (Huang, 1994). Considering 10-fold higher concentration in local areas, 10-30 μ M might be a physiologically relevant level of 9E,11E-CLA. Large-scale and economically efficient separation of 9E,11E-CLA is difficult due to its minimal existence in the CLA mixture. Kishino *et al.* (2002, 2003) reported the bacterial production of 9Z,11E- and 9E,11E-CLA from ricinoleic acid (12-hydroxy-cis-9-octadecaenoic acid) and linoleic acid. This biological system for 9E,11E-CLA production may promise the large-scale and selective preparation of 9E,11E-CLA.

Conclusion

In spite of intensive studies of CLAs performed over the last two decades, the mechanism of action of each CLA isomer is not well-appreciated yet. In this review, various health-related effects of CLAs were discussed with a focus on superior and/or distinctive characteristics of 9E,11E-CLA. The limited numbers of recent studies of 9E,11E-CLA in *in vitro* and *in vivo* studies suggest that this specific CLA isomer appears to have the anti-carcinogenic, anti-inflammatory, and potential anti-atherogenic effects. Although it requires more *in vivo* and clinical studies before making a solid conclusion, understanding the effects of the individual CLA isomer helps us to make a combination of the CLA isomers in dietary supplements to maximize its healthful effects and minimize its harmful effects.

Literature cited

- Albers R, van der Wielen RP, Brink EJ, Hendriks HF, Dorovska-Taran VN & Mohede IC (2003). Effects of cis-9, trans-11 and trans-10, cis-12 conjugated linoleic acid (CLA) isomers on immune function in healthy men. *Eur J Clin Nutr* 57:595-603.
- Al-Madaney MM, Kramer JK, Deng Z & Vanderhoek JY (2003). Effects of lipid-esterified conjugated linoleic acid isomers on platelet function: evidence for stimulation of platelet phospholipase activity. *Biochim Biophys Acta* 1635:75-82.
- Arbones-Mainar JM, Navarro MA, Guzman MA, Arnal C, Surra JC, Acin S, Carnicer R, Osada J & Roche HM (2006). Selective effect of conjugated linoleic acid isomers on atherosclerotic lesion development in apolipoprotein E knockout mice. *Atherosclerosis* 189:318-327.
- Benito P, Nelson GJ, Kelley DS, Bartolini G, Schmidt PC & Simon V (2001). The effect of conjugated linoleic acid on plasma lipoproteins and tissue fatty acid composition in humans. *Lipids* 36:229-236.
- Beppu F, Hosokawa M, Tanaka L, Kohno H, Tanaka T & Miyashita K (2006). Potent inhibitory effect of trans9, trans11 isomer of conjugated linoleic acid on the growth of human colon cancer cells. *J Nutr Biochem* 17:830-836.
- Bhattacharya A, Banu J, Rahman M, Causey J & Fernandes G (2006). Biological effects of conjugated linoleic acids in health and disease. *J Nutr Biochem* 17:789-810.
- Calder PC & Grimble RF (2002). Polyunsaturated fatty acids, inflammation and immunity. *Eur J Clin Nutr* 56:S14-S19.
- Cheng WL, Lii CK, Chen HW, Lin TH & Liu KL (2004). Contribution of conjugated linoleic acid to the suppression of inflammatory responses through the regulation of the NF-kappaB pathway. *J Agric Food Chem* 52:71-78.
- Coen P, Cummins P, Birney Y, Devery R & Cahill P (2004). Modulation of nitric oxide and 6-keto-prostaglandin F(1alpha) production in bovine aortic endothelial cells by conjugated linoleic acid. *Endothelium* 11:211-220.
- Cook ME, Miller CC, Park Y & Pariza M (1993). Immune modulation by altered nutrient metabolism: nutritional control of immune-induced growth depression. *Poult Sci* 72:1301-1305.
- Cuzzocrea S, Pisano B, Dugo L, Ianaro A, Maffia P, Patel NS, Di Paola R, Ialenti A, Genovese T, Chatterjee PK, Di Rosa M, Caputi AP & Thiemermann C (2004). Rosiglitazone, a ligand of the peroxisome proliferator-activated receptor-gamma, reduces acute inflammation. *Eur J Pharmacol* 483:79-93.
- Desroches S, Chouinard PY, Galibois I, Corneau L, Delisle J, Lamarche B, Couture P & Bergeron N (2005). Lack of effect of dietary conjugated linoleic acids naturally incorporated into butter on the lipid profile and body composition of overweight and obese men. *Am J Clin Nutr* 82:309-319.
- Ecker J, Langmann T, Moehle C & Schmitz G (2007). Isomer specific effects of Conjugated Linoleic Acid on macrophage ABCG1 transcription by a SREBP-1c dependent mechanism. *Biochem Biophys Res Commun* 352:805-811.
- Evans M, Brown J & McIntosh M (2002). Isomer-specific effects of conjugated linoleic acid (CLA) on adiposity and lipid metabolism. *J Nutr Biochem* 13:508.
- Gaullier JM, Berven G, Blankson H & Gudmundsen O (2002). Clinical trial results support a preference for using CLA preparations enriched with two isomers rather than four isomers in human studies. *Lipids* 37:1019-1025.
- Gaullier JM, Halse J, Hoye K, Kristiansen K, Fagertun H, Vik H & Gudmundsen O (2004). Conjugated linoleic acid supplementation for 1 y reduces body fat mass in healthy overweight humans. *Am J Clin Nutr* 79:1118-1125.
- Hayek MG, Han SN, Wu D, Watkins BA, Meydani M, Dorsey JL, Smith DE & Meydani SN (1999). Dietary conjugated linoleic acid influences the immune response of young and old C57BL/6NCrIBR mice. *J Nutr* 129:32-38.
- Huang YC, Luedecke LO & Shultz TD (1994). Effect of cheddar cheese consumption on plasma conjugated linoleic acid concentrations in men. *Nutr Res* 14:373-386.
- Ip C, Scimeca JA & Thompson HJ (1994). Conjugated linoleic acid. A powerful anticarcinogen from animal fat sources. *Cancer* 74:1050-1054.
- Jaudszus A, Foerster M, Kroegel C, Wolf I & Jahreis G (2005). Cis-9,Trans-11-CLA exerts anti-inflammatory effects in human bronchial epithelial cells and eosinophils: Comparison to Trans-10, Cis-12-CLA and to linoleic acid. *Biochim Biophys Acta* 1737:111-1118.
- Kelley DS, Taylor PC, Rudolph IL, Benito P, Nelson GJ, Mackey BE & Erickson KL (2000). Dietary conjugated linoleic acid did not alter immune status in young healthy women. *Lipids* 35:1065-1071.
- Kelley DS, Warren JM, Simon VA, Bartolini G, Mackey BE & Erickson KL (2002). Similar effects of c9,t11-CLA and t10, c12-CLA on immune cell functions in mice. *Lipids* 37:725-728.
- Kelly GS (2001). Conjugated linoleic acid: a review. *Altern Med Rev* 6:367-382.
- Kishino S, Ogawa J, Ando A, Iwashita T, Fujita T, Kawashima H & Shimizu S (2003). Structural analysis of conjugated linoleic acid produced by *Lactobacillus plantarum*, and factors affecting isomer production. *Biosci Biotechnol Biochem* 67:179-182.
- Kishino S, Ogawa J, Ando A, Omura Y & Shimizu S (2002). Ricinoleic acid and castor oil as substrates for conjugated linoleic acid production by washed cells of *Lactobacillus plantarum*. *Biosci Biotechnol Biochem* 66:2283-2286.
- Kramer JK, Parodi PW, Jensen RG, Mossoba MM, Yurawecz MP & Adlof RO (1998). Ruminic acid: a proposed common name for the major conjugated linoleic acid isomer found in natural products. *Lipids* 33:835.
- Kritchevsky D, Tepper SA, Wright S, Tso P & Czarnecki SK (2000). Influence of conjugated linoleic acid (CLA) on establishment and progression of atherosclerosis in rabbits. *J Am Coll Nutr* 19:472S-477S.
- Lai KL, Torres-Duarte AP & Vanderhoek JY (2005). 9-trans, 11-trans-CLA: antiproliferative and proapoptotic effects on bovine endothelial cells. *Lipids* 40:1107-1116.
- Lee KN, Kritchevsky D & Pariza MW (1994). Conjugated linoleic acid and atherosclerosis in rabbits. *Atherosclerosis* 108:19-25.
- Lee Y, Thompson JT, de Lera AR & Vanden Heuvel JP (2008). Isomer-specific effects of conjugated linoleic acid on gene expression in RAW 264.7. *J Nutr Biochem* 5:[Epub ahead of print].
- Li Y & Watkins BA (1998). Conjugated linoleic acids alter bone fatty acid composition and reduce ex vivo prostaglandin E2 biosynthesis in rats fed n-6 or n-3 fatty acids. *Lipids* 33:417-425.
- Loscher CE, Draper E, Leavy O, Kelleher D, Mills KH & Roche HM (2005). Conjugated linoleic acid suppresses NF-kappa B activation and IL-12 production in dendritic cells through ERK-mediated IL-10 induction. *J Immunol* 175:4990-4998.
- Luongo D, Bergamo P & Rossi M (2003). Effects of conjugated linoleic acid on growth and cytokine expression in Jurkat T cells.

- Immunol Lett* 90:195-201.
- Mitchell PL, Langille MA, Currie DL & McLeod RS (2005). Effect of conjugated linoleic acid isomers on lipoproteins and atherosclerosis in the Syrian Golden hamster. *Biochim Biophys Acta* 1734:269-276.
- Mougiou V, Matsakas A, Petridou A, Ring S, Sagredos A, Melissopoulou A, Tsigilis N & Nikolaidis M (2001). Effect of supplementation with conjugated linoleic acid on human serum lipids and body fat. *J Nutr Biochem* 12:585-594.
- Munday JS, Thompson KG & James KA (1999). Dietary conjugated linoleic acids promote fatty streak formation in the C57BL/6 mouse atherosclerosis model. *Br J Nutr* 81:251-255.
- Nestel P, Fujii A & Allen T (2006). The cis-9,trans-11 isomer of conjugated linoleic acid (CLA) lowers plasma triglyceride and raises HDL cholesterol concentrations but does not suppress aortic atherosclerosis in diabetic apoE-deficient mice. *Atherosclerosis* 189:282-287.
- Nicolosi RJ, Rogers EJ, Kritchevsky D, Scimeca JA & Huth PJ (1997). Dietary conjugated linoleic acid reduces plasma lipoproteins and early aortic atherosclerosis in hypercholesterolemic hamsters. *Artery* 22:266-277.
- Nugent AP, Roche HM, Noone EJ, Long A, Kelleher DK & Gibney MJ (2005). The effects of conjugated linoleic acid supplementation on immune function in healthy volunteers. *Eur J Clin Nutr* 59:742-750.
- O'Hagan S & Menzel A (2003). A subchronic 90-day oral rat toxicity study and in vitro genotoxicity studies with a conjugated linoleic acid product. *Food Chem Toxicol* 41:1749-1760.
- Olefsky JM (2001). Nuclear receptor minireview series. *J Biol Chem* 276:36863-36864.
- Pariza MW (2004). Perspective on the safety and effectiveness of conjugated linoleic acid. *Am J Clin Nutr* 79:1132S-1136S.
- Pariza MW & Ha YL (1990). Conjugated dienoic derivatives of linoleic acid: a new class of anticarcinogens. *Med Oncol Tumor Pharmacother* 7:169-171.
- Pariza MW, Park Y & Cook ME (2001). The biologically active isomers of conjugated linoleic acid. *Prog Lipid Res* 40:283-298.
- Petridou A, Mougiou V & Sagredos A (2003). Supplementation with CLA: isomer incorporation into serum lipids and effect on body fat of women. *Lipids* 38:805-811.
- Rahman MM, Bhattacharya A, Banu J & Fernandes G (2007). Conjugated linoleic acid protects against age-associated bone loss in C57BL/6 female mice. *J Nutr Biochem* 18:467-474.
- Rainer L & Heiss CJ (2004). Conjugated linoleic acid: health implications and effects on body composition. *J Am Diet Assoc* 104:963-968.
- Ramakrishna JD, Plat J, Sebedio JL & Mensink RP (2005). Effects of the individual isomers cis-9,trans-11 vs. trans-10,cis-12 of conjugated linoleic acid (CLA) on inflammation parameters in moderately overweight subjects with LDL-phenotype B. *Lipids* 40:909-918.
- Ringseis R, Muller A, Herter C, Gahler S, Steinhart H & Eder K (2006). CLA isomers inhibit TNF α -induced eicosanoid release from human vascular smooth muscle cells via a PPAR γ ligand-like action. *Biochim Biophys Acta* 1760:290-300.
- Ritzenthaler KL, McGuire MK, McGuire MA, Shultz TD, Koepf AE, Lueddecke LO, Hanson TW, Dasgupta N & Chew BP (2005). Consumption of conjugated linoleic acid (CLA) from CLA-enriched cheese does not alter milk fat or immunity in lactating women. *J Nutr* 135:422-430.
- Ross R (1999). Atherosclerosis--an inflammatory disease. *N Engl J Med* 340:115-126.
- Schlesler S, Ringseis R & Eder K (2006). Conjugated linoleic acids have no effect on TNF α -induced adhesion molecule expression, U937 monocyte adhesion, and chemokine release in human aortic endothelial cells. *Atherosclerosis* 186:337-344.
- Scimeca JA (1998). Toxicological evaluation of dietary conjugated linoleic acid in male Fischer 344 rats. *Food Chem Toxicol* 36:391-395.
- Sneddon AA, McLeod E, Wahle KW & Arthur JR (2006). Cytokine-induced monocyte adhesion to endothelial cells involves platelet-activating factor: suppression by conjugated linoleic acid. *Biochim Biophys Acta* 1761:793-801.
- Song HJ, Grant I, Rotondo D, Mohede I, Sattar N, Heys SD & Wahle KW (2005). Effect of CLA supplementation on immune function in young healthy volunteers. *Eur J Clin Nutr* 59:508-517.
- Sugano M, Tsujita A, Yamasaki M, Noguchi M & Yamada K (1998). Conjugated linoleic acid modulates tissue levels of chemical mediators and immunoglobulins in rats. *Lipids* 33:521-527.
- Toomey S, Harhen B, Roche HM, Fitzgerald D & Belton O (2006). Profound resolution of early atherosclerosis with conjugated linoleic acid. *Atherosclerosis* 187:40-49.
- Tricon S, Burdge GC, Kew S, Banerjee T, Russell JJ, Grimble RF, Williams CM, Calder PC & Yaqoob P (2004a). Effects of cis-9, trans-11 and trans-10,cis-12 conjugated linoleic acid on immune cell function in healthy humans. *Am J Clin Nutr* 80:1626-1633.
- Tricon S, Burdge GC, Kew S, Banerjee T, Russell JJ, Jones EL, Grimble RF, Williams CM, Yaqoob P & Calder PC (2004b). Opposing effects of cis-9,trans-11 and trans-10,cis-12 conjugated linoleic acid on blood lipids in healthy humans. *Am J Clin Nutr* 80:614-620.
- Wahle KW, Heys SD & Rotondo D (2004). Conjugated linoleic acids: are they beneficial or detrimental to health? *Prog Lipid Res* 43: 553-587.
- Weldon S, Mitchell S, Kelleher D, Gibney MJ & Roche HM (2004). Conjugated linoleic acid and atherosclerosis: no effect on molecular markers of cholesterol homeostasis in THP-1 macrophages. *Atherosclerosis* 174:261-273.
- Whigham LD, Cook ME & Atkinson RL (2000). Conjugated linoleic acid: implications for human health. *Pharmacol Res* 42:503-510.
- Wilson TA, Nicolosi RJ, Saati A, Kotyla T & Kritchevsky D (2006). Conjugated linoleic acid isomers reduce blood cholesterol levels but not aortic cholesterol accumulation in hypercholesterolemic hamsters. *Lipids* 41:41-48.
- Wong MW, Chew BP, Wong TS, Hosick HL, Boylston TD & Shultz TD (1997). Effects of dietary conjugated linoleic acid on lymphocyte function and growth of mammary tumors in mice. *Anticancer Res* 17:987-993.
- Yasui Y, Suzuki R, Kohno H, Miyamoto S, Beppu F, Hosokawa M, Miyashita K & Tanaka T (2007). 9trans,11trans conjugated linoleic acid inhibits the development of azoxymethane-induced colonic aberrant crypt foci in rats. *Nutr Cancer* 59:82-91.
- Yu Y, Correll PH & Vanden Heuvel JP (2002). Conjugated linoleic acid decreases production of pro-inflammatory products in macrophages: evidence for a PPAR γ -dependent mechanism. *Biochim Biophys Acta* 1581:89-99.
- Yurawecz MP (1999). *Advances in conjugated linoleic acid research*. Champaign, Ill.: AOCS Press. p.154-179. Champaign, IL. USA
- Zulet MA, Marti A, Parra MD & Martinez JA (2005). Inflammation and conjugated linoleic acid: mechanisms of action and implications for human health. *J Physiol Biochem* 61:483-494.