

Hypoxia inducible factors regulate pluripotency and proliferation in human embryonic stem cells cultured at reduced oxygen tensions

Catherine E Forristal^{1,2}, Kate L Wright^{1,2}, Neil A Hanley^{1,2}, Richard O C Oreffo^{1,3} and Franchesca D Houghton^{1,2}

¹Centre for Human Development, Stem Cells and Regeneration, ²Human Genetics Division and ³Developmental Origins of Health & Disease Division, School of Medicine, University of Southampton, Southampton General Hospital, Duthie Building (MP 808), Tremona Road, Southampton, SO16 6YD, UK

Correspondence should be addressed to F D Houghton; Email: f.d.houghton@soton.ac.uk

N A Hanley is now at Endocrine Sciences Research Group, Manchester Academic Health Science Centre, University of Manchester, AV Hill Building, Oxford Road, Manchester, M13 9PT, UK

Abstract

Human embryonic stem (hES) cells are routinely cultured under atmospheric, 20% oxygen tensions but are derived from embryos which reside in a 3–5% oxygen (hypoxic) environment. Maintenance of oxygen homeostasis is critical to ensure sufficient levels for oxygen-dependent processes. This study investigates the importance of specific hypoxia inducible factors (HIFs) in regulating the hypoxic responses of hES cells. We report that culture at 20% oxygen decreased hES cell proliferation and resulted in a significantly reduced expression of *SOX2*, *NANOG* and *POU5F1* (*OCT4*) mRNA as well as *POU5F1* protein compared with hypoxic conditions. HIF1A protein was not expressed at 20% oxygen and displayed only a transient, nuclear localisation at 5% oxygen. HIF2A (EPAS1) and HIF3A displayed a cytoplasmic localisation during initial hypoxic culture but translocated to the nucleus following long-term culture at 5% oxygen and were significantly upregulated compared with cells cultured at 20% oxygen. Silencing of HIF2A resulted in a significant decrease in both hES cell proliferation and *POU5F1*, *SOX2* and *NANOG* protein expression while the early differentiation marker, *SSEA1*, was concomitantly increased. HIF3A upregulated HIF2A and prevented HIF1A expression with the knockdown of HIF3A resulting in the reappearance of HIF1A protein. In summary, these data demonstrate that a low oxygen tension is preferential for the maintenance of a highly proliferative, pluripotent population of hES cells. While HIF3A was found to regulate the expression of both HIF1A and HIF2A, it is HIF2A which regulates hES cell pluripotency as well as proliferation under hypoxic conditions.

Reproduction (2010) **139** 85–97

Introduction

Human embryonic stem (hES) cells, typically derived from the inner cell mass (ICM) of the blastocyst, proliferate by self renewal and have the potential to differentiate into all cells of the body (Thomson *et al.* 1998). Thus, hES cells provide an excellent model to investigate developmental mechanisms and are a potentially unlimited source of cells for transplantation in degenerative disease. However, hES cells are notoriously difficult to maintain in culture due to their propensity for spontaneous differentiation, an effect likely caused by a suboptimal culture environment. Clues to improve the culture of hES cells may be gained by investigating the environment from which these cells

are derived. *In vivo*, preimplantation embryos develop in the secretions of the reproductive tract which are characterised by low, 1.5–8% oxygen tensions (Fischer & Bavister 1993), an environment which has been shown to improve the *in vitro* embryo development of several species including human (Dumoulin *et al.* 1999, Petersen *et al.* 2005, Kovacic & Vlaisavljevic 2008), mouse (Orsi & Leese 2001) and bovine (Thompson *et al.* 1990, Olson & Seidel 2000). Moreover, mouse embryos cultured at 5% oxygen rather than 20% oxygen displayed a global gene expression pattern which more closely resembled their *in vivo* counterparts (Rinaudo *et al.* 2006). Harvey *et al.* (2004), also observed that culture under 2% oxygen rather than atmospheric oxygen significantly increased the proportion of ICM

cells compared with trophectoderm cells, findings consistent with pluripotent cells displaying increased proliferation when cultured under reduced oxygen tensions. Thus, mimicking the *in vivo*, physiological oxygen concentration may be beneficial for the propagation of hES cells *in vitro*. This is supported by data from rat mesenchymal and neural crest stem cells, where culture under physiological oxygen tensions showed an increase in the rate of proliferation compared with those maintained at 20% oxygen (Morrison *et al.* 2000, Lennon *et al.* 2001). These combined data highlight the benefit that culture under reduced oxygen tension has on the ability to expand scarce stem/precursor cell populations.

In terms of hES cells, atmospheric oxygen remains the culture environment of routine use. There is emerging evidence to suggest that reducing the oxygen concentration towards physiological levels is beneficial for the *in vitro* maintenance of hES cells in terms of decreasing the amount of spontaneous differentiation, supporting self-renewal (Ezashi *et al.* 2005, Ludwig *et al.* 2006, Westfall *et al.* 2008), and reducing spontaneous chromosomal aberrations (Forsyth *et al.* 2006). Thus, while hES cells can be maintained under atmospheric oxygen tensions, lowering the oxygen tension to 2–5% appears beneficial for the propagation of a highly proliferative, pluripotent population of cells. However, controversy still remains as a recent report suggests that there are no significant advantages of culturing hES cells under reduced oxygen tension (Chen *et al.* 2009).

Upon exposure to hypoxic conditions, cells mount a physiological response to ensure sufficient levels for oxygen-dependent processes. This response is regulated by hypoxia inducible factors (HIFs) which regulate the expression of over 200 genes including those involved in erythropoiesis, apoptosis and proliferation (Semenza 2000). HIFs are transcription factors consisting of three oxygen-dependent α subunits: HIF1A, HIF2A (also known as EPAS1) and HIF3A, and a constitutively expressed β subunit, HIF1B (also known as ARNT). Under atmospheric oxygen tensions, HIF1A protein is rapidly degraded due to hydroxylation by prolyl hydroxylase proteins (PHDs). These hydroxylated proteins are then recognised by the Von Hippel–Lindau (VHL) protein which targets them for proteosomal degradation. Under hypoxic conditions, PHDs are unable to hydroxylate HIF1A which therefore cannot be targeted for degradation by the VHL protein (Semenza 2003). Stabilised HIF1A subunits translocate from the cytoplasm to the nucleus where they bind with HIF1B to activate target genes (Wenger 2002). HIF1A, the first HIF- α subunit described, is thought to be the global regulator of the hypoxic response (Semenza & Wang 1992). However, the contribution of HIF2A and HIF3A, which are believed to be regulated in a similar manner (Ivan *et al.* 2001, Masson *et al.* 2001), remain to be fully characterised. All three α subunits share many sequence

similarities (reviewed by Lee *et al.* (2004)); they all possess the Per, Arnt/HIF1B and Sim domain essential for binding to HIF1B and the oxygen-dependent degradation domain, the target for degradation under normoxic conditions. HIF3A differs from HIF1A and HIF2A as it lacks the C-terminal activation domain required for co-activator binding. Thus, HIF3A is unable to recruit co-transcriptional regulators and basal transcriptional machinery to gene targets. HIF- α subunits have been found to activate genes that contain the hypoxia response element sequence located in the promoter region of hypoxia responsive genes (Wenger 2002). HIF2A can target this sequence independently of HIF1A, suggesting that they have functionally diverse roles. For example, HIF1A, originally thought to be the main oxygen sensing subunit, predominantly regulates glycolytic genes (Hu *et al.* 2003), whereas HIF2A is the main regulator of hypoxia-induced erythropoietin in tissues which express both HIF1A and HIF2A (Warnecke *et al.* 2004, Rankin *et al.* 2007). HIF2A has also been shown to promote cell cycle progression in hypoxic renal clear cell carcinoma cells (Gordan *et al.* 2007). To date, little is known about which genes are targeted by HIF3A.

The functional significance of HIF genes has been highlighted using targeted gene inactivation. HIF1A^{-/-} mice are non-viable, displaying developmental arrest by E9.0 with significant mesenchymal cell death and impaired vascular development (Kotch *et al.* 1999). HIF1B null mice are embryonic lethal by E10.5, displaying yolk sac and placental deficiencies and decreased numbers of haematopoietic progenitors (Nishi *et al.* 2004). HIF2A^{-/-} mice develop severe vascular defects and show developmental arrest between E9.5 and E12.5 with variability depending on the genetic background (Compernelle *et al.* 2002, Nishi *et al.* 2004).

In mouse ES cells, both HIF1A and HIF2A are expressed but HIF1A appears to be central to regulating hypoxic responses, since it targets many oxygen-dependent genes that are not regulated by HIF2A (Hu *et al.* 2006). However, HIF2A has been found to be a direct upstream regulator of POU5F1 (OCT4) in mouse ES cells, suggesting that HIF2A is involved in the regulation of stem cell maintenance (Covello *et al.* 2006). In hES cells, the mechanism of hypoxic regulation appears to differ since HIF1A protein is only transiently expressed for ~48 h following exposure to low oxygen tension (Cameron *et al.* 2008).

While there has been controversy in the literature, this study aims to prove that hES cell culture is improved under physiological oxygen concentrations compared with ambient, atmospheric oxygen tension. We investigate the effect of oxygen tension on hES cell morphology, pluripotency and proliferation and determine the functional significance and potential hierarchy of HIFs in regulating these hypoxic responses.

Results

Cell morphology

Under hypoxic conditions, morphology was altered (Fig. 1a); colonies appeared more compact with clearly defined borders. At 20% oxygen, the cells began to show peripheral, spontaneous differentiation by day 4 post-passage, characterised by enlarged, flattened cells with translucent cytoplasm (Fig. 1aF). As a consequence, hES cells cultured at 20% oxygen required more frequent passaging to limit spontaneous differentiation. In contrast, hES cells maintained at 5% oxygen were passaged due to space constraints, rather than to remove differentiated areas. Thus, colony diameter was ~50% larger at 5% oxygen than those cultured under 20% oxygen on each day post-passage ($P < 0.01$ – $P < 0.001$;

Fig. 1b). In total, hES cells were maintained under both 5% and 20% oxygen for ~10–12 months, or 36–40 passages and the same morphological differences were maintained throughout this period. To further quantify the increase in colony size, equal numbers of cells were seeded on Matrigel coated plates on day 0 under both oxygen tensions and total hES cell number determined on each subsequent day post-passage. A significant increase in cell number was apparent by 48 h post-passage under 5% oxygen compared with 20% oxygen and was maintained on days 3 and 4 ($P < 0.001$; Fig. 1c). Using Ki67 labelling of hES cells on Matrigel on day 3 post-passage, there was a dramatic decrease in the number of proliferating cells at 20% oxygen compared with 5% oxygen with ~50% of hES cells cultured under 20% oxygen being Ki67 positive compared with virtually




Figure 1 (a) Phase contrast images of Hues 7 hES colonies cultured on MEFs under 5% (A, C and E) and 20% (B, D and F) oxygen on days 2 (A and B), 3 (C and D) and 4 (E and F) post-passage. Area of differentiation highlighted with arrow. Scale bar, 500 µm. (b) Average maximum hES cell colony diameter at 5 and 20% oxygen on days 2–4 post-passage ** $P < 0.01$, *** $P < 0.001$ significantly different from 5% oxygen ($n = 6$ for each day). (c) Average hES cell number at 5% and 20% oxygen on days 1–4 post-passage. Cell numbers were normalised on day 0 by passaging equal numbers of cells under both oxygen tensions. *** $P < 0.001$ significantly different from 5% oxygen ($n = 21$ for each day). (d) Ki67 (green) and DAPI (blue) labelling of Hues 7 hES cells at 5% (A–C) and 20% (D–F) oxygen on day 3 post-passage, phase contrast (C and F). Negative control, secondary antibody only (G). Scale bar, 250 µm.

all cells at 5% oxygen (Fig. 1d). Similar results were obtained for hES cells cultured on mouse embryonic fibroblasts (MEFs; data not shown).

Pluripotency marker expression

To investigate the effect of oxygen tension on the expression of pluripotency markers, hES cells cultured under either 20% or 5% oxygen were obtained on day 3 post-passage, prior to the onset of any overt differentiation observed in cells cultured at 20% oxygen. There was a significant ($P < 0.05$) reduction in *POU5F1* ($80 \pm 20\%$), *SOX2* ($75 \pm 18\%$) and *NANOG* ($60 \pm 10\%$) mRNA expression in cells maintained at 20% oxygen compared with those cultured at 5% oxygen (Fig. 2a). Although immunocytochemistry detected similar immunoreactivity for *POU5F1*, *SOX2*, TRA-1-60 and TRA-1-81 (Fig. 2b), using western blotting, *POU5F1* was significantly decreased by $\sim 40\%$ in hES cells cultured at 20% oxygen compared with 5% oxygen ($P < 0.01$; Fig. 2c and d).

HIF expression

Real-time RT-PCR was used to investigate the effect of environmental oxygen tension on the differential expression of *HIF1A*, *HIF1B*, *HIF2A* and *HIF3A* subunits. All HIFs were expressed in hES cells cultured under both 20% and 5% oxygen, but there was no

significant difference in the mRNA expression of the α subunits with respect to oxygen tension. Surprisingly, *HIF1B* was significantly upregulated under hypoxic conditions (Fig. 3a).

Western blotting was performed on day 3 post-passage to quantify the level of HIF protein expression in hES cells (Fig. 3b). *HIF1A* was not expressed following long-term culture under 5% or 20% oxygen tension and there was no significant difference in the expression of *HIF1B*. However, *HIF2A* and *HIF3A* were significantly upregulated under 5% oxygen compared with 20% oxygen ($P < 0.05$; Fig. 3c).

Using immunocytochemistry, *HIF1A* was found to be only transiently expressed being present in the nucleus of hES cells cultured under 5% oxygen for 48 h but absent after long-term culture under hypoxic conditions (Fig. 3d). *HIF1A* was also not expressed in hES cells cultured under 20% oxygen. *HIF2A* and *HIF3A* both displayed a largely cytoplasmic expression following 48 h of culture at 5% oxygen. However, following long-term hypoxic culture they both displayed a predominantly nuclear localisation. Under 20% oxygen, *HIF2A* and *HIF3A* remained cytoplasmic. *HIF1B*, which is known to be constitutively expressed, was present in the nuclei of hES cells when cultured under both 20% and 5% oxygen tensions. It was therefore not necessary to investigate the localisation of *HIF1B* in hES cells cultured in 5% oxygen for 48 h.




Figure 2 (a) mRNA expression of *POU5F1*, *SOX2* and *NANOG* cultured under 5% and 20% oxygen using relative quantification real-time RT-PCR. All data have been normalised to *UBC* and to 1 for 5% oxygen ** $P < 0.01$ ($n = 6$). (b) Immunocytochemistry of *POU5F1* (A and E), *SOX2* (B and F), TRA-1-60 (C and H) and TRA-1-81 (D and G) of hES cells cultured under 5% (A–D) and 20% (E–H) oxygen. Scale bar = 100 μ m. (c) Representative western blot of *POU5F1* expression in hES cells cultured under 5% and 20% oxygen. (d) Quantification of *POU5F1* western blots. Data were normalised to β -actin and to 1 for 5% oxygen. ** $P < 0.01$ significantly different to 5% oxygen ($n = 3$).




Figure 3 (a) mRNA expression of HIFs in hES cells at 5% and 20% oxygen. All data have been normalised to *UBC* and to 1 for 20% oxygen. * $P < 0.05$ significantly different to 20% oxygen ($n = 3$). (b) Representative western blots of HIFs after long-term culture at 5% or 20% oxygen (C) represents positive control protein. (c) Quantification of HIF western blots. Data were normalised to β -actin and to 1 for 20% oxygen. * $P < 0.05$ significantly different to 20% oxygen ($n = 3$). (d) Protein expression of HIFs by immunocytochemistry in hES cells cultured under 5% oxygen for 48 h (5% 48 h; A and B, G and H, M and N), 5% oxygen for long-term (5% LT = more than three passages; C and D, I and J, O and P, S and T) and 20% oxygen (E and F, K and L, Q and R, U and V). HIF1A (green; A–F), HIF2A (green; G–L), HIF3A (red; M–R) and HIF1A (green; S–V). DAPI (blue). Negative controls = FITC secondary only (W) and Texas Red secondary only (X). Scale bar = 40 μ m.

HIF knockdown

Following siRNA real-time RT-PCR confirmed an 83%, 86% and 90% knockdown of *HIF1A*, *HIF2A* and *HIF3A* respectively compared with transfection control siRNA (Fig. 4a, c and e). Initial studies examined the effect of knocking down individual HIF- α subunits for 48 h. When *HIF2A* was silenced *HIF1A* mRNA expression was not affected (Fig. 4a). However, the knockdown of *HIF3A* significantly upregulated *HIF1A* mRNA expression ($P < 0.05$, Fig. 4a). As expected, HIF1A protein was not expressed in hES cells maintained at 5% oxygen and was not induced by silencing *HIF2A*. Interestingly, HIF1A protein was induced when *HIF3A* was silenced (Fig. 4b). When both *HIF2A* and *HIF3A* were silenced, HIF1A was expressed but at a significantly reduced level compared with when *HIF3A* alone was knocked down ($P < 0.001$, Fig. 4b).

When *HIF1A* was silenced, *HIF2A* mRNA expression were unaffected. However, there was a significant reduction of HIF2A mRNA ($P < 0.01$) and protein ($P < 0.05$) expression observed when *HIF3A* was silenced (Fig. 4c and d).

HIF3A mRNA (Fig. 4e) and protein (Fig. 4f) expression were found to be significantly upregulated when *HIF1A* and *HIF2A* were knocked down independently.

Effect of HIFs on pluripotency marker expression

Using real-time RT-PCR there was a significant reduction in *POU5F1* (Fig. 5a; $P < 0.001$), *SOX2* (Fig. 5c; $P < 0.001$, $P < 0.05$) and *NANOG* (Fig. 5e; $P < 0.001$) when *HIF2A* and *HIF3A* were silenced independently. As expected, silencing of *HIF1A* did not alter the mRNA expression of *POU5F1*, *SOX2* and *NANOG* compared with transfection control siRNA (Fig. 5a, c and e). At the protein level, POU5F1, SOX2 and NANOG were significantly reduced when *HIF2A* ($P < 0.001$) and *HIF3A* ($P < 0.05$) were silenced (Fig. 5b, d and f).

Effect of HIF expression on hES cells morphology

Silencing *HIF1A* or *HIF3A* did not affect hES cell morphology (Fig. 6a) and following knockdown these cells could be maintained in culture, remaining TRA-1-60 and POU5F1 positive 48 h (Fig. 6b) and two passages (Fig. 6c) post transfection. Moreover, *HIF1A* and *HIF3A* silenced colonies contained similar levels of SSEA1 expression as the transfection controls (Fig. 6b). However, when *HIF2A* expression was knocked down colonies appeared to have less clearly defined borders and large areas of differentiation (Fig. 6a). These cells failed to maintain pluripotency, being SSEA1 positive




Figure 4 mRNA expression levels of (a) *HIF1A*, (c) *HIF2A* and (e) *HIF3A* when each HIF- α isoform was silenced in hES cells cultured under 5% oxygen for 48 h. All data has been normalised to UBC and to 1 for the transfection control. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ significantly different to transfection control ($n = 3$). (b) Representative HIF1A protein expression and quantification when *HIF2A* and *HIF3A* were silenced separately and simultaneously. *** $P < 0.001$ significantly different to *HIF3A* siRNA. Representative (d) HIF2A and (f) HIF3A protein expression and quantification when each HIF- α isoform was silenced * $P < 0.05$, ** $P < 0.01$, * $P < 0.001$ significantly different to transfection control ($n = 3$). Protein was collected for all samples 48 h post-transfection.




Figure 5 mRNA expression levels of (a) *POU5F1*, (c) *SOX2* and (e) *NANOG* when HIF- α subunits were silenced in hES cells cultured under 5% oxygen for 48 h. Data normalised to UBC and to 1 for the transfection control. * $P < 0.05$, *** $P < 0.001$ significantly different to transfection control ($n = 3$). Representative (b) *POU5F1*, (d) *SOX2* and (f) *NANOG* protein expression and quantification by western blotting when HIF- α subunits were silenced. * $P < 0.05$, *** $P < 0.001$ significantly different to transfection control ($n = 3$). Protein was collected for all samples 48 h post-transfection.

and displaying large areas that were TRA-1-60 and *POU5F1* negative (Fig. 6b) and failed to proliferate during culture. Consequently, *HIF2A* silenced hES cells that were cultured for two passages post transfection were negative for TRA-1-60 and *POU5F1* (Fig. 6c). Double knockdowns combining *HIF1A* and either *HIF2A* or *HIF3A* showed significant areas of differentiation and the cells failed to form colonies (Fig. 6a). When *HIF2A* and *HIF3A* were both silenced simultaneously, hES cells were capable of forming colonies but possessed large areas of differentiation. In contrast, triple HIF- α knockdown hES cells did not form colonies and did not survive in culture.

Effect of HIF expression on hES cells proliferation

At 48 h post-transfection, hES cell number (Fig. 7a) and colony size (Fig. 7b) were significantly ($P < 0.001$) reduced by $\sim 30\%$ when *HIF2A* expression was knocked down and displayed a further decrease ($P < 0.01$) when *HIF2A* and *HIF3A* were silenced simultaneously. There was no additional increase in the diameter of *HIF2A* silenced cells 72 h post-transfection (data not shown). Colony size and cell number were unaffected by *HIF1A* or *HIF3A* knockdown. It was not possible to measure the

colony size of double knockdown combinations of *HIF1A* and *HIF2A* or *HIF1A* and *HIF3A* or the triple knockdown due to the absence of colony formation occurring in these populations. Virtually all cells were positive for Ki67 when either *HIF1A* or *HIF3A* expression was silenced whereas Ki67 expression decreased to $\sim 85\%$ when *HIF2A* was knocked down ($P < 0.001$, Fig. 7c and d).

Discussion

There is emerging evidence to suggest that culture under more physiological oxygen conditions (2–5% oxygen) decreases the amount of spontaneous hES cell differentiation (Ezashi *et al.* 2005, Westfall *et al.* 2008). Similarly, we found that spontaneous differentiation was apparent by day 4 post-passage in hES cells cultured under atmospheric conditions, whereas cells cultured at 5% oxygen maintained pluripotency. As a result of this, hES cells cultured under 20% oxygen required more frequent passaging prior to confluency to remove differentiated areas, whereas hES cells maintained at 5% oxygen were only passaged due to space constraints on the plate. These findings suggest that environmental oxygen tension has a critical role in the maintenance of hES

cell pluripotency. To investigate differences between the two oxygen concentrations, results were obtained on day 3 post-passage, prior to overt cell differentiation.

Culture at 5% oxygen, whether on MEFs or feeder-free on Matrigel, increased the rate of cell proliferation which agrees with data from Ludwig *et al.* (2006), producing significantly larger hES cell colonies. This may be due to the observed increase in NANOG expression levels, since it has recently been shown that NANOG regulates S-phase entry in hES cells and that overexpression of NANOG significantly increases proliferation by binding to regulatory regions of CDK6 and CDC25A, two important cell cycle regulators (Neganova *et al.* 2009, Zhang *et al.* 2009). Similar increased rates of proliferation have also been observed in many other primitive populations including mesenchymal stem cells and neural progenitor cells when cultured under low oxygen

tensions (Grayson *et al.* 2007, Zhao *et al.* 2008). Data from mesenchymal stem cells suggest the involvement of HIFs, specifically HIF2A (Grayson *et al.* 2007). This is supported in hypoxic carcinoma cells where HIF2A has been shown to promote cell cycle progression (Gordan *et al.* 2007).

In agreement with Forsyth *et al.* (2008) and Westfall *et al.* (2008), there was no significant increase in the mRNA expression of the HIF- α subunits under hypoxia suggesting that altered protein levels are due to post-transcriptional regulation as found in other cell types (Huang *et al.* 1998, Lang *et al.* 2002). However, in contrast to Westfall *et al.* (2008), *HIF1B* mRNA expression was upregulated under hypoxia which may reflect differences in both the methodology and cell lines used. To date, most research has focused on the HIF1A subunit which is thought to be the master regulator of the




Figure 6 (a) Representative phase contrast images of hES cells cultured under 5% oxygen 48 h after transfection with HIF- α siRNA. Controls contain the same volume and concentration of transfection reagent and AllStars control siRNA as each of the knockdowns. Scale bar=100 μ m. (b) Protein expression of pluripotency markers TRA-1-60 (A, G, M and S), merged with DAPI (B, H, N and T), POU5F1 (C, I, O and U), merged with DAPI (D, J, P and V) and differentiation marker SSEA1 (E, K, Q and W), merged with DAPI (F, L, R and X) in hES cells 48 h following siRNA transfection of HIF- α subunits. Scale bar=100 μ m. (c) Phase contrast and protein expression of POU5F1 and TRA-1-60 in hES cells two passages following HIF- α gene silencing. Scale bar=100 μ m.




Figure 7 Effect of silencing HIF- α subunits in hES cells cultured under 5% oxygen 48 h following siRNA transfection on average cell number ($n=3$). Cell numbers were normalised on day 0 by passaging an equal number of cells. (b) Effect of HIF- α silencing 48 h following siRNA transfection on average maximum colony diameter ($n=6$). $***P<0.001$ significantly different from transfection control. Bars with the same superscript are significantly different; ^a $P<0.01$. (c) Ki67 (green) and DAPI (blue) labelling of HIF- α silenced hES cells and transfection control. Scale bar = 100 μm . (d) Percentage of HIF- α silenced Ki67 positive hES cells. $***P<0.001$ significantly different from transfection control.

hypoxic response (Mazure *et al.* 2004). However, our data suggests this is not the case, at least in hES cells since HIF1A was only transiently expressed in the nucleus for ~ 48 h following exposure to hypoxia. These findings are in agreement with recent data from Cameron *et al.* (2008) using a 2% environmental oxygen tension, although contrary to those of Westfall *et al.* (2008), who found HIF1A protein to be expressed in hES cells after 10 days of culture under both 5 and 20% oxygen. The reason for this discrepancy is unknown although HIF1A would be expected to be rapidly degraded by PHDs under atmospheric conditions. Moreover, the HIF1A staining observed by the latter investigators was cytoplasmic and therefore would not be expected to regulate the expression of hypoxia regulated genes. Our study suggests that HIF1A may play a role in the initial adaptation of hES cells to hypoxia but some other factor(s) must be responsible for maintaining the hypoxic response. We propose that HIF2A or HIF3A, both of which are translocated from the cytoplasm to the nucleus and are upregulated following

culture under hypoxic conditions, may take over the initial, transient role of HIF1A. This is the first time that HIF3A and HIF1B have been demonstrated in hES cells.

hES cell colonies cultured at 5% or 20% oxygen were both positive for pluripotency markers albeit at a significantly decreased level at 20% oxygen. These results were not apparent using immunocytochemistry and highlight the importance of performing quantitative techniques when assessing pluripotency. The current results suggest that although hES cells are capable of being maintained under atmospheric conditions pluripotency is reduced before the appearance of morphological differentiation. This shows that the manual removal of differentiated cells at the time of passage, a method often used during routine culture at 20% oxygen is not effective in ensuring the maintenance of a highly pluripotent population. However, culturing hES cells at 5% oxygen does not require the routine removal of differentiated areas and can therefore be considered a much more effective method of maintaining pluripotency during long term culture. Interestingly, our data

active as observed in the initial adaptation of hES cells to hypoxia.

In conclusion, these studies demonstrate that the culture of hES cells is preferential under a reduced oxygen environment. In contrast to other cell types, HIF1A is only responsible for the initial adaptation of cells to hypoxia, whereas HIF2A regulates the long-term hypoxic response by controlling hES cell pluripotency and proliferation. HIF3A acts to regulate the expression level of both HIF1A and HIF2A. These data provide a greater understanding of the mechanisms which regulate hES cell function and the pluripotent state.

Materials and Methods

Culture of hES cells

Hues-7 hES cells (D Melton, Howard Hughes Medical Institute/Harvard University) were maintained on γ -irradiated MEFs in knockout (KO) DMEM (Invitrogen) supplemented with 10% KO serum replacement (Invitrogen), 1 mM L-glutamine, 50 mM β -mercaptoethanol, 1% non-essential amino acids, 10 ng/ml basic FGF factor (Peprotech Ltd, London, UK) and 100 μ g/ml penicillin/streptomycin before being transferred on to plates pre-coated with Matrigel (BD Biosciences, San Diego, CA, USA), in medium that had been cultured over-night on γ -irradiated MEFs. hES cells were initially cultured under atmospheric (~20%) oxygen before half were transferred into 5% O₂, 5% CO₂ and balanced nitrogen. Unless stated, cells were cultured at 5% oxygen for a minimum of three passages prior to use.

Quantitative real-time RT-PCR

mRNA was isolated from hES cells cultured on Matrigel on day 3 post-passage using TriReagent (Sigma) and RNA (2 μ g) was reverse transcribed into cDNA using Moloney murine leukaemia virus reverse transcriptase (Promega). The standard curve method of relative quantification real-time PCR was performed using Applied Biosystems reagents in 20 μ l reactions containing 4 μ g of cDNA, 10 μ l 2 \times Taqman Universal PCR Master Mix, 1 μ l Probes and Primer Mix (*HIF1A*: Hs00936368_m1; *HIF1B*: Hs00231048_m1; *HIF2A*: Hs01026142_m1; *HIF3A*: Hs00541709_m1; *POU5F1*: Hs01895061_u1; *NANOG*: Hs02387400_g1; *SOX2*: Hs00602736_s1; ubiquitin C (*UBC*): Hs00824723_m1) using a 7500 Real-Time PCR System. The following conditions were used; 50 °C for 2 min, 95 °C for 10 min, 45 cycles at 95 °C for 15 s and 60 °C for 1 min. Placental cDNA (0–10 ng) was used to create a standard curve for each gene of interest as well as for *UBC*, used as an endogenous control. All target transcripts were analysed in duplicate and normalised to *UBC*.

Western blotting

hES cells cultured on Matrigel were rinsed with ice-cold PBS and treated with ice-cold radio immunoprecipitation assay lysis buffer (50 mM Tris-HCl (pH 7.4), 1% nonyl

phenoxy polyethoxy ethanol, 0.25% Na-deoxycholate and complete protease inhibitors (Roche) supplemented with 1 mM phenylmethylsulphonyl fluoride, 1 mM Na₃VO₄ and 1 mM NaF) for 30 min with gentle agitation. Protein concentration was quantified using the Bradford assay (Bradford 1976) and lysates (75 μ g) were resolved on an 8% SDS bisacrylamide gel. Positive controls were run for each protein of interest; hypoxic colorectal carcinoma cell protein for HIF1A, foetal heart (obtained following ethical approval from the Southampton & South West Hampshire Local Research Ethics Committee, under guidelines issued by the Polkinghorne Committee) for HIF1B and NT2 cells treated with 130 μ M cobalt chloride for 4 h for HIF2A and HIF3A. The protein was electro-transferred to a nitrocellulose membrane and blocked in PBS containing 5% non-fat powdered milk and 0.1% Tween-20 for 1 h at room temperature. Primary antibodies were diluted in blocking buffer and incubated with the membrane overnight at 4 °C; HIF1A (BD Biosciences) 1:250, HIF1B (Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA) 1:250, HIF2A (Novus Biologicals, Cambridge, UK) 1:500, HIF3A (Santa Cruz) 1:1000, POU5F1 (Santa Cruz) 1:1000, SOX2 (Millipore, Billerica, MA, USA) 1:000 and NANOG (Abcam, Cambridge, UK) 1:1000. Membranes were washed and incubated with either peroxidase labelled anti-mouse antibody (Amersham) 1:50 000, or anti-goat antibody (Sigma) 1:200 000, in blocking buffer for 1 h at room temperature. Protein expression was quantified relative to β -actin expression which was detected with mouse anti- β -actin peroxidase-conjugated antibody (1:50 000; Sigma). Membranes were developed using ECL advanced western blotting detection kit (Amersham).

Immunocytochemistry

hES cells cultured on Matrigel were fixed in 4% paraformaldehyde for 20 min. Non-specific antibody binding was blocked with 3% donkey serum and where necessary cells were permeabilised with 0.1% tritonX-100 for 1 h before the addition of primary antibodies diluted in PBS and 3% donkey serum. Primary antibodies used were TRA-1-60 (Santa Cruz) 1:100, TRA-1-81 (gift from P Andrews) 1:50, SSEA1 (Santa Cruz) 1:100, POU5F1 1:100, SOX2 (Chemicon) 1:150, HIF1A 1:250, HIF1B 1:100, HIF2A 1:100, HIF3A 1:50, Ki67 (Novocastra Laboratories, Newcastle upon Tyne, UK) 1:100 and incubated overnight in a humidified chamber at 4 °C. Secondary antibody staining was performed with anti-mouse IgG conjugated-FITC 1:100, anti-mouse IgM conjugated-FITC 1:200 (Sigma) or anti-goat IgG Alexa Fluor 594 1:100 (Sigma) for 1 h in a humidified chamber. Nuclei were labelled with DAPI (Vecta Laboratories, Peterborough, UK).

siRNA

siRNA experiments were carried out on hES cells cultured on Matrigel under 5% oxygen for a minimum of three passages. hES cells cultured under 5% oxygen were passaged and incubated overnight. For each transfection 50 nM siRNA (*HIF1A*: Hs_HIF1A_5; *HIF2A*: Hs_EPAS1_5_HP; *HIF3A*: HsHIF3A_1_HP; Qiagen) along with 12 μ l HiPerfect transfection reagent (Qiagen) were mixed into 200 μ l of KO-DMEM

and added in a dropwise manner to 1-well of a 6-well plate. At 48 h after transfection, cells were harvested and extracts were prepared for mRNA and protein. For *HIF3A*, siRNA targeting exon 5 was used which is present in all the major known *HIF3A* isoforms (Maynard *et al.* 2003). AllStars Negative Control (Transfection control, Qiagen) siRNA that has no homology to any known mammalian gene was used as a negative control for each transfection. For double and triple knockdowns 50 nM siRNA and 12 μ l HiPerfect transfection reagent were added in 600 μ l of KO-DMEM. Twice or three times the volume of Allstars Negative control was added to controls for double and triple knockdowns respectively. hES cells were analysed for knockdown using relative quantification real-time RT-PCR.

Statistical analysis

All data were analysed to determine whether they were normally distributed, using the Anderson–Darling normality test. Differences between oxygen tension and maximum colony diameter and cell number were analysed using a Student's *t*-test. Differences in relative gene and protein expression between cells cultured at 5 and 20% oxygen were analysed using a 1-sample *t*-test. Differences between negative control and knockdown results were analysed using a Student's *t*-test. All data represent at least three independent experiments. A value of $P < 0.05$ was considered significant. Data are presented as mean \pm S.E.M.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was funded by the Gerald Kerkut Charitable Trust, a Wellcome Trust Research Career Development Fellowship (WT066492MA), the MRC (G0701153), the Wellcome Trust in partnership with JDRF (WT074320MA). N A H receives support from Manchester NIHR BRC.

References

- Bradford MM 1976 A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72** 248–254.
- Cameron CM, Harding F, Hu WS & Kaufman DS 2008 Activation of hypoxic response in human embryonic stem cell-derived embryoid bodies. *Experimental Biology and Medicine* **233** 1044–1057.
- Chen HF, Kuo HC, Chen W, Wu FC, Yang YS & Ho HN 2009 A reduced oxygen tension (5%) is not beneficial for maintaining human embryonic stem cells in the undifferentiated state with short splitting intervals. *Human Reproduction* **1** 71–80.
- Compennolle V, Brusselmans K, Acker T, Hoet P, Tjwa M, Beck H, Plaisance S, Dor Y, Keshet E, Lupu F *et al.* 2002 Loss of HIF-2 α and inhibition of VEGF impair fetal lung maturation, whereas treatment with VEGF prevents fatal respiratory distress in premature mice. *Nature Medicine* **8** 702–710.
- Covello KL, Kehler J, Yu H, Gordan JD, Arsham AM, Hu CJ, Labosky PA, Simon MC & Keith B 2006 HIF-2 α regulates Oct-4: effects of hypoxia on stem cell function, embryonic development, and tumor growth. *Genes and Development* **20** 557–570.
- Dumoulin JC, Meijers CJ, Bras M, Coonen E, Geraedts JP & Evers JL 1999 Effect of oxygen concentration on human *in vitro* fertilisation and embryo culture. *Human Reproduction* **14** 465–469.
- Ezashi T, Das P & Roberts RM 2005 Low O₂ tensions and the prevention of differentiation of hES cells. *PNAS* **102** 4783–4788.
- Fischer B & Bavister BD 1993 Oxygen tension in the oviduct and uterus of rhesus monkeys, hamsters and rabbits. *Journal of Reproduction and Fertility* **99** 673–679.
- Forsyth NR, Musio A, Vezzoni P, Simpson AH, Noble BS & McWhir J 2006 Physiologic oxygen enhances human embryonic stem cell clonal recovery and reduces chromosomal abnormalities. *Cloning and Stem Cells* **8** 16–23.
- Forsyth NR, Kay A, Hampson K, Downing A, Talbot R & McWhir J 2008 Transcriptome alterations due to physiological normoxic (2% O₂) culture of human embryonic stem cells. *Regenerative Medicine* **3** 817–833.
- Gordan JD, Bertout JA, Hu C-J, Diehl JA & Simon MC 2007 HIF-2 α promotes hypoxic cell proliferation by enhancing c-myc transcriptional activity. *Cancer Cell* **11** 335–347.
- Grayson WL, Zhao F, Bunnell B & Ma T 2007 Hypoxia enhances proliferation and tissue formation of human mesenchymal stem cells. *Biochemical and Biophysical Research Communications* **358** 948–953.
- Hara S, Hamada J, Kobayashi C, Kondo Y & Imura N 2001 Expression and characterization of hypoxia-inducible factor (HIF)-3 α in human kidney: suppression of HIF-mediated gene expression by HIF-3 α . *Biochemical and Biophysical Research Communications* **287** 808–813.
- Harvey AJ, Kind KL, Pantaleon M, Armstrong DT & Thompson JG 2004 Oxygen-regulated gene expression in bovine blastocysts. *Biology of Reproduction* **71** 1108–1119.
- Hu CJ, Wang LY, Chodosh LA, Keith B & Simon MC 2003 Differential roles of hypoxia-inducible factor 1 α (HIF-1 α) and HIF-2 α in hypoxic gene regulation. *Molecular and Cellular Biology* **23** 9361–9374.
- Hu CJ, Iyer S, Sataur S, Covello KL, Chodosh LA & Simon MC 2006 Differential regulation of the transcriptional activities of hypoxia-inducible factor 1 α (HIF-1 α) and HIF-2 α in stem cells. *Molecular Cell Biology* **26** 3514–3526.
- Huang LE, Gu J, Schau M & Bunn HF 1998 Regulation of hypoxia-inducible factor 1 α is mediated by an O₂-dependent degradation domain via the ubiquitin-proteasome pathway. *PNAS* **95** 7987–7992.
- Ivan M, Kondo K, Yang H, Kim W, Valiando J, Ohh M, Salic A, Asara JM, Lane WS & Kaelin WG 2001 HIF α targeted for VHL-mediated destruction by proline hydroxylation: implications for O₂ sensing. *Science* **292** 464–468.
- Jeong CH, Lee HJ, Cha JH, Kim JH, Kim KR, Kim JH, Yoon DK & Kim KW 2007 Hypoxia-inducible factor-1 α inhibits self-renewal of mouse embryonic stem cells *in vitro* via negative regulation of the leukemia inhibitory factor-STAT3 pathway. *Journal of Biological Chemistry* **282** 13672–13679.
- Kotch LE, Iyer NV, Laughner E & Semenza GL 1999 Defective vascularization of HIF-1 α -null embryos is not associated with VEGF deficiency but with mesenchymal cell death. *Developmental Biology* **209** 254–267.
- Kovacic B & Vlasisvljevic V 2008 Influence of atmospheric versus reduced oxygen concentration on development of human blastocysts *in vitro*: a prospective study on sibling oocytes. *Reproductive Biomedicine Online* **17** 229–236.
- Lang KJD, Kappel A & Goodall GJ 2002 Hypoxia-inducible factor-1 α mRNA contains an internal ribosome entry site that allows efficient translation during normoxia and hypoxia. *Molecular Biology of the Cell* **13** 1792–1801.
- Lee J-W, Bae S-H, Jeong J-W, Kim S-H & Kim K-W 2004 Hypoxia-inducible factor (HIF)-1 α : its protein stability and biological functions. *Experimental and Molecular Medicine* **36** 1–12.
- Lennon DP, Edmison JM & Caplan AI 2001 Cultivation of rat marrow-derived mesenchymal stem cells in reduced oxygen tension: effects on *in vitro* and *in vivo* osteochondrogenesis. *Journal of Cellular Physiology* **187** 345–355.

- Ludwig TE, Levenstein ME, Jones JM, Berggren WT, Mitchen ER, Frane JL, Crandall LJ, Daigh CA, Conard KR, Piekarczyk MS *et al.* 2006 Derivation of human embryonic stem cells in defined conditions. *Nature Biotechnology* **24** 185–187.
- Masson N, Willam C, Maxwell PH, Pugh CW & Ratcliffe PJ 2001 Independent function of two destruction domains in hypoxia-inducible factor- α chains activated by prolyl hydroxylation. *EMBO Journal* **20** 5197–5206.
- Maynard MA, Qi H, Chung J, Lee EH, Kondon Y, Hara S, Conaway RC, Conaway JW & Ohh M 2003 Multiple splice variants of the human HIF-3 α locus are targets of the von Hippel–Lindau E3 ubiquitin ligase complex. *Journal of Biological Chemistry* **278** 11032–11040.
- Mazure NM, Brahimi-Horn MC, Berta MA, Benizri E, Bilton RL, Dayan F, Ginouves A, Berra E & Pouyssegur J 2004 HIF-1: master and commander of the hypoxic world. A pharmacological approach to its regulation by siRNAs. *Biochemical Pharmacology* **68** 971–980.
- Morrison SJ, Csete M, Groves AK, Melega W, Wold B & Anderson DJ 2000 Culture in reduced levels of oxygen promotes clonogenic sympathoadrenal differentiation by isolated neural crest stem cells. *Journal of Neuroscience* **20** 7370–7376.
- Neganova I, Zhang X, Atkinson S & Lako M 2009 Expression and functional analysis of G1 to S regulatory components reveals an important role for CDK2 in cell cycle regulation in human embryonic stem cells. *Oncogene* **28** 20–30.
- Nishi H, Nakada T, Kyo S, Inoue M, Shay JW & Isaka K 2004 Hypoxia-inducible factor 1 mediates upregulation of telomerase (hTERT). *Molecular Cell Biology* **24** 6076–6083.
- Olson SE & Seidel GE 2000 Reduced oxygen tension and EDTA improve bovine zygote development in a chemically defined medium. *Journal of Animal Science* **78** 152–157.
- Orsi NM & Leese HJ 2001 Protection against reactive oxygen species during mouse preimplantation embryo development: role of EDTA, oxygen tension, catalase, superoxide dismutase and pyruvate. *Molecular Reproduction and Development* **59** 44–53.
- Petersen A, Mikkelsen AL & Lindenberg S 2005 The impact of oxygen tension on developmental competence of post-thaw human embryos. *Acta Obstetrica et Gynecologica Scandinavica* **84** 1181–1184.
- Rankin EB, Biju MP, Liu Q, Unger TL, Rha J, Johnson RS, Simon MC, Keith B & Haase VH 2007 Hypoxia-inducible factor-2 (HIF-2) regulates hepatic erythropoietin *in vivo*. *Journal of Clinical Investigation* **117** 1068–1077.
- Rinaudo PF, Giritharan G, Talbi S, Dobson AT & Schultz RM 2006 Effects of oxygen tension on gene expression in preimplantation mouse embryos. *Fertility and Sterility* **86** 1252–1265.
- Schöler HR, Balling R, Hatzopoulos AK, Suzuki N & Gruss P 1989 Octamer binding proteins confer transcriptional activity in early mouse embryogenesis. *EMBO Journal* **8** 2551–2557.
- Semenza GL 2000 HIF-1 and human disease: one highly involved factor. *Genes and Development* **14** 1983–1991.
- Semenza GL 2003 Targeting HIF-1 for cancer therapy. *Nature Reviews. Cancer* **3** 721–732.
- Semenza GL & Wang GL 1992 A nuclear factor induced by hypoxia via *de novo* protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation. *Molecular and Cellular Biology* **12** 5447–5454.
- Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS & Jones JM 1998 Embryonic stem cell lines derived from human blastocysts. *Science* **282** 1145–1147.
- Thompson JC, Simpson AC, Pugh PA, Donnelly PE & Tervit HR 1990 Effect of oxygen concentration on *in vitro* development of preimplantation sheep and cattle embryos. *Journal of Reproduction and Fertility* **89** 573–578.
- Warnecke C, Zaborowska Z, Kurreck J, Erdmann VA, Frei U, Wiesener M & Eckardt KU 2004 Differentiating the functional role of hypoxia-inducible factor (HIF)-1 α and HIF-2 α (EPAS-1) by the use of RNA interference: erythropoietin is a HIF-2 α target gene in Hep3B and Kelly cells. *FASEB Journal* **18** 1462–1464.
- Wenger RH 2002 Cellular adaptation to hypoxia: O₂-sensing protein hydroxylases, hypoxia-inducible transcription factors, and O₂-regulated gene expression. *FASEB Journal* **16** 1151–1162.
- Westfall SD, Sachdev S, Das P, Hearne LB, Hannink M, Roberts RM & Ezashi T 2008 Identification of oxygen-sensitive transcriptional programs in human embryonic stem cells. *Stem Cells and Development* **17** 869–882.
- Zhang X, Neganova I, Przyborski S, Yang C, Cooke M, Atkinson SP, Anyfantis G, Fenyk S, Keith WN, Hoare SF *et al.* 2009 A role for NANOG in G1 to S transition in human embryonic stem cells through direct binding of CDK6 and CDC25A. *Journal of Cell Biology* **184** 67–82.
- Zhao T, Zhang CP, Liu ZH, Wu LY, Huang X, Wu HT, Xiong L, Wang X, Wang XM, Zhu LL *et al.* 2008 Hypoxia-driven proliferation of embryonic neural stem/progenitor cells—role of hypoxia-inducible transcription factor-1 α . *FEBS Journal* **275** 1824–1834.

Received 16 July 2009

First decision 18 August 2009

Revised manuscript received 8 September 2009

Accepted 14 September 2009