



Published in final edited form as:

Comp Biochem Physiol B Biochem Mol Biol. 2010 February ; 155(2): 201–209. doi:10.1016/j.cbpb.2009.11.012.

Characterization of expression, activity and role in antibacterial immunity of *Anopheles gambiae* lysozyme c-1

Mayur K. Kajla, Olga Andreeva, Thomas M. Gilbreath III^a, and Susan M. Paskewitz^{*}

Department of Entomology, University of Wisconsin, Madison, Wisconsin, USA

Abstract

There are eight lysozyme genes in the *Anopheles gambiae* genome. Transcripts of one of these genes, *LYSC-1*, increased in *Anopheles gambiae* cell line 4a3B by 24 h after exposure to heat-killed *Micrococcus luteus*. Lysozyme activity was also identified in conditioned media from the cell line from which the protein was purified to homogeneity using ion exchange and gel filtration. Mass spectrometric analysis of the purified protein showed 100% identity to lysozyme c-1. Purified lysozyme c-1 was tested against non-mosquito derived as well as culturable bacteria isolated from mosquito midguts. Lysozyme c-1 had negligible effects on the growth of most mosquito-derived bacteria *in vitro* but did inhibit the growth of *M. luteus*. Although Lys c-1 did not directly kill most bacteria, knockdown of *LYSC-1* resulted in significant mortality in mosquitoes subjected to hemocoelic infections with *Escherichia coli* but not *M. luteus* thus suggesting that this protein plays an important role in antibacterial defense against selected bacteria.

Keywords

insect immunity; mosquito; antibacterial protein; muramidase

1. Introduction

Lysozymes (EC 3.2.1.17) hydrolyze the β -1, 4-glycosidic linkage between N-acetylmuramic acid and N-acetylglucosamine of peptidoglycan, a cell wall component of bacteria. They are widespread in nature, occurring in bacteriophage, plants and animals. The role of this enzyme as a defense molecule has been well documented in vertebrates (Jolles and Jolles, 1984; Markart et al., 2004) and insects (Dunn, 1986; Hultmark, 1996; Powning and Davidson, 1973; Jolles et al., 1979; Hultmark et al., 1980; Abraham et al., 1995). In both vertebrates and invertebrates, lysozymes may act through multiple pathways to affect bacterial killing. First, these enzymes directly inhibit or kill a number of bacterial species through both enzymatic and nonenzymatic mechanisms (Nash et al. 2006). Lysozyme generally exhibits greater microbiocidal activity against Gram positive than Gram negative bacteria. However, the degree of activity against Gram negative bacteria varies interspecifically (Mai and Hu 2009; Yu et al. 2002) and is affected by osmolarity, ionic concentration and the presence of synergistic co-factors (Skerritt

© 2009 Elsevier Inc. All rights reserved.

Corresponding author: Dr. Susan M. Paskewitz, 237 Russell Labs, 1630 Linden Dr., Department of Entomology, Madison, WI 53706, USA. Tel. (608) 262-1269/ Fax (608) 262-3322 paskewit@entomology.wisc.edu.

^aCurrent address: Department of Health Sciences, University of California, Hewitt Hall, Room 3501, Mail Code: 4050, Irvine, CA 92697, USA

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

2004). Second, lysozymes may play broader roles in regulating the overall response to bacteria. For example, it has been suggested that the generation of small peptidoglycan fragments by lysozyme is a key aspect of the pattern recognition process that functions to initiate antibacterial responses in insects (Boman and Hultmark, 1987; Boman, 1991; Dunn et al., 1994). Indeed, inhibition of lysozyme resulted in reduced activation of the melanization pathway in *Tenebrio molitor* (Park et al 2007). Conversely, lysozymes may be essential to downregulation of the inflammatory response in some situations. Knockout of a mouse lysozyme resulted in the prolonged upregulation of an inflammatory response to infection by *Micrococcus luteus*, suggesting that the degradation of peptidoglycan eliminates a key signal of infection (Markart et al., 2004). Many invertebrates have more than one lysozyme and it is possible that these actions are carried out by separate enzymes with specific peptidoglycan degrading activities or temporal and spatial expression profiles.

In dipteran insects, a large expansion of the c-type lysozyme gene family has occurred (Daffre et al., 1994; Li et al., 2005; Ursic-Bedoya et al., 2005). In the mosquito *Anopheles gambiae*, there are eight c-type lysozymes with different tissue and stage expression profiles, biochemical properties (pI and molecular weight), and with varying conservation of amino acids critical for function (Li et al., 2005). One of these proteins, lysozyme c-1, is a typical basic lysozyme with the required amino acids for muramidase activity. It is constitutively expressed in most tissues of adult mosquitoes but is upregulated during bacterial infections (Li et al., 2005; Dong et al., 2006; Dong et al., 2009). We previously reported detection of this lysozyme in conditioned medium from *An. gambiae* derived 4a3B cells (Li and Paskewitz, 2006). To begin to clarify the roles of lysozymes in mosquito immunity, we now report on purification of Lys c-1 from this source and tests of its activity against selected bacteria as well as bacteria isolated from midguts of *An. gambiae* and *An. stephensi* females. We also conducted lysozyme c-1 knockdown experiments in *An. gambiae* female mosquitoes and analyzed the survival of mosquitoes after challenge with bacteria. Results of the Lys c-1 purification, its antibacterial activity and whole insect responses after gene silencing are presented.

2. Materials and methods

2.1. Mosquitoes, bacteria, and cell line

The G3 and 4a rr strains of *Anopheles gambiae* were acquired through the Malaria Research and Reference Reagent Resource Center (MR4, American Type Culture Collection Manassas, VA, USA) and the Center for Disease Control and Prevention and have been in culture for at least 5 years at the Department of Entomology, UW-Madison. A colony of *An. stephensi* was acquired through Johns Hopkins University (gift of Prof. Marcelo Jacobs Lorena) and has also been in culture for 5 years at UW. Mosquitoes were reared as described previously (Paskewitz et al., 1999). These mosquito colonies were used for the isolation of midgut bacteria. The G3 strain was used for gene silencing experiments.

A list of selected bacteria used in this study and their sources is given in Table 2. Some strains of bacteria used in this study were selected on the basis of previous reports of mosquito midgut bacterioflora (Lindh et al., 2005; Favia et al., 2007; Gusmao et al., 2007; Dillon and Dillon, 2004; Azambuja et al., 2005; Demaio et al., 1996). These included *Bacillus subtilis*, *Enterobacter sp.*, *Asaia sp.* (gifts from Dr. Nichole Broderick, Dept. of Plant Pathology, UW Madison). Bacteria were maintained on Luria Broth (LB) plates as well on tryptic soy broth (TSA) plates. Glycerol stocks were maintained at -80°C .

Anopheles gambiae cell line 4a3B was provided by Dr. Hans-Michael Mueller (Müller et al., 1999). This cell line has been in culture since 2000. Cells are routinely cultured in Schneider's Drosophila medium (Lonza, Walkersville, MD, USA) supplemented with 5-10% fetal calf serum and an antibiotic/antimycotic mixture containing 10,000 units of penicillin (base),

10,000 µg of streptomycin (base), and 25 µg of amphotericin B/mL utilizing penicillin G (sodium salt), streptomycin sulfate, and amphotericin B as Fungizone® Antimycotic in 0.85% saline (Invitrogen Corporation, Carlsbad, CA, USA) used at a final dilution of 1:100. For the purpose of Lys c-1 purification, cells were cultured in serum-free medium to confluence before conditioned medium was harvested. Conditioned medium was collected after removal of cell debris by centrifugation and then stored at -20 °C.

2.2. Isolation and identification of culturable bacteria from mosquito midguts

For the isolation of culturable bacteria, 2-3 day old female mosquitoes were cold anaesthetized for 5 min on ice and subsequently surface sterilized twice in 70% ethanol for 2 min. Mosquitoes were transferred to sterile saline (0.9% NaCl w/v in water) and were dissected under aseptic conditions. Ten midguts were transferred to 500 µL sterile saline and were homogenized using sterile pestles with a motor driven hand homogenizer (Kimble/Kontes, Vineland, New Jersey, USA).

Homogenates were sonicated in a water bath sonicator which operated at 125 watts for 2 min to facilitate the release of bacteria from the gut tissues. Serial dilutions (from 10^{-1} to 10^{-5}) from this homogenate were prepared in 0.9% sterile saline and 100 µl from each dilution were plated onto tryptic soy agar (3 g/L tryptic soy broth and 1.5% agar w/v). Plates were incubated at 28 °C for 48 h. Bacteria were analyzed for their morphology, color, texture and size. Bacterial colonies with unique morphologies were picked from different dilutions and streaked on TSA plates to isolate pure cultures. After three serial isolation steps, the bacteria were stored as glycerol stocks at -80 °C. For the identification of bacteria, genomic DNA from each isolated bacteria was prepared using the Master Pure Gram Positive DNA purification kit (Epicenter Biotechnologies, Madison, WI). Approximately 50-100 ng genomic DNA was used as a template for PCR amplification of the 16S rRNA gene using 27F and 1492R primers (gift of Dr. Jo Handlesman, Dept. of Bacteriology, UW Madison, WI). The sequences for these primers are: 27F: 5-AGR GTT TGA TYM TGG CTC AG-3; 1492R: 5-GGY TAC CTT GTT ACG ACT T-3. Amplification conditions were as follows: 94 °C - 4 min; followed by 30 cycles of 94 °C - 45 s; 58 °C - 45 s; 72 °C - 1.5 min; final extension of 72 °C - 10 min. PCR products were analyzed on 1% agarose gels stained with ethidium bromide. A DNA band of approximately 1.4 kb was excised from the gels and purified using a QIAquick gel extraction kit (QIAGEN Sciences, Maryland, USA). The purified PCR product was sequenced with the same primer sets used for PCR amplification (Biotech Center, UW Madison, WI, USA) and analyzed by FinchTV software (Geospiza Inc., Seattle, WA). Each manually corrected sequence over the entire length of the 16s RNA gene (1327-1411bp) was blasted against NCBI nr/nt database. Searches yielding the highest identity scores are reported as the closest matches for each respective bacterial sequence. Sequences from midgut derived bacteria were submitted to GenBank and the accession numbers are given in Table 2.

2.3. One dimensional SDS-PAGE and immunoblotting

Purified Lys c-1 was fractionated by 10% SDS-PAGE gels (NuPage, Invitrogen, USA). The gels were silver stained or used for immunoblotting. Polyclonal antibodies were produced to a peptide specific to Lys c-1 and do not cross react with other lysozymes (Li et al., 2005). In all experiments, the anti-Lys c-1 antibody (9122) was used at a 1:500 and blots were incubated overnight at 4 °C. Bound antibodies were detected by a peroxidase labeled goat anti-rabbit IgG (Kirkegaard and Perry Labs, Maryland, USA) at 1:5000 using a diaminobenzidine reagent (DAB; Sigma-Aldrich) as per supplied instructions.

2.4. Immune challenge to the cell line

The 4a3B cells were cultured in Schneider's Drosophila medium as described above. For immune challenge, *M. luteus* was grown overnight in tryptic soy broth medium (30 g/L). This

culture was then heat inactivated by autoclaving (121 °C for 15 min). The cells were pelleted by centrifugation at 7800 g for 10 min at room temperature and the pellet was resuspended in 0.1 M MOPS, pH-6.5. Five hundred microliters of a bacterial cell suspension (A_{600} of 1.0) were added to each T25 flask containing 10 mL of 4a3B cells three days after passaging and were further incubated at 28 °C for 24 or 72 h. Conditioned medium was collected from three different flasks and lysozyme activity was monitored from each of three flasks by turbidometric assays.

2.5. Purification and identification of Lys c-1

Frozen conditioned medium was thawed at room temperature. The pH of the medium was raised to 8.0 with 0.1 N NaOH without any loss of lysozyme activity. Changing the pH of the conditioned medium caused the medium to become turbid. The precipitates were removed by filtration through 0.8 μ m membrane filters. Clean filtrate was allowed to pass through an ion exchange column (1 \times 10 cm) packed with CM C-25 Sephadex beads pre-equilibrated with 0.1 M Tris-Cl pH-8.0 containing 0.15 M NaCl. The column was washed with 20 mL of 0.1 M Tris-Cl pH 8.0, 0.15 M NaCl. Lysozyme activity was eluted with 0.6 M NaCl in 0.1 M Tris-Cl pH 8.0. The fractions with highest lysozyme activity were pooled and 300 μ L of this mixed sample was loaded onto a pre-equilibrated Sephadex G-75 (Sigma) gel filtration column (1 \times 20 cm). Lysozyme was eluted with 0.1 M Tris-Cl, pH 8.0 containing 0.15 M NaCl and 0.01 % Triton X-100. Each fraction was assessed for lysozyme activity via turbidity assays and for purity on SDS-PAGE. Protein concentration was measured by Biorad's protein assay reagent (Biorad) and/or EZQ™ Protein Quantitation Kit (Invitrogen). About 200 μ L (0.01 mg/mL) of the purified sample was loaded on a 12.5% preparative polyacrylamide gel (NuPage, Invitrogen, USA). The protein band at the molecular mass of 14 kDa was excised from the gel using a sterile scalpel blade and processed for MS analysis. In gel digestion and mass spectrometric analysis was done at the Mass Spectrometry Facility (Biotechnology Center, University of Wisconsin-Madison). The digestion was performed as outlined on the website: <http://www.biotech.wisc.edu/ServicesResearch/MassSpec/ingel.htm>. The digested peptides were subjected to Matrix-Assisted Laser Desorption/Ionization-Time of Flight-Time of Flight (MALDI TOF-TOF) mass spectrometer (Applied Biosystems Inc., Foster City, CA). The peptide fingerprint was subjected to subsequent tandem MS analysis. Raw data was deconvoluted using GPS Explorer™ software (Applied Biosystems) and submitted for peptide mapping and MS/MS ion search analysis against non-redundant NCBI database with an in-house licensed Mascot search engine (Matrix Science, London, UK).

2.6. dsRNA mediated silencing of Lys c-1 and bacterial infections in the KD mosquitoes

An *in vitro* transcription template was produced using a one-step PCR protocol. The primers used to prepare the templates for *LYSC-1* and *GFP* contains T7 sequence on both forward and reverse primers at the 5'-end. The primer sequences are: (Fwd 5'-TAATACGACTCACTATAGGGATGAAAGTGTTCACAGTTTTG-3', Rev 5'-TAATACGACTCACTATAGGGAAAAACAGGAGCTAACATTCGG-3'). These primers were used to amplify a product from mosquito cDNA. The unrelated *GFP* sequences (Fwd: 5'-TAATACGACTCACTATAGGGCGTGATCAAGCCCGACA-3', Rev: 5'-TAATACGACTCACTATAGGGCTTCGGCGTGCTC-3') were used to amplify a product from *phMGFP* vector (Promega). The PCR product was purified using a QIAquick gel extraction kit (QIAGEN) and 1-2 μ g of the product were used as template for transcription. The MEGAscript™ RNAi kit (Ambion, Austin, Texas) was used for transcription and the production of dsRNA following the instruction manual. All dsRNA preparations were quantified by measuring absorbance at 260 nm, checked for integrity on an agarose gel, and stored at -80 °C.

One day old *An. gambiae* G3 females were injected with 1.4 μg *dsLYSC-1* RNA or *dsGFP* RNA (as control). After 24 h the mosquitoes were injected with 0.1 μL live *E. coli* ($A_{600}=0.004$) or *M. luteus* ($A_{600}=0.4$). The survival rate of the mosquitoes was monitored for seven days and specificity and efficiency of the knockdown were examined using RT-PCR. Survival of the mosquitoes after infection with bacteria were analyzed using Log-rank and Gehan-Breslow-Wilcoxon tests (GraphPad Software Inc., CA, USA). At least 45 mosquitoes were alive at the start of the monitoring period (beginning 24 h after bacterial challenge) for each experiment. A subset of mosquitoes were also examined for melanotic nodules following *M. luteus* inoculation. The results of two independent experiments are presented. Expression of *LYSC-2*, *LYSC-8*, defensin (*DEF1*) and cecropin (*CECI*) was also monitored in bacteria challenged mosquitoes after *LYSC-1* KD using RT-PCR.

2.7. Real time quantitative (QPCR) and semiquantitative RT-PCR

Real time PCR was used to compare relative levels of *LYSC-1* transcripts in the cell line before and after treatment with heat killed *M. luteus*. The protocol used for real time PCR has been described previously (Paskewitz et al., 2008). Briefly, total RNA was isolated from 1 mL cells ($\sim 1.5 \times 10^6$ cells/mL) using MasterPure™ Complete DNA and RNA Purification Kit (Epicentre Biotechnologies, Madison, WI, USA) according to the supplied instructions. Total RNA was treated with RQ1 DNase (Promega) at concentration of 1 unit/ μg RNA preparation to remove genomic DNA. RNA concentration was measured with micro-spectrophotometry (Nanodrop NT1000, Thermo Fisher Scientific, Waltham, MA). RNA quality was assessed on an Agilent 2100 bioanalyzer (Agilent Technologies, Inc. Santa Clara, CA, USA) using the Agilent RNA 6000 Nano Kit. Total RNA was also checked for genomic DNA contamination with each set of primers used and across all conditions/replicates. Threshold cycle (Ct) values above 32 in the QPCR amplification were considered free of significant genomic DNA contamination. Five hundred nanograms of total RNA were used to synthesize cDNA using a High-Capacity cDNA Archive Kit (Applied Biosystems). The resulting cDNA was stored at -20°C .

Primers based on *An. gambiae* cDNA sequences for internal reference genes (ribosomal protein *S7* (Salazar et al., 1993) actin and *RPL31*) and *LYSC-1* were designed using the Beacon designer software (Premier Biosoft International, Palo Alto, CA, USA). The primers were synthesized by IDT Technologies (Coralville, IA, USA). The treatments are compared to the calibrator as control cells (minus *M. luteus*) at zero hour time point versus 24 and 72 h post challenge with *M. luteus*.

Semiquantitative RT-PCR was performed to analyze transcript abundance in adult female mosquitoes following RNAi and bacterial challenge. Total RNA was isolated from five mosquitoes from respective treatments as described above. One hundred nanograms of total RNA were used to synthesize cDNA using a High-Capacity cDNA Archive Kit. The resulting cDNA was stored at -20°C until use. Primers based on *An. gambiae* cDNA sequences for immune effector genes *LYSC-2*, *LYSC-8*, cecropin (*CECI*) and defensin (*DEF1*) were designed using the Beacon designer software; other primers were as described above. Sources and sequences of gene specific primer sets are given in Table 1. Final concentrations of reagents used for RT-PCR were 1X buffer, 200 μM dNTPs, 0.5 μM each primer, 0.2 μL (4 units) Advantage Taq polymerase (Clontech Laboratories Inc., CA, USA). PCR cycle conditions were: an initial denaturation at 95°C for 3 min; then repeated cycles of 95°C for 30 s, 60°C for 30 s, 72°C for 30 s; and a final extension step at 72°C for 10 min. For *LYSC-1*, *LYSC-2*, *LYSC-8*, *CECI* and *DEF1*, 35 cycles were used while 25 cycles were used for *S7*.

2.8. Lysozyme activity assays

2.8.1. Turbidometric assay—For turbidometric assays, 30 μ L of the protein (0.01 mg/mL) sample was incubated with 970 μ L of *M. luteus* cells (0.3 mg/mL of purchased *Micrococcus lysodeikticus* cells; Sigma) prepared in 0.1 M MOPS, pH 6.5. Decreases in turbidity were monitored at 475 nm over a period of 60 s using a preset program in a Genesys 5 Spectrophotometer (Spectronic Instruments Inc. Rochester, NY, USA). For simplicity the absorbance values versus time were plotted directly.

2.8.2. Zonal inhibition assay—The zonal inhibition assay was carried out as described previously (Nasr et al., 2003). A solution of 1% agarose was prepared in 0.1 M sodium phosphate buffer pH 6.5 containing 0.05 M NaCl. The melted agarose was mixed with lyophilized powder of *M. luteus* cells to give a final concentration of 0.5 mg/mL cells and then poured in sterile petri-plates. After solidification, 1.5 mm wells were made and 10 μ L (0.01 mg/mL) of test samples were added to the wells. Plates were incubated at 37°C overnight. Buffer served as negative control. In addition to lyophilized *M. luteus*, Lys c-1 activity was tested against growing bacteria isolated from mosquitoes as well as representative non-mosquito derived Gram-positive and Gram-negative bacteria. Each bacterium was grown on TSA (Tryptic Soy Broth 30g/L containing 1.5% agar) plates. Wells (1.5 mm in diameter) were punched in the agar using sterile plastic tip and 10 μ L (of 0.01 mg/mL) of the purified lysozyme preparation was applied to wells. Plates were incubated at 30 °C for overnight and the zone of clearance was measured.

3. Results

3.1. Lysozyme activity in the 4a3B cell line following bacterial challenge

The abundance of *LYSC-1* transcripts was measured by real time quantitative PCR before and after treatment of 4a3B cells with heat-killed *M. luteus* (Fig. 1A). Expression of the *LYSC-1* transcripts increased dramatically (15.8 fold; average of three replicates) 24 h after treatment in three replicate experiments and remained significantly elevated at 72 h post treatment (10 fold).

To monitor muramidase activity in conditioned medium from the 4a3B cell line, turbidity assays were performed. Results (Fig. 1B) indicated that lysozyme activity increases in conditioned medium for a prolonged period following exposure to heat-killed *M. luteus* and remains elevated through 120 h post challenge. Correspondence of this activity with elevated levels of Lys c-1 (rather than another of the *An. gambiae* lysozymes) in the conditioned medium 24 h post challenge was confirmed by Western blotting with anti-Lys-c-1 antibodies (Fig. 1C).

3.2. Purification and identification of lysozyme c-1 from the conditioned medium

Lysozyme was purified from the cell culture medium of 4a3B cell line using a two step purification protocol. Almost all of the lysozyme activity could be recovered following the first ion exchange step (Fig. 2A). Gel filtration on Sephadex-G75 of fractions exhibiting the highest lysozyme activity (Fig. 2B) from the ion exchange step yielded one major purified band of about 14 kDa and a very faint band (probably in the femtograms range) running below the lysozyme band (Fig. 2C). We confirmed this upper protein as Lys c-1 (GenBank accession no. [AAC47326](#)) by ESI-MS and MS/MS (Biotech Center, UW Madison). The lower protein band could not be identified as even a 10X concentrated protein sample was below the detection limit of the MS instrument.

3.3. Zonal inhibition assays against midgut-derived bacterial isolates and other bacteria

Because several lysozymes have been reported as active against Gram-negative bacteria (Abraham et al., 1995; Ibrahim et al., 2001), we tested the mosquito enzyme for its activity against Gram-negative as well as Gram-positive bacteria. For these assays, we chose bacteria that were isolated from the midguts of laboratory colonies *An. gambiae* and *An. stephensi*, as well as other bacteria from various sources. The respective bacteria were grown on tryptic soy broth plates and the purified enzyme was applied to the wells. The results are described in Table 2. We tested antibacterial activity in two samples, one from the peak fraction of the ion exchange step and second in the purified fraction after second step of gel filtration. The antibacterial activity was higher in the partially purified fraction of the first ion exchange step as compared to the pure Lys c-1 indicating that additional antibacterial proteins are secreted in the conditioned medium that exert antibacterial effect in concert. In this partially purified fraction, the growth of most bacteria was inhibited except mosquito-derived *Elizabethkingia meningoseptica*, *Asaia krungthepensis*, non-mosquito derived *Asaia sp.*, *Serratia marcescens* and *Pseudomonas aeruginosa* (data not shown). By contrast, purified Lys c-1 was strongly active only against *M. luteus* and *Bacillus subtilis*. It had trace activity against two mosquito-derived bacteria (GS-3 and GS-4; Table 2). Unexpectedly, Lys c-1 was not active against a mosquito-derived strain of *M. luteus* from *An. gambiae* G3.

3.4. Lysozyme c-1 knockdown reduces the survival of mosquitoes upon infection with bacteria

Since Lys c-1 did not directly kill many bacteria but is strongly induced by bacteria in hemolymph from whole insects (Fig 3A) and in 4a3B cells (Fig. 1), we tested whether gene knockdown would affect survival after exposure to mildly pathogenic bacterial strains. RT-PCR results documented efficient KD of *LYSC-1* that persisted to at least seven days after injection of *E. coli*. The survival of mosquitoes injected with *dsLYSC-1* was compared to control *dsGFP* injected mosquitoes in two independent experiments using Log-Rank and Gehan-Breslow-Wilcoxon test of survival curves. Silencing of *LYSC-1* caused significant mortality in the mosquitoes infected with *E. coli* as compared to a control group ($p < 0.0001$ experiment 1 and $p < 0.005$ in experiment 2; Fig. 3B and C). The majority of the mortality was concentrated in the first 3 days after administration of the bacteria.

M. luteus infection in the *LYSC-1* KD mosquitoes did not have any significant effects on their survival (Fig. 4A). These bacteria are rapidly melanized in *A. gambiae* and *LYSC-1* KD did not alter that response. However, in contrast to the results for *E. coli*, *LYSC-1* transcripts re-appeared 96 h after *M. luteus* injection of KD mosquitoes (Fig. 4B). This result was repeated in three independent experiments, although the timing of re-appearance of the transcript varied somewhat.

3.5. Effects of LYSC-1 knockdown on transcript abundance of defensin, cecropin, LYSC-2 and LYSC-8 in mosquitoes

Since Lys c-1 does not kill bacteria directly but does enhance survival of the insects after bacterial exposure, we hypothesized that this protein might produce small peptidoglycan fragments which are needed to upregulate signaling cascades resulting in the production of antimicrobial peptides. We measured transcript abundance for *LYSC-2*, *LYSC-8*, *DEF1* and *CEC1* following *LYSC-1* knockdown and after the introduction of *E. coli*. These four genes were selected because they are all induced by bacterial inoculation. This was compared with *dsGFP* mosquitoes that were challenged with bacteria and with *dsLYSC-1* mosquitoes that were not challenged. We found no evidence that *LYSC-1* knockdown affected transcript abundance for the other immune genes after bacterial injection (Fig. 5).

4. Discussion

Expression of eight lysozyme genes in *An. gambiae* has been documented (Li et al., 2005; Kang et al., 1996) but the functional roles of these proteins are unknown. Transcripts of several lysozymes were more abundant in adult *An. gambiae* mosquitoes following infection with bacteria or malaria parasites (Dong et al., 2006; Li et al., 2005). *LYSC-1* and *LYSC-2* transcript abundance increased when adult female mosquitoes were challenged with *E. coli* or *M. luteus* (Li et al., 2005) which suggested a possible role for these proteins in immunity. Upregulation of *LYSC-1* was even more striking in the *An. gambiae* 4a3B cell line after bacterial challenge and we used this finding to further investigate the activity of this enzyme and its role in mosquito immunity.

Lysozymes are defined by their ability to cleave peptidoglycan in the cell wall of bacteria and this generally limits direct growth inhibition to Gram-positive bacteria (Nakimbugwe et al., 2006) although there are a growing number of exceptions (Clarke and DuPont, 1992; Yu et al., 2002; Gandhe et al., 2007). We investigated whether Lys c-1 can affect the growth of mosquito midgut-derived bacteria since the midgut is likely to be the most common site of entry in natural settings. Most of the bacteria identified in this study have been previously reported from mosquitoes, including some insects that had been field-collected (Straif et al., 1998; Gonzalez-Ceron et al., 2003; Favia et al., 2007; Lindh et al., 2008; Dong et al., 2009). Nine out of eleven mosquito-derived bacteria belonged to the Gram-negative group suggesting that the midgut environment is favorable for this type of bacteria. Gram-negative bacteria are generally considered resistant to lysozyme, due to the outer layer of lipopolysaccharide and protein which shields the peptidoglycan layer (Masschalck and Michiels, 2003). Secretion of lysozyme inhibitors also has been proposed as a resistance mechanism of several bacteria (Masschalck and Michiels, 2003). In keeping with these strong defense mechanisms, purified Lys c-1 did not show growth inhibitory attributes against most bacteria when cultured under neutral conditions *in vitro*. Perhaps most surprisingly, no activity was detected against a mosquito-derived *M. luteus* strain (Table 2) suggesting that this Gram positive strain may exhibit unique adaptations for life in the mosquito gut. Lys c-1 is expressed in the midgut and in the salivary glands so resident gut bacteria are likely to be exposed to this protein. Purified Lys c-1 did exhibit activity against non-mosquito derived strains of *M. luteus* and *Bacillus subtilis*, against a commercially available preparation of lyophilized *M. luteus*, and against two mosquito-derived bacteria, *Klebsiella oxytoca* (GS-4) and GS-3 (most closely related to an uncultured bacterium from an ant lion (Dunn and Stabb, 2005).

Although purified Lys c-1 from the 4a3B cell line did not show significant growth inhibitory activity against most mosquito-derived bacteria, the protein increased in conditioned media after exposure to heat-killed bacteria and in hemolymph from adult insects after injection of live *M. luteus*. An increase in c-type lysozyme gene expression following immune challenge has been reported for several Lepidoptera (Fujimoto et al., 2001; Lee & Brey, 1995; Sun et al., 1991; Mulnix and Dunn, 1994) and for adult mosquitoes (Li et al., 2005) and cultured cells from *Aedes aegypti* or *A. albopictus* (Gao & Fallon, 2000; Hernandez et al., 2003). The strong increase in Lys c-1 activity after immune challenge with Gram-negative or Gram-positive bacteria is intriguing since Lys c-1 can directly kill only a small number of Gram-positive bacterial species *in vitro*. To verify that Lys c-1 was important for mosquito survival after bacterial challenge, we silenced the gene through RNA interference. The impact of Lys c-1 depletion varied by bacterial species demonstrating specificity in the response to the two bacteria. Silencing of *LYSC-1* in adult mosquitoes reduced their survival upon subsequent infection with *E. coli*. In the absence of *LYSC-1*, the mosquitoes died at a rapid rate during the first three days after infection but thereafter the mortality curve plateaued. By contrast, no significant mortality was observed after *M. luteus* infection in the KD mosquitoes although this bacterium is susceptible to Lys c-1 activity *in vitro*. Redundancy in antibacterial

mechanisms could account for this observation. *M. luteus* is melanized within the hemocoel within a few minutes or hours after introduction. This response appears to be quite effective since *M. luteus* is rapidly cleared from the insect, unlike *E. coli* which persists for many days after infection (Gorman and Paskewitz, 2000). In the beetle, *Tenebrio molitor*, melanization is regulated by lysozyme which is necessary for the generation of peptidoglycan fragments that are bound by a peptidoglycan recognition protein (Park et al. 2007). However, knockdown of *A. gambiae* *LYSC-1* did not affect rapid melanization of *M. luteus*.

We also considered the possibility that these effects (plateau and failure to affect survival after inoculation with *M. luteus*) resulted from lack of persistence of the silencing phenotype. When we examined mosquitoes that had been silenced and then injected with *E. coli*, no transcripts were detected at any time out to 7 days after the bacterial inoculation. However, *LYSC-1* transcripts reappeared 48-96 h after *M. luteus* infection. This result indicates that silencing is not systemic and that *LYSC-1* may be expressed in a unique subset of cells only after *M. luteus* introduction. Since extensive melanization of this bacteria occurs along the dorsal vessel (unpub. obs.) and in hemocytes (Baton et al. 2009), we hypothesize that the generation of replacements for the hemocytes involved in this reaction occurs and that these new cells now express Lys c-1. Finally, we note that studies that rely on gene silencing to examine functional roles of proteins over extended timeframes will need to incorporate controls to verify the persistence of silencing during the entire observational period.

Since Lys c-1 did not directly kill *E. coli* in *in vitro* assays but did affect survival after exposure to this organism, we considered three other possible roles for the enzyme. First, Lys c-1 might produce peptidoglycan fragments that could activate transcription pathways that result in the production of antimicrobial peptides, in a manner similar to that identified for Gram positive bacteria and the TOLL pathway by Park and colleagues (2007). Second, Lys c-1 might assist in the removal of peptidoglycan, thereby downregulating an antimicrobial response that is likely to be costly to the insect. Finally, this lysozyme might act synergistically with other antimicrobial mechanisms to destroy bacteria.

We tested the hypothesis that the loss of Lys c-1 might affect the expression of other antimicrobial proteins which are responsible for clearing bacteria. Bacterially-induced upregulation of AMP (antimicrobial peptide) gene expression might not occur if a failure to produce small peptidoglycan fragments resulted in inefficient activation of the antibacterial peptide signaling cascades. Conversely, AMP gene expression might be hyper-elevated if most peptidoglycan fragments were not rendered non-immunogenic. In *Drosophila*, this latter function is attributed to catalytic peptidoglycan recognition proteins (PGRPs) that hydrolyze amide bonds between N-acetylmuramic acid and L-alanine moieties of PGN (peptidoglycan), thereby “scavenging” PGN and downregulating AMP transcription pathways (Mellroth et al., 2003; Zaidman et al., 2006). We examined transcript levels of *CEC1*, *DEF1* or *LYSC-2/8* in *LYSC-1* KD and *E. coli* challenged adult female mosquitoes. However, no significant differences in transcript levels were noted in comparison with control insects following introduction of *E. coli*.

It remains possible that Lys c-1 acts synergistically with other antimicrobial mechanisms, and/or affects key immune functions. Chalk et al. (1995) reported synergy between a chicken lysozyme and insect cecropin (but not defensin) at physiological concentrations against *E. coli*. Human peptidoglycan recognition proteins (i.e. PGLYRP) have been documented to act synergistically against both Gram-positive and Gram-negative bacteria in conjunction with lysozyme and lysostaphin (Lu et al., 2006; Cho et al., 2005; Wang et al., 2007). This synergism is likely to occur *in vivo* because both PGLYRPs and lysozyme are found within granules in the most abundant phagocyte, the polymorphonuclear leukocyte (Lu et al., 2006; Wang et al., 2007). Phagocytosis by granulocytic hemocytes is important for clearance of bacteria in insects

(Strand, 2008; Hillyer et al., 2003; 2004) and some insect PGRPs as well as Lys c-1 (Castillo et al. 2006) are produced by hemocytes. Therefore, it is possible that Lys c-1 depletion may cause impairment in removal of bacterial loads through synergy with other antibacterial peptides in the hemolymph plasma or following phagocytosis by granulocytes.

In conclusion we have isolated and purified a lysozyme corresponding to Lys c-1 from a mosquito-derived cell line 4a3B. Lys c-1 had limited direct effects on the growth of culturable bacteria isolated from the community in mosquito midguts. However its depletion through RNAi led to increased mortality in adult mosquitoes following challenge with Gram-negative bacteria. We found this system a useful tool to produce, isolate and characterize an important but neglected immune protein *in vitro* given the challenge in isolating it from whole mosquitoes in sufficient amounts for biochemical studies. Future studies will investigate the kinetics of Lys c-1 against specific targets under various conditions. Combining the cell culture system and selective gene inhibition by RNA interference will allow further investigation of the role of lysozyme c-1 in antibacterial immune responses.

Acknowledgments

We thank Dr. Nichole Broderick and Jo Handelsman for sharing protocol for bacteria isolation experiments from mosquitoes and for providing bacterial strains; Dr. John Lindquist for supplying bacteria; Erica Yashiro for helping with sequence analysis; Heather Sage and Ellie Walker for immune challenge to cell lines; Patrick Irwin for statistical analyses and Beth Schadd and Eric J. Shelley for rearing mosquitoes used in these experiments. This work was supported by NIH grant RO1 AI 037083 to SMP.

References

- Abraham EG, Nagaraju J, Salunke D, Gupta HM, Dalta RK. Purification and partial characterization of an induced antibacterial protein in the silkworm, *Bombyx mori*. *J. Invertebr. Pathol* 1995;65:17–24. [PubMed: 7876591]
- Azambuja P, Garcia ES, Ratcliffe NA. Gut microbiota and parasite transmission by insect vectors. *Trends Parasitol* 2005;21:568–572. [PubMed: 16226491]
- Boman HG, Hultmark D. Cell-free immunity in insects. *Ann. Rev. Microbiol* 1987;41:103–126. [PubMed: 3318666]
- Boman HG. Antibacterial peptides: key components needed in immunity. *Cell* 1991;65:205–207. [PubMed: 2015623]
- Castillo JC, Robertson AE, Strand MR. Characterization of hemocytes from the mosquitoes *Aedes aegypti* and *Anopheles gambiae*. *Insect Biochem. Mol. Biol* 2006;36:891–903. [PubMed: 17098164]
- Chalk R, Albuquerque CMR, Ham PJ, Townson H. Full sequence and characterization of two insect defensins: immune peptides from the mosquito *Aedes aegypti*. *Proc. R. Soc. Lond. B* 1995;261:217–221.
- Cho JH, Fraser IP, Fukase K, Kusumoto Yukari, Fujimoto Shoichi, Stahl GL, Ezekowitz RAB. Human peptidoglycan recognition protein S is an effector of neutrophil-mediated innate immunity. *Blood* 2005;106:2551–2558. [PubMed: 15956276]
- Clark A, Dupont C. O-acetylated peptidoglycan: its occurrence, pathological significance and biosynthesis. *Can. J. Microbiol* 1992;38:85–91. [PubMed: 1521192]
- Daffre S, Kylsten P, Samakovlis C, Hultmark D. The lysozyme locus in *Drosophila melanogaster*: an expanded gene family adapted for expression in the digestive tract. *Mol. Gen. Genet* 1994;242:152–162. [PubMed: 8159165]
- Demaio J, Pumpuni CB, Kent M, Beier JC. The midgut bacterial flora of wild *Aedes triseriatus*, *Culex pipiens* and *Psorophora columbiae* mosquitoes. *Am. J. Trop. Med. Hyg* 1996;54:219–223. [PubMed: 8619452]
- Dziarski R. Peptidoglycan recognition proteins (PGRPs). *Mol. Immunol* 2004;40:877–886. [PubMed: 14698226]

- Dillon RJ, Dillon VM. The gut bacteria of insects: nonpathogenic interactions. *Annu. Rev. Entomol* 2004;49:71–92. [PubMed: 14651457]
- Dimopoulos G, Christophides GK, Meister S, Schultz J, White KP, Barillas-Mury C, Kafatos FC. Genome expression analysis of *Anopheles gambiae*: responses to injury, bacterial challenge, and malaria infection. *Proc. Natl. Acad. Sci. USA* 2002;99:8814–8819. [PubMed: 12077297]
- Dong Y, Aguilar R, Xi Z, Warr E, Mongin E, Dimopoulos G. *Anopheles gambiae* immune responses to human and rodent *Plasmodium* parasite species. *PLoS Pathog* 2006;2:e52. [PubMed: 16789837]
- Dong Y, Manfredini F, Dimopoulos G. Implication of the mosquito midgut microbiota in the defense against malaria parasites. *PLoS Pathog* 2009;5:e1000423. [PubMed: 19424427]
- Dunn PE. Biochemical aspects of insect immunology. *Ann. Rev. Entomol* 1986;31:321–339.
- Dunn PE, Bohnert TJ, Russell V. Regulation of antibacterial protein synthesis following infection and during metamorphosis of *Manduca sexta*. *Ann. N.Y. Acad. Sci* 1994;15:117–130. [PubMed: 8192326]
- Dunn AK, Stabb EV. Culture-independent characterization of the microbiota of the ant lion *Myrmeleon mobilis* (Neuroptera: Myrmeleontidae). *Appl Environ Microbiol* 2005;71:8784–8794. [PubMed: 16332874]
- Favia G, Ricci I, Damiani C, Raddadi N, Crotti E, Marzorati M, Rizzi A, Urso R, Brusetti L, Borin S, Mora D, Scuppa P, Pasqualini L, Clementi E, Genchi M, Corona S, Negri I, Grandi G, Alma A, Kramer L, Esposito F, Bandi C, Sacchi L, Daffonchio D. Bacteria of the genus *Asaia* stably associate with *Anopheles stephensi*, an Asian malarial mosquito vector. *Proc. Natl. Acad. Sci. USA* 2007;104:9047–9051. [PubMed: 17502606]
- Fujimoto S, Tsuda I, Kishimoto K, Yamano Y, Morishima I. Protein purification, cDNA cloning and gene expression of lysozyme from eri-silkworm, *Samia cynthia ricini*. *Comp. Biochem. Physiol. B* 2001;128:709–718. [PubMed: 11290453]
- Gandhe AS, Janardhan G, Nagaraju J. Immune upregulation of novel antibacterial proteins from silkworms (Lepidoptera) that resemble lysozymes but lack muramidase activity. *Insect Biochem. Mol. Biol* 2007;37:655–666. [PubMed: 17550822]
- Gao Y, Fallon AM. Immune activation upregulates lysozyme gene expression in *Aedes aegypti* mosquito cell culture. *Insect Biochem. Mol. Biol* 2000;9:553–558.
- Gonzalez-Ceron L, Santillan F, Rodriguez MH, Mendez D, Hernandez-Avila JE. Bacteria in midguts of field-collected *Anopheles albimanus* block *Plasmodium vivax* sporogonic development. *J. Med. Entomol* 2003;40:371–374. [PubMed: 12943119]
- Guan R, Roychowdhury A, Ember B, Kumar S, Boons GJ, Mariuzza RA. Structural basis for peptidoglycan binding by peptidoglycan recognition proteins. *Proc. Natl. Acad. Sci. USA* 2004;101:7168–7173.
- Gusmão DS, Santos AV, Marini DC, Russo-Ede S, Peixoto AM, Bacci-Júnior M, Berbert-Molina MA, Lemos FJ. First isolation of microorganisms from the gut diverticulum of *Aedes aegypti* (Diptera: Culicidae): new perspectives for an insect-bacteria association. *Mem. Inst. Oswaldo. Cruz* 2007;102:919–924. [PubMed: 18209929]
- Hernandez VP, Higgins L, Fallon AM. Characterization and cDNA cloning of an immune-induced lysozyme from cultured *Aedes albopictus* mosquito cells. *Dev. Comp. Immunol* 2003;27:11–20. [PubMed: 12477497]
- Hillyer JF, Schmidt SL, Christensen BM. Hemocyte-mediated phagocytosis and melanization in the mosquito *Armigeres subalbatus* following immune challenge by bacteria. *Cell Tissue Res* 2003;313:117–127. [PubMed: 12838409]
- Hillyer JF, Schmidt SL, Christensen BM. The antibacterial innate immune response by the mosquito *Aedes aegypti* is mediated by hemocytes and independent of Gram type and pathogenicity. *Microbes Infect* 2004;6:448–459. [PubMed: 15109959]
- Hoffmann JA. The immune response of *Drosophila*. *Nature* 2003;426:33–38. [PubMed: 14603309]
- Hultmark D. Insect lysozymes *EXS* 1996;75:87–102.
- Hultmark D, Steiner H, Rasmuson T, Boman HG. Insect immunity. Purification and properties of three inducible bactericidal proteins from hemolymph of immunized pupae of *Hyalophora cecropia*. *Eur. J. Biochem* 1980;106:7–16. [PubMed: 7341234]

- Ibrahim HR, Thomas U, Pellegrini A. A helix-loop-helix peptide at the upper lip of the active site cleft of lysozyme confers potent antimicrobial activity with membrane permeabilization action. *J. Biol. Chem* 2001;276:43767–43774. [PubMed: 11560930]
- Jollès J, Schoentgen F, Croizier G, Croizier L, Jollès P. Insect lysozymes from three species of Lepidoptera: their structural relatedness to the C (chicken) type lysozyme. *J. Mol. Evol* 1979;14:267–721. [PubMed: 537106]
- Jolles P, Jolles J. What is new in lysozyme research. Always a model system, today as yesterday. *Mol. Cell. Biochem* 1984;63:165–189. [PubMed: 6387440]
- Kang D, Romans P, Lee JY. Analysis of a lysozyme gene from the malaria vector mosquito, *Anopheles gambiae*. *Gene* 1996;174:239–244. [PubMed: 8890741]
- Lee WJ, Brey PT. Isolation and characterization of the lysozyme-encoding gene from the silkworm *Bombyx mori*. *Gene* 1995;161:199–203. [PubMed: 7665079]
- Lindh JM, Terenius O, Faye I. 16S rRNA gene-based identification of midgut bacteria from field-caught *Anopheles gambiae* sensu lato and *A. funestus* mosquitoes reveals new species related to known insect symbionts. *Appl. Environ. Microbiol* 2005;71:7217–7223. [PubMed: 16269761]
- Li B, Calvo E, Marinotti O, James AA, Paskewitz SM. Characterization of the c-type lysozyme gene family in *Anopheles gambiae*. *Gene* 2005;360:131–139. [PubMed: 16137842]
- Li B, Paskewitz SM. A role for lysozyme in melanization of Sephadex beads in *Anopheles gambiae*. *J. Insect. Physiol* 2006;52:936–942. [PubMed: 16876189]
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 2001;25:402–408. [PubMed: 11846609]
- Lu X, Wang M, Qi J, Wang H, Li X, Gupta D, Dziarski R. Peptidoglycan recognition proteins are a new class of human bactericidal proteins. *J. Biol. Chem* 2006;281:5895–5907. [PubMed: 16354652]
- Mai W, Hu C. cDNA cloning, expression and antibacterial activity of lysozyme C in the blue shrimp (*Litopenaeus stylirostris*). *Prog. Natural Sci* 2009;19:837–844.
- Markart P, Korfhagen TR, Weaver TE, Akinbi HT. Mouse lysozyme M is important in pulmonary host defense against *Klebsiella pneumoniae* infection. *Am. J. Respir. Crit. Care Med* 2004;169:454–458. [PubMed: 14617511]
- Masschalck B, Michiels CW. Antimicrobial properties of lysozyme in relation to foodborne vegetative bacteria. *Crit. Rev. Microbiol* 2003;29:191–214. [PubMed: 14582617]
- Mellroth P, Karlsson J, Steiner H. A scavenger functions for a *Drosophila* peptidoglycan recognition protein. *J. Biol. Chem* 2003;278:7059–7064. [PubMed: 12496260]
- Müller HM, Dimopoulos G, Blass C, Kafatos FC. A hemocyte-like cell line established from the malaria Vector *Anopheles gambiae* expresses six prophenoloxidase genes. *J. Biol. Chem* 1999;274:11727–11735. [PubMed: 10206988]
- Mulnix AB, Dunn PE. Structure and induction of a lysozyme gene from the tobacco hornworm, *Manduca sexta*. *Insect Biochem. Mol. Biol* 1994;24:271–281. [PubMed: 7517269]
- Nakimbugave D, Masschalck B, Atanassova M, Zewdie-Bosüher A, Michiels CW. Comparison of bactericidal activity of six lysozymes at atmospheric pressure and under high hydrostatic pressure. *Int. J. Food. Microbiol* 2006;108:355–363. [PubMed: 16487612]
- Nasr NM, Fallon AM. Detection of lysozyme-like enzymatic activity secreted by an immune-responsive mosquito cell line. *J. Invertebr. Pathol* 2003;82:162–166. [PubMed: 12676552]
- Park JW, Kim CH, Kim JH, Je BR, Roh KB, Kim SJ, Lee HH, Ryu JH, Lim JH, Oh BH, Lee WJ, Ha NC, Lee BL. Clustering of peptidoglycan recognition protein-SA is required for sensing lysine-type peptidoglycan in insects. *Proc. Natl. Acad. Sci. USA* 2007;104:6602–6607. [PubMed: 17409189]
- Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper-Excel-based tool using pairwise correlations. *Biotechnol. Lett* 2004;26:509–515. [PubMed: 15127793]
- Powning RF, Davison WJ. Studies on insect bacteriolytic enzymes. I. Lysozyme in haemolymph of *Galleria mellonella* and *Bombyx mori*. *Comp. Biochem. Physiol. B* 1973;45:669–686.
- Paskewitz SM, Reese-Stardy S, Gorman MJ. An easter-like serine protease from *Anopheles gambiae* exhibits changes in transcript abundance following immune challenge. *Insect Mol. Biol* 1999;8:329–337. [PubMed: 10469250]

- Paskewitz SM, Li B, Kajla MK. Cloning and molecular characterization of two invertebrate-type lysozymes from *Anopheles gambiae*. *Insect Mol Biol* 2008;17:217–225. [PubMed: 18397277]
- Royet J, Dziarski R. Peptidoglycan recognition proteins: pleiotropic sensors and effectors of antimicrobial defences. *Nat. Rev. Microbiol* 2007;5:264–277. [PubMed: 17363965]
- Salazar CE, Mills-Hamm D, Kumar V, Collins FH. Sequence of a cDNA from the mosquito *Anopheles gambiae* encoding a homologue of human ribosomal protein S7. *Nucleic Acids Res* 1993;21:4147. [PubMed: 8371989]
- Skerrett SJ. Lysozyme in pulmonary host defense. *Am. J. Resp. Crit. Care Med* 2004;169:435–436. [PubMed: 14766658]
- Sun SC, Asling B, Faye I. Organization and expression of the immunoresponsive lysozyme gene in the giant silk moth, *Hyalophora cecropia*. *J. Biol. Chem* 1991;266:6644–6649. [PubMed: 2007608]
- Straif SC, Mbogo CNM, Toure AM, Walker ED, Kaufman M, Toure YT, Beier JC. Midgut Bacteria in *Anopheles gambiae* and *An. funestus* (Diptera: Culicidae) from Kenya and Mali. *J. Med. Entomol* 1998;35:222–226. [PubMed: 9615538]
- Strand, MR. Insect hemocytes and their role in immunity. In: Beckage, NE., editor. *Insect Immunity*. Academic Press; San Diego, CA: 2008. p. 25-48.
- Ursic-Bedoya RJ, Mitzey AM, Obratsova M, Lowenberger C. Molecular cloning and transcriptional activation of lysozyme-encoding cDNAs in the mosquito *Aedes aegypti*. *Insect Mol. Biol* 2005;14:89–94. [PubMed: 15663778]
- Wang M, Liu LH, Wang S, Li X, Lu X, Gupta D, Dziarski R. Human peptidoglycan recognition proteins require zinc to kill both Gram-positive and Gram-negative bacteria and are synergistic with antibacterial peptides. *J. Immunol* 2007;178:3116–3125. [PubMed: 17312159]
- Yu KH, Kim KN, Lee JH, Lee HS, Kim SH, ChoKY-Nam MH, Lee HI. Comparative study on characteristics of lysozymes from the hemolymph of three lepidopteran larvae, *Galleria mellonella*, *Bombyx mori*, *Agrius convolvuli*. *Dev. Comp. Immunol* 2002;26:707–713. [PubMed: 12206834]
- Zaidman-Rémy A, Hervé M, Poidevin M, Pili-Floury S, Kim MS, Blanot D, Oh BH, Ueda R, Mengin-Lecreulx D, Lemaitre B. The *Drosophila* amidase PGRP-LB modulates the immune response to bacterial infection. *Immunity* 2006;24:463–473. [PubMed: 16618604]




Figure 1.

Lysozyme expression in the *Anopheles gambiae* 4a3B cell line. A. Quantitative PCR for *LYSC-1* expression in 4a3b cell line following exposure to heat killed *M. luteus*. 0 TP as Calibrator=Zero time point as calibrator. B. Muramidase activity in the conditioned medium of control and bacteria challenged cells. Lysozyme activity was assayed via turbidometric assay from 0 to 120 h in three replicates. Error bars represent standard deviation of the mean of three independent measurements. C. Western blot of conditioned media from 4a3b cell line before (lane 1) and 24 h after (lane 2) *M. luteus* induction.




Figure 2. Purification of lysozyme c-1 from conditioned medium of 4a3B cells. A. Ion-exchange chromatography of the crude cell culture medium. B. Gel filtration chromatography of peak fractions from A. on Sephadex G-75. C. Silver stained gel from ion-exchange peak and G-75 fractions; Lane IE- ion exchange peak, lanes marked 19-25 are the respective fractions from B. Fraction 22-24 were used for characterization of activity under varied conditions. Lysozyme band is seen at an apparent molecular mass of 14 kDa marked by an arrow.






Figure 3. Induction and silencing of lysozyme c-1 in adult mosquitoes. A. Western blot demonstrating induction of lysozyme c-1 in mosquito hemolymph following inoculation with *M. luteus* (C=control; hpi= hour post infection; M=molecular mass marker. B and C. Kaplan Meier survival curves of mosquitoes showing effect of *LYSC-1* silencing after *E. coli* injection in two independent experiments. One day old *An. gambiae* females were injected with 1.4 μ g *dsLYSC-1* or an equal concentration of *dsGFP* as control. After 24 h the mosquitoes were injected with *E. coli*. The mortality was monitored over seven days.

A



B

**Figure 4.**

Effect of *LYSC-1* KD on survival of mosquitoes after *M. luteus* injection. A. Mortality curve following *M. luteus* infection 24 h after *LYSC-1* KD. Results of four independent experiments are shown. Error bars depicts standard deviation. B. Semi-quantitative RT-PCR analysis of *LYSC-1* transcripts following injection of *M. luteus*.




Figure 5. Semiquantitative RT-PCR analysis of selective immunity gene transcripts in *LYSC-1* KD and *E. coli* injected mosquitoes over 72 h.

Table 1

Sources and sequences of primers used for semiquantitative RT-PCR and real time PCR.

Gene	GenBank Accession no.	Sense primer 5'-3'	Antisense primer 5'-3'
<i>LYSC-1</i> (RT-PCR)	DQ007317	ATGAAAGTGTTTCCACAGTTTTG	AAAAACAGGAGCTAACATTCCG
<i>LYSC-1</i> (real time PCR)	DQ007317	AATGTGAACTGGCAAAGG	AGTCGTTTGAACCGTAGC
<i>LYSC-2</i>	AY659929	CTACCATCAGAAAGCAAATCAG	TAGCCTCACCAACGGAAC
<i>LYSC-8</i>	DQ004402	AACGAGAGCCGATACGATAC	CGCACCAGTAGTAGTTGTTG
Cecropin (<i>CEC1</i>)	AF200686	CCAACAACAATGAACTTCTCC	CTGCTGCCTTGAACACTC
Defensin (<i>DEF1</i>)	X93562	AAGAGGGTGTGCCGTTC	GCCTGTGTGTAAACTATTATCC
<i>RPL31</i>	XP_320630	TGGACAGCATTGGGACGATTC	TCACCTGGCAACGAAGTAAAGC
<i>RPS7</i>	L20837	CGCTATGGTGTTCGGTTCC	TGCTGCAAACCTTCGGCTAT

Table 2

Muramidase activity of Lys c-1 against various bacteria in lysoplate assays. Mosquito derived bacteria were identified by BLAST search based on 16s rRNA sequences. Purified Lys c-1 (10 µL of 0.01 mg/mL) was incubated with various bacteria derived from female mosquitoes as well as stock bacteria.

Bacterial isolates from	GenBank accession no.	Identification (top matches) ⁺⁺	Sequence length/query coverage/identity%	Muramidase activity in lysoplate assays ^{**}	Gram-type	Purified Lys c-1
An. gambiae G3						
G3-1	<u>FJ816020</u>	<i>Elizabethkingia meningoseptica</i> *	1402/99/99	nd	Gram ⁻	nd
G3-3	<u>FJ816021</u>	<i>Asaia krunghepensis</i>	1361/99/99	nd	Gram ⁻	nd
G3-6	<u>FJ816022</u>	<i>Micrococcus luteus</i>	1387/99/100	nd	Gram ⁺	nd
An. gambiae 4a rr						
GS-1	<u>FJ816023</u>	<i>Erwinia sp.</i>	1403/100/100	nd	Gram ⁻	nd
GS-2	<u>FJ816024</u>	<i>Serratia sp.</i>	1350/99/99	nd	Gram ⁻	nd
GS-3	<u>FJ816025</u>	Uncultured bacterium	1385/100/99	5.5	Gram ⁺	5.5
GS-4	<u>FJ816026</u>	<i>Klebsiella oxytoca</i>	1400/100/99	tr	Gram ⁻	tr
GS-6	<u>FJ816027</u>	Uncultured bacterium	1411/100/99	nd	Gram ⁻	nd
GS-7	<u>FJ816028</u>	<i>Elizabethkingia meningoseptica</i> *	1374/100/99	nd	Gram ⁻	nd
An. Stephens						
ST-2	<u>FJ816029</u>	<i>Serratia sp.</i>	1335/100/99	nd	Gram ⁻	nd
ST-3	<u>FJ816030</u>	<i>Bacterium_spbL1type1-2</i>	1327/100/100	nd	Gram ⁻	nd
Non-mosquito derived bacteria						
<i>Micrococcus luteus</i>				6	Gram ⁺	6
<i>Serratia marcescens</i>				nd	Gram ⁻	nd
<i>Enterobacter cloacae</i>				nd	Gram ⁻	nd
<i>Staphylococcus epidermidis</i>				nd	Gram ⁺	nd
<i>Bacillus subtilis</i>				7	Gram ⁺	7
<i>Bacillus thuringiensis</i>				nd	Gram ⁺	nd
<i>Asaia sp.</i>				nd	Gram ⁻	nd
<i>Pseudomonas aeruginosa</i>				nd	Gram ⁻	nd
<i>Enterobacter sp.</i>				nd	Gram ⁻	nd
<i>Micrococcus lysodactylicus</i> (lyophilized)				14	Gram ⁺	14

Bacterial isolates from	GenBank accession no.	Identification (top matches) ⁺⁺	Sequence length/query coverage/identity%	Muramidase activity in lysoplate assays ^{**}
<i>Escherichia coli</i> DH5 α				Gram ⁻ nd

⁺⁺ Top score from NCBI BLAST database nr/nt; Query coverage is over the entire length of sequenced PCR product as indicated

^{*} also known as *Chryseobacterium meningosepticum*; Zone of clearance measured in mm; Measurements were made after O/N incubation of the bacteria with lysozyme sample; nd-absence of detectable zone of clearance; tr-trace zone of clearance (below 5 mm).