

Published in final edited form as:

J Am Chem Soc. 2009 December 23; 131(50): 18044–18045. doi:10.1021/ja908372b.

Unnatural amino acid substitution as a probe of the allosteric coupling pathway in a mycobacterial Cu(I) sensor

Zhen Ma^{‡,§}, Darin M. Cowart[¶], Brian P. Ward[‡], Randy J. Arnold[‡], Richard D. DiMarchi[‡], Limei Zhang[†], Graham N. George[†], Robert A. Scott[¶], and David P. Giedroc^{*,‡}


[‡]Department of Chemistry, Indiana University, Bloomington, IN 47405-7102 USA.

[§]Department of Biochemistry and Biophysics, Texas A&M University, College Station, TX 77843-2128 USA

[¶]Departments of Chemistry and of Biochemistry and Molecular Biology, University of Georgia, Athens, GA 30602 USA.

[†]Department of Geological Sciences, University of Saskatchewan, Saskatoon, S7N 5E2, Canada.

Abstract



The Cu(I) sensor *Mycobacterium tuberculosis* CsoR is a founding member of a new metalloregulatory protein family. Here we show that two "atom" substitutions of the Nε2 face of a Cu(I) coordinating histidine-61 allosterically uncouple Cu(I) and DNA binding, with no effect on Cu(I) binding affinity and coordination structure. A model analogous to the allosteric switch mechanism in *Staphylococcus aureus* CzrA, a zinc sensor protein with a completely different fold, is proposed.

Elucidation of structural and dynamical pathways of communication or coupling between ligand binding sites, *i.e.*, classical allostery, is of topical interest and fundamental importance in biological regulation,^{1–3} but remains a challenge to detect and quantify.^{4–6}

Metalloregulatory proteins are specialized allosteric proteins that control the intracellular availability of essential transition metal ions by binding a specific metal, which in turn,

giedroc@indiana.edu.

Supporting Information Available: Supplementary Methods, Tables (S1–S3) and Figures (S1–S9) are available free of charge via the Internet at <http://pubs.acs.org>.

activates or inhibits operator DNA binding, thus programming a defined transcriptional output.^{7,8} In this study, we target the "second coordination shell" of the Cu(I) sensor *M. tuberculosis* CsoR and successfully abrogate allosteric coupling of Cu(I) and DNA binding.

Mtb CsoR is a Cu(I) regulatory protein distantly related to the Ni(II)/Co(II) sensor *E. coli* RcnR,¹⁰ in which Cu(I)-binding allosterically decreases operator DNA binding affinity. This leads to transcriptional derepression of the copper-sensing operon (*cs0*) in response to Cu(I) stress.⁹ The Cu(I) ion in CsoR occupies a pocket at the protomer interface, where the S γ atoms of Cys36' and Cys65, and the N δ 1 atom of His61 form a trigonal coordination complex (Figure 1).⁹ The structure further reveals that the N ϵ 2 face of His61 points toward the side chains of two conserved "second coordination shell" residues, Tyr35 and Glu81 (Figure 1). This structure suggests that Tyr35, Glu81 and His61 form a hydrogen bonding network that stabilizes the allosterically inhibited Cu(I) bound state.^{9,11} Substitution of the Cu(I) ligand His61 with Ala changes the Cu(I) coordination number from three to two, resulting in a linear *bis*-thiolato Cu(I) complex.⁹ This substitution also greatly reduces the *cs0* operator binding affinity by $\geq 10,000$ -fold to undetectable levels, despite adopting a folded, α -helical tetrameric assembly state (Figure S1). Thus, His61 is a multifunctional residue, the distinct physicochemical roles of which cannot be directly probed by conventional mutagenesis.⁹ We therefore employed native chemical ligation to perform an "atom" substitution of His61 by replacement with N ϵ 2-methyl-histidine (MeH) or 4-(thiazolyl)-L-alanine (Thz) in order to directly test this allosteric coupling model (Figure 1). Both substitutions are expected to preserve the Cu(I) coordination and DNA binding properties of His61 while blocking the ability of the opposite face of His61 to donate hydrogen bonds to Tyr35' and/or Glu81.

The Cu(I) binding affinity ($\log K_{Cu}$) of wild-type *Mtb* CsoR was determined to be 18.0 (± 0.2) from both a conventional titration of Cu(I) into a mixture of CsoR and BCS (Figure S2), as well as an assay that employs varying BCS concentrations, rather than Cu(I) concentrations (Table 1 and Supplementary Methods).^{9,12,13} The coupling free energy (ΔG_c) was obtained from the ratio of the macroscopic CsoR DNA binding affinities in the presence (A_2^{Cu}) and absence (A_2^{apo}) of bound Cu(I) measured by fluorescence anisotropy⁷ (Supplementary Methods and Figures S3–S5). Cu(I) binding to wild-type CsoR significantly decreases the *cs0* operator DNA binding affinity, resulting in a ΔG_c of +3.6 (± 0.2) kcal/mol (pH 7.0, 0.2 M NaCl, 2 mM DTT, 25 °C) (Table 1), with no effect on the assembly state of the CsoR tetramer.¹³

A semisynthetic route using an intein fusion strategy was used to introduce MeH and Thz analogs of His61 (Figure 1 and Supplementary Methods). To increase the efficiency of such a strategy, L58C CsoR 1–106 was chosen as the parent molecule to incorporate both H61MeH and H61Thz substitutions by chemical ligation, since it showed no significant differences from wild-type CsoR in Cu(I) binding affinity (Table 1), Cu(I) coordination geometry (Figure 2 and Table S2), DNA binding affinity and negative allosteric regulation by Cu(I) (Table 1 and Figure S5). The integrity of the resulting H61MeH and H61Thz *Mtb* CsoRs was confirmed by trypsin digestion and tandem LC-MS/MS sequencing (Figure S6).

Both H61MeH and H61Thz *Mtb* CsoRs bind Cu(I) with an affinity comparable to the parent L58C CsoR 1–106 ($\log K_{Cu}=17.3$) (Table 1) and 100-1000-fold higher affinity than that of H61A CsoR, which forms a digonal S_2 complex.⁹ This suggests that H61MeH and H61Thz CsoRs form trigonal coordination complexes with Cu(I), a result confirmed by the Cu(I) near-edge region of the x-ray absorption spectrum (Figure 2, Table S2 and Figure S7). Quantitative analysis of the EXAFS spectrum of H61MeH CsoR reveals a coordination geometry that is identical to wild-type CsoR (Table S2).⁹ Both H61MeH and H61Thz CsoRs bind the *cs0* operator with near wild-type affinity in their apo-forms which reveals the semisynthetic proteins are folded correctly and likely form stable tetramers (Figure S5 and Table S1).¹³

Strikingly, the DNA binding affinities of both H61MeH and H61Thz CsoRs are not strongly regulated by Cu(I), with $\Delta G_c = 0 (\pm 0.4)$ and $0.5 (\pm 0.2)$ kcal mol⁻¹, respectively (Table 1). These findings clearly establish that the N ϵ 2 face of His61 is specifically required to link thermodynamically and functionally Cu(I) binding to DNA binding, with no significant influence on the magnitude of the Cu(I) binding affinity or coordination geometry.

To further test this structural linkage model (Figure 1), we targeted the other "second coordination shell" residues, Tyr35 and Glu81, using conventional site-directed mutagenesis. Y35F and all E81 mutant CsoRs retained wild-type Cu(I) binding affinity and trigonal S₂N Cu (I) coordination geometries by x-ray absorption spectroscopy (Table 1, Table S3 and Figure S8). Y35F CsoR is characterized by a 33% decrease in ΔG_c , while *cso* operator DNA binding by E81A CsoR is nearly refractory to Cu(I) regulation ($\Delta G_c = 0.6$ kcal mol⁻¹) (Table 1 and Figure S9). More conservative substitutions of Glu81, *i.e.*, with Gln, Asp or Asn, also give rise to significant decreases in ΔG_c , to about 50% of wild-type CsoR, while Cu(I) binding affinities remain unaffected (Table 1). Inspection of the magnitude of ΔG_c for a double mutant, Y35F/E81Q CsoR, reveals a near additivity of ΔG_c measured for the component single mutants (Table 1). Thus, these residues play important roles, but only weakly interact energetically with one another to drive allosteric negative regulation of CsoR by Cu(I). This suggests that hydrogen bonds from Tyr35 and Glu81 to His61, and not to each other (Figure 1), are most critical for metalloregulation in this system.

We propose that the heteroaromatic character of His61 contacts the DNA in a way that does not require the N ϵ 2 atom of the imidazole ring. Cu(I) coordination to the N δ 1 atom is necessary but not sufficient to drive allosteric switching, but critically, initiates formation of a hydrogen-bonding network involving the N ϵ 2 face of the same histidine, and the side chains of Y35 and E81, that ultimately leads to dissociation of the CsoR tetramer from the DNA. It is remarkable that opposite N δ 1 and N ϵ 2 faces of a key metal coordinating histidine residue, albeit one far from the DNA binding site,¹⁴ are proposed to play exactly analogous roles in the arsenic repressor (ArsR) family zinc sensor CzcA, despite the completely unrelated protein fold and different metal that is sensed.^{14–16} This speaks to the evolutionary generality of a mechanism of allosteric regulation that exploits the unique physicochemical features of a key histidine side chain to orchestrate cellular metal homeostasis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by grants from the National Institutes of Health to D. P. G. (GM042569) and R. A. S. (GM042025). Portions of this research were carried out at the Stanford Synchrotron Radiation Lightsource (SSRL). Work at the University of Saskatchewan was supported by NSERC (Canada), the CIHR and a CRC award (to G.N.G.). R.J.A. acknowledges support from the US National Institutes of Health National Center for Glycomics & Glycoproteomics (P41 RR018942) and Eksigent Technologies for providing the nanoLC used in this work.

References

1. Kuriyan J, Eisenberg D. *Nature* 2007;450:983–990. [PubMed: 18075577]
2. Swint-Kruse L, Matthews KS. *Curr. Opin. Microbiol* 2009;12:129–137. [PubMed: 19269243]
3. Popovych N, Tzeng SR, Tonelli M, Ebright RH, Kalodimos CG. *Proc. Natl. Acad. Sci* 2009;106:6927–6932. [PubMed: 19359484]
4. Goodey NM, Benkovic SJ. *Nat. Chem. Biol* 2008;4:474–482. [PubMed: 18641628]
5. Kern D, Zuiderweg ER. *Curr. Opin. Struct. Biol* 2003;13:748–757. [PubMed: 14675554]

6. Suel GM, Lockless SW, Wall MA, Ranganathan R. *Nat. Struct. Biol* 2003;10:59–69. [PubMed: 12483203]
7. Giedroc DP, Arunkumar AI. *Dalton Trans* 2007:3107–3120. [PubMed: 17637984]
8. Ma Z, Jacobsen FE, Giedroc DP. *Chem. Rev* 2009;109:4644–4681. [PubMed: 19788177]
9. Liu T, Ramesh A, Ma Z, Ward SK, Zhang L, George GN, Talaat AM, Sacchettini JC, Giedroc DP. *Nat. Chem. Biol* 2007;3:60–68. [PubMed: 17143269]
10. Iwig JS, Leitch S, Herbst RW, Maroney MJ, Chivers PT. *J. Am. Chem. Soc* 2008;130:7592–7606. [PubMed: 18505253]
11. Eiamphungporn W, Soonsanga S, Lee JW, Helmann JD. *Nucleic Acids Res* 2009;37:1174–1181. [PubMed: 19129220]
12. Xiao Z, Donnelly PS, Zimmermann M, Wedd AG. *Inorg. Chem* 2008;47:4338–4347. [PubMed: 18412332]
13. Ma Z, Cowart DM, Scott RA, Giedroc DP. *Biochemistry* 2009;48:3325–3334. [PubMed: 19249860]
14. Arunkumar AI, Campanello GC, Giedroc DP. *Proc. Natl. Acad. Sci* 2009;106:18177–18182. [PubMed: 19822742]
15. Eicken C, Pennella MA, Chen X, Koshlap KM, VanZile ML, Sacchettini JC, Giedroc DP. *J. Mol. Biol* 2003;333:683–695. [PubMed: 14568530]
16. Pennella MA, Arunkumar AI, Giedroc DP. *J. Mol. Biol* 2006;356:1124–1136. [PubMed: 16406068]




Figure 1.

Structure around the Cu(I) binding region of the *Mtb* CsoR dimer interface (protomers shaded *slate* and *salmon*)⁹ and space filling and chemical structures of histidine, Nε2-methyl-histidine (MeH) and 4-(thiazolyl)-L-alanine (Thz) used in this study. Note the Nδ1 nitrogen of His61 forms a Cu(I)-N coordination bond while the Nε2 face is within hydrogen bonding distance of Y35' (*salmon* protomer) and E81.




Figure 2.

Cu X-ray absorption edge (a), EXAFS (b, *inset*), and FT (b; $k = 2\text{--}13 \text{ \AA}^{-1}$, k^3 weighting) for WT 1-106 (*black dashed*), L58C (*black solid*), and H61MeH (*red dashed*) CsoRs. The best fit of H61MeH CsoR based on a $\text{CuS}_2(\text{imid})_1$ model is shown in *bold red* in (b). Fitting parameters are compiled in Table S2.

Table 1Cu(I) affinities and allosteric coupling free energies for various *Mtb* CsoRsa^a

<i>Mtb</i> CsoR	log K_{Cu} ^b	ΔG_c (kcal/mol) ^d
Wild-type	18.0 (± 0.2)	3.6 (± 0.2)
WT 1–106	17.3 (± 0.4)	3.0 (± 0.3)
L58C 1–106	17.3 (± 0.3)	3.0 (± 0.4)
H61MeH	18.4 (± 0.2)	0 (± 0.4)
H61Thz	16.7 (± 0.1)	0.5 (± 0.2)
H61A	14.5 (± 0.3)	n.d. ^e
	14.9 (± 0.4) ^c	
E81A	17.7 (± 0.4)	0.6 (± 0.3)
E81Q	17.6 (± 0.5)	1.9 (± 0.3)
E81D	17.8 (± 0.4)	1.5 (± 0.3)
E81N	17.3 (± 0.4)	1.6 (± 0.4)
Y35F	16.7 (± 0.4)	2.4 (± 0.2)
Y35F/E81Q	17.6 (± 0.3)	0.8 (± 0.3) ^f

^a Conditions: 10mM HEPES, 0.2 M NaCl, pH 7.0, 25 °C, with 2 mM DTT present only in the DNA binding experiments.^b Determined by competition with the chromophoric complex Cu^I(BCS)₂ (log $\beta_2=19.8$) or^c Cu^I(BCA)₂ (log $\beta_2=17.2$) as described in Supplementary Methods.^d Determined using fluorescence anisotropy-based titrations like those shown in Figure S5, with detailed fitting parameters compiled in Table S1.^e n.d., not detected, $A_2 \leq 10^{10} M^{-1}$ under these conditions.^f $\delta = \Delta \Delta G_c^{Y35F/E81Q} - (\Delta \Delta G_c^{Y35F} + \Delta \Delta G_c^{E81Q}) = +0.1$ kcal/mol where $\Delta \Delta G_c^{mutant} = \Delta G_c^{mutant} - \Delta G_c^{WT}$.