



Published in final edited form as:

Anesthesiology. 2009 December ; 111(6): 1271–1278. doi:10.1097/ALN.0b013e3181bf1d79.

## Propofol Facilitates Glutamatergic Transmission to Neurons of the Ventrolateral Preoptic Nucleus

Ke Y. Li, M.D., Ph.D<sup>1,\*</sup>, Yan-zhong Guan, MD, Ph.D<sup>1,\*</sup>, Kresimir Krnjević, MD, Ph.D<sup>2,†</sup>, and Jiang H. Ye, MD, MSc<sup>1,†</sup>

<sup>1</sup>Department of Anesthesiology, Pharmacology and Physiology, University of Medicine and Dentistry of New Jersey, New Jersey Medical School, Newark, New Jersey.

<sup>2</sup>Department of Physiology, McGill University, Montréal, QC, Canada

### Abstract

**Background**—There is much evidence that the sedative component of anesthesia is mediated by  $\gamma$ -aminobutyric acid A receptors on hypothalamic neurons responsible for arousal, notably in the tuberomammillary nucleus. These  $\gamma$ -aminobutyric acid A receptors are targeted by GABAergic neurons in the ventrolateral preoptic area (VLPO): when these neurons become active, they inhibit the arousal-producing nuclei and induce sleep. According to recent studies, propofol induces sedation by enhancing VLPO-induced synaptic inhibition, making the target cells more responsive to  $\gamma$ -aminobutyric acid A. We explored the possibility that propofol also promotes sedation less directly by facilitating excitatory inputs to the VLPO GABAergic neurons.

**Methods**—Spontaneous excitatory postsynaptic currents were recorded from VLPO cells – principally mechanically isolated, but also in slices from rats.

**Results**—In isolated VLPO GABAergic neurons, propofol increased the frequency of glutamatergic spontaneous excitatory postsynaptic currents without affecting their mean amplitude. Propofol's action was mimicked by muscimol and prevented by gabazine, respectively a specific agonist and antagonist at  $\gamma$ -aminobutyric acid type-A receptors. It was also suppressed by bumetanide, a blocker of  $\text{Na}^+\text{-K}^+\text{-Cl}^-$  cotransporter-mediated inward  $\text{Cl}^-$  transport. In slices, propofol also increased the frequency of spontaneous excitatory postsynaptic currents and, at low doses, accelerated firing of VLPO cells.

**Conclusion**—Propofol induces sedation, at least in part, by increasing firing of GABAergic neurons in ventrolateral preoptic area, indirectly by activation of  $\gamma$ -aminobutyric acid type-A receptors on glutamatergic afferents: because these axons/terminals have a relatively high internal  $\text{Cl}^-$  concentration, they are depolarized by GABAergic agents such as propofol which thus enhance glutamate release.

### Introduction

THE CELLULAR and molecular mechanisms underlying the effects of general anesthetics are not well understood. The sedative component of anesthesia appears to be mediated by type A

---

Corresponding author: Jiang Hong Ye, Tel: 973-972-1866, Fax: 973-972-4172 ye@umdnj.edu Department of Anesthesiology, UMDNJ, New Jersey Medical School.

\*postdoctoral fellows

†Professor

Parts of the contents have been presented at the Annual meeting, American Society of Anesthesiology, October 20, 2008, Orlando, FL.

Summary Statement: Propofol enhances glutamate release onto neurons of the ventrolateral preoptic area by potentiating the function of  $\gamma$ -aminobutyric acid A receptors on the glutamatergic afferents. These effects may contribute to its sedative action.

$\gamma$ -aminobutyric acid receptors (GABA<sub>A</sub>Rs) in an endogenous sleep pathway<sup>1,2</sup>, the relevant GABAergic neurons being located in the ventrolateral preoptic area (VLPO)<sup>3–8</sup>. Recently, Zecharia et al.<sup>2</sup> showed that GABAergic agents such as muscimol and propofol potentiate the GABA-mediated inhibition of orexinergic perifornical<sup>9</sup> and histaminergic tuberomammillary neurons<sup>6</sup>. Together with other monoaminergic and cholinergic projections to cortex, the neurons in tuberomammillary nuclei directly promote cortical arousal<sup>10,11</sup>; and all are activated by orexinergic neurons<sup>12</sup>. Thus, their inhibition by VLPO neurons is pivotal in the initiation of sleep<sup>13,14</sup>. Although the activity of VLPO neurons is strongly influenced by circadian fluctuations in input from suprachiasmatic cells<sup>15</sup>, the detailed mechanisms underlying activation of VLPO neurons are largely unexplored<sup>16</sup>.

General anesthetics are known to have marked effects on synaptic transmission. Most previous studies found that propofol enhances the function of inhibitory GABA<sub>A</sub>Rs<sup>17–20</sup> and may depress the release of glutamate, the major excitatory neurotransmitter<sup>21</sup>. However, in vitro electrophysiological studies have shown that, in several brain areas, activation of presynaptic GABA<sub>A</sub>Rs have a depolarizing effect which enhances glutamatergic transmission<sup>22–25</sup>. We hypothesized that by potentiating the depolarizing action of presynaptic GABA<sub>A</sub>Rs, propofol may increase glutamatergic transmission in the VLPO, and thus increase the activity of VLPO GABAergic neurons. To test this hypothesis, we recorded spontaneous excitatory postsynaptic currents (sEPSCs) in the rat VLPO GABAergic neurons, either in slices, or isolated by a mechanical, enzyme-free procedure. These neurons preserve some functional glutamate-releasing terminals after isolation.

## Materials and Methods

All experiments were performed in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals, and they were approved by the Institutional Animal Care and Use Committee of the University of Medicine and Dentistry of New Jersey (Newark, NJ). The experiments were done on brains from Sprague-Dawley rats.

### Slice Preparation and Mechanical Dissociation

The hypothalamic slices were prepared as described previously<sup>26,27</sup>. Briefly, Sprague–Dawley rats (aged 10–28 postnatal days) were anesthetized and then decapitated. Coronal slices (300  $\mu$ m thick) were cut using a VF-200 slicer (Precisionary Instruments Inc., Greenville, NC). They were prepared in an ice-cold glycerol-based artificial cerebrospinal fluid containing 250 mM glycerol, 1.6 mM KCl, 1.2 mM NaH<sub>2</sub>PO<sub>4</sub>, 1.2 mM MgCl<sub>2</sub>, 2.4 mM CaCl<sub>2</sub>, 25 mM NaHCO<sub>3</sub>, and 11 mM glucose, and saturated with 95% O<sub>2</sub>/5% CO<sub>2</sub> (carbogen)<sup>28</sup>. Slices were allowed to recover for at least 1 h in a holding chamber at 32°C in carbogen-saturated regular artificial cerebrospinal fluid, which has the same composition as glycerol-based artificial cerebrospinal fluid, except that glycerol was replaced by 125 mM NaCl.

Neurons with functional presynaptic terminals attached were obtained by mechanical dissociation (Fig. 1A), as previously described<sup>29</sup>, with some modifications<sup>28,30</sup>. A slice containing the VLPO was transferred to a 35 mm culture dish (Falcon, Rutherford, NJ) and held down with a flat U-shape wire. The dish was filled with standard external solution containing 140 mM NaCl, 5 mM KCl, 2 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub>, 10 mM HEPES, and 10 mM glucose (320 mOsm, pH adjusted to 7.3 with Tris base). Under an inverted microscope (Nikon, Tokyo, Japan), we identified VLPO by its stereotaxic coordinates<sup>31</sup>, and located the VLPO cells that project to the noradrenergic locus coeruleus and the histaminergic tuberomammillary nucleus<sup>3,6,32</sup>. A fire-polished pipette, held by a micromanipulator, lightly touched the surface of the VLPO region and vibrated horizontally at 15–20 Hz for 2–5 min. After 20 min, isolated neurons that adhered to the bottom of the dish were kept for electrophysiological recordings at room temperature (21–22 °C). These mechanically

dissociated neurons often preserve functional nerve terminals, including some that release glutamate<sup>29,30</sup>.

### Electrophysiological Recordings

Electrical signals were obtained in whole-cell configurations with an Axon 200B amplifier or a MultiClamp 700A (Molecular Devices Co., Union City, CA), a Digidata 1320A A/D converter (Molecular Devices Co.), and pCLAMP 9.2 software (Molecular Devices Co.). Data were filtered at 1 kHz and sampled at 5 kHz. The patch electrodes had a resistance of 2–5 M $\Omega$  when filled with the pipette solution containing: 135 mM CsF, 5 mM KCl, 2 mM MgCl<sub>2</sub>, 10 mM HEPES, 2 mM Mg Adenosine-5'-triphosphate, and 0.2 mM Guanosine-5'-triphosphate (for voltage clamp). For current clamp recordings, the CsF in the above pipette solution was replaced by K-gluconate. The pH of the pipette solution was adjusted to 7.2 with Tris base. Neurons were voltage clamped at –60 mV to record  $\alpha$ -amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptor-mediated spontaneous and miniature excitatory postsynaptic currents (sEPSCs and mEPSCs).

In several experiments, a single coronal slice was transferred into a 0.4 ml recording chamber where it was held down by a platinum ring. Warm carbogenated artificial cerebrospinal fluid flowed through the bath (1.5–2.0 ml/min). Under infrared video microscopy (E600FN; Nikon), we recorded from neurons located in the core of VLPO, where a cluster of sleep-active neurons were originally identified, as illustrated in Fig. 1<sup>5,6</sup>. Cells were deemed of the sleep promoting type according to previously established criteria<sup>3</sup>: first, they were triangular in shape (Fig. 1A1), which were in contrast to the fusiform, non-low-threshold spike (non-LTS) type (Fig. 1A2); second, when recorded in current-clamp mode, they generated characteristic low-threshold spikes in response to a depolarizing pulse or after a hyperpolarizing current step (Fig. 1B1, C) (Fig. 1B2 shows the high-threshold spikes of non-LTS cells); and finally, they were inhibited by noradrenaline (Fig. 1D1). Triangular-shaped cells inhibited by noradrenaline always displayed low-threshold spikes. By contrast, noradrenaline excited a non-LTS neuron (Fig. 1D2). The series resistance (15–30 M $\Omega$ ) or input resistance (300–500 M $\Omega$ ) was monitored throughout the whole-cell recording, and data were discarded if the resistance changed by more than 20%. All these recordings were made at 32°C, maintained by an automatic temperature controller (Warner Instruments, Hamden, CT).

### Chemicals and Applications

The chemicals, including 2, 6-diisopropylphenol (propofol), gabazine (SR 95531), 2,3-dihydroxy-6,7-dinitroquinoxaline, tetrodotoxin, bicuculline and bumetanide were obtained from Sigma-Aldrich Chemical Company (St Louis, MO). Drugs were added to the superfusate or applied to a cell by a fast perfusion system (Y tube). Solutions in the vicinity of a neuron could be completely exchanged within 40 ms without damaging the seal<sup>33</sup>.

### Statistical Analysis

sEPSCs and mEPSCs were counted and analyzed with Clampfit 9.2 (Molecular Devices Co.); sEPSCs and mEPSCs were screened automatically (6 pA amplitude threshold), checked visually, and accepted or rejected according to their rise and decay times. The frequency and amplitude of all events, during and after drug application, were compared to the mean of the values observed during the initial control period. Cumulative probability plots of the incidence of various interevent intervals and amplitudes, recorded under different conditions from the same neuron, were subjected to the Kolmogorov–Smirnov (K-S) test. For other plots, data obtained over one min periods at the peak of a drug response were compared to the average values of frequency (or amplitude) of the sEPSC (or mEPSC) recorded during the initial control period (1–4 min). Data are expressed as means ( $\pm$  SEM). sEPSCs were fitted with a standard single exponential equation (Clampfit 9.2, Molecular Devices Co.) to determine the time

constant of decay from the EPSC peak. The statistical significance of drug effects was assessed by a paired two-tailed *t*-test on normalized data with Sigma Plot (Systat Software Inc., San Jose, CA). Values of  $P < 0.05$  were considered significant.

## Results

### Spontaneous EPSCs recorded in neurons mechanically dissociated from VLPO

In VLPO, two thirds of the neurons have a characteristic triangular and multipolar shape and low-threshold-spikes; moreover, as these cells contain the GABA synthesizing enzyme glutamic acid decarboxylase and are inhibited by noradrenaline and carbachol, they correspond to the GABAergic VLPO cells known to be active during sleep<sup>3,8</sup>. In the current study, we recorded from such multipolar, triangular shaped neurons within VLPO (Fig. 1). Most of the experiments were done on mechanically dissociated neurons (Fig. 1A1), which combine several advantages: good space clamp, preservation of functioning synaptic terminals, including some that release glutamate, as well as better control of the surrounding solution<sup>29, 30,34,35</sup>. The traces in Fig. 1 B-D were recorded under current-clamp. The voltage traces in the Fig. 1B1, Fig. 1C, and Fig. 1D1 were from a triangular-shaped, multipolar VLPO neuron. Depolarization from a relatively hyperpolarized level was followed by the characteristic depolarizing hump and burst of firing (inset in Fig. 1B1, and Fig. 1C), and noradrenaline induced membrane hyperpolarization (Fig. 1D1). Conversely, in the bipolar, fusiform VLPO neuron (Fig. 1A2), in response to a depolarization pulse there was a slow voltage rise towards the first action potential (Fig. 1B2, arrow) and noradrenaline induced membrane depolarization and firing (Fig. 1D2). Under voltage-clamp, sEPSCs were recorded with the CsF-based internal solution at a holding potential of  $-60$  mV. Under these conditions, postsynaptic responses to GABA or glycine are suppressed<sup>36</sup>, allowing us to monitor changes in frequency and amplitude of the isolated EPSCs<sup>30,37,38</sup>. 2,3-dihydroxy-6,7-dinitroquinoxaline ( $10$   $\mu$ M), an antagonist of  $\alpha$ -amino-3-hydroxy-5-methylisoxazole-4-propionic acid (AMPA) /kainate receptor reversibly blocked the spontaneous synaptic events (Fig. 1E), confirming that these VLPO neurons receive glutamatergic excitatory inputs<sup>15</sup>.

### Propofol raises sEPSC frequency in neurons mechanically dissociated from VLPO

Propofol ( $1$ – $100$   $\mu$ M) was applied to isolated VLPO neurons by a fast perfusion system (Y-tube)<sup>39</sup>. While propofol ( $10$   $\mu$ M) did not significantly change the holding current, it robustly but reversibly increased the frequency of sEPSCs (Fig. 2). A second application of propofol had a comparable effect ( $88 \pm 7\%$ , of first response;  $n = 8$ ,  $P > 0.05$ , Fig. 2A, B). As depicted by averaged traces recorded before and during propofol application (Fig. 2C) and the dose-response data in Fig. 2E, propofol ( $10$   $\mu$ M) did not significantly alter the mean amplitude of sEPSCs ( $7.6 \pm 1.2$  pA in control and  $10.3 \pm 1.8$  pA in presence of propofol,  $P > 0.05$ ,  $n = 30$  events). The superimposed normalized mean traces at right in Fig. 2C illustrate the absence of any change in sEPSC kinetics: the decay time constant of sEPSCs was not significantly altered (control:  $3.9 \pm 1.3$  ms, propofol:  $2.3 \pm 0.4$  ms,  $P = 0.13$ , by *t*-test). The effect of propofol on sEPSC frequency depended on its concentration, as shown in Fig. 2D. The apparent  $EC_{50}$  for propofol was  $4.2$   $\mu$ M. These results are consistent with a presynaptic mechanism of propofol action, enhancing glutamatergic transmission by increasing glutamate release.

### Propofol acts only on sEPSCs that are suppressed by tetrodotoxin and $Cd^{2+}$

We next examined how block of synaptic transmission affects the propofol-induced facilitation of sEPSC frequency. First, we examined the effect of  $1$   $\mu$ M tetrodotoxin, a blocker of voltage-dependent  $Na^+$  channels. Tetrodotoxin ( $1$   $\mu$ M) sharply decreased sEPSC frequency to  $38 \pm 10\%$  of control (from  $3.4 \pm 1.2$  Hz in control to  $1.1 \pm 0.6$  Hz in tetrodotoxin;  $n = 5$ ,  $P < 0.01$ ) but a moderate reduction in sEPSC amplitude was not significant (to  $75 \pm 7\%$  of control, from  $17 \pm 5$  pA to  $13 \pm 4$  pA;  $n = 5$ ,  $P > 0.05$ ). In the presence of  $1$   $\mu$ M tetrodotoxin, propofol ( $10$

$\mu\text{M}$ ) did not significantly alter either the frequency or the amplitude of the remaining EPSCs (presumably mEPSCs) (Fig. 3A-C). These results indicate that propofol acts only on sEPSCs generated by ongoing activity of tetrodotoxin-sensitive  $\text{Na}^+$  channels.

In further experiments, we applied  $\text{Cd}^{2+}$ , a general blocker of voltage-dependent  $\text{Ca}^{2+}$  channels, which also suppresses synaptic transmission. Indeed, acting somewhat like tetrodotoxin,  $100 \mu\text{M}$   $\text{Cd}^{2+}$  reduced sEPSC frequency by  $36 \pm 10\%$  ( $P = 0.03$ ,  $n = 5$ ) and amplitude by  $32 \pm 9\%$  ( $P = 0.02$ ,  $n = 5$ ); and in its presence,  $10 \mu\text{M}$  propofol did not significantly alter the frequency of the remaining EPSCs (Fig. 3D-F). Judging by these results, propofol targets only EPSCs generated by  $\text{Na}^+$  channel dependent terminal depolarization and the resulting  $\text{Ca}^{2+}$  influx.

### **The propofol-induced increase in glutamatergic activity is eliminated by a GABA<sub>A</sub>R antagonist and mimicked by a GABA<sub>A</sub>R agonist**

It is well known that propofol enhances the function of GABA<sub>A</sub>Rs, in particular those containing  $\beta 2$  and  $\beta 3$  subunits<sup>40</sup>, and that activation of presynaptic GABA<sub>A</sub>Rs enhances glutamate release in several brain areas<sup>22-25</sup>. To test for a possible involvement of GABA<sub>A</sub>Rs in the acceleration of sEPSC frequency, we applied gabazine ( $10 \mu\text{M}$ ), a specific antagonist of GABA<sub>A</sub>Rs. As shown in Fig. 4A-C, in the presence of gabazine,  $10 \mu\text{M}$  propofol caused a mild reduction in sEPSC frequency ( $-17.9 \pm 2.1\%$ ,  $n = 6$ ), presumably mediated by its GABA<sub>A</sub>R independent depressant effect on glutamate release<sup>21</sup>. To further confirm the role of GABA<sub>A</sub>Rs in the facilitation of sEPSCs, we applied muscimol, the specific GABA<sub>A</sub>R agonist. In 5 cells tested,  $10 \mu\text{M}$  muscimol significantly increased sEPSC frequency, but not amplitude (Fig. 4C-E).

### **The propofol-induced increase in glutamatergic activity is attenuated by an antagonist of $\text{Na}^+\text{-K}^+\text{-2Cl}^-$ cotransporters (NKCC)**

We have shown that voltage-dependent  $\text{Na}^+$  and  $\text{Ca}^{2+}$  channels, as well as GABA<sub>A</sub>Rs, are involved in the propofol-induced facilitation of sEPSC frequency. These data suggest that the activation of presynaptic GABA<sub>A</sub>Rs depolarizes some glutamatergic nerve terminals by generating  $\text{Cl}^-$  efflux. This implies that the intraterminal  $\text{Cl}^-$  concentration ( $[\text{Cl}^-]$ ) is relatively high, perhaps owing to inwardly directed  $\text{Cl}^-$  transport systems via NKCC and/or the  $\text{Cl}^-$ - $\text{HCO}_3^-$  exchanger<sup>41-43</sup>. In peripheral sensory axons, and immature or injured CNS neurons, NKCC raises intracellular  $[\text{Cl}^-]$ ; hence, in these cells GABA induces depolarization<sup>43</sup>. Indeed, in some other hypothalamic neurons NKCC is responsible for a relatively high  $[\text{Cl}^-]$  in glutamatergic nerve terminals<sup>44,45</sup>. To determine whether a high internal  $[\text{Cl}^-]$ , maintained by NKCC, is essential for the propofol-induced facilitation of sEPSC frequency, we tested propofol in the presence of  $10 \mu\text{M}$  bumetanide, which selectively blocks NKCC<sup>43,46</sup> (but not GABA receptors<sup>47</sup>). As illustrated in Fig. 5 by original traces in A, the plot in B and mean data in C, after 5 min of  $10 \mu\text{M}$  bumetanide application,  $10 \mu\text{M}$  propofol did not significantly alter the frequency of sEPSCs. Propofol's effect slowly recovered to control values after washing out bumetanide (data not shown). This result suggests that functional NKCC are present on the juxta-terminal portion of glutamatergic axons, where they maintain a relatively high internal  $[\text{Cl}^-]$ .

### **Propofol raises the frequency both of sEPSCs and of ongoing firing of VLPO neurons in brain slices**

We confirmed that propofol modulates glutamate release onto VLPO neurons under more physiological conditions in slices (Fig. 6). Propofol was tested on multipolar VLPO neurons, identified by the standard criteria (Fig. 6A). As illustrated by the traces (Fig. 6B), plot (Fig. 6C) and histograms (Fig. 6D),  $2 \mu\text{M}$  propofol robustly increased the frequency (but not amplitude) of glutamatergic sEPSCs in these VLPO neurons. Having established that propofol ( $2 \mu\text{M}$ ) facilitates glutamatergic transmission in slices, we next assessed the physiological



consequences of propofol's action by examining the ongoing discharge of VLPO neurons. As shown in Fig. 6E and Fig. 6F, 0.1  $\mu\text{M}$  propofol significantly raised the frequency of spontaneous firing of VLPO neurons in brain slices. Interestingly, as illustrated in Fig. 6G and Fig. 6H, at a higher concentration (2  $\mu\text{M}$ ) propofol had a sharp depolarizing action, which inactivated firing.

## Discussion

The central finding of this study is that propofol – at concentrations (1–10  $\mu\text{M}$ ) similar to those found effective in potentiating GABA<sub>A</sub>Rs in previous studies (e.g. 48–51) – raises the frequency of spontaneous glutamatergic transmission to VLPO neurons, both isolated and in slices. This is the first electrophysiological demonstration of the potentiating effect of propofol on glutamatergic transmission at the single cell level. An interesting parallel observation is that propofol, over a range of moderate concentrations (10–13  $\mu\text{M}$ ), selectively increases Ca-dependent glutamate release from cortical synaptosomes<sup>52</sup>. The VLPO neurons under study were multipolar in shape, generated low threshold spikes and were inhibited by noradrenaline, and therefore belonged to the GABAergic population<sup>3</sup> that projects to and inhibits the principal deep nuclei responsible for cortical and behavioural arousal<sup>8</sup>. According to previous studies<sup>2,20</sup>, the sedative action of propofol is mediated especially by enhanced GABAergic inhibition of the cells of the tuberomammillary nuclei. Specifically, Zecharia and colleagues<sup>2</sup> recently found that propofol enhances GABAergic inhibition of these cells by increasing the duration of inhibitory synaptic responses. This did not exclude the possibility that propofol also increases the excitability of VLPO neurons in some indirect manner, for example by disinhibition<sup>1</sup>. The present findings show that propofol increases ongoing excitation of the GABAergic VLPO neurons by enhancing glutamate release. Clearly, this mechanism could be a significant component of the sedative action of propofol.

Our recordings of AMPA receptor-mediated glutamatergic sEPSCs in the VLPO neurons are in keeping with a previous report<sup>15</sup>. A presynaptic target for propofol's action was indicated by the increase in sEPSC frequency and insignificant changes in the mean amplitude of sEPSCs and mEPSCs. The presence of GABA<sub>A</sub>Rs on or near the glutamatergic terminals was supported by the demonstration that propofol's effect was mimicked by muscimol, a GABA<sub>A</sub>R agonist, and blocked by gabazine, a GABA<sub>A</sub>R antagonist. The fact that activation of these GABA<sub>A</sub>Rs enhances glutamate release is readily explained if the Cl<sup>-</sup> concentration in these glutamatergic axons/terminals is relatively high; in which case, opening the presynaptic Cl<sup>-</sup> channels would induce Cl<sup>-</sup> efflux and thus membrane depolarization. The higher internal Cl<sup>-</sup> concentration appears to be the result of the activity of axonal NKCC-mediated inward Cl<sup>-</sup> transport, since bumetanide, - a selective inhibitor of Cl<sup>-</sup> transport that does not block GABA<sub>A</sub> receptors<sup>47</sup> - prevented the action of propofol. On the other hand, in view of the block of propofol's effect by tetrodotoxin, the depolarization directly caused by presynaptic GABA<sub>A</sub>R –induced Cl<sup>-</sup> efflux was unable to induce glutamate release; either because it was of insufficient magnitude to activate Ca<sup>2+</sup> influx into the terminal or because the operative GABA<sub>A</sub>Rs are situated at some distance, requiring the conduction of tetrodotoxin-sensitive action potentials: in either case, the participation of voltage-dependent Na<sup>+</sup> channels was crucial.

Previous studies on spinal cord, hippocampus, thalamus, and neocortex have not reported an excitatory effect of propofol on glutamatergic EPSCs. The simplest explanation could be that in contrast to previous authors, we recorded EPSCs in the absence of GABA<sub>A</sub>R antagonists – which eliminate this propofol action (as shown by our observations in the presence of gabazine).

Previous studies suggest that some GABA<sub>A</sub>R agonists promote sleep by inhibiting the histaminergic cells in the tuberomammillary nucleus and weakly activating the VLPO via

agonist binding to the  $\alpha 1$  subunit of GABA<sub>A</sub>Rs; whereas, gaboxadol (THIP; 4,5,6,7-tetrahydroisoxazolo[5,4-c]pyridin-3-ol) binds to the  $\alpha 4\delta$ -subunits, potentially promoting sleep by activation of the VLPO<sup>16</sup>. However, how activation of GABA<sub>A</sub>Rs could excite VLPO neurons was not determined. Our study may provide an answer to this question. Future research will define the relevant subunits of GABA<sub>A</sub>Rs, and whether the  $\alpha 4\delta$ -subunits in particular mediate the propofol-induced potentiation of sEPSCs of VLPO neurons.

In the current study, we tested 2  $\mu$ M propofol on both mechanically dissociated neurons and slices. Its effects on sEPSCs (in whole cell voltage clamp recordings) were similar in both preparations; but the facilitation of sEPSCs occurred more quickly in mechanically dissociated neurons than in slices, as might be expected in view of the very slow diffusion of propofol in slices<sup>53</sup>. On the other hand, in current clamp recordings of ongoing firing in slices, we found that even 0.1  $\mu$ M propofol significantly raised the firing rate of some VLPO neurons. Interestingly, at 2  $\mu$ M, propofol induced a sufficiently large depolarization that firing was inactivated. A possible explanation is that the indirect potentiation of glutamate release produced by propofol is especially pronounced in slices, where many more glutamatergic inputs to VLPO neurons are likely to be preserved.

## Conclusion

Our results show that chloride channel and GABA<sub>A</sub> receptors exist on the glutamatergic axons/terminals which make synapses on the VLPO neurons. By enhancing Cl<sup>-</sup> efflux from these axons, propofol depolarizes these terminals and stimulates the release of glutamate, which increases the activity of VLPO neurons, and thus potentiates GABAergic inhibition of arousal systems. This indirect mode of propofol action can be expected to contribute to its known effectiveness as a general anesthetic.

## Acknowledgments

Financial supports: This work is made possible by the Foundation of the University of Medicine and Dentistry of New Jersey, Newark, NJ; and National Institute of Health, Bethesda, MD, grant: AA016964.

## References


1. Nelson LE, Guo TZ, Lu J, Saper CB, Franks NP, Maze M. The sedative component of anesthesia is mediated by GABA(A) receptors in an endogenous sleep pathway. *Nat Neurosci* 2002;5:979–984. [PubMed: 12195434]
2. Zecharia AY, Nelson LE, Gent TC, Schumacher M, Jurd R, Rudolph U, Brickley SG, Maze M, Franks NP. The involvement of hypothalamic sleep pathways in general anesthesia: Testing the hypothesis using the GABAA receptor beta3N265M knock-in mouse. *J Neurosci* 2009;29:2177–2187. [PubMed: 19228970]
3. Gallopin T, Fort P, Eggermann E, Cauli B, Luppi PH, Rossier J, Audinat E, Muhlethaler M, Serafin M. Identification of sleep-promoting neurons in vitro. *Nature* 2000;404:992–995. [PubMed: 10801127]
4. Szymusiak R. Magnocellular nuclei of the basal forebrain: Substrates of sleep and arousal regulation. *Sleep* 1995;18:478–500. [PubMed: 7481420]
5. Sherin JE, Shiromani PJ, McCarley RW, Saper CB. Activation of ventrolateral preoptic neurons during sleep. *Science* 1996;271:216–219. [PubMed: 8539624]
6. Sherin JE, Elmquist JK, Torrealba F, Saper CB. Innervation of histaminergic tuberomammillary neurons by GABAergic and galaninergic neurons in the ventrolateral preoptic nucleus of the rat. *J Neurosci* 1998;18:4705–4721. [PubMed: 9614245]
7. Szymusiak R, Alam N, Steininger TL, McGinty D. Sleep-waking discharge patterns of ventrolateral preoptic/anterior hypothalamic neurons in rats. *Brain Res* 1998;803:178–188. [PubMed: 9729371]

8. Saper CB, Scammell TE, Lu J. Hypothalamic regulation of sleep and circadian rhythms. *Nature* 2005;437:1257–1263. [PubMed: 16251950]
9. Alam MN, Kumar S, Bashir T, Suntsova N, Methippara MM, Szymusiak R, McGinty D. GABA-mediated control of hypocretin- but not melanin-concentrating hormone-immunoreactive neurones during sleep in rats. *J Physiol* 2005;563:569–582. [PubMed: 15613374]
10. Lin JS, Sakai K, Jouvet M. Evidence for histaminergic arousal mechanisms in the hypothalamus of cat. *Neuropharmacology* 1988;27:111–122. [PubMed: 2965315]
11. Lin JS, Hou Y, Sakai K, Jouvet M. Histaminergic descending inputs to the mesopontine tegmentum and their role in the control of cortical activation and wakefulness in the cat. *J Neurosci* 1996;16:1523–1537. [PubMed: 8778302]
12. Sakurai T. The neural circuit of orexin (hypocretin): Maintaining sleep and wakefulness. *Nat Rev Neurosci* 2007;8:171–181. [PubMed: 17299454]
13. Ericson H, Kohler C, Blomqvist A. GABA-like immunoreactivity in the tuberomammillary nucleus: An electron microscopic study in the rat. *J Comp Neurol* 1991;305:462–469. [PubMed: 2037716]
14. Steininger TL, Gong H, McGinty D, Szymusiak R. Subregional organization of preoptic area/anterior hypothalamic projections to arousal-related monoaminergic cell groups. *J Comp Neurol* 2001;429:638–653. [PubMed: 11135241]
15. Sun X, Whitefield S, Rusak B, Semba K. Electrophysiological analysis of suprachiasmatic nucleus projections to the ventrolateral preoptic area in the rat. *Eur J Neurosci* 2001;14:1257–1274. [PubMed: 11703455]
16. Lu J, Greco MA. Sleep circuitry and the hypnotic mechanism of GABAA drugs. *J Clin Sleep Med* 2006;2:S19–S26. [PubMed: 17557503]
17. Trapani G, Altomare C, Liso G, Sanna E, Biggio G. Propofol in anesthesia. Mechanism of action, structure-activity relationships, and drug delivery. *Curr Med Chem* 2000;7:249–271. [PubMed: 10637364]
18. Bieda MC, MacIver MB. Major role for tonic GABAA conductances in anesthetic suppression of intrinsic neuronal excitability. *J Neurophysiol* 2004;92:1658–1667. [PubMed: 15140905]
19. Kotani Y, Shimazawa M, Yoshimura S, Iwama T, Hara H. The experimental and clinical pharmacology of propofol, an anesthetic agent with neuroprotective properties. *CNS Neurosci Ther* 2008;14:95–106. [PubMed: 18482023]
20. Zeller A, Jurd R, Lambert S, Arras M, Drexler B, Grashoff C, Antkowiak B, Rudolph U. Inhibitory ligand-gated ion channels as substrates for general anesthetic actions. *Handb Exp Pharmacol* 2008;31–51. [PubMed: 18175085]
21. Lingamaneni R, Birch ML, Hemmings HC Jr. Widespread inhibition of sodium channel-dependent glutamate release from isolated nerve terminals by isoflurane and propofol. *Anesthesiology* 2001;95:1460–1466. [PubMed: 11748406]
22. Turecek R, Trussell LO. Reciprocal developmental regulation of presynaptic ionotropic receptors. *Proc Natl Acad Sci U S A* 2002;99:13884–13889. [PubMed: 12370408]
23. Jang IS, Nakamura M, Ito Y, Akaike N. Presynaptic GABAA receptors facilitate spontaneous glutamate release from presynaptic terminals on mechanically dissociated rat CA3 pyramidal neurons. *Neuroscience* 2006;138:25–35. [PubMed: 16378694]
24. Stell BM, Rostaing P, Triller A, Marty A. Activation of presynaptic GABA(A) receptors induces glutamate release from parallel fiber synapses. *J Neurosci* 2007;27:9022–9031. [PubMed: 17715339]
25. Koga H, Ishibashi H, Shimada H, Jang IS, Nakamura TY, Nabekura J. Activation of presynaptic GABAA receptors increases spontaneous glutamate release onto noradrenergic neurons of the rat locus coeruleus. *Brain Res* 2005;1046:24–31. [PubMed: 15896724]
26. Ye JH, McArdle JJ. Excitatory amino acid induced currents of isolated murine hypothalamic neurons and their suppression by 2,3-butanedione monoxime. *Neuropharmacology* 1995;34:1259–1272. [PubMed: 8570023]
27. Dittman JS, Regehr WG. Mechanism and kinetics of heterosynaptic depression at a cerebellar synapse. *J Neurosci* 1997;17:9048–9059. [PubMed: 9364051]
28. Ye JH, Zhang J, Xiao C, Kong JQ. Patch-clamp studies in the CNS illustrate a simple new method for obtaining viable neurons in rat brain slices: Glycerol replacement of NaCl protects CNS neurons. *J Neurosci Methods* 2006;158:251–259. [PubMed: 16842860]




29. Akaike N, Moorhouse AJ. Techniques: Applications of the nerve-bouton preparation in neuropharmacology. *Trends Pharmacol Sci* 2003;24:44–47. [PubMed: 12498731]
30. Deng C, Li KY, Zhou C, Ye JH. Ethanol enhances glutamate transmission by retrograde dopamine signaling in a postsynaptic neuron/synaptic bouton preparation from the ventral tegmental area. *Neuropsychopharmacology* 2009;34:1233–1244. [PubMed: 18784647]
31. Paxinos, G.; Watson, C. *The Rat brain in stereotaxic coordinates* 6th edition. Paxinos, G.; Watson, C., editors. Academic press; 2007. Publisher: Elsevier Inc., Chapter page range 33–38.
32. Lu J, Greco MA, Shiromani P, Saper CB. Effect of lesions of the ventrolateral preoptic nucleus on NREM and REM sleep. *J Neurosci* 2000;20:3830–3842. [PubMed: 10804223]
33. Zhou C, Xiao C, Commissiong JW, Krnjevic K, Ye JH. Mesencephalic astrocyte-derived neurotrophic factor enhances nigral gamma-aminobutyric acid release. *Neuroreport* 2006;17:293–297. [PubMed: 16462600]
34. Ye JH, Wang F, Krnjevic K, Wang W, Xiong ZG, Zhang J. Presynaptic glycine receptors on GABAergic terminals facilitate discharge of dopaminergic neurons in ventral tegmental area. *J Neurosci* 2004;24:8961–8974. [PubMed: 15483115]
35. Matsuo S, Jang IS, Nabekura J, Akaike N. alpha 2-Adrenoceptor-mediated presynaptic modulation of GABAergic transmission in mechanically dissociated rat ventrolateral preoptic neurons. *J Neurophysiol* 2003;89:1640–1648. [PubMed: 12626630]
36. Bormann J, Hamill OP, Sakmann B. Mechanism of anion permeation through channels gated by glycine and gamma-aminobutyric acid in mouse cultured spinal neurones. *J Physiol* 1987;385:243–286. [PubMed: 2443667]
37. Turecek R, Trussell LO. Presynaptic glycine receptors enhance transmitter release at a mammalian central synapse. *Nature* 2001;411:587–590. [PubMed: 11385573]
38. Xiao C, Shao XM, Olive MF, Griffin WC 3rd, Li KY, Krnjevic K, Zhou C, Ye JH. Ethanol facilitates glutamatergic transmission to dopamine neurons in the ventral tegmental area. *Neuropsychopharmacology* 2009;34:307–318. [PubMed: 18596684]
39. Zhou C, Xiao C, McArdle JJ, Ye JH. Mefloquine enhances nigral gamma-aminobutyric acid release via inhibition of cholinesterase. *J Pharmacol Exp Ther* 2006;317:1155–1160. [PubMed: 16501066]
40. Rudolph U, Antkowiak B. Molecular and neuronal substrates for general anaesthetics. *Nat Rev Neurosci* 2004;5:709–720. [PubMed: 15322529]
41. Kaila K. Ionic basis of GABAA receptor channel function in the nervous system. *Prog Neurobiol* 1994;42:489–537. [PubMed: 7522334]
42. Plotkin MD, Kaplan MR, Peterson LN, Gullans SR, Hebert SC, Delpire E. Expression of the Na(+)-K(+)-2Cl- cotransporter BSC2 in the nervous system. *Am J Physiol* 1997;272:C173–C183. [PubMed: 9038823]
43. Russell JM. Sodium-potassium-chloride cotransport. *Physiol Rev* 2000;80:211–276. [PubMed: 10617769]
44. Jang IS, Jeong HJ, Akaike N. Contribution of the Na-K-Cl cotransporter on GABA(A) receptor-mediated presynaptic depolarization in excitatory nerve terminals. *J Neurosci* 2001;21:5962–5972. [PubMed: 11487619]
45. Choi HJ, Lee CJ, Schroeder A, Kim YS, Jung SH, Kim JS, Kim do Y, Son EJ, Han HC, Hong SK, Colwell CS, Kim YI. Excitatory actions of GABA in the suprachiasmatic nucleus. *J Neurosci* 2008;28:5450–5459. [PubMed: 18495878]
46. Haas M, Harrison JH Jr. Stimulation of K-Cl cotransport in rat red cells by a hemolytic anemia-producing metabolite of dapsone. *Am J Physiol* 1989;256:C265–C272. [PubMed: 2919657]
47. Dzhala VI, Talos DM, Sdrulla DA, Brumback AC, Mathews GC, Benke TA, Delpire E, Jensen FE, Staley KJ. NKCC1 transporter facilitates seizures in the developing brain. *Nat Med* 2005;11:1205–1213. [PubMed: 16227993]
48. Hales TG, Lambert JJ. The actions of propofol on inhibitory amino acid receptors of bovine adrenomedullary chromaffin cells and rodent central neurones. *Br J Pharmacol* 1991;104:619–628. [PubMed: 1665745]
49. Lam DW, Reynolds JN. Modulatory and direct effects of propofol on recombinant GABAA receptors expressed in xenopus oocytes: influence of alpha- and gamma2-subunits. *Brain Res* 1998;784:179–187. [PubMed: 9518600]

50. Patten D, Foxon GR, Martin KF, Halliwell RF. An electrophysiological study of the effects of propofol on native neuronal ligand-gated ion channels. *Clin Exp Pharmacol Physiol* 2001;28:451–458. [PubMed: 11380521]
51. Martella G, De Persis C, Bonsi P, Natoli S, Cuomo D, Bernardi G, Calabresi P, Pisani A. Inhibition of persistent sodium current fraction and voltage-gated L-type calcium current by propofol in cortical neurons: implications for its antiepileptic activity. *Epilepsia* 2005;46:624–635. [PubMed: 15857426]
52. Westphalen RI, Hemmings HC Jr. Selective depression by general anesthetics of glutamate versus GABA release from isolated cortical nerve terminals. *J Pharmacol Exp Ther* 2003;304:1188–1196. [PubMed: 12604696]
53. Gredell JA, Turnquist PA, Maciver MB, Pearce RA. Determination of diffusion and partition coefficients of propofol in rat brain tissue: Implications for studies of drug action in vitro. *Br J Anaesth* 2004;93:810–817. [PubMed: 15377586]



**Fig. 1.** Mechanically dissociated two types of ventrolateral preoptic nucleus (VLPO) neurons and glutamatergic excitatory postsynaptic currents (sEPSCs). (A) Typical mechanically dissociated VLPO neuron: note triangular shape characteristic of GABAergic cells with (LTS) low-threshold spikes<sup>3</sup> (A1, with patch pipette in place), scale bar: 20  $\mu\text{m}$ , and bipolar fusiform non-LTS neuron (A2), scale bar: 15  $\mu\text{m}$ . Low-threshold spikes occurred when the multipolar cell was depolarized from a hyperpolarizing level (at \* in inset in B1, and in trace C). This same neuron was hyperpolarized by 100  $\mu\text{M}$  noradrenaline (NA, D1). (B2), a slow voltage response toward the first action potential (arrow) recorded from a bipolar, fusiform non-LTS neuron (A2). This neuron was depolarized, and firing was induced by 100  $\mu\text{M}$  noradrenaline (NA,


D2). (E) sEPSCs recorded in a multipolar neuron were completely blocked by 2,3-dihydroxy-6,7-dinitroquinoxaline (DNQX, 10  $\mu$ M), indicating these events were mediated by  $\alpha$ -amino-3-hydroxy-5-methylisoxazole-4-propionic acid (AMPA) /Kainate-type glutamate receptors. Some traces are shown on an expanded time scale. For all figures, horizontal bars indicate period of drug applications.



**Fig. 2.** Propofol increases the frequency (but not amplitude) of spontaneous excitatory postsynaptic currents (sEPSCs) recorded in mechanically dissociated multipolar ventrolateral preoptic nucleus neurons. (A) Increased incidence of sEPSCs during two applications of 10  $\mu$ M propofol. (B) Time course of propofol-induced enhancement of sEPSC frequency (same data as illustrated in A). (C) Averaged traces show only minor change in amplitude, which was not significant (see E). Overlay of normalized traces indicates no change in sEPSC kinetics in presence of propofol (10  $\mu$ M) ( $n = 30$  events). Dose-response relationship for propofol-induced changes in sEPSC frequency (D) and amplitude (E). Solid line in D is least square fit of the frequency data to the Michaelis-Menten equation:  $P = (P_{MAX} * C^n) / (C^n + EC_{50}^n)$ , where P,




$P_{MAX}$ ,  $C$ ,  $EC_{50}$  and  $n$  are percent potentiation, maximal potentiation, propofol concentration, the concentration of propofol at which the potentiation of sEPSC frequency is 50% of maximum, and the Hill coefficient, respectively. The  $EC_{50}$  was  $4.2 \mu M$ . Each circle represents the mean  $\pm$  SEM of results from 3 to 14 cells. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$  by paired t test for propofol versus pre-propofol.



**Fig. 3.** Miniature excitatory postsynaptic currents, in mechanically isolated ventrolateral preoptic nucleus neurons recorded in presence of tetrodotoxin (TTX) or Cd<sup>2+</sup>, are not sensitive to propofol. (A-C) effects of propofol on spontaneous excitatory postsynaptic currents (sEPSCs) recorded in absence and presence of 1  $\mu$ M tetrodotoxin. (A) Original traces of spontaneous EPSCs obtained at times indicated by small letters in B. (B) time course of changes in sEPSC frequency. (C) Cumulative probability plots of intervals between sEPSCs show that sharp effect of propofol disappears in the presence of tetrodotoxin (n = 5). Insets are mean changes ( $\pm$  SEM); \*\*, P < 0.01 by paired t test for propofol versus pre-propofol. (D-F) effects of propofol on sEPSCs recorded in absence and presence of 100  $\mu$ M Cd<sup>2+</sup>. (D) Original traces of sEPSCs;


(E) time course of changes in sEPSC frequency. (F) Cumulative probability plots of intervals between sEPSCs: as indicated by inset histograms, (means  $\pm$  SEM, n = 5), effect of propofol was no longer significant in presence of 100  $\mu$ M Cd<sup>2+</sup>. \*\*, P < 0.01 by paired t test for propofol versus pre-propofol values.




**Fig. 4.** The action of propofol is mediated via  $\gamma$ -aminobutyric acid A receptors ( $GABA_A$ R): propofol-induced increase in frequency of spontaneous excitatory postsynaptic currents (sEPSCs) was abolished by selective  $GABA_A$ R antagonist gabazine and mimicked by the selective  $GABA_A$ R agonist muscimol. (A-C) Gabazine effects. (A) Traces from a neuron isolated from ventrolateral preoptic nucleus show that propofol (10  $\mu$ M) had no effect on sEPSCs in the presence of gabazine (10  $\mu$ M). (B) Time course of above data. (C-E) Like propofol, muscimol increases sEPSC frequency but not amplitude. (C) Traces recorded before (a, control), during (b, 10  $\mu$ M Muscimol) and after washout (c). (D) Time course of effect of muscimol on one cell. (E) Cumulative plots and the insets (means  $\pm$  SEM from five cells) show increase in sEPSC

frequency (left panel) but not amplitude (right panel). \*,  $P < 0.05$  by paired t test for muscimol versus control ( $n = 5$ ).





**Fig. 5.** Suppression of outward  $\text{Cl}^-$  gradient eliminates excitatory action of propofol. (A) Traces from a dissociated ventrolateral preoptic nucleus neuron show much smaller effect of propofol (10  $\mu\text{M}$ ) on the frequency of spontaneous excitatory postsynaptic currents (sEPSCs) in presence of 10  $\mu\text{M}$  bumetanide, a blocker of  $\text{Na}^+\text{-K}^+\text{-Cl}^-$  co-transporter. (B) Time course of changes in frequency. (C) Means  $\pm$  SEM from four cells. \*,  $P < 0.05$  by paired t test for propofol versus pre-propofol data.



**Fig. 6.** Propofol increases the frequency of spontaneous excitatory postsynaptic currents (sEPSCs) and firings in multipolar neurons in ventrolateral preoptic nucleus in slices. (A) A triangular-shaped neuron in ventrolateral preoptic nucleus with patch pipette in place. Scale bar, 20  $\mu$ m. (B) 2  $\mu$ M propofol significantly increased the frequency of sEPSCs in such a multipolar neuron in ventrolateral preoptic nucleus. (C), Time course of propofol-induced enhancement of sEPSC frequency. (D) Mean effects ( $\pm$  SEM,  $n = 4$  cells) of 2  $\mu$ M propofol on frequency and amplitude of sEPSCs. (E) Typical current traces and mean effects ( $\pm$  SEM,  $n = 3$  cells) (F) of spontaneous action potential (AP) frequency of neurons in ventrolateral preoptic nucleus show that 0.1  $\mu$ M propofol significantly increases the firing rate. (G) Typical current traces and mean effects ( $\pm$

SEM,  $n = 5$  cells, H) of neurons in ventrolateral preoptic nucleus show that  $2 \mu\text{M}$  propofol significantly depolarizes the neuron. \*,  $P < 0.05$ , \*\*,  $P < 0.01$ , respectively, by paired t test for propofol versus pre-propofol data.