

Published in final edited form as:

*Curr Opin Clin Nutr Metab Care*. 2008 January ; 11(1): 45–49. doi:10.1097/MCO.0b013e3282f2a592.

## Amino acid metabolism and regulatory effects in aging

Kyle L. Timmerman and Elena Volpi

Division of Geriatric Medicine, University of Texas Medical Branch, Galveston, Texas, USA

### Abstract

**Purpose of review**—To examine recent discoveries related to the amino acid metabolism and regulatory effects in aging, focusing on the development and treatment of age-related muscle loss (sarcopenia).

**Recent findings**—While basal amino acid metabolism may be unaffected by age, elderly subjects appear to have a decreased ability to respond to anabolic stimuli such as insulin and, to a lesser extent, amino acids. Specifically, compared to young subjects, the stimulation of muscle protein synthesis is attenuated in elderly subjects following the administration of mixed meals due to insulin resistance. In addition, the anabolic effect of amino acids appears blunted at low doses. Recent studies, however, have highlighted that these age-related alterations in amino acid metabolism may be overcome by provision of excess leucine, changes in the daily protein intake pattern or exercise, which improve activation of translation initiation and muscle protein synthesis.

**Summary**—Muscle loss with aging is associated with significant changes in amino acid metabolism, which can be acutely reversed using nutritional manipulations and exercise. Long-term, large clinical trials are, however, needed to determine the clinical significance of these findings in the elderly population, and to establish if nutritional and exercise interventions can help prevent and treat sarcopenia.

### Keywords

aging; amino acid; leucine; metabolism; muscle protein; sarcopenia

### Introduction

The main amino acid reservoir in the body is skeletal muscle, which contains 50–75% of all proteins in the human body [1,2]. In addition to its role in movement and posture, regulation of metabolism, and storage of energy and nitrogen, skeletal muscle becomes a vital supplier of amino acids to be used as a fuel by the brain and immune system, and as a substrate for wound healing during malnutrition, starvation, injury and disease [3]. The maintenance of body protein mass is critical not only to remain physically independent, but also for survival. The loss of approximately 30% of the body proteins results in impaired respiration and circulation due to muscle weakness, reduced immune function due to lack of nutrients, and inadequate barrier effect of the epithelia, which eventually will result in death [2].

Senescence in humans is characterized by an involuntary loss of muscle mass and function, termed sarcopenia. This degenerative loss of skeletal muscle occurs at a rate of 3–8% per decade after the age of 30 and accelerates with advancing age [4,5]. Sarcopenia is associated with decreased metabolic rate [6], decreased strength [7,8], increased risk of falls and

fractures [9], increased morbidity [7,10], and loss of independence [7]. When defining sarcopenia as appendicular skeletal muscle mass/height<sup>2</sup> less than 2 standard deviations below the mean for young, healthy reference populations, a quarter to a half of men and women aged 65 and older are likely sarcopenic [8,11]. Given our rapidly aging population, research designed to better understand the development, progression and treatment of sarcopenia is of substantial importance.

The mechanisms underlying the development of sarcopenia are not completely understood and likely numerous, but significant progress has been made over the past few years to identify some of the major contributors to the development of this condition. Here, we will review recent studies related to the regulatory effects and the role of amino acid metabolism in the development and treatment of age-related muscle loss. Following a logical progression of the discoveries in this area, we will begin with a discussion of amino acid and protein metabolism in the basal, postabsorptive state. Subsequently, we will outline the effects of nutrients and, in particular, amino acid on muscle metabolism with aging.

### **Basal amino acid and protein metabolism in aging**

Although the mechanisms leading to sarcopenia are likely numerous, a disproportionate rate of muscle protein breakdown compared to muscle protein synthesis clearly plays a role. Such an imbalance between breakdown and synthesis is smaller in size than that observed in wasting conditions, such as infections or traumatic injuries; however, when protracted over time it can lead to gradual and significant loss of muscle. Since muscle protein degradation has been consistently reported to remain essentially unchanged with advancing age [12-16], there has been an emphasis on studies examining the influence of age on muscle protein synthesis in the basal (postabsorptive) and fed (post-prandial) state. While some researchers have reported a decrease in basal muscle protein synthesis rate with age [17,18], others [14,19] could not confirm those findings in older individuals exhibiting a reduction in muscle mass. The reasons for these discrepancies are still unclear, but it is likely that differences in the health, nutritional status and physical activity level of the different older cohorts enrolled in the various studies [14,19] may have played a significant role. Furthermore, because in the studies reporting a reduced muscle protein synthesis with aging muscle protein breakdown had only been indirectly estimated using whole-body approaches, it is not possible to establish if the subjects were truly experiencing a reduction in net muscle protein balance with aging (i.e. net muscle loss). For example, if a slower muscle protein synthesis was accompanied by a concomitant decrease in breakdown (i.e. decreased turnover) the protein net balance would not change and muscle would not be lost. If there is no age-related difference in basal protein net balance, then it may be inferred that the events contributing to the development of sarcopenia are active outside of the postabsorptive period.

### **Nutrient intake on amino acid metabolism in aging**

The most important anabolic stimulus for muscle proteins is nutrient intake because it allows for replacement of the essential amino acids (EAAs) lost through oxidation. There is clear evidence that increased amino acid or protein availability can enhance muscle protein synthesis and anabolism in young and older subjects [19-21,22\*]. It has, however, been suggested that the recommended dietary allowance for protein (0.8 g/kg/day) may not be sufficient for older adults to maintain their muscle mass. In fact, some researchers have reported that the elderly should consume up to 1.2 g/kg/day [23]. In partial support of this conjecture, Thalacker-Mercer *et al.* [24\*] reported that inadequate protein intake (0.5 g/kg/day) by older adults resulted in a significant downregulation of muscle transcripts associated with synthesis, energy metabolism and proliferation compared to older adults consuming

adequate amounts of dietary protein (1.2 g/kg/day). No measures of muscle protein synthesis or balance were, however, available to determine the net effect of these protein intakes on muscle mass.

Despite suggestions that the elderly should consume more protein [23], the use of high-protein diets alone to increase muscle mass and strength in the elderly has been mostly ineffective [25,26]. There are a number of reasons why these nutritional interventions may have failed to produce positive results. First, when subjects are given nutritional supplements, there is evidence to suggest that they naturally compensate by consuming fewer calories as part of their *ad libitum* diet [25,27] and thus negate any anabolic effects associated with protein supplementation. Second, it is also possible that older adults have a diminished ability to respond to the anabolic effects of the supplements analogous to that observed in old animals. The latter hypothesis is corroborated by the finding that ingestion of an amino acid/glucose mixture stimulated muscle protein synthesis in young, but not older adults [13]. These data have since been confirmed by Guillet *et al.* [28] through the use of a hyperinsulemic/euglycemic clamp while intravenously administering amino acids to emulate the postprandial state.

The existence of insulin resistance of muscle protein metabolism with aging, independent of glucose tolerance, has been further demonstrated in older, healthy and nondiabetic subjects [29]. This defect appears associated with the age-related reduction in endothelium-dependent vasodilation [29], and can be reversed by aerobic exercise through improvements in endothelial function and insulin-induced vasodilation, and insulin signaling [30]. These data suggest that vasodilation and nutrient flow to the muscle are important regulators of the muscle anabolic response during hyperinsulinemia [29] and during feeding [13]. This hypothesis is further supported by recent data obtained in young individuals where various levels of physiological hyperinsulinemia were induced in the absence of amino acid replacement [31]. In this experiment, the muscle protein anabolic response was mainly related to insulin-induced changes in blood flow and amino acid delivery to the muscle, rather than the absolute insulin level. In other words, in order for hyperinsulinemia to stimulate muscle protein anabolism it must increase capillary recruitment and amino acid flow to the muscle. Altogether, the studies described above highlight the importance of an adequate amino acid supply to the muscle tissue in order to initiate and sustain muscle protein anabolism both in young and older persons.

## Amino acids and regulation of muscle metabolism in healthy aging

A number of studies have shown that pure amino acids can stimulate muscle protein synthesis and improve net protein balance in older as well as in younger individuals [19,32-35]. Although older persons have an increased splanchnic extraction of orally administered amino acids at first pass (i.e. immediately after absorption) [34,36], this does not appear to influence the systemic amino acid concentration, which normally increases in the elderly as well as in the young, and consequently the muscle anabolic effect of amino acids [34]. EAAs, in particular, are able to stimulate muscle protein synthesis in the elderly, whereas non-EAAs do not appear to provide any additional benefit with regard to muscle protein deposition and anabolism [35]. Among the EAAs, the branched-chain amino acid (BCAA) leucine has been shown to be a key regulator of muscle protein synthesis in both humans and rats [37-39]. Leucine activates translation initiation in the skeletal muscle cells by increasing the phosphorylation of several signaling proteins including the mammalian target of rapamycin, 70-kDa ribosomal protein S6 kinase and eukaryotic initiation factor 4E-binding protein-1 [37,38,40,41\*].

Although large amounts of EAAs exert similar effects in young and older persons [19,34], age-related differences in the muscle anabolic response to submaximal amino acid doses have been recently uncovered. Katsanos *et al.* [42] reported that older subjects had significantly less muscle protein accretion than younger subjects following the ingestion of a 7-g EAA bolus. In a more recent study, the same authors [43\*\*] found that while both a 26% (1.721 g leucine) and a 41% (2.79 g leucine) leucine EAA bolus increased muscle protein synthesis in young men, only the 41% leucine EAA bolus was effective in elderly men. These data are consistent with Paddon-Jones *et al.* [44\*\*] who reported that the iso-caloric ingestion of EAAs containing 2.79 g leucine increased phenylalanine uptake and muscle protein synthesis to a significantly greater degree in older adults than did ingestion of the same amount of a whole protein supplement, whey protein, containing only 1.75 g leucine. Based on these studies it is possible to speculate that aged muscle may be slightly less sensitive to the anabolic effects of leucine than young muscle, but that this age-related difference could be overcome by increased leucine intake.

Recent studies have examined the influence of leucine supplementation as part of a meal on protein synthesis in older muscle, and confirmed that supplementation with leucine dramatically improves postprandial muscle protein synthesis in both old rats [45,46,47\*] and humans [48\*\*]. Data from Koopman *et al.* [49\*\*] further highlight that coingestion of protein and leucine along with carbohydrate increased the muscle protein synthesis rate in young and old men to a similar degree. Rieu *et al.* [47\*] also examined the influence of meals supplemented with various milk proteins, containing varying amounts of leucine, on postprandial muscle protein synthesis in old rats. They found that protein supplements with the highest leucine content (e.g.  $\beta$ -lactoglobulin, 14.5% leucine) elicited a significantly greater postprandial response of muscle protein synthesis than protein supplements with lower leucine proportion (e.g. casein, 10% leucine). Thus, strategies capable of significantly increasing plasma levels of leucine appear capable of restoring the stimulatory effects of a meal in aging. This phenomenon may partly explain reports [23,50\*,51] that older adults require a higher protein intake or a protein 'pulse feeding' pattern, i.e. daily protein intake concentrated on one of the daily meals [52,53], to improve their nitrogen and amino acid balance.

## Amino acids and muscle metabolism in chronic disease

The ability of amino acid ingestion to stimulate muscle protein synthesis has also recently been demonstrated in clinical populations [54\*\*,55\*]. Killewich *et al.* [55\*] measured muscle protein synthesis in the calf muscles of older patients with peripheral arterial disease and sex-matched controls before and after the ingestion of 15 g EAAs. Despite the fact that the peripheral arterial disease patients had reduced leg blood flow and, thus, presumably decreased delivery of EAAs, they experienced a significant increase in muscle protein synthesis comparable to healthy controls. This led the authors to speculate that the decreased muscle perfusion in these patients was not sufficient to influence delivery of EAAs to muscle or to attenuate their anabolic effects. In elderly subjects with chronic obstructive pulmonary disease, Engelen *et al.* [54\*\*] reported that soy protein added with BCAAs stimulated whole-body protein synthesis to a greater degree than did soy protein alone. These data are consistent with previous reports that EAAs, particularly BCAAs, are predominantly responsible for the anabolic effects of amino acid ingestion. Together, these data imply that both healthy and diseased elderly may benefit from amino acid supplementation, and that composition, timing and dosage of the supplement should be carefully considered.

Caution is, however, warranted given reports that high physiologic levels of amino acids can induce insulin resistance in humans [56,57]. Specifically, Tremblay *et al.* [57] have reported

that increased availability of amino acids impairs the ability of insulin to attenuate glucose production and the ability of muscle to dispose of excess glucose. Their data suggest that the mechanisms underlying amino acid-induced insulin resistance include overactivation of mammalian target of rapamycin and 70-kDa ribosomal protein S6 kinase, along with the inhibition of insulin receptor substrate-1 via serine phosphorylation. Other than the obvious complications that arise from increased insulin resistance (e.g. metabolic syndrome and type 2 diabetes), a recent study [58] suggests that insulin resistance may accelerate muscle protein degradation. Thus, studies are necessary to determine not only the minimum amount of amino acid and protein intake necessary to maintain an adequate muscle mass in aging, but also the upper limit beyond which side effects could occur.

## Conclusion

In summary, aging is associated with a progressive loss of muscle mass, which is at least in part due to negative changes in protein and amino acid homeostasis. While older adults may still exhibit normal basal muscle protein synthesis, recent data imply that there may be an age-related decrease in the ability of aged muscle to respond to various anabolic stimuli, including insulin, mixed meals containing amino acids and carbohydrate, and, to some extent, amino acids themselves. Consequently, there is a clear need for strategies that are effective at maximizing muscle protein synthesis and anabolism in the elderly. Based on results from the most recent studies, such strategies may include nutritional supplementation with protein or amino acids, particularly leucine, pulse protein feeding and exercise. Two important points must be noted, however: (i) many of the studies published in the literature have been acute in nature and small in size, and (ii) high physiologic levels of amino acids may potentially induce insulin resistance. As such, recommendations regarding specific dietary and/or exercise interventions await large longitudinal, randomized clinical trials.

## Acknowledgments

Supported by the National Institute on Aging grants R01 AG18311 and P30 AG024832 (UTMB Claude D. Pepper Older Americans Independence Center).

## References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (p. 85).

1. Welle, S. Human protein metabolism. Springer; New York: 1999.
2. Matthews, DE. Proteins and amino acids.. In: Shils, ME., editor. Modern nutrition in health and disease. Williams & Wilkins; Baltimore: 1999. p. 11-48.
3. Reeds PJ, Fjeld CR, Jahoor F. Do the differences between the amino acid compositions of acute-phase and muscle proteins have a bearing on nitrogen loss in traumatic states? *J Nutr.* 1994; 124:906-910. [PubMed: 7515956]
4. Holloszy JO. The biology of aging. *Mayo Clin Proc.* 2000; 75(Suppl):S3-S8. [PubMed: 10959208]
5. Melton LJ III, Khosla S, Riggs BL. Epidemiology of sarcopenia. *Mayo Clin Proc.* 2000; 75(Suppl):S10-S12. [PubMed: 10959209]
6. Karakelides H, Sreekumaran NK. Sarcopenia of aging and its metabolic impact. *Curr Top Dev Biol.* 2005; 68:123-148. [PubMed: 16124998]

7. Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc.* 2002; 50:889–896. [PubMed: 12028177]
8. Iannuzzi-Sucich M, Prestwood KM, Kenny AM. Prevalence of sarcopenia and predictors of skeletal muscle mass in healthy, older men and women. *J Gerontol A Biol Sci Med Sci.* 2002; 57:M772–M777. [PubMed: 12456735]
9. Sayer AA, Syddall HE, Martin HJ, et al. Falls, sarcopenia, and growth in early life: findings from the Hertfordshire cohort study. *Am J Epidemiol.* 2006; 164:665–671. [PubMed: 16905644]
10. Janssen I, Baumgartner RN, Ross R, et al. Skeletal muscle cutpoints associated with elevated physical disability risk in older men and women. *Am J Epidemiol.* 2004; 159:413–421. [PubMed: 14769646]
11. Baumgartner RN, Koehler KM, Gallagher D, et al. Epidemiology of sarcopenia among the elderly in New Mexico. *Am J Epidemiol.* 1998; 147:755–763. [PubMed: 9554417]
12. Hasten DL, Pak-Loduca J, Obert KA, Yarasheski KE. Resistance exercise acutely increases MHC and mixed muscle protein synthesis rates in 78–84 and 23–32 yr olds. *Am J Physiol Endocrinol Metab.* 2000; 278:E620–E626. [PubMed: 10751194]
13. Volpi E, Mittendorfer B, Rasmussen BB, Wolfe RR. The response of muscle protein anabolism to combined hyperaminoacidemia and glucose-induced hyperinsulinemia is impaired in the elderly. *J Clin Endocrinol Metab.* 2000; 85:4481–4490. [PubMed: 11134097]
14. Volpi E, Sheffield-Moore M, Rasmussen BB, Wolfe RR. Basal muscle amino acid kinetics and protein synthesis in healthy young and older men. *JAMA.* 2001; 286:1206–1212. [PubMed: 11559266]
15. Welle S, Thornton C, Statt M. Myofibrillar protein synthesis in young and old human subjects after three months of resistance training. *Am J Physiol.* 1995; 268(3 Pt 1):E422–E427. [PubMed: 7900788]
16. Yarasheski KE, Zachwieja JJ, Bier DM. Acute effects of resistance exercise on muscle protein synthesis rate in young and elderly men and women. *Am J Physiol.* 1993; 265(2 Pt 1):E210–E214. [PubMed: 8368290]
17. Balagopal P, Rooyackers OE, Adey DB, et al. Effects of aging on *in vivo* synthesis of skeletal muscle myosin heavy-chain and sarcoplasmic protein in humans. *Am J Physiol.* 1997; 273(4 Pt 1):E790–E800. [PubMed: 9357810]
18. Welle S, Thornton C, Jozefowicz R, Statt M. Myofibrillar protein synthesis in young and old men. *Am J Physiol.* 1993; 264(5 Pt 1):E693–E698. [PubMed: 8498491]
19. Paddon-Jones D, Sheffield-Moore M, Zhang XJ, et al. Amino acid ingestion improves muscle protein synthesis in the young and elderly. *Am J Physiol Endocrinol Metab.* 2004; 286:E321–E328. [PubMed: 14583440]
20. Rennie MJ, Edwards RH, Halliday D, et al. Muscle protein synthesis measured by stable isotope techniques in man: the effects of feeding and fasting. *Clin Sci (Lond).* 1982; 63:519–523. [PubMed: 6181926]
21. Biolo G, Tipton KD, Klein S, Wolfe RR. An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *Am J Physiol.* 1997; 273(1 Pt 1):E122–E129. [PubMed: 9252488]
- 22•. Symons TB, Schutzler SE, Cocke TL, et al. Aging does not impair the anabolic response to a protein-rich meal. *Am J Clin Nutr.* 2007; 86:451–456. [PubMed: 17684218] [• of special interestAging does not impair the ability of a high-quality protein source (beef) to stimulate muscle protein synthesis.]
23. Campbell WW, Trappe TA, Wolfe RR, Evans WJ. The recommended dietary allowance for protein may not be adequate for older people to maintain skeletal muscle. *J Gerontol A Biol Sci Med Sci.* 2001; 56:M373–M380. [PubMed: 11382798]
- 24•. Thalacker-Mercer AE, Fleet JC, Craig BA, et al. Inadequate protein intake affects skeletal muscle transcript profiles in older humans. *Am J Clin Nutr.* 2007; 85:1344–1352. [PubMed: 17490972] [• of special interestShort-term inadequate protein intake elicits changes in muscle protein transcripts that may contribute to reduced muscle protein synthesis and muscle loss.]



25. Fiatarone MA, O'Neill EF, Ryan ND, et al. Exercise training and nutritional supplementation for physical frailty in very elderly people. *N Engl J Med*. 1994; 330:1769–1775. [PubMed: 8190152]
26. Welle S, Thornton CA. High-protein meals do not enhance myofibrillar synthesis after resistance exercise in 62- to 75-yr-old men and women. *Am J Physiol*. 1998; 274(4 Pt 1):E677–E683. [PubMed: 9575829]
27. Fiatarone Singh MA, Bernstein MA, Ryan AD, et al. The effect of oral nutritional supplements on habitual dietary quality and quantity in frail elders. *J Nutr Health Aging*. 2000; 4:5–12. [PubMed: 10828934]
28. Guillet C, Prod'homme M, Balage M, et al. Impaired anabolic response of muscle protein synthesis is associated with S6K1 dysregulation in elderly humans. *FASEB J*. 2004; 18:1586–1587. [PubMed: 15319361]
29. Rasmussen BB, Fujita S, Wolfe RR, et al. Insulin resistance of muscle protein metabolism in aging. *FASEB J*. 2006; 20:768–769. [PubMed: 16464955]
30. Fujita S, Rasmussen BB, Cadenas JG, et al. Aerobic exercise overcomes the age-related insulin resistance of muscle protein metabolism by improving endothelial function and Akt/mammalian target of rapamycin signaling. *Diabetes*. 2007; 56:1615–1622. [PubMed: 17351147] [• of special interestA single bout of aerobic exercise restores the anabolic response of muscle to insulin by improving endothelial function, anabolic signaling and amino acid availability for the muscle.]
31. Fujita S, Rasmussen BB, Cadenas JG, et al. Effect of insulin on human skeletal muscle protein synthesis is modulated by insulin-induced changes in muscle blood flow and amino acid availability. *Am J Physiol Endocrinol Metab*. 2006; 291:E745–E754. [PubMed: 16705054]
32. Rasmussen BB, Wolfe RR, Volpi E. Oral and intravenously administered amino acids produce similar effects on muscle protein synthesis in the elderly. *J Nutr Health Aging*. 2002; 6:358–362. [PubMed: 12459885]
33. Volpi E, Ferrando AA, Yeckel CW, et al. Exogenous amino acids stimulate net muscle protein synthesis in the elderly. *J Clin Invest*. 1998; 101:2000–2007. [PubMed: 9576765]
34. Volpi E, Mittendorfer B, Wolf SE, Wolfe RR. Oral amino acids stimulate muscle protein anabolism in the elderly despite higher first-pass splanchnic extraction. *Am J Physiol*. 1999; 277(3 Pt 1):E513–E520. [PubMed: 10484364]
35. Volpi E, Kobayashi H, Sheffield-Moore M, et al. Essential amino acids are primarily responsible for the amino acid stimulation of muscle protein anabolism in healthy elderly adults. *Am J Clin Nutr*. 2003; 78:250–258. [PubMed: 12885705]
36. Boirie Y, Gachon P, Beaufrere B. Splanchnic and whole-body leucine kinetics in young and elderly men. *Am J Clin Nutr*. 1997; 65:489–495. [PubMed: 9022534]
37. Anthony JC, Yoshizawa F, Anthony TG, et al. Leucine stimulates translation initiation in skeletal muscle of postabsorptive rats via a rapamycin-sensitive pathway. *J Nutr*. 2000; 130:2413–2419. [PubMed: 11015466]
38. Anthony JC, Anthony TG, Kimball SR, et al. Orally administered leucine stimulates protein synthesis in skeletal muscle of postabsorptive rats in association with increased eIF4F formation. *J Nutr*. 2000; 130:139–145. [PubMed: 10720160]
39. Garlick PJ. The role of leucine in the regulation of protein metabolism. *J Nutr*. 2005; 135(6 Suppl): 1553S–1556S. [PubMed: 15930468]
40. Anthony TG, Anthony JC, Yoshizawa F, et al. Oral administration of leucine stimulates ribosomal protein mRNA translation but not global rates of protein synthesis in the liver of rats. *J Nutr*. 2001; 131:1171–1176. [PubMed: 11285321]
41. Fujita S, Dreyer HC, Drummond MJ, et al. Nutrient signalling in the regulation of human muscle protein synthesis. *J Physiol*. 2007; 582(Pt 2):813–823. [PubMed: 17478528] [• of special interestIngestion of a leucine-enriched EAA/carbohydrate mixture promotes muscle protein synthesis via both an increased initiation of translation and an increased translation elongation.]
42. Katsanos CS, Kobayashi H, Sheffield-Moore M, et al. Aging is associated with diminished accretion of muscle proteins after the ingestion of a small bolus of essential amino acids. *Am J Clin Nutr*. 2005; 82:1065–1073. [PubMed: 16280440]
43. Katsanos CS, Kobayashi H, Sheffield-Moore M, et al. A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the

- elderly. *Am J Physiol Endocrinol Metab.* 2006; 291:E381–E387. [PubMed: 16507602] [•• of outstanding interestThe muscle protein anabolic response to a small (submaximal) bolus of EAAs is blunted in older subjects, but can be restored by changing the EAA proportion in favor of leucine. Conversely, excess leucine has no additional anabolic effect in young subjects.]
- 44••. Paddon-Jones D, Sheffield-Moore M, Katsanos CS, et al. Differential stimulation of muscle protein synthesis in elderly humans following isocaloric ingestion of amino acids or whey protein. *Exp Gerontol.* 2006; 41:215–219. [PubMed: 16310330] [•• of outstanding interestIngestion of EAAs stimulates muscle protein synthesis to a greater extent than ingestion of the same amount of whey protein, which contains non-EAAs. This confirms and highlights that EAAs are mainly responsible for the anabolic action of proteins.]
45. Dardevet D, Sornet C, Bayle G, et al. Postprandial stimulation of muscle protein synthesis in old rats can be restored by a leucine-supplemented meal. *J Nutr.* 2002; 132:95–100. [PubMed: 11773514]
46. Rieu I, Sornet C, Bayle G, et al. Leucine-supplemented meal feeding for ten days beneficially affects postprandial muscle protein synthesis in old rats. *J Nutr.* 2003; 133:1198–1205. [PubMed: 12672943]
- 47•. Rieu I, Balage M, Sornet C, et al. Increased availability of leucine with leucine-rich whey proteins improves postprandial muscle protein synthesis in aging rats. *Nutrition.* 2007; 23:323–331. [PubMed: 17367997] [• of special interestProteins rich in leucine provide a more potent stimulus for muscle protein synthesis than protein sources with a lower percentage of leucine in rats.]
- 48••. Rieu I, Balage M, Sornet C, et al. Leucine supplementation improves muscle protein synthesis in elderly men independently of hyperaminoacidaemia. *J Physiol.* 2006; 575(Pt 1):305–315. [PubMed: 16777941] [•• of outstanding interestLeucine supplementation improves muscle protein synthesis in elderly men during feeding regardless of increases in the concentrations of other amino acids. This highlights how leucine can at least partially overcome the resistance of muscle proteins to the anabolic action of mixed feeding.]
- 49••. Koopman R, Verdijk L, Manders RJ, et al. Co-ingestion of protein and leucine stimulates muscle protein synthesis rates to the same extent in young and elderly lean men. *Am J Clin Nutr.* 2006; 84:623–632. [PubMed: 16960178] [•• of outstanding interestCo-ingestion of protein and leucine with carbohydrates stimulates muscle protein synthesis to the same extent in old as in young subjects.]
- 50•. Morais JA, Chevalier S, Gougeon R. Protein turnover and requirements in the healthy and frail elderly. *J Nutr Health Aging.* 2006; 10:272–283. [PubMed: 16886097] [• of special interestThis is a recent review on protein requirements for healthy and frail older adults.]
51. Morse MH, Haub MD, Evans WJ, Campbell WW. Protein requirement of elderly women: nitrogen balance responses to three levels of protein intake. *J Gerontol A Biol Sci Med Sci.* 2001; 56:M724–M730. [PubMed: 11682582]
52. Arnal MA, Mosoni L, Boirie Y, et al. Protein pulse feeding improves protein retention in elderly women. *Am J Clin Nutr.* 1999; 69:1202–1208. [PubMed: 10357740]
53. Arnal MA, Mosoni L, Dardevet D, et al. Pulse protein feeding pattern restores stimulation of muscle protein synthesis during the feeding period in old rats. *J Nutr.* 2002; 132:1002–1008. [PubMed: 11983828]
- 54••. Engelen MP, Rutten EP, De Castro CL, et al. Supplementation of soy protein with branched-chain amino acids alters protein metabolism in healthy elderly and even more in patients with chronic obstructive pulmonary disease. *Am J Clin Nutr.* 2007; 85:431–439. [PubMed: 17284740] [•• of outstanding interestBCAA supplementation of soy protein improves the anabolic effect of soy protein on whole-body metabolism, particularly in elderly patients with chronic obstructive pulmonary disease.]
- 55•. Killewich LA, Tuvdendorj D, Bahadorani J, et al. Amino acids stimulate leg muscle protein synthesis in peripheral arterial disease. *J Vasc Surg.* 2007; 45:554–559. [PubMed: 17321342] [• of special interestDespite decreased blood flow, patients with peripheral arterial disease respond to the stimulatory effects of amino acids to the same extent as healthy elderly subjects.]
56. Krebs M, Krssak M, Bernroider E, et al. Mechanism of amino acid-induced skeletal muscle insulin resistance in humans. *Diabetes.* 2002; 51:599–605. [PubMed: 11872656]



57. Tremblay F, Krebs M, Dombrowski L, et al. Overactivation of S6 kinase 1 as a cause of human insulin resistance during increased amino acid availability. *Diabetes*. 2005; 54:2674–2684. [PubMed: 16123357]
58. Wang X, Hu Z, Hu J, et al. Insulin resistance accelerates muscle protein degradation: activation of the ubiquitin–proteasome pathway by defects in muscle cell signaling. *Endocrinology*. 2006; 147:4160–4168. [PubMed: 16777975]