

Research

Open Access

The 12-item General Health Questionnaire (GHQ-12): translation and validation study of the Iranian version

Ali Montazeri*, Amir Mahmood Harirchi, Mohammad Shariati, Gholamreza Garmaroudi, Mehdi Ebadi and Abolfazl Fateh

Address: Iranian Institute for Health Sciences Research, Tehran, Iran

Email: Ali Montazeri* - ali@jdcord.jd.ac.ir; Amir Mahmood Harirchi - hariram@hbi.or.ir; Mohammad Shariati - IHSR@jdcord.jd.ac.ir; Gholamreza Garmaroudi - IHSR@jdcord.jd.ac.ir; Mehdi Ebadi - IHSR@jdcord.jd.ac.ir; Abolfazl Fateh - IHSR@jdcord.jd.ac.ir

* Corresponding author

Published: 13 November 2003

Received: 13 October 2003

Health and Quality of Life Outcomes 2003, **1**:66

Accepted: 13 November 2003

This article is available from: <http://www.hqlo.com/content/1/1/66>

© 2003 Montazeri et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: The objective of this study was to translate and to test the reliability and validity of the 12-item General Health Questionnaire (GHQ-12) in Iran.

Methods: Using a standard 'forward-backward' translation procedure, the English language version of the questionnaire was translated into Persian (Iranian language). Then a sample of young people aged 18 to 25 years old completed the questionnaire. In addition, a short questionnaire containing demographic questions and a single measure of global quality of life was administered. To test reliability the internal consistency was assessed by Cronbach's alpha coefficient. Validity was performed using convergent validity. Finally, the factor structure of the questionnaire was extracted by performing principal component analysis using oblique factor solution.

Results: In all 748 young people entered into the study. The mean age of respondents was 21.1 (SD = 2.1) years. Employing the recommended method of scoring (ranging from 0 to 12), the mean GHQ score was 3.7 (SD = 3.5). Reliability analysis showed satisfactory result (Cronbach's alpha coefficient = 0.87). Convergent validity indicated a significant negative correlation between the GHQ-12 and global quality of life scores as expected ($r = -0.56$, $P < 0.0001$). The principal component analysis with oblique rotation solution showed that the GHQ-12 was a measure of psychological morbidity with two-factor structure that jointly accounted for 51% of the variance.

Conclusion: The study findings showed that the Iranian version of the GHQ-12 has a good structural characteristic and is a reliable and valid instrument that can be used for measuring psychological well being in Iran.

Background

The General Health Questionnaire (GHQ) is a measure of current mental health and since its development by Goldberg in the 1970s it has been extensively used in different settings and different cultures [1–5]. The questionnaire was originally developed as a 60-item instrument but at present a range of shortened versions of the questionnaire

including the GHQ-30, the GHQ-28, the GHQ-20, and the GHQ-12 is available. The scale asks whether the respondent has experienced a particular symptom or behavior recently. Each item is rated on a four-point scale (less than usual, no more than usual, rather more than usual, or much more than usual); and for example when using the GHQ-12 it gives a total score of 36 or 12 based

on the selected scoring methods. The most common scoring methods are bi-modal (0-0-1-1) and Likert scoring styles (0-1-2-3). Since the GHQ-12 is a brief, simple, easy to complete, and its application in research settings as a screening tool is well documented; it was decided to translate the GHQ-12 into Persian (the Iranian language) and to examine the psychometric properties of the questionnaire in a sample of young Iranian adolescents. This was to develop the Iranian version of the GHQ-12 to meet the increasing demand for the questionnaire. There is evidence that the GHQ-12 is a consistent and reliable instrument when used in general population samples [6].

Methods

The standard "forward-backward" procedure was applied to translate the questionnaire from English into Persian. Two independent health professionals translated the items and two others translated the response categories and a provisional version was provided. Subsequently it was back translated into English and following a careful cultural adaptation the final version was provided. Finally after pilot testing, the translated questionnaire was administered to a sample of healthy young people aged 18 to 25 years old. The sample was recruited through a national family journal and from two higher education institutes. The participants who were recruited through the journal returned the completed questionnaire by post and those who were recruited from two higher education institutes were tested in several groups while the assessors were present. In addition each respondent was asked to complete a short questionnaire containing demographic questions and to rate a measure of global quality of life; a subscale derived from the validated Iranian version of the European Organization for Research and Treatment of Cancer Quality of Life Questionnaire (EORTC QLQ-C30)[7]. Global quality of life subscale contains 2 items and each item is rated on a seven-point scale. A linear transformation was performed to standardize the row scores. Scores range from 0 to 100 and the higher value indicates a higher level of global quality of life [8].

To test the reliability, the internal consistency of the questionnaire was assessed by Cronbach's alpha coefficient and alpha equal to or greater than 0.70 was considered satisfactory [9]. Validity was performed using convergent validity to demonstrate the extent to which the GHQ-12 correlates with global quality of life. It was expected that the GHQ-12 would correlate negatively with global quality of life. This was assessed by the Pearson product moment statistic (Pearson's correlation coefficient = r) and r equal to 0.40 or above was considered satisfactory. Furthermore the factor structure of the questionnaire was extracted by performing principal component analysis using oblique factor solution. The study used Goldberg's original scoring method. In this method response categories

score 0, 0, 1, and 1 respectively. This gives scores ranging from 0 to 12 [10].

Results

Descriptive findings

In all 748 young people aged 18 to 25 years entered into the study. The descriptive findings are presented in Table 1. The mean age of respondents was 21.1 (SD = 2.1) years and most were female (76%), single (84%) and college/university students (50%). Employing the bi-modal method of scoring (ranging from 0 to 12) the mean GHQ score was found to be 3.7 (SD = 3.5). Forty-four percent scored above the mean GHQ score for the whole population of the respondents. The mean global quality of life score was 65.1 (SD = 21.5).

Table 1: The characteristics of respondents and descriptive findings (n = 748)

	No.	%
Age (group)		
18–19	209	28
20–22	335	45
23–25	204	27
Mean (SD)	21.1 (2.1)	
Gender		
Female	567	76
Male	181	24
Educational level (n = 730)		
Primary/ Secondary	310	41
College/ University	420	56
Marital status		
Single	632	84
Married	105	14
Widowed	11	2
Employment status (n = 556)		
Employed	114	22
Student	280	50
Housewife	69	12
Unemployed	93	16
GHQ score		
Mean (SD)	3.7 (3.5)	
Range	0–12	
Global quality of life (n = 743)		
Mean (SD)	65.1 (21.5)	
Range	0–100	

Reliability

To test the reliability the internal consistency of the questionnaire was measured using Cronbach's alpha coefficient. The alpha for the whole sample was found to be 0.87 and was the same for both males and females indicating satisfactory results.

Validity

Validity of the instrument was performed using convergent validity. When the correlation between the GHQ-12 and global quality of life scores was investigated, as expected a significant negative correlation emerged ($r = -0.56, P < 0.0001$) indicating that those who were more distressed showed lower levels of global quality of life.

Factor structure

The principal component analysis with oblique rotation solution was performed and a two-factor structure was loaded that jointly accounted for 51% of the variance. The results are shown in Table 2. Apart from item seven (enjoy normal activities), other items loaded in two distinct factors producing the factors of 'psychological distress' and 'social dysfunction'.

Table 2: Factor structure of the GHQ-12 using principal component analysis with oblique rotation solution

	Factor 1	Factor 2
GHQ-12 Items		
1. Able to concentrate	0.57	-
2. Lost much sleep	-	0.69
3. Playing useful part	0.81	-
4. Capable of making decisions	0.56	-
5. Under stress	-	0.68
6. Could not overcome difficulties	-	0.62
7. Enjoy normal activities	0.57	0.46
8. Face up to problems	0.64	-
9. Feeling unhappy and depressed	-	0.64
10. Losing confidence	0.62	-
11. Thinking of self as worthless	0.79	-
12. Feeling reasonably happy	-	0.56

Discussion

The GHQ is a well-known instrument for measuring minor psychological distress and has been translated into a variety of languages [11–16]. However, it is not a tool for indicating a specific diagnosis. This study reports data from a validation study of the 12-item GHQ in Iran. In general, the findings showed promising results and were comparable with most research findings throughout the world. Our two-factor solution was similar to those reported in the WHO study of psychological disorders in general health care [17]. Reliability was assessed by internal consistency of the questionnaire reporting Cronbach's alpha coefficient and validity was examined by convergent validity performing correlation between the GHQ-12 and global quality of life scores and both showed satisfactory results.

It is argued that although the GHQ-12 was originally developed as a unitary screening measure for psychologi-

cal problems, there have also been efforts to identify whether it has a multidimensional structure [18]. The World Health Organization study of psychological disorders in general health care in 15 different centers indicated that for the GHQ-12 substantial factor variation between centers exist. However, the study reported that after rotation two factors expressing depression and social dysfunction could be identified [17]. The findings from present study showed that the Iranian GHQ-12 is a valid measure of psychological distress and the factor structure of the questionnaire was very similar to that of the original language. Interestingly a recent findings from a Japanese study reported that the GHQ-12 could be used as an internally reliable and homogenous scale that produces mainly the factors of psychological distress and social dysfunction [16].

We used the bi-modal fashion of scoring and the results indicated that this method in Iran appears to be useful. Evidence suggests that there is no tendency for the GHQ to work less efficiently in developing countries [19]. The mean GHQ-12 score in this study was 3.7 (SD = 3.5). It is recommended that the mean GHQ score for the whole population of respondents provides a rough guide to the best cut-off threshold [20]. Thus considering people who scored above the mean, the findings from the present study indicated that 44% of the respondents showed an indication of mental health problems [21]. This clearly suggests that if investigators wish to use a screening instrument as a case detector, the shorter GHQ is remarkably robust and works as well as the longer instrument [19]. Similar studies among young adolescents reported that the GHQ-12 is a particularly useful measure with adolescents where there are likely to be a number of different threats to their psychological health, such as poor self-esteem, that may not necessarily constitute a formal psychiatric condition [22]. In contrast, studies have shown that the GHQ-12 is not a suitable instrument for some special populations such as elderly patients [23].

Iran has a very young population. In general, the findings from this study indicated that mental health in young people in Iran is poor and it is strongly associated with their quality of life. Since mental health in young adolescents could be regarded as a risk factor for psychological disorders such as antisocial behavior, criminal activity, suicidal behavior, substance abuse, depression, and eating disorders [24], improving quality of life in this age group becomes very important task. However, in interpreting the study findings it should be noted that the young people in the sample were a selected sample and thus it cannot be generalized to the whole population of young adolescents in Iran.

Conclusion

The findings suggest that the Iranian version of the 12-item GHQ is a reliable and valid instrument to measure minor psychological distress in young people and has a good factor structure.

List of abbreviations

GHQ: General Health Questionnaire; SD = standard deviation; WHO: World Health Organization.

Competing interest

None declared.

Authors' contribution

AM was the main investigator, analyzed the data, and wrote the paper. AMH, MSh GhG, ME, and AF all contributed to the study design, the translation procedure, data collection and first draft of the paper in Persian.

References

- Goldberg DP and Blackwell B: **Psychiatric illness in general practice. A detailed study using a new method of case identification.** *Br Med J* 1970, **1**:439-443.
- Golderberg D and Williams P: **A user's guide to the General Health questionnaire.** Windsor, UK: NFER-Nelson; 1988.
- Schmitz N, Kruse J and Tress W: **Psychometric properties of the General Health Questionnaire (GHQ)-12 in a German primary care.** *Acta Psychiatr Scand* 1999, **100**:462-468.
- Jacob KS, Bhugra D and Mann AH: **The validation of the 12-item General Health questionnaire in ethnic Indian women living in the United Kingdom.** *Psychol Med* 1997, **27**:1215-1217.
- Donath S: **The validity of the 12-item General Health Questionnaire in Australia: a comparison between three scoring methods.** *Aust N Z J Psychiatry* 2001, **35**:231-235.
- Pevalin DJ: **Multiple applications of the GHQ-12 in a general population sample: an investigation of long-term retest effects.** *Soc Psychiatry Psychiatr Epidemiol* 2000, **35**:508-512.
- Montazeri A, Harirchi I and Vahdani M et al.: **The European Organization for Research and Treatment of Cancer: translation and validation study of the Iranian version.** *Support Care Cancer* 1999, **7**:400-406.
- Fayers P, Aaronson N, Bjordal K, Groenvold M, Curran D and Bottomley A: **The EORTC QLQ-C30 Scoring Manual.** 3rd edition. EORTC Quality of Life Group, Brussels; 2001.
- Nunnally JC and Bernstein IH: **Psychometric Theory.** 3rd edition. New York: McGraw-Hill; 1994.
- Goldberg D: **General Health Questionnaire (GHQ-12).** Windsor, UK: NFER-Nelson; 1992.
- Politi PL, Piccinelli M and Wilkinson G: **Reliability, validity and factor structure of the 12-item General Health Questionnaire among young males in Italy.** *Acta Psychiatr Scand* 1994, **90**:432-437.
- Daradkeh TK, Ghubash R and el-rufai OE: **Reliability, validity, and factor structure of the Arabic version of the 12-item General Health Questionnaire.** *Psychol Rep* 2001, **89**:85-94.
- Quek KF, Low WY, Razack AH and Loh CS: **Reliability and validity of the General Health Questionnaire (GHQ-12) among urological patients: A Malaysian study.** *Psychiatry Clin Neurosci* 2001, **55**:509-513.
- Kilic C, Rezaki M, Rezaki B, Kaplan I, Ozgen G, Sagduyu A and Ozturk MO: **General Health Questionnaire (GHQ-12, GHQ-28): psychometric properties and factor structure of the scales in a Turkish primary care sample.** *Soc Psychiatry Psychiatr Epidemiol* 1997, **32**:327-331.
- Chan DW: **The Chinese General Health Questionnaire in a psychiatric setting. The development of the Chinese scaled version.** *Soc Psychiatry Psychiatr Epidemiol* 1993, **28**:124-129.
- Doi Y and Minowa M: **Factor structure of the 12-item General Health Questionnaire in the Japanese general adult population.** *Psychiatry Clin Neurosci* 2003, **57**:379-383.
- Werneke U, Goldberg DP, Yalcin I and Ustun BT: **The stability of the factor structure of the General Health Questionnaire.** *Psychol Med* 2000, **30**:823-829.
- Campbell A, Walker J and Farrell G: **Confirmatory factor analysis of the GHQ-12: can I see that again?** *Aust N Z J Psychiatry* 2003, **37**:475-483.
- Goldberg DP, Gater R, Sartorius N, Ustun TB, Piccinelli M, Gureje O and Rutter C: **The validity of two version of the GHQ in the WHO study of mental illness in general health care.** *Psychol Med* 1997, **27**:191-197.
- Goldberg DP, Oldhinkel T and Ormel J: **Why GHQ threshold varies from one place to another.** *Psychol Med* 1998, **28**:915-921.
- Montazeri A, Hosseini Kh and Sadighi J et al.: **Quality of life and mental health in young Iranian population [abstract].** *Quality of Life Research* 2001, **10**:274.
- Tait RJ, French DJ and Hulse GK: **Validity and psychometric properties of the General Health Questionnaire-12 in young Australian adolescents.** *Aust N Z J Psychiatry* 2003, **37**:374-381.
- Heun R, Muller H, Freyberger HJ and Maier W: **Reliability of interview information in a family study in the elderly.** *Soc Psychiatry Psychiatr Epidemiol* 1998, **33**:140-144.
- Pearce J: **Psychological disturbances in young people: challenges for prevention.** *Br Med J* 1996, **312**:453.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

