

Respiratory syncytial virus infection in children with acute respiratory infections in Zambia

M. SAIJO^{1*}, H. TERUNUMA¹, K. MIZUTA^{1,2}, E. M. MPABALWANI¹, M. MONZE¹,
H. OSHITANI³, N. LUO¹, H. SUZUKI³ AND Y. NUMAZAKI^{1,2}

¹ *The Virology Laboratory, Department of Microbiology and Pathology, University Teaching Hospital, Lusaka, Zambia*

² *Virus Research Center, Sendai National Hospital, Sendai, Japan*

³ *Department of Public Health, Niigata University School of Medicine, Niigata, Japan*

(Accepted 23 April 1998)

SUMMARY

Epidemiological research on respiratory syncytial virus (RSV) infections in children was carried out at the Virology Laboratory, University Teaching Hospital (UTH), in Lusaka, Zambia, from January–December 1996. Specimens including 736 nasal washings and 2424 throat swabs were collected from children with acute respiratory infections (ARI) and tested for RSV by enzyme immunoassay and by virus isolation. RSV was isolated in 62 (4·1%) of 1496 throat swabs collected from March to September and was detected in 99 (16·3%) of 609 nasal washings from March to November. The average RSV isolation rate was 2·6% and the average RSV detection rate was 13·5%. The highest RSV isolation (8·1%) and detection (30·5%) rates were in June 1996. RSV antibody in the 278 serum specimens collected from Zambian children, who were hospitalized in the paediatric ward, UTH, was detected using a standard neutralization test. The antibody positive rate was 60–80% in children > 4 years. It is evident that RSV is one of the main causal agents of ARI in children in Zambia.

INTRODUCTION

Acute lower respiratory tract infections are a major cause of morbidity and mortality in children in developing countries [1–3]. Viruses such as respiratory syncytial virus (RSV), influenza virus, parainfluenza virus, and adenovirus play an important role in respiratory infections in children [2, 4]. However, there are few reports on the epidemiology of viral respiratory infections in developing countries because of the limitations in diagnostic technology.

RSV is the commonest cause of acute respiratory infections (ARI) in young children in temperate regions [5–9]. It may cause life-threatening lower respiratory tract infections [10, 11], and so it is important to monitor outbreaks of RSV infection in each region. We studied the incidence of RSV in ARI

in Zambian children using cell culture techniques and an enzyme immunoassay (EIA). This is the first report documenting RSV as one of the important agents causing ARI in children in Zambia, a sub-Saharan African country.

MATERIALS AND METHODS

Study period

An epidemiological study on RSV infection in Zambia was carried out from January to December 1996.

Patients and specimens

Patients were recruited from two urban health centres in Lusaka. Throat swab specimens of children with ARI who visited the two centres were collected for viral isolation during the study period. The specimen

* Author for correspondence.

collection and handling methods have already been described [12, 13]. Nasal washings were also collected for RSV antigen detection using EIA, from January to December 1996 [14]. Clinical data on each patient were collected using standard questionnaires including sex, age, clinical manifestations, clinical diagnosis and complications. Serum samples from 278 children who were hospitalized in a paediatric ward and submitted to the paediatric laboratory for various diagnostic tests were randomly selected and supplied to the Virology laboratory for RSV antibody detection.

Virus isolation and identification

Virus isolation from the throat swab specimens were performed using the previously described microplate method at the Virology Laboratory, University Teaching Hospital (UTH), Lusaka, Zambia [12, 13]. Identification of RSV isolates was performed by direct immunofluorescence [12].

RSV antigen detection

RSV antigen in the nasal washings was detected with the EIA using TESTPACK RSVTM (Abbott Laboratories, IL, USA) [15, 16].

RSV neutralizing antibody detection

RSV antibody in the serum specimens was detected using a standard neutralization test. Briefly, Long strain virus and HEp-2 cells were used. A virus suspension containing 10 µl of 10 TCID₅₀ was mixed for 60 min with fourfold serial dilutions of inactivated serum in phosphate buffered saline (10 µl). The virus mixture was inoculated onto duplicate HEp-2 cell monolayers seeded in 96-well microplates. The specific cytopathic effect (CPE) of RSV was observed and antibody titre was evaluated after 3 days incubation period.

RESULTS

Patients and specimens

During the one-year study period, throat swabs were collected from 2424 children with ARI, the average number was 202 per month (range 150–240). Nasal washings were collected from 736 children with ARI from January to December 1996, the average number was 61 per month (range 39–84 per month from January to November 1996, but and 10 in December 1996). The male to female ratio was 1:1. The age

Table 1. Number of RSV isolates and the isolation rates in children by age group. The throat swabs were collected between January 1995 and December 1996

Age (year)	Number of samples	Number of isolates	Rate (%)
< 1	1160	38	3.3
1	713	15	2.1
2	281	5	1.8
3	170	2	1.2
4	90	2	3.3
= B 5	10	0	0
Total	2424	62	2.6

Table 2. Number of RSV positives by EIA by age group between January and September 1996

Age (year)	Number of samples	Number of isolates	Rate (%)
< 1	370	48	13.0
1	337	46	13.6
2	27	5	17.2
Total	731	99	13.5




Fig. 1. Monthly distribution of RSV cases recognized by cell culture [■] and by EIA [□], January–December 1996.

distribution of children from whom throat swabs and nasal washings were collected is shown in Tables 1 and 2, respectively.

RSV isolation and RSV antigen detection

RSV was isolated from 62 (4.1%) out of 1496 specimens collected from March to September. The highest RSV isolation rate was 8.1% in June 1996. RSV antigen was detected in 99 (16.3%) out of 609 nasal washings collected from March to November 1996




Fig. 2. Monthly RSV isolation rate (●) and antigen positive rate (■), January–December 1996.




Fig. 3. RSV neutralizing antibody prevalence among Zambian children. The figures on the top of the bars indicate the number of specimens.

(Fig. 1). The RSV antigen positive rate by EIA was 13.5%, and the highest monthly rate was 30.5% in June. The monthly distribution of isolation and antigen positive rates is shown in Fig. 2. The age distribution of children, from whom RSV was isolated or was detected by EIA, is shown in Tables 1 and 2.

RSV neutralizing antibody prevalence

The RSV neutralizing antibody prevalence in Zambian children is shown in Fig. 3. The lowest RSV antibody positive rate was at 1 year of age, and it increased with age. The antibody positive rate was 60–80% among children > 4 years.

DISCUSSION

RSV is one of the main causes of ARI in children in Zambia, and seems to have a seasonal pattern. The

number of RSV isolates and antigen positives identified per month increased gradually from March, peaked in June and declined thereafter. The peak months of influenza virus infections were July and August [13]. This means that in Zambia outbreaks of RSV infection among children are followed by outbreaks of influenza virus infection. The high prevalence of RSV neutralizing antibody by 4 years of age suggests that RSV is commonly transmitted among small children in Zambia.

Gbaldero and colleagues [17] and Johnson and colleagues [18] report that RSV is one of the most important viral agents, and causes bronchiolitis and asthma in Nigeria. Greenwood [19] reported that RSV was one of the most important causes of acute lower respiratory infection in the Gambia, West Africa, together with *Haemophilus influenzae* type *b*. However, the reports on RSV epidemiology are very few from developing African countries and we believe that our report presents valuable information on the understanding of RSV epidemiology in Africa.

In this study, there was a difference in the rates between RSV isolation and RSV antigen detection. The difference in the rates can be explained by the difficulty associated with virus isolation. RSV is easily inactivated in a warm environment, and specimens must be inoculated onto cells as soon as possible [14]. However, there was a time lag between taking throat swabs and inoculation. Further, throat swabs are inferior to nasal washings for RSV isolation. Throat swabs were collected for RSV isolation in our study. So the RSV isolation rate by cell culture was lower than RSV antigen detection rate by EIA.

We did not analyse the clinical manifestations and outcome of RSV infections in Zambian children in this study. Further study is needed to make them clear. RSV infections in small children are severe, and life-threatening in some cases [10, 11]. The mortality rate of RSV infections in human immunodeficiency virus (HIV)-infected children is reported to be higher than that of immunocompetent children [20]. The HIV-seroprevalence in Zambian children, who were hospitalized in the paediatric ward, UTH, was reported to be 30.4% in 1995 [21]. Also, the mortality of lower respiratory tract infections was significantly higher among HIV-seropositives than among HIV-seronegatives [21]. Malnutrition in Zambian children also seems to be a co-factor for a high mortality of lower respiratory infections. So, we speculate that the mortality rate of RSV infections may be high. To clarify this speculation, further study is needed. We

conclude that RSV is one of the major causes of ARI in Zambian children.

ACKNOWLEDGEMENTS

We thank the staff of the two participating health centres in Lusaka for specimen collection and clinical investigation. We also thank the staff of the Virology Laboratory, UTH, for virus isolation and identification. This study was performed as part of a technical cooperation project between the governments of Japan and Zambia through the Japan International Cooperation Agency (JICA).

REFERENCES

- Glezen WP, Couch RB. Influenza viruses. In: Evans AS, ed. *Viral infections of humans: Epidemiology and Control*, 3rd ed. New York: Plenum Publishing, 1991; 419–50.
- Bulla A, Hitze KI. Acute respiratory infections: a review. *Bull WHO* 1978; **56**: 481–98.
- World Health Organization. Program for the control of acute respiratory infections. *Weekly Epidemiol Rec* 1993; **68**: 353–7.
- Walter EB, Shurin PA. Acute respiratory infections. In: Krugman S, ed. *Infectious diseases of children*, 9th ed. St. Louis: Mosby Year Book, 1992; 329–76.
- Saijo M, Ishii T, Kokubo M, Takimoto M, Takahashi Y. Respiratory syncytial virus infection in lower respiratory tract and asthma attack in hospitalized children in North Hokkaido, Japan. *Acta Paediatr Jpn* 1993; **35**: 233–7.
- Saijo M, Takahashi S, Kokubo M, et al. The role of respiratory syncytial virus in acute bronchiolitis in small children in northern Japan. *Acta Paediatr Jpn* 1994; **36**: 371–4.
- Orstavik I, Carlsen K-H, Halvolson K. Respiratory syncytial virus infection in Oslo 1972–1978. *Acta Paediatr Scand* 1980; **69**: 717–22.
- Avendano LF, Larranaga C, Palomino MA, et al. Community- and hospital-acquired respiratory syncytial infection in Chile. *Pediatr Infect Dis J* 1991; **10**: 564–8.
- Popow-Kraupp T, Kunz C, Huber E. Respiratory virus infection in hospitalized children in Austria 1979–1982. *Infection* 1984; **12**: 185–9.
- Disney ME, Sandiford BR, Cragg J, Wolff J. Epidemic bronchiolitis in infants. *BMJ* 1960; **1**: 1407–11.
- Heycock JB, Noble TC. 1230 cases of acute bronchiolitis in infancy. *BMJ* 1962; **2**: 879–81.
- Numazaki Y, Oshima T, Ohmi A, et al. A microplate method for isolation of viruses from infants and children with acute respiratory infections. *Microbiol Immunol* 1987; **31**: 1085–95.
- Mizuta K, Oshitani H, Saijo M, et al. Epidemiology of influenza virus infections in children with acute respiratory infections in Zambia. *Ann Trop Paediatr* 1997; **17**: 115–9.
- Caroline B, Douglas RG Jr. Clinically useful method for the isolation of respiratory syncytial virus. *J Clin Microbiol* 1975; **131**: 1–5.
- Swierkosz EM, Flanders R, Melvin L, Miller JD, Kline MW. Evaluation of the Abbott TestPack RSV enzyme immunoassay for the detection of respiratory syncytial virus in nasopharyngeal swab specimens. *J Clin Microbiol* 1989; **27**: 1151–4.
- Thomas EE, Book LE. Comparison of two rapid methods for detection of respiratory syncytial virus (RSV) (Testpack RSV and Ortho RSV ELISA) with direct immunofluorescence and virus isolation for the diagnosis of pediatric RSV infection. *J Clin Microbiol* 1990; **29**: 632–5.
- Gbadero DA, Johnson AW, Aderole WI, Olaleye OD. Microbiol inciters of acute asthma in urban Nigerian children. *Thorax* 1995; **50**: 739–45.
- Johnson AW, Aderole WI, Osinusi K, Gbadero DA, Fagbami AH, Rotowa NA. Acute bronchiolitis in tropical Africa: a hospital-based perspective in Ibadan, Nigeria. *Pediatr Pulmonol* 1996; **22**: 236–47.
- Greenwood B. Epidemiology of acute lower respiratory tract infections, especially those due to *Haemophilus influenzae* type b, in The Gambia, West Africa. *J Infect Dis* 1992; **165**: S26–8.
- Chandwani S, Borkowsky W, Krasinski K, Lawrence R, Welliver R. Respiratory syncytial virus infection in human immunodeficiency virus-infected children. *J Pediatr* 1990; **117**: 251–4.
- Baht G, Mawenyeho S, Luo C, Terunuma H, Numazaki Y. HIV seroprevalence and childhood morbidity and mortality in Lusaka, Zambia (O-58-4). In: *Abstracts of the 14th International Congress for Tropical Medicine and Malaria*. Nagasaki, Japan: International Federation for Tropical Medicine, 1996: 297.