

Pronounced Enhancement of $\cdot\text{NO}$ -Dependent Antimicrobial Action by an $\cdot\text{NO}$ -Oxidizing Agent, Imidazolineoxyl *N*-Oxide

KAZUKO YOSHIDA,^{1,2} TAKA AKAIKE,¹ TOSHINORI DOI,^{1,2} KEIZO SATO,¹ SUMIKO IJIRI,¹ MORITAKA SUGA,² MASAYUKI ANDO,² AND HIROSHI MAEDA^{1*}

Department of Microbiology¹ and The First Department of Internal Medicine,² Kumamoto University School of Medicine, Kumamoto 860, Japan

Received 22 March 1993/Accepted 22 May 1993

The antimicrobial action of $\cdot\text{NO}$ against *Cryptococcus neoformans* was investigated by using imidazolineoxyl *N*-oxide, which we recently reported removes $\cdot\text{NO}$ via oxidation (T. Akaike, M. Yoshida, Y. Miyamoto, K. Sato, M. Kohno, K. Sasamoto, K. Miyazaki, S. Ueda, and H. Maeda, *Biochemistry* 32:827-832, 1993). No appreciable fungicidal activity was observed in neutral $\cdot\text{NO}$ solutions. Imidazolineoxyl *N*-oxide induced or enhanced fungicidal action in neutral or acidic $\cdot\text{NO}$ solutions, respectively. Our results provide convincing evidence that $\cdot\text{NO}$ is not a microbicidal molecular species.

Nitric oxide ($\cdot\text{NO}$) has been implicated as an antitumor or antimicrobial molecule that is generated by activated macrophages (23). A number of studies have clearly demonstrated the cytostatic action of $\cdot\text{NO}$ on tumor cells and cryptococci through various metabolic inhibitions in the cells (3, 7, 8a, 9, 10, 15, 22). However, $\cdot\text{NO}$ is not as reactive and toxic as its derivatives (5, 8, 13, 14, 21, 23), and the cytotoxic (cytolytic) mechanism of $\cdot\text{NO}$ remains obscure (8a, 16). Therefore, investigating the biological activities of $\cdot\text{NO}$ in more detail, focusing on its antimicrobial activity, is worthwhile. To identify the biological significance of $\cdot\text{NO}$, it is essential to examine the effects of specific inhibitors of $\cdot\text{NO}$ and their antimicrobial actions. Until now, however, very few compounds with specific inhibitory action against $\cdot\text{NO}$ have been available. Only L-arginine analogs, such as *N*^ω-monomethyl-L-arginine, are widely used as inhibitors of $\cdot\text{NO}$ synthase to block $\cdot\text{NO}$ generation and analyze the effect of $\cdot\text{NO}$, but L-arginine analogs do not antagonize $\cdot\text{NO}$ directly. In contrast, we have found that the imidazolineoxyl *N*-oxide derivative 2-phenyl-4,4,5,5-tetramethylimidazoline-3-oxide-1-oxyl (PTIO) acts as an oxygen donor for $\cdot\text{NO}$, forming $\cdot\text{NO}_2$, and thus scavenges $\cdot\text{NO}$ (Fig. 1) (2). Therefore, it now becomes possible to investigate the role of $\cdot\text{NO}$ as a cytotoxic molecular species that is known to be generated by activated macrophages.

PTIO was synthesized and purified as reported recently (2). 2-Phenyl-4,4,5,5-tetramethylimidazoline-1-oxyl (PTI) was prepared by deoxidation of PTIO with $\cdot\text{NO}$ and was purified by silica gel column chromatography (2). PTIO is a stable radical compound, reacts with $\cdot\text{NO}$ more than 10-fold faster than does molecular oxygen, and yields $\cdot\text{NO}_2$ and PTI in a stoichiometric manner in solutions with a rate constant of $\sim 10^4 \text{ M}^{-1} \text{ s}^{-1}$. We also demonstrated that PTIO strongly inhibited the vasorelaxation induced by endothelium-derived relaxing factor. The inhibitory potential of PTIO is comparable to that of the $\cdot\text{NO}$ synthesis inhibitor *N*^ω-monomethyl-L-arginine (2).

Cryptococcus neoformans (serotype A or D), originally obtained from clinical isolates, was used for the assays. The serotype A strain (TIMM1855) was a gift from K. Uchida (Research Center for Medical Mycology, Teikyo University,

Tokyo, Japan), and the serotype D strain was a gift from T. Shinoda (Department of Microbiology, Meiji College of Pharmacy, Tokyo, Japan). Yeast cells were grown on Sabouraud agar plates, harvested in log-phase growth, and washed with 0.45% saline three times before use. Cells were then added to the reaction mixtures at 10^6 CFU/ml (final concentration).

Washed cells of *C. neoformans* serotype D were added to the reaction mixtures, which contained 1 mM NaNO_2 in 80 mM succinate buffer (pH 4.0) and generated $\cdot\text{NO}$ chemically (3, 20), in the presence or absence of PTIO (Fig. 2). The number of viable cells in each reaction mixture was then quantitated by colony formation on the Sabouraud agar plates after incubation for indicated periods at room temperature. NaNO_2 , succinate buffer, and PTIO did not exhibit any fungicidal action at the concentrations used. The $\cdot\text{NO}$ -generating mixture showed fungicidal action (open circles). PTIO enhanced the cytotoxic action in a dose-dependent manner: in the presence of 200 μM PTIO (closed circles), a 10^5 -fold increase in cytotoxic activity was observed during a 3-h incubation period.

An $\cdot\text{NO}$ -saturated solution was prepared by bubbling $\cdot\text{NO}$ gas (99% purity; Nippon Sanso Co., Ltd., Tokyo, Japan) into pure water, which had been degassed and bubbled with helium gas. The content of $\cdot\text{NO}$ was quantitated by electron spin resonance spectroscopy as described recently (2). An aliquot of $\cdot\text{NO}$ -saturated aqueous solution was added to the suspension of *C. neoformans* in 200 mM sodium phosphate buffer (pH 7.0) or in 200 mM succinate buffer (pH 4.0). $\cdot\text{NO}$ (380 μM) without PTIO did not kill *C. neoformans* within a 24-h incubation period at pH 7.0. At pH 7.0, in the presence of PTIO, $\cdot\text{NO}$ (130 μM) did not induce fungicidal action within 8 min (Fig. 3), but after 3 h of incubation with 130 μM $\cdot\text{NO}$ and 200 μM PTIO, 92% of the cells were killed (data not shown). The viability of *C. neoformans* was not affected by either PTI (200 μM) or PTIO (500 μM) alone. At pH 4.0, $\cdot\text{NO}$ (130 μM) without PTIO did not induce cytotoxicity against *C. neoformans* within 8 min (Fig. 3), but after 3 h of incubation, 99% of the cells were killed (data not shown). In contrast, as demonstrated in Fig. 3, PTIO significantly enhanced fungicidal action under acidic conditions. Under these cytotoxic conditions generated by $\cdot\text{NO}$ plus PTIO, almost all $\cdot\text{NO}$ was converted to $\cdot\text{NO}_2$, as judged by electron

* Corresponding author.




FIG. 1. L-Arginine-nitrite-nitrate flow (11) and the use of PTIO to determine the biological significance of [•]NO. Until now, it has been impossible to distinguish the biological significance of [•]NO from that of its reactive derivatives by conventional reagents or methods (e.g., the nitric oxide synthase [NOS] inhibitor N^ω-monomethyl-L-arginine [L-NMMA], the [•]NO scavenger HbO₂ or its photometric assay, and measurements of nitrate and nitrite or L-citrulline). However, we did so by using PTIO. The pH-dependent nitrite cycle (20) was modified according to recent reports (5, 8, 13, 14, 17, 19).




FIG. 2. Fungicidal action against *C. neoformans* of a system chemically generating [•]NO. *C. neoformans*, serotype D, was added to either the reaction mixture containing 1 mM NaNO₂ in 80 mM succinate buffer (pH 4.0) with various concentrations of PTIO or to the succinate buffer alone. The number of viable cells after the indicated incubation periods was measured by the colony-forming assay. Each datum point represents the mean value for three experiments. Symbols: ○, 0 μM PTIO; ▲, 2 μM PTIO; □, 20 μM PTIO; ●, 200 μM PTIO; △, succinate buffer only.

spin resonance spectroscopy by our recently reported method (2) (data not shown).

The cytotoxic effects of various concentrations of [•]NO and PTIO were examined with different yeast strains at pH 4.0 (Fig. 4). Dose-dependent fungicidal action was observed with both [•]NO and PTIO.

The effects of [•]NO and PTIO on the viability of *Staphylococcus aureus* (ATCC 25923) were also examined, and a strong enhancement of [•]NO-dependent cytotoxicity similar to the enhancement of fungicidal action of PTIO (data not shown) was observed.

We compared the actions of PTIO with those of oxyhemoglobin (HbO₂), which entraps and oxidizes [•]NO as follows: [•]NO + HbO₂ → Hb⁺ + NO₃⁻ (11). HbO₂ was purified from healthy human erythrocytes as described previously (1). [•]NO (130 μM) killed 99% of the cells after incubation for 3 h at pH 4.0, and HbO₂ (100 or 300 μM) reduced the fungicidal action to 89 or 26%, respectively. These results can be interpreted as HbO₂ converting [•]NO to the stable compound, nitrate, thus bypassing the nitrite cycle, so that the fungicidal effects of toxic nitrogen oxide derivatives (Fig. 1) are no longer exerted.

The antimicrobial effect of the [•]NO-generating system has been known for years (3, 4, 6, 20). The mechanisms for this effect have been studied extensively, originally because of the worldwide use of nitrite in meat curing and now as a model of antimicrobial action of reactive nitrogen oxides within activated macrophages (3, 6). A pH-dependent nitrite cycle had been proposed, and the reactive nitrogen species responsible for antimicrobial action was suggested to be [•]NO, [•]NO₂, N₂O₃, N₂O₄, or HNO₂ (Fig. 1) (4, 6, 17, 20).




FIG. 3. Fungicidal action of authentic NO against *C. neoformans* in buffer (pH 7.0 or 4.0). NO -saturated solution was added to the suspension of *C. neoformans* in 200 mM sodium phosphate buffer (pH 7.0) or in 200 mM succinate buffer (pH 4.0) to yield 130 μM NO . The number of viable cells was measured by the colony-forming assay after the indicated incubation periods at room temperature. Symbols: \circ , 0 μM PTIO; \blacktriangle , 200 μM PTIO; \bullet , 500 μM PTIO. Each value shown represents the mean value for two experiments.

Recently, Stamler et al. (21) proposed the possible involvement of various redox-activated forms of nitrogen monoxides such as nitrosonium cation (NO^+) and nitroxyl anion (NO^-) in a number of pathophysiological phenomena induced by NO . N_2O_3 and N_2O_4 are known to possess potent chemical reactivities equivalent to that of NO^+ , which confers nitration of various nucleophiles (21). However, a good tool for distinguishing the biological reactivity of each nitrogen oxide involved in this cycle has not been available (4-6, 17, 20).

In this study, we used PTIO to identify the antimicrobial action of NO , and our results provide convincing evidence that (i) NO is not a fungicidal molecule and (ii) the NO -dependent fungicidal action was enhanced by PTIO. Therefore, it had been suggested that one or more of the nitrogen oxides (such as NO_2 , N_2O_4 , HNO_2 , and N_2O_3) generated in the nitrite cycle would be cytotoxic against various microbes, whereas we have demonstrated unequivocally that NO has little cytotoxic potency (Fig. 1). This conclusion is consistent with the observations of other researchers who have studied the cytotoxic activities of NO for mammalian cells and cryptococci (3, 8-10, 20-22).




FIG. 4. Dose-dependent fungicidal action of PTIO and NO . *C. neoformans*, serotype A or D, was incubated in 200 mM succinate buffer (pH 4.0) at room temperature with various concentrations of PTIO and/or NO , and the number of viable cells was measured by the colony-forming assay after 8-min incubation periods at room temperature. Each value shown represents the mean values for two experiments. Symbols: \blacktriangle , 0 μM NO ; \circ , 60 μM NO ; \bullet , 120 μM NO ; \square , 240 μM NO .

Previously, Kosaka et al. (13, 14) reported that mutagenic action of NO_2 , N_2O_4 , and N_2O_3 against both *Salmonella typhimurium* and *Escherichia coli* was more potent than that of NO ; NO had only negligible action against these bacteria. Although mutagenesis may not necessarily be related to direct cytotoxic action of nitrogen oxides, these observations seem to be consistent with our finding that NO_2 , N_2O_4 , and N_2O_3 would be the principal cytotoxic agents rather than NO . Recently, Rockett et al. (19) reported that a saturated solution of NO was not cytotoxic against *Plasmodium falciparum*. This finding also seems to be compatible with our results, except that they did not test the toxic effects of NO_2 , N_2O_4 , and/or N_2O_3 .

The cytotoxic potential of peroxyntirite (ONOO^-), which is formed with NO and O_2^- , has been another current focus in research on NO -dependent bactericidal action (12, 18, 23). However, as revealed in this study, it has become apparent that strong fungicidal activity could be exhibited through the NO oxidation process without the involvement of ONOO^- generation. Therefore, it is crucially important to clarify the involvement of toxic nitrogen oxides (NO_2 , N_2O_4 , N_2O_3 , and HNO_2) other than NO and ONOO^- (ONOOH) in the antimicrobial activity of activated macrophages.

This work was supported by a Grant-in-Aid for Scientific Research from Monbusho (Ministry of Education, Science and Culture) of Japan (to T. Akaike and H. Maeda), by funds from the Okukubo Memorial Fund for Medical Research in Kumamoto University School of Medicine (to T. Akaike), and by a grant from Yakult Honsha Co., Tokyo, Japan.

We thank Judith B. Gandy for editorial work and R. Yoshimoto for typing the manuscript. We also thank Hiroaki Kosaka, Department of Physiology, Osaka University Medical School, for stimulating discussions.

REFERENCES

1. Akaike, T., K. Sato, S. Ijiri, Y. Miyamoto, M. Kohno, M. Ando, and H. Maeda. 1992. Bactericidal activity of alkyl peroxy radicals generated by heme-iron-catalyzed decomposition of organic peroxides. *Arch. Biochem. Biophys.* **294**:55-63.
2. Akaike, T., M. Yoshida, Y. Miyamoto, K. Sato, M. Kohno, K. Sasamoto, K. Miyazaki, S. Ueda, and H. Maeda. 1993. Antagonistic action of imidazolineoxyl *N*-oxides against endothelium-derived relaxing factor/NO through a radical reaction. *Biochemistry* **32**:827-832.
3. Alspaugh, J. A., and D. L. Granger. 1991. Inhibition of *Cryptococcus neoformans* replication by nitrogen oxides supports the role of these molecules as effectors of macrophage-mediated cytostasis. *Infect. Immun.* **59**:2291-2296.
4. Castellani, A. G., and C. F. Niven, Jr. 1955. Factors affecting the bacteriostatic action of sodium nitrite. *Appl. Microbiol.* **3**:154-159.
5. Challis, B. C. 1981. The chemistry of formation of N-nitroso compounds, p. 16-55. In G. G. Gibson, and C. Ioannides (ed.), *Safety evaluation of nitrosatable drugs and chemicals*. Taylor & Francis, Ltd., London.
6. Chan, B. J., Y. Xing, R. S. Magliozzo, and B. R. Bloom. 1992. Killing of virulent *Mycobacterium tuberculosis* by reactive nitrogen intermediates produced by activated murine macrophages. *J. Exp. Med.* **175**:1111-1122.
7. Curran, R. D., F. K. Ferrari, P. H. Kispert, J. Stadler, D. J. Stuehr, R. L. Simmons, and T. R. Billiar. 1991. Nitric oxide and nitric oxide-generating compounds inhibit hepatocyte protein synthesis. *FASEB J.* **5**:2085-2092.
8. Douglass, M. L., B. L. Kabakoff, G. A. Anderson, and M. C. Cheng. 1978. The chemistry of nitrosamine formation, inhibition and destruction. *J. Soc. Cosmet. Chem.* **29**:581-606.
- 8a. Drapier, J.-C. 1991. L-Arginine-derived nitric oxide and the cell-mediated immune response. *Res. Immunol.* **142**:551-602.
9. Drapier, J.-C., and J. B. Hibbs, Jr. 1986. Murine cytotoxic activated macrophages inhibit aconitase in tumor cells. Inhibition involves the iron-sulfur prosthetic group and is reversible. *J. Clin. Invest.* **78**:790-797.
10. Granger, D. L., J. B. Hibbs, Jr., J. R. Perfect, and D. T. Durack. 1990. Metabolic fate of L-arginine in relation to microbiostatic capability of murine macrophages. *J. Clin. Invest.* **85**:264-273.
11. Ignarro, L. J. 1990. Biosynthesis and metabolism of endothelium-derived nitric oxide. *Annu. Rev. Pharmacol. Toxicol.* **30**:535-560.
12. Koppenol, W. H., J. J. Moreno, W. A. Pryor, H. Ischiropoulos, and J. S. Beckman. 1992. Peroxynitrite, a cloaked oxidant formed by nitric oxide and superoxide. *Chem. Res. Toxicol.* **5**:834-842.
13. Kosaka, H., Y. Oda, and M. Uozumi. 1985. Induction of *umuC* gene expression by nitrogen dioxide in *Salmonella typhimurium*. *Mutat. Res.* **142**:99-102.
14. Kosaka, H., K. Yamamoto, Y. Oda, and M. Uozumi. 1986. Induction of SOS functions by nitrogen dioxide in *Escherichia coli* with different DNA-repair capacities. *Mutat. Res.* **162**:1-5.
15. Lepoivre, M., J.-M. Flaman, and Y. Henry. 1992. Early loss of the tyrosyl radical in ribonucleotide reductase of adenocarcinoma cells producing nitric oxide. *J. Biol. Chem.* **267**:22994-23000.
16. Levitz, S. M., and D. J. DiBenedetto. 1988. Differential stimulation of murine resident peritoneal cells by selectively opsonized encapsulated and acapsular *Cryptococcus neoformans*. *Infect. Immun.* **56**:2544-2551.
17. Marletta, M. A., P. S. Yoon, R. Iyengar, C. D. Leaf, and J. S. Wishnok. 1988. Macrophage oxidation of L-arginine to nitrite and nitrate: nitric oxide is an intermediate. *Biochemistry* **27**:8706-8711.
18. Moreno, J. J., and W. A. Pryor. 1992. Inactivation of α_1 -proteinase inhibitor by peroxynitrate. *Chem. Res. Toxicol.* **5**:425-431.
19. Rockett, K. A., M. M. Awburn, W. B. Cowden, and I. A. Clark. 1991. Killing of *Plasmodium falciparum* in vitro by nitric oxide derivatives. *Infect. Immun.* **59**:3280-3283.
20. Shank, J. L., J. H. Silliker, and R. H. Harper. 1962. The effect of nitric oxide on bacteria. *Appl. Microbiol.* **10**:185-189.
21. Stamler, J. S., D. J. Singel, and J. Loscalzo. 1992. Biochemistry of nitric oxide and its redox-activated forms. *Science* **258**:1898-1902.
22. Stuehr, D. J., and C. F. Nathan. 1989. A macrophage product responsible for cytostasis and respiratory inhibition in tumor target cells. *J. Exp. Med.* **169**:1543-1555.
23. Zhu, L., C. Gunn, and J. S. Beckman. 1992. Bactericidal activity of peroxynitrite. *Arch. Biochem. Biophys.* **298**:452-457.