
A rabies serosurvey of domestic dogs in rural Tanzania: results of a rapid fluorescent focus inhibition test (RFFIT) and a liquid-phase blocking ELISA used in parallel

S. CLEAVELAND^{1,2*}, J. BARRAT³, M.-J. BARRAT³, M. SELVE³, M. KAARE⁴
AND J. ESTERHUYSEN⁵

¹ *Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK*

² *Institute of Zoology, Zoological Society of London, Regent's Park, London NW1 4RY, UK*

³ *Centre National d'Études Vétérinaires et Alimentaires (CNEVA), Boîte Postale 9, Domaine de Pixérécourt, Malzeville 54220, France*

⁴ *Ministry of Agriculture, P.O. Box 162, Mugumu, Serengeti District, Tanzania*

⁵ *Onderstepoort Institute for Exotic Diseases, Private Bag X6, Onderstepoort 0110, South Africa*

(Accepted 9 March 1999)

SUMMARY

During a serosurvey of domestic dogs in Tanzania, a rapid fluorescent focus inhibition test (RFFIT) and a liquid-phase blocking ELISA (LPBE) were used to measure rabies antibodies in vaccinated and unvaccinated dogs. Post-vaccination titres measured by LPBE correlated closely with those found by RFFIT. Of 567 unvaccinated dogs tested using the LPBE, 42 (7·4%) were seropositive, with titres exceeding 32. Of this group, 233 dogs were tested using the RFFIT and 115 (49·4%) were seropositive, with titres exceeding 0·5 IU/ml. Two lines of evidence pointed to the greater specificity of the LPBE when measuring rabies antibodies induced by natural infections: (a) no seropositive dogs were detected among the 162 unvaccinated dogs from the rabies-free island of Pemba, Tanzania, when using LPBE, whereas 15/145 (10·3%) dogs of the same group were seropositive using RFFIT; (b) among Tanzanian dogs there was a close association between the location of rabies cases and location of seropositive dogs when using LPBE, but not when using RFFIT. These results suggest that LPBE may be of value in rabies seroepidemiological studies and could be developed as a reference technique for the detection of rabies antibody in domestic dogs.

INTRODUCTION

Throughout much of sub-Saharan Africa, case incidence data on canine rabies have often been fragmentary and official figures reported to the World Health Organization (WHO) provide a poor indication of both the magnitude of the rabies problem and the trends in disease incidence. In Tanzania, for

example, a total of only 42 cases of dog rabies were officially reported from 1990–7 [1, 2] despite unofficial reports that the disease is a widespread problem throughout the country. With so few rabies cases reported, there has clearly been a need to develop epidemiological approaches that do not rely on case incidence data. The aim of this investigation was therefore to explore the use of serological assays for epidemiological studies of rabies in rural African domestic dog populations.

Serological surveys are widely used to investigate the dynamics of viral infections in natural pop-

* Author for correspondence. Current address: Centre for Tropical Veterinary Medicine, Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush, Roslin, Midlothian EH25 9RG, UK.

ulations. However, serosurveys have generally been considered to be of little value for the study of rabies epidemiology because, classically, antibodies are detected only in the terminal stages of infection and shortly before death. Serological tests have therefore been developed principally to measure post-vaccination antibody titres in people and animals and not to detect antibodies induced by natural infections.

The general explanation for the late appearance of antibodies in rabies infections is that the intrinsic neurotropism of the rabies virus protects it from immune surveillance and virus must multiply in the central nervous system before sufficient antigen is released to induce serum neutralizing antibody [3]. This assumption is challenged by the detection of rabies antibody in a proportion of healthy domestic dogs [4–9] and wild animals [5, 10–21] in rabies-endemic areas in various parts of the world. However, interpretation of such serological data is problematic because of the wide range of assays adopted, the variation in cut-off points and, in many cases, the lack of validation of test specificity for the species in question.

Although several rabies serological tests have been described [22], at the time of this study, only the mouse neutralization test (MNT) [23] and the rapid fluorescent focus inhibition test (RFFIT) [24] were recognized as reference techniques by the WHO. These recommendations were based on the fact that both tests measure inhibition of infectivity, a biological property usually correlated with protection. However, both tests have practical drawbacks, requiring specialized facilities, use of infective virus, several days to complete and, in the case of the MNT, large numbers of mice. A further problem with the RFFIT has been a lack of consistency between laboratories as a result of numerous modifications to the original RFFIT protocol [25, 26].

Enzyme-immunosorbent assays (ELISA) are gaining increasing popularity as a rapid and sensitive method of detecting rabies antibody. Several new techniques have been described using competitive systems [27] or enzyme-linked protein A [28] to allow sera of all species to be tested. The development of competitive assays that use monoclonal antibodies [29] also overcomes the need to produce highly purified antigen. Despite these advances, no ELISA has yet been validated as a reference technique for measuring rabies antibodies in dogs.

In this paper, we compare results of a rabies serological survey in rural domestic dog populations

in Tanzania that were obtained using a RFFIT and a liquid-phase blocking ELISA (LPBE). Given the uncertainties about the interpretation of rabies seropositivity in natural infections, we adopted two approaches to assess the specificity of our results. The first was to compare titres of Tanzanian mainland dogs with those from a negative control population on the rabies-free island of Pemba, Tanzania [1]. Previous studies have addressed the specificity of RFFIT and LPBE through analysis of sera from dogs in rabies-free countries, such as UK, Mauritius and Hawaii [30, 31]. However, none of these countries provides an ideal negative control group for Tanzanian dogs because cross-reactivity to other Lyssavirus genotypes, such as Lagos Bat and Mokola virus [32], cannot be ruled out.

In the second analysis, we adopt an epidemiological approach to the assessment of test specificity, investigating the association between the location of rabies seropositive dogs and the location of rabies cases. We assume that, if non-specific or cross-reactions are the cause of rabies seropositivity in dogs, we would not expect to find a significant association between the location of seropositive dogs and rabies cases. The results are discussed with respect to the potential value of these assays for rabies seroepidemiological studies in natural populations. The significance of these findings in terms of rabies epidemiology has been discussed elsewhere [31].

MATERIALS AND METHODS

Field studies

Blood sample collection from unvaccinated dogs

Blood samples were collected from 567 domestic dogs from two Districts in north-western Tanzania; Ngorongoro District to the east of the Serengeti National Park (35°–36° E, 1° 30′–3° 7′ S), and Serengeti District to the west of the park [33]. Blood samples were collected from September–December 1993 and from August–December 1994 during systematic surveys of households within 19 villages bordering the park (Fig. 1). Details of household surveys are given elsewhere [33]. Dogs were manually restrained and muzzled while blood was collected from the cephalic vein. Blood samples were centrifuged within 24 h of collection, serum was stored for up to 5 months at between –5 °C and –20 °C and transported to laboratories on dry ice.




Fig. 1. Map of the study area showing location of villages (dark circles) where domestic dogs were sampled. Ngorongoro crater is shown by the large open circle.

Survivorship studies

In 1994, 76/193 (39.3%) households visited in 1993 were re-visited to determine survival of seropositive and seronegative dogs sampled previously. If a dog had died, information was collected from the owner on the date of death, clinical history and whether rabies was a suspected cause of death. Although results relied on owners identifying rabies as a cause of death, the disease is well-recognized in rural Tanzania. In this study area, villagers were able to detect rabies in domestic dogs with a diagnostic specificity of 83.3% and sensitivity of 86.5%, based on an analysis

of 139 cases where dogs had died and samples had been sent for laboratory diagnosis.

Post-vaccination samples

From September–December 1993, 240 dogs from 6 villages were vaccinated against rabies, by a single subcutaneous injection of 1 ml Rabisin (Rhône-Poulenc, Nairobi). Ninety-one vaccinated dogs were re-bled between 28–31 days after vaccination to assess post-vaccinal titres; 62 vaccinated dogs were also re-bled 1 year after vaccination.

Rabies case incidence data

Records of reported rabies cases for the three study areas were obtained from veterinary offices in Mugumu (Serengeti District), Loliondo and Ngorongoro (Ngorongoro District). Suspected rabies cases were reported by village extension officers stationed in each village. Wherever possible, samples from suspect rabies cases were examined for rabies antigen using immunofluorescence diagnostic tests [34] carried out on brain stem samples collected in WHO collection kits containing 50% glycerol-saline with 0.01% methiolate as preservative [35]. Tests were carried out at the WHO Collaborating Centre for Zoonoses, Centre National d'Études Vétérinaires et Alimentaires (CNEVA), France.

Negative control population

From November–December 1996, 166 serum samples were collected from domestic dogs on Pemba by sampling every household within four rural villages, Fundo Island, Kizambani, Mchanga Mdogo, Gombani, and one urban location Bahrein/Masipa. Samples were obtained and treated as described above.

Serological tests*Liquid phase blocking ELISA*

A total of 729 serum samples were analysed with the liquid phase blocking ELISA (LPBE) at the Onderstepoort Institute for Exotic Diseases, South Africa, using a technique adapted for the measurement of antibodies to foot-and-mouth disease virus [36, 37]. Details of the protocol and methods of calculating serum titres have been described fully elsewhere [27]. In this study, the protocol differed from that described previously only in the use of CVS strain of rabies virus as test antigen in place of Flury HEP virus. Sera were initially screened at dilutions 1:16–1:128. Any serum showing a titre > 1:16 was re-titrated at least once using dilutions from 1:16–1:1024 and the geometric mean titre adopted as the value. Titres were expressed as the \log_{10} of the reciprocal dilution (referred to as log dilution). Previous studies have demonstrated specificity of the LPBE at a threshold > log dilution 1.5 [27, 31] and we adopt this cut-off point to define seropositivity in this analysis.

Rapid Fluorescent Focus Inhibition Test (RFFIT)

A modification of the original RFFIT technique [24] was used to measure neutralizing antibodies to rabies

virus in 378 serum samples. This technique was developed at CNEVA, Nancy, using an adaptation for use of microtitre plates, based on the techniques of Zalan and colleagues [38] and Perrin and colleagues [39]. The protocol is described in detail elsewhere [30]. The titre of the serum was determined as the dilution at which 50% of fluorescence was inhibited (D_{50}) according to the Spearman–Karber method [40]. The antibody titre in International Units (IUs)/ml was determined by comparison with the WHO standard immunoglobulin. Titres were also expressed as the \log_{10} of the reciprocal D_{50} (referred to as log dilution) for comparison with LPBE titres. In this study, we use the international reference threshold of 0.5 IU/ml (equivalent to log dilution 1.5) to define seropositivity.

Data analysis

Frequency distributions of Serengeti and negative control populations were compared by grouping values into four categories in contingency tables and analysed using a χ^2 test. Associations between the location of seropositive dogs and the location of rabies cases were investigated by calculating odds ratios (OR) expressed with 95% confidence limits using Epi-Info 5.0 [41].

RESULTS**Sample sizes**

The number of serum samples tested with each serological test is shown in Table 1. Eleven dogs were sampled in both 1993 and 1994, and sera analysed using the LPBE. Of these dogs, only one sample was included for each individual to avoid pseudo-replication. There was no significant difference in the distribution of titres that included only the first samples and the distribution of titres that included only the second samples ($\chi^2_{(3)} = 0.04$, $P > 0.05$). For consistency, results including the first sample only are used in subsequent analyses.

Frequency distributions

The frequency distributions of LPBE values are shown for Serengeti and Pemba dogs in Figure 2*a*. The distribution of titres of Pemba dogs differed significantly from the distribution of Serengeti dogs ($\chi^2_{(3)} = 12.7$, $P < 0.01$). None of 162 Pemba dogs was seropositive. In the Serengeti, 21/286 (7.34%) unvaccinated dogs were seropositive in 1993 and 21/281 (7.47%) seropositive in 1994.

Table 1. The number of serum samples analysed by each of the serological tests

Origin of samples	LPBE (Onderstepoort)	RFFIT (CNEVA, Nancy)	Samples analysed on both LPBE and RFFIT
Serengeti, 1993	286	134	126
Serengeti, 1994	281	99	90
Pemba, 1996	162	145	141




Fig. 2. Distribution of rabies antibody in unvaccinated domestic dogs from Pemba and Serengeti measured by (a) liquid phase blocking ELISA (LPBE) and (b) rapid fluorescent focus inhibition test (RFFIT).

The frequency distributions of RFFIT titres from unvaccinated dogs is shown in Figure 2b. The distribution of Pemba dogs was significantly different from Serengeti dogs ($\chi^2_{(3)} = 56.8, P < 0.001$). 15/145 (10.34%) dogs from Pemba were seropositive. In 1993, 71/134 (53.0%) Serengeti dogs were seropositive and in 1994, 44/99 (44.4%) Serengeti dogs were seropositive.

Survivorship

Six of 14 (42.9%) dogs that were defined as seropositive by the LPBE were alive 1 year later. There was no significant difference in survival between




Fig. 3. Correlation between LPBE and RFFIT titres in (a) unvaccinated Serengeti domestic dogs and (b) vaccinated Serengeti domestic dogs.

seropositive and seronegative dogs ($\chi^2_{(1)} = 1.28, P > 0.05, n = 135$). No seropositive dogs were reported to have died with clinical signs consistent with rabies.

Of 32 dogs defined as seropositive by the RFFIT, 19 (59.4%) were alive 1 year later. Survival of seropositive and seronegative dogs was very similar ($\chi^2_{(1)} = 0, P > 0.05$). No seropositive individuals were reported to have died with clinical signs consistent with rabies.

Correlation analysis

There was only a weak correlation between LPBE and RFFIT titres in unvaccinated Serengeti dogs ($r = 0.18, n = 216, P < 0.05$) (Fig. 3a). There was a much closer agreement between LPBE and RFFIT titres in

vaccinated dogs ($r = 0.84$, $n = 131$, $P < 0.001$) (Fig. 3*b*).

Geographical associations

Dogs defined as seropositive on the LPBE were significantly more likely to be found in villages where rabies had been confirmed during the past 12 months than in villages where it had not; OR = 5.16 (1.45–18.05), $P < 0.01$, $n = 216$. In contrast, there was no significant association between the location of RFFIT seropositives and confirmed rabies cases; OR = 1.26 (0.53–2.97), $P > 0.05$, $n = 216$.

DISCUSSION

The results of this study demonstrate that a proportion of unvaccinated domestic dogs from a rabies-endemic area of Africa had detectable levels of rabies antibodies when measured by a serum neutralization test (RFFIT) and by a liquid-phase blocking ELISA (LPBE). No seropositive dogs were reported to have died of suspect rabies and a proportion remained alive for at least a year. Our findings thus suggest that seropositive dogs were unlikely to be incubating rabies, but that seropositivity was consistent with aborted infection [42]. Although this outcome differs from that classically described for rabies, much of our understanding about the pathogenesis of rabies is derived from experimental, not natural, infections. In natural populations, the potential variation in infection routes, infective dose and virus characteristics may lead to a broader spectrum of outcomes than occurs in most experimental situations.

In this study, uncertainties arose in the interpretation of seropositivity because the two serological tests identified different dogs as seropositive. A key issue was therefore to investigate the relative specificity of the LPBE and the RFFIT to assess whether seropositivity reflected genuine exposure to rabies virus. Results of both analyses suggested that the LPBE was the more specific test when detecting rabies antibody induced by natural infection. Thus, fewer Pemba (rabies-negative) dogs were defined as seropositive ('false positives') when using the LPBE (0%) than when using the RFFIT (10.3%). Furthermore, the location of LPBE seropositive dogs was significantly associated with villages where rabies had been confirmed, whereas there was no significant association between the location of RFFIT seropositive dogs and rabies cases.

The close correlation between the location of LPBE seropositive dogs and rabies cases demonstrates the potential value of this technique for detecting rabies infection in unvaccinated populations. We have previously shown that rabies antibodies induced by natural infection are short-lived [31], which raises the possibility that seropositivity may also provide a useful comparative measure of the incidence of infection. The application of rabies serology may be of particular value in areas where incidence data are difficult to obtain, as occurs throughout much of rural Africa. Even in areas where active surveillance measures have significantly improved rabies detection rates [43], the high costs involved may favour rabies serology as a more cost-effective alternative for monitoring epidemiological trends. Still required, however, are more detailed longitudinal data comparing trends in both disease incidence and rabies seropositivity.

For wildlife populations, in which only a small fraction of rabies cases are ever detected, sero-epidemiological approaches clearly also have potential value. However, the conclusions of our study regarding the specificity of rabies seropositivity in natural infections applies only to domestic dogs and cannot be freely extrapolated to other species. Even where serological tests have been validated for wildlife populations, seroprevalence data may be difficult to interpret. It is likely that seropositivity in wildlife populations varies both with host species and virus strain. In highly susceptible hosts, such as red foxes and jackals, only very low seroprevalences have been recorded [10, 11, 13, 44], presumably because most infected animals die of the disease. However, virus strain also plays a role; in experimental infections, for example, red foxes infected with domestic dog isolates develop an immune response more often than when infected with red fox isolates [45].

We cannot explain why RFFIT and LPBE tests produce similar results when measuring antibody induced by a known dose of immunogenic antigen (rabies vaccine), but showed a very poor correlation when measuring antibody resulting from possible natural exposure. This is not just a matter of titre range, since unvaccinated dogs with high titres on one test had low titres on the other, whereas even low post-vaccinal titres correlated more closely. These results may simply reflect variability in the test systems, but biological differences in anti-N (detected by the LPBE) or anti-G responses (detected by the RFFIT) in non-lethal infections could also give rise to

discrepancies in titre. To our knowledge, however, there are no data comparing the kinetics of anti-G and anti-N antibody in infection.

In summary, the results of this study demonstrate the existence of rabies seropositivity in a proportion of healthy, unvaccinated domestic dogs in rural Africa and support the view that the liquid phase blocking ELISA is more specific than the RFFIT for detection of rabies antibody in unvaccinated dogs. ELISAs are intrinsically more easily standardized among laboratories than neutralization tests and allow simple and more rapid large-scale processing of sera. These practical advantages, coupled with the relatively high specificity demonstrated in this study, should encourage the development and validation of ELISAs for use in rabies epidemiological studies and as a reference technique for the detection of rabies antibody in domestic dogs.

ACKNOWLEDGEMENT

SC was supported by grants from the BBSRC and the Department for International Development (formerly ODA/NRRD Animal Health Programme). We thank Tanzania National Parks, Ngorongoro Conservation Area Authority and Serengeti Wildlife Research Institute for permission to work in the Serengeti and Dr F. Gulland, Dr S. Mkumbo, Dr P. Loomu, Mr P. Tiringa, Mr Mosha and Mr Magoto for field assistance. The Frankfurt Zoological Society and British Development Division, East Africa provided logistic support for which we are most grateful. Dr Chris Dye provided valuable advice and comments on earlier drafts of the manuscript.

REFERENCES

1. National Rabies Control Programme. Dar es Salaam: Tanzania Ministry of Agriculture and Cooperatives, 1994.
2. Loretu K. Rabies in Tanzania. In: King AA, ed. Proceedings of the International Conference on Epidemiology, Control and Prevention of Rabies in Eastern and Southern Africa, Lusaka, Zambia, 1992: Fondation Marcel Merieux, Lyon, 1993: 17–18.
3. King AA, Turner GS. Rabies: a review. *J Comp Path* 1993; **108**: 1–39.
4. Andral L, Serie C. Études expérimentales sur la rage en Ethiopie. *Ann Inst Pasteur* 1957; **93**: 475–88.
5. Mebatsion T, Frost JW, Krauss H. ELISA using staphylococcal protein A for the measurement of rabies antibody in various species. *J Vet Med B* 1989; **36**: 532–6.
6. Yasmuth C, Nelson KE, Laima T, Supawadee J, Thaiyanant P. Prevalence of abortive canine rabies in Chiang Mai, Thailand. *J Med Assoc Thailand* 1983; **66**: 169–75.
7. Ogunkoya AB, Beran GW, Umoh JU, Gomwalk NE, Abdulkadir A. Serological evidence of infection of dogs and man in Nigeria by lyssaviruses (family Rhabdoviridae). *Trans Roy Soc Trop Med Hyg* 1990; **84**: 842–5.
8. Alexander KA, Smith JS, Macharia MJ, King AA. Rabies in the Masai Mara, Kenya; preliminary report. *Ond J Vet Res* 1993; **60**: 411–4.
9. Laurenson K, Esterhuysen J, Stander P, van Heerden J. Aspects of rabies seroepidemiology in Tsumkwe District, Namibia. *Ond J Vet Res* 1997; **64**: 39–45.
10. Carey AB, McLean RG. Rabies antibody prevalence and virus tissue tropism in wild carnivores in Virginia. *J Wild Dis* 1978; **14**: 487–91.
11. Foggini CM. Rabies and rabies-related viruses in Zimbabwe: historical, virological and ecological aspects [PhD thesis]. Harare, Zimbabwe: University of Zimbabwe, 1988.
12. Gascoyne SC, King AA, Laurenson MK, Borner M, Schildger B, Barrat J. Aspects of rabies infection and control in the conservation of the African wild dog (*Lycaon pictus*) in the Serengeti Region, Tanzania. *Ond J Vet Res* 1993; **60**: 415–20.
13. Wandeler AI, Wachendörfer G, Förster U, Krekel H, Müller J, Steck F. Rabies in wild carnivores in central Europe II. Virological and serological examinations. *Zbl Vet Med B* 1974; **21**: 757–64.
14. McLean RG. Raccoon rabies. In: Baer GM, ed. The natural history of rabies, vol. 2. New York: Academic Press, 1991: 53–77.
15. Bigler WJ, Hoff GL, Smith JS, McLean RG, Trevino HA. Persistence of rabies antibody in free-ranging raccoons. *J Infect Dis* 1983; **148**: 610.
16. Jenkins SR, Perry BD, Winkler WG. The ecology and epidemiology of raccoon rabies. *Rev Infect Dis* 1988; **10**: S620–5.
17. Ferguson DV, Heidt GA. Survey for rabies, leptospirosis, toxoplasmosis and tularemia in striped skunks (*Mephitis mephitis*) from three public use areas in northwestern Arkansas. *J Wild Dis* 1981; **17**: 515–9.
18. Carey AB, McLean RG. The ecology of rabies: evidence of co-adaptation. *J App Ecol* 1983; **20**: 777–800.
19. Rosatte RC, Gunson JR. Presence of neutralizing antibodies to rabies virus in striped skunks from areas free of skunk rabies in Alberta. *J Wild Dis* 1984; **20**: 171–6.
20. Everard COR, Baer GM, James A. Epidemiology of mongoose rabies in Grenada. *J Wild Dis* 1974; **10**: 190–6.
21. Everard COR, Baer GM, Alls ME, Moore, SA. Rabies serum neutralizing antibody in mongooses from Grenada. *Trans Roy Soc Trop Med Hyg* 1981; **75**: 654–66.
22. Smith J. Rabies serology. In: Baer GM, ed. The natural history of rabies, 2nd ed. Boca Raton: CRC Press, 1991: 235–52.

23. Atanasiu P. Quantitative assay and potency test of antirabies serum. In: Kaplan MM, Koprowski H, eds. *Laboratory techniques in rabies*, 2nd ed. Geneva: World Health Organization, 1966: 167–72. (WHO Monograph Series: no. 23).
24. Smith JS, Yager PA, Baer GM. A rapid reproducible test for determining rabies neutralizing antibody. *Bull WHO* 1973; **48**: 535–41.
25. Cliquet F, Aubert M, Sagne F. Comparison of the virus neutralization protocols in cells used by the participating laboratories. In: *Proceedings of the workshop on rabies serum neutralization tests in European laboratories*. Nancy, France: WHO/OIE Collaborating Centre, Centre National d'Études Vétérinaires et Alimentaires, 1995.
26. Barrat J, Cleaveland S, Cliquet F, Esterhuysen J, Aubert M. Rabies sero-antibody measurement. In: Bingham J, Bishop GC, King AA, eds. *Proceedings of the 3rd International Meeting of the Southern and Eastern African Rabies Group*, Harare, March 1995. Harare, Zimbabwe: Southern and Eastern African Rabies Group, 1995; 178–83.
27. Esterhuysen JJ, Prehaud C, Thomson GR. A liquid-phase blocking ELISA for the detection of antibodies to rabies virus. *J Virol Meth* 1995; **51**: 31–42.
28. Mebatsion T, Sillero-Zubiri C, Gottelli D, Cox JH. Detection of rabies antibody by ELISA and RFFIT in unvaccinated dogs and in the endangered Simien jackal (*Canis simensis*) of Ethiopia. *J Vet Med B* 1992; **39**: 233–5.
29. Elmgren LD, Wandeler AI. Competitive ELISA for the detection of rabies virus-neutralizing antibodies. In: Meslin FX, Kaplan MM, Koprowski H, eds. *Laboratory techniques in rabies*, 4th ed. Geneva, Switzerland: World Health Organization, 1996: 200–8.
30. Cliquet F, Aubert M, Sagne L. Development of a fluorescent antibody virus neutralisation test (FAVN) for the quantitation of rabies-neutralising antibody. *J Imm Meth* 1998; **212**: 79–87.
31. Cleaveland S, Dye C. Maintenance of a microparasite infecting several host species: rabies in the Serengeti. *Parasitology* 1995; **111**: S33–S47.
32. King AA, Meredith CD, Thomson GR. The biology of Southern African Lyssavirus variants. In: Rupprecht CE, Dietzschold B, Koprowski H, eds. *Lyssaviruses*. Berlin: Springer-Verlag, 1994: 267–295.
33. Cleaveland S. The epidemiology of rabies and canine distemper in the Serengeti, Tanzania [PhD Thesis], London: University of London, 1996.
34. Kaplan MM, Koprowski H. *La rage – techniques de laboratoire*, 3rd ed. Geneva, Switzerland: World Health Organization, 1973: 75–87 (Monograph Series no. 23).
35. Barrat J, Blancou J. Simplified technique for the collection, storage and shipment of brain specimens for rabies diagnosis. Geneva, Switzerland: World Health Organization, 1988 (WHO Rabies Research 88.27).
36. McCullough KC, Crowther JR, Butcher RN. A liquid-phase ELISA and its use in the identification of epitopes of foot-and-mouth disease virus antigens. *J Virol Meth* 1985; **11**: 39–338.
37. Hamblin C, Barnett ITR, Hedger RS. A new enzyme-linked immunosorbent assay (ELISA) for the detection of antibodies against foot-and-mouth disease virus. I. Development and method of ELISA. *J Immunol Meth* 1986; **93**: 115–21.
38. Zalan E, Wilson C, Pukitis D. A microtest for the quantitation of rabies virus neutralizing antibodies. *J Biol Stand* 1979; **7**: 213–220.
39. Perrin P, Lafon M, Versmisse P, Sureau P. Application d'une methode immunoenzymatique au titrage des anticorps rabiques neutralisants en culture cellulaire. *J Biol Stand* 1985; **13**: 35–42.
40. Lorenz RG, Bogel K. Calculation of titres and their significance. A. Spearman-Kärber, B. Method of Reed and Muench, C. Infectivity titration based on plaque counts. In: Kaplan MM, Koprowski H, eds. *Laboratory techniques in rabies*, 3rd ed. Geneva, Switzerland: World Health Organization, 1973: 321–35.
41. Dean AG, Dean JA, Burton AH, et al. *Epi Info Version 5: A word processing, database and statistics program for epidemiology on microcomputers*. Stone Mountain, Georgia, USA: USD, Incorporated, 1990.
42. Fekadu M. Latency and aborted rabies. In: Baer GM, ed. *The natural history of rabies*, 2nd ed. Boca Raton: CRC Press, 1991: 191–8.
43. Kitala PM, McDermott JJ, Kyule MN, et al. Investigation of rabies incidence and control by active surveillance in Machakos District, Kenya, Kenya Veterinarian 1994; **18**: 528–30.
44. Baradel JM, Barrat J, Blancou J, et al. Results of a serological survey of wild mammals in France. *Rev Sci Tech Off Int Epi* 1988; **7**: 873–83.
45. Blancou J. Ecology and epidemiology of fox rabies. *Rev Infect Dis* 1988; **10**: S606–14.