

---

## Levels of virulence are not determined by genomic lineage of *Salmonella enterica* serotype Enteritidis strains

---

J. E. OLSEN\*, T. TIAINEN AND D. J. BROWN

Department of Veterinary Microbiology, The Royal Veterinary and Agricultural University, Stigbøjlen 4, DK-1870 Frederiksberg C., Denmark

(Accepted 16 August 1999)

### SUMMARY

Mouse virulence and the ability to adhere to, and invade cultured MDCK cells were investigated in 38 phage type reference strains of *Salmonella enterica* serotype Enteritidis and correlated with genomic lineage. The genomic lineage of 11 of the strains was determined in the present study; one IS200 and one ribotype pattern that had not been reported previously were observed. Log c.f.u. in the spleen 10 days post intraperitoneal (i.p.) infection with  $3 \times 10^3$  bacteria (log<sub>10</sub>VC10) varied between 2.9 and 8.7. The reference strains of PT7 and PT23 were found to be semi-rough and were of low virulence. All other strains possessed smooth LPS. Within each of the two major clonal lines, as well as among phage types outside these, both highly virulent and moderate to low virulent strains were present. While all strains of PT1, PT2 and PT8 were highly virulent, low virulent strains were detected in PT4 and PT13. The ability to adhere to, and invade MDCK cells varied between phage types (adherence between 13 and 61% of the inocula and invasion between 4 and 151% of the adherent cells). The results of the cell culture experiments did not correlate with the results of mouse virulence tests. No correlation between clonal lineage and virulence was found within *S. Enteritidis*. It seems most likely that some strains have lost some of the essential factors enabling this serotype to cause successful systemic infection.

### INTRODUCTION

*Salmonella enterica* serotype Enteritidis is the most common cause of human salmonellosis in many countries [1], often associated with consumption of eggs and poultry products [2, 3]. In chickens, *S. Enteritidis* does not normally cause disease, although internal organs, such as liver, spleen, heart and gallbladder may be colonized at a low level for a short period. Of particular significance to the epidemiology of human infections is the fact that the intestine and cloaca of infected birds remains culture-positive for a prolonged period, and that oviduct tissues may be

colonized [4], increasing the risk of crop contamination during slaughter and the production of shell-contaminated, as well as internally contaminated eggs.

Strains of *S. Enteritidis* may differ with respect to virulence [5–7]. Within phage type 4, which has been dominating in Europe [1], virulence variation has been linked to the possession of a virulence plasmid [8, 9], as well as to tolerance of a number of environmental conditions [10]. Recently, strains isolated from clinical cases were shown to have lower LD<sub>50</sub> in chicken than strains isolated from the environment or from food products [11].

From a public health point of view, it is essential to be able to differentiate between virulent and avirulent

\* Author for correspondence.

Table 1. *Mouse virulence and LPS-profile of strains of S. Enteritidis in relation to clonal line*

Strain	Phage type	Clonal line*	LPS R/S†	Ribotype‡	LogVC10	$\bar{x}$ LogVC10
E2341	1	I	S	I	8.3 ± 1.3	7.5 ± 1.4**
P66040	3	I	S	III	8.3 ± 0.4	
E2187	4	I	S	II	8.3 ± 0.7	
P99764	4a	I	S	II	8.0 ± 1.3	
P70001	5	I	S	I	8.5 ± 0.5	
Ent5a	5a§	I	S	II	n.d.	
P99327	6	I	S	II	5.1 ± 0.9	
E2408	6a	I	S	I	5.7 ± 0.6	
E1492	7	I	R	II	4.6 ± 0.3	
E2402	9	I	S	II	8.7 ± 0.3	
Ent9a	9a§	I	S	IX	n.d.	
P95661	12	I	S	II	7.0 ± 2.1	
Ent11A	11a§	I	S	III	n.d.	
P95940	17	I	S	II	7.9 ± 0.8	
E1949	19	I	S	III	8.5 ± 0	
Ent20a	20a§	I	S	II	n.d.	
P72580	21	I	S	II	8.3 ± 1.3	
P100613	25	I	S	II	7.2 ± 2.0	
Ent31	31§	I	S	II	n.d.	
Ent32	32§	I	S	I	n.d.	
Ent33	34§	I	S	II	n.d.	
E2452	2	II	S	II	8.7 ± 0.3	7.0 ± 2.3**
E2468	8	II	S	II	8.5 ± 0	
E3945	10	II	S	III	4.0 ± 0.2	
E464	13	II	S	II	7.6 ± 2.1	
Ent13A	13a§	II	S	VIII	n.d.	
P84357	22	II	S	II	8.4 ± 0.2	
P88255	23	II	R	II	3.5 ± 1.4	
P99768	24	II	S	II	8.5 ± 0	
Ent28	28†	II	S	II	n.d.	
Ent9b	9b§	Other	S	II	n.d.	6.6 ± 2.8
E2109	11	Other	S	I	8.3 ± 0.8	
E2384	14	Other	S	IV	2.9 ± 1.2	
Ent14b	14b§	Other	S	II	n.d.	
E2400	15	Other	S	I	8.5 ± 0	
E866	16	Other	S	VII	3.0 ± 1.4	
P89448	18	Other	S	II	8.5 ± 0	
P68147	20	Other	S	I	8.5 ± 0	

\* Clonality defined based on IS200 typing according to Stanley and colleagues (1991) with grouping according to Olsen and colleagues (1994).

† S, smooth; R, rough LPS profile.

‡ *Sma*I ribotype with probe complementary to 16S and 23S rRNA.

§ IS200 and ribotyping results from this study.

\*\*  $\bar{x}$ LogVC10 without the rough strains of PT7 and PT23 were 7.2 ± 1.2 for clonal line I and 7.6 ± 1.8 for clonal line II.

strains, preferably without the use of animal models. *S. Enteritidis* contains more than one clonal line as identified by multi locus enzyme electrophoresis (three lines) [12], the copy number and the location of insertion sequence IS200 (at least 3 lines) [13, 14], and typing with a variety of DNA based typing methods (two major and several minor lines) [14]. In the

present study, mouse virulence of strains of *S. Enteritidis* from different clonal lines was compared with adhesion and invasion in cell culture in order to evaluate whether levels of virulence could be predicted by the use of cell culture, and to determine whether virulence was influenced by genomic lineage of the *S. Enteritidis* strains.

## MATERIALS AND METHODS

### Bacterial strains

The 38 phage type (PT) reference strains of *S. Enteritidis* used are listed in Table 1. They consisted of 27 strains selected based on reports on genomic typing results [14] and 11 phage type reference strains which had not previously been allocated to a clonal line. The strains were kindly provided by Dr L. R. Ward, Central Public Health Laboratory (CPHL), London, UK. For PT1, PT2, PT4, PT8 and PT13, additional wild-type strains were included to test the generality of the results obtained by challenge of mice (Table 2). These strains were kindly provided by Dr E. J. Threlfall, CPHL, London, UK. *S. Typhimurium* 13/74 and its isogenic *invH*<sup>-</sup> mutant [15] were used as positive and negative controls in tissue culture experiments. Strains were cultured and maintained as previously reported [14].

### Allocation of strains into clonal lines based on genotype

Strains were allocated to a clonal line by the method of Stanley and colleagues [13] based on IS200 profiles using the restriction enzymes *PvuII* and *PstI* (Boehringer–Mannheim), and further characterized by ribotyping using the enzyme *SmaI*. Isolation of DNA, restriction endonuclease digestions, agarose gel electrophoresis, preparation of probes, Southern analysis, detection of hybridization results and allocation to IS200-type and ribotype were carried out as previously reported [14].

### Profiling of LPS

LPS molecules were selectively isolated by the proteinase K method, separated by SDS–Page and demonstrated by silver staining as previously reported [16].

### Mouse virulence

Strains were tested for mouse virulence following intraperitoneal challenge of groups of five 21 g female Balb/C mice with suspensions made from overnight cultures and containing  $3 \times 10^3$  c.f.u. in Brain Heart Infusion broth (BHI) (Oxoid) as previously reported [17]. Five mice, which received BHI only, served as controls. These were all negative for salmonella upon

Table 2. Variation of virulence within selected phage types of *S. Enteritidis*

Phage type	Strain	LogVC10	$\bar{x}$ LogVC10
PT1	E2341	8.3 ± 1.3	8.8 ± 0.6
	P79582	9.4 ± 0.3	
	P82987	8.7 ± 0.5	
PT2	E2452	8.7 ± 0.3	8.6 ± 0.1
	P79578	8.5 ± 0	
	P88343	8.7 ± 0.3	
PT4	E2187	8.3 ± 0.4	7.3 ± 1.7
	P49120	5.7 ± 1.6	
	P132344	6.5 ± 0.6	
	P125592	8.5 ± 0	
PT8	E2468	8.5 ± 0	8.6 ± 0.1
	P87975	8.5 ± 0	
	P87888	8.7 ± 0.6	
	P127527	8.5 ± 0	
PT13	E464	7.6 ± 2.1	7.2 ± 2.1
	P106547	8.5 ± 0	
	P143029	8.5 ± 0	
	P141006	4.2 ± 0.9	

the termination of the testings. The log<sub>1</sub> c.f.u. in spleens 10 days post infection (LogVC10) was used as a measure of virulence. For statistical purposes, mice that were killed in the terminal stages for animal welfare reasons were allotted a value of 8.5, provided the spleen contained a pure culture of salmonella. The detection level of the assay was 10<sup>2</sup> c.f.u. per spleen. The minimum logVC10 was therefore taken as 2.0.

### In vitro cell assays

The ability of the bacteria to adhere to, and to invade cultured MDCK (Manine–Darby Canine Kidney) cells was determined essentially as reported by Fields and colleagues [18]. A detailed description of methods used has been given elsewhere [17].


### Statistics

LogVC10 values and c.f.u. associated with cell monolayers were compared by Bonferroni's pairwise comparison of means using the computer software Statistix version 3.

## RESULTS AND DISCUSSION

### Allocation of strains into clonal line

Eleven of the phage type reference strains had not previously been allocated to a clonal line. These



**Fig. 1.** IS200 and ribotype patterns that had not previously been demonstrated among strains of *S. Enteritidis*. (a) IS200 patterns. Lanes 1, 5: molecular size marker phage lambda *Hind*III; lanes 2–4 show *Pst*I-IS200 patterns: 2, 3: the commonly observed pattern I (strains of PT1 and PT5a); lane 4: PT14a (at least 10 copies of IS200), lanes 6–8 shows *Pvu*II-IS200 patterns of the same strains. (b) *Sma*I-ribotype patterns. Lane 1: molecular size marker phage lambda *Hind*III; lanes 2 and 3: type III and type II according to [14] (strains P66040 and E2187 of PT3 and PT4; lane 4: PT9a (new type IX)).

strains were shown to be of 5 IS200 types when the RFLP-patterns of both *Pvu*II and *Pst*I were considered (Table 1). One IS200 and one ribotyping pattern that had not previously been observed among *S. Enteritidis* strains are shown in Figure 1.

The strains of PT5a, PT9a, 11a, 20a, 31, 32 and 34 were of *Pvu*II and *Pst*I pattern I, and the strains are therefore included in *S. Enteritidis* clonal line I together with the type strains of PT1, 3, 4, 4a, 5, 6, 6a, 7, 9, 12, 17, 19, 20, 21 and 25. In support of this observation, Olsen and colleagues [14] typed clinical isolates of 20a, 31 and 32 and allocated these strains to the same clonal line, as has now been shown for the reference strains. The strains of PT13a and PT28 were allocated to clonal line II based on the presence of IS200 type II with both enzymes. These phage types belong to the same clonal line as PT2, 8, 10, 13, 22, 23 and 24. The result with the PT13a type strain confirms the allocation of this phage type to clonal line II as previously suggested by a clinical PT13a isolate [14].

Olsen and colleagues [14] observed that a wild-type strain of PT14b contained at least 7 copies of IS200. With the phage type reference strain, the *Pvu*II pattern differed from the pattern in the wild type strain, and the strain was found to contain at least 10 copies of

the insertion element (Fig. 1). The strain of PT9b showed a pattern identical to one previously demonstrated in a clinical isolate of this phage type [14].

Based on previous studies [13, 14] and the results reported here, IS200 typing is shown to divide strains of *S. Enteritidis* into two major clonal lines, each characterized by a unique IS200 pattern. In addition, several minor groups are formed outside these two lines. Human cases may be caused by strains of all lines, but isolates from Europe have been predominantly of clonal line I, while clonal line II strains have been the most common on the American continent [1, 19]. There is, however, now a tendency towards a more complex picture, as for example PT4 outbreaks are reported from the USA [20].

### LPS profiles

Thirty-seven of the 39 phage type reference strains possessed fully smooth LPS, while the strains representing phage type 7 and phage type 23 showed a shortened profile (data not shown) and were scored as semi-rough. The roughness of the strains was not evident from a simple visual examination of the colony morphology on agar plates. Strains of phage

type 7 have been reported previously to be rough and of reduced virulence [21]. Phage type 7 strains may be derived from other phage types such as PT1, PT4 and PT6 under natural conditions, and such phage type conversions have been demonstrated apparently to be associated with loss of lipopolysaccharide [22]. Phage type conversion in *S. Enteritidis* has also been reported to be caused by introduction of extra chromosomal DNA [23, 24].

### Mouse challenge experiments

Log<sub>10</sub>VC10 values of the phage type reference strains of *S. Enteritidis* are shown in Table 1 in relation to the genomic line. Within the two major clonal lines, both strains of high (log<sub>10</sub>VC10 > 7.5) and medium (7.5 > log<sub>10</sub>VC10 > 5) to low (log<sub>10</sub>VC10 < 5) virulence were present. The strains with high values had log<sub>10</sub>VC10 values in the range 7.6–8.7 and included PT1, PT3, PT4, PT4a, PT5, PT9, PT17, PT19 and PT21 of clonal line I and PT2, PT8, PT13, PT22 and PT24 of clonal line II. The log<sub>10</sub>VC10 values of these strains were not statistically different from each other. The strains of PT6, PT6a, PT12 and PT25 showed values of 5.1, 5.7, 7.0 and 7.2, while the two semi-rough strains, i.e. PT7 of clonal line I and PT23 of clonal line II, and the strain of PT10 from clonal line II were of low virulence, and log<sub>10</sub>VC10 values of 4.6, 3.5 and 4.0, respectively, were obtained.

The mean log<sub>10</sub>VC10 values of strains belonging to the three groups (clonal line I and II and other strains) were not statistically different, irrespective of whether or not the semi-rough strains were included in the calculations, indicating that IS200 typing is not a useful method to distinguish between high and low virulence strains and that essential virulence factors are probably a common feature of all genomic lines of *S. Enteritidis*. The latter suggestion is supported by studies of population genetics within salmonella. The *spv* operon is believed to have been acquired early in *S. enterica* evolution, as the genes are present in the chromosome of strains belonging to subspecies II, IIIa and IV (and the Selander group VII), are carried on extrachromosomal elements of several of the more common serotypes of *S. enterica* subspecies I, but are not present in strains of *S. bongori* [24]. Other important genes for salmonella virulence have also most likely been acquired from an external source early in the evolution but after the split between salmonella and *E. coli*, i.e. genes belonging to the so-called pathogenicity islands (SPIs) [26]. The best

characterized are SPI-1, which encodes genes involved in intestinal invasion and macrophage cytotoxicity [27] and SPI-2, which encodes genes required for survival within macrophages [28]. SPI-1 is present in all salmonellas while SPI-2 is absent in strains of *S. bongori* [26]. Thus it seems unlikely that high levels of virulence have been acquired separately by strains belonging to the different clonal lineages of *S. Enteritidis*, and it is a reasonable assumption that strains of moderate to low virulence in the mouse model have lost some of the essential factors needed to cause a systemic infection.

Ribotyping may be used as an alternative to IS200 typing for determining genomic types of *S. Enteritidis* [14], and virulence was also assessed in relation to grouping determined by this method. Strains of PT3 and PT19 could be separated from the remaining strains of clonal line I that were tested in mice, based on a unique ribotype (Table 1). The type strains of these phage types both had log<sub>10</sub>VC10 values in the high level. The type strain of PT10 (clonal line II) was of the same ribotype. The strain does not carry a virulence plasmid [29], and as expected, it was of low virulence. Strains of PT1, PT5, PT11, PT15 and PT20 are of one ribotype [14]. These strains were all shown to be highly virulent in the mouse model. However, the strain of PT6a showed the same ribotype, and this strain, although still in possession of its virulence plasmid, was only moderately virulent (Table 1). As with IS200 typing, ribotyping was not a predictable indicator of virulence in *S. Enteritidis*.

In order to assess the variability of virulence within a phage type, more strains of PT1, PT2, PT4, PT8 and PT13 were tested (Table 2). All strains of PT1, PT2 and PT8 were highly virulent (log<sub>10</sub>VC10 > 8.3) while medium to low virulent strains (log<sub>10</sub>VC10 < 5.7) were observed among the PT4 and PT13 strains. This indicates that neither phage type, nor genomic typing is a predictable indicator of the virulence of *S. Enteritidis* strains. Strains of *S. Enteritidis* have been reported previously to vary in virulence, as have strains of the same phage type, and it has been noted that differences observed between strains in one animal model may not be detected in another animal model [7], indicating that different virulence factors are important in different animal species.

The reason for the variation in virulence observed in the present study is unknown, but may differ between strains. Mouse virulence drops dramatically when the virulence associated plasmid is removed [30, 31], but the demonstration of low virulent strains

Table 3. Adherence and invasion of strains of *S. Enteritidis* into MDCK cells

Strain	Phage type	Clonal line	Adherence*	Invasion†
E2341	1	I	20 ± 3	5 ± 4
P66040	3	I	17 ± 10	5 ± 4
E2187	4	I	21 ± 5	7 ± 8
P99764	4a	I	25 ± 6	4 ± 5
P70001	5	I	20 ± 5	9 ± 3
P99327	6	I	13 ± 1	11 ± 10
E2408	6a	I	14 ± 5	2 ± 1
E1492	7	I	18 ± 9	11 ± 10
E2402	9	I	17 ± 3	25 ± 22
P95661	12	I	16 ± 3	11 ± 8
P95940	17	I	37 ± 16	9 ± 1
E1949	19	I	23 ± 10	8 ± 4
P72580	21	I	24 ± 8	38 ± 23
E2457	2	II	17 ± 3	26 ± 35
E2468	8	II	22 ± 5	24 ± 22
E3945	10	II	19 ± 11	31 ± 34
E464	13	II	20 ± 4	7 ± 0
P84357	22	II	18 ± 6	13 ± 10
P88255	23	II	61 ± 4	4 ± 1
P99768	24	II	28 ± 9	151 ± 34
E2109	11	Other	28 ± 9	33 ± 41
E2384	14	Other	22 ± 9	11 ± 3
E2400	15	Other	32 ± 6	16 ± 11
E866	16	Other	43 ± 6	25 ± 19
P89448	18	Other	40 ± 18	106 ± 6
P68147	20	Other	17 ± 2	16 ± 16

\* Adhesion defined as associated bacteria as a per cent of inoculum.

† Invasion defined as intra cellular bacteria as a per cent of associated bacteria.

cannot be due simply to the absence of *spv* genes, as virulence plasmids and *spv*-genes have been demonstrated in all phage type reference strains tested, except PT10 [29]. environmental conditions often control expression of virulence factors in pathogenic bacteria [32], and PT4 strains that show a generally impaired ability to adapt to changing environmental condition have also been demonstrated to be essentially avirulent in mice by the oral infection route [10]. The strains have a variant *rpoS*-gene with a deletion, and in a direct comparison, a strain with the variant gene was less virulent in mice following oral challenge than a non-isogenic strain with the normal gene [33], while such strains do not differ in the ability to invade and colonize internal organs following oral challenge of one day old chickens [34]. Since the strains were not isogenic, however, other underlying factors may have caused the difference in virulence. Also, different sequence variations in the *E. coli katF*

(*rpoS*)-gene have been shown to have different effects on the phenotype [35], and this may also be a possibility for similar mutations in strains of salmonella.

Strains originating from clinical cases seem to be more virulent than those from the environment [11], and it may be that strains of *S. Enteritidis* of low and medium virulence are common in nature but that animal and human hosts select for the more virulent variants. As the majority of studies dealing with virulence characterization have used clinical isolates this may give an unbalanced picture of the virulence variation with the serotype.

#### Adhesion to, and invasion of cultured cells

Results from *in vitro* adhesion and invasion assays with selected strains are summarized in Table 3. After 1 h of incubation, the number of c.f.u.s associated with the MDCK cells were highest for the semi-rough strain of 23. The per cent association of this strain was significantly higher than for all other strains except PT16, PT17 and PT18. This result is in accordance with previous observations that LPS-mutants adhere strongly to cells and other substances in *in vitro* assays [36]. Among the remaining strains, phage types 6 and 6a showed the lowest ability to adhere to cultured MDCK cells but the values observed were not statistically different from the values of other phage types, with the noted exception of phage type 23.

In an attempt to correlate virulence with an *in vitro* feature of strains of *S. Enteritidis*, Solano and colleagues [11] showed that when the bacteria were starved, only virulent strains were able to adhere to the wall of glass tubes, forming a biofilm. No direct comparison with the present study was possible since different strains were tested and since bacteria in the present investigation were not starved before the ability to adhere to cells was tested.

After incubation with gentamicin for 2 h (intracellular bacteria only), 4% of the c.f.u.s associated with cells after 1 h could be detected in the positive control (*S. Typhimurium* 13/74), while the negative control showed 0% (*S. Typhimurium* invH<sup>-</sup>) intracellular location. The values for most of *S. Enteritidis* strains varied between 5 and 40%. An apparent intracellular replication, however, was observed with PT24 (c.f.u. more than 150% of associated cells) and with PT18 (c.f.u. 106% of associated cells).

In previous testing, cells of *S. Enteritidis* had been shown to be capable of adhering to, and entering cultured epithelial cells [37]. Based upon the present results it can be concluded that this ability seems to vary greatly between different strains of the serotype. However, due to the small number of tests performed with each strain, the standard deviation was high, and only a few strains differed significantly.

The tissue culture results did not correlate with the results of i.p. mouse virulence tests, possibly because the i.p. challenge model does not require invasion of the gut. However, for some of the strains included in the present study, studies of invasion in a chicken loop test model have been performed recently. In this model, PT4 was significantly better at invading than strains of other phage types (Aabo, personal communication). This was not reflected in the present *in vitro* adherence and invasion assays. Although it is desirable to reduce the use of experimental animal infections to a minimum, our results indicate that cell culture assays cannot be used to predict the outcome of the challenge of the mouse with *S. Enteritidis*.

#### ACKNOWLEDGEMENTS

This study was supported by the Danish Agricultural and Veterinary Research Council (grant no. 13-4538-1 and 95-01056). We are grateful to Charlotte Rasmussen and Gitte Christensen for skilful technical assistance.

#### REFERENCES

1. Fisher I. Salm/Enter-Net records a resurgence in *Salmonella enteritidis* infections throughout the European. Eurosurveill Wkly, 26 June 1997.
2. Anonymous. Annual Report on Zoonosis in Denmark 1998. Ministry of Food, Agriculture and Fisheries, 1999.
3. Rampling A, Anderson JR, Upton R, Peters E, Ward LR, Rowe B. *Salmonella enteritidis* phage type 4 infection of broiler chickens: a hazard to public health. *Lancet* 1989; **ii**: 436–8.
4. Keller LH, Benson CE, Krotec K, Eckroade RJ. *Salmonella enteritidis* colonization of the reproductive tract and forming and freshly laid eggs of chickens. *Infect Immun* 1995; **63**: 2443–9.
5. Gast RK, Benson ST. The comparative virulence for chicks of *Salmonella enteritidis* phage type 4 isolates and isolates of phage types commonly found in the United States. *Avian Dis* 1995; **39**: 567–74.
6. Hinton M, Threlfall EJ, Rowe B. The invasive potential of *Salmonella enteritidis* phage types for young chickens. *Lett Appl Microbiol* 1990; **10**: 237–9.
7. Poppe C, Demczuk W, McFadden K, Johnson RP. Virulence of *Salmonella enteritidis* phage types 4, 8 and 13 and other *Salmonella* spp. for day-old chicks, hens and mice. *Can J Vet Res* 1993; **57**: 281–7.
8. Chart H, Threlfall EJ, Rowe B. Virulence of *Salmonella enteritidis* phage type 4 is related to the possession of a 38 MDa plasmid. *FEMS Microbiol Lett* 1989; **58**: 299–304.
9. Chart H, Threlfall EJ, Rowe B. Virulence studies of *Salmonella enteritidis* phage types. *Lett Appl Microbiol* 1991; **12**: 188–91.
10. Humphrey TJ, Williams A, McAlpine K, Lever MS, Guard-Peter J, Cox JM. Isolates of *Salmonella enteritidis* Enteritidis PT4 with enhanced heat acid tolerance are more virulent in mice and more invasive in chickens. *Epidemiol Infect* 1996; **117**: 79–88.
11. Solano C, Sesma B, Alvarez M, Humphrey TJ, Thorns CJ, Gamazo C. Discrimination of strains of *Salmonella enteritidis* with different levels of virulence by an *in vitro* glass adherence test. *J Clin Microbiol* 1998; **36**: 674–8.
12. Beltran P, Musser JM, Helmuth R, et al. Toward a population genetic analysis of *Salmonella*: genetic diversity and relationships among strains of serotypes *S. choleraesuis*, *S. derby*, *S. dublin*, *S. enteritidis*, *S. heidelberg*, *S. infantis*, *S. newport* and *S. typhimurium*. *Proc Natl Acad Sci USA* 1988; **85**: 7753–7.
13. Stanley J, Jones C, Threlfall EJ. Evolutionary lines among *Salmonella enteritidis* phage types are identified by insertions sequence IS200 distribution. *FEMS Microbiol Lett* 1991; **82**: 83–90.
14. Olsen JE, Skov MN, Threlfall EJ, Brown DJ. Clonal lines of *Salmonella enterica* serotype Enteritidis documented by IS200-, ribo-, pulsed field gel electrophoresis- and RFLP-typing. *J Med Microbiol* 1994; **40**: 15–22.
15. Watson PR, Paulin SM, Bland AP, Jones PW, Wallis TS. Characterization of intestinal invasion by *Salmonella typhimurium* and *Salmonella dublin* and the effect of a mutation in the *invH* gene. *Infect Immun* 1995; **63**: 2743–54.
16. Tiainen T, Pedersen K, Larsen JL. *Vibrio anguillarum* serogroup O3 and *V. anguillarum*-like serogroup O3 cross-reactive species – comparison and characterization. *J Appl Microbiol* 1997; **82**: 211–8.
17. Olsen JE, Skov MM, Brown DJ, Christensen JP, Bisgaard M. Virulence and genotype stability of *Salmonella enterica* serovar Berta during a natural outbreak. *Epidemiol Infect* 1996; **116**: 267–74.
18. Fields PI, Swanson RV, Hardaris CG, Heffron F. Mutants of *Salmonella typhimurium* that cannot survive within the macrophage are avirulent. *Proc Natl Acad Sci USA* 1986; **83**: 5189–93.
19. Rodrigue DC, Tauxe RV, Rowe B. International increase in *Salmonella enteritidis*: a new pandemic? *Epidemiol Infect* 1990; **105**: 21–7.
20. Boyes TG, Koo D, Swerdlow DL, et al. Recurrent outbreaks of *Salmonella* Enteritidis infections in a Texas restaurant: phage type 4 arrives in the United States. *Epidemiol Infect* 1996; **117**: 29–34.
21. Chart H, Rowe B, Threlfall EJ, Ward LR. Conversion of *Salmonella enteritidis* phage type 4 to phage type 7

- involves loss of lipopolysaccharide and with concomitant loss of virulence. FEMS Microbiol Lett 1989; **60**: 37–40.
22. Baggesen DL, Wegener HC, Madsen M. Correlation of conversion of *Salmonella enterica* serotype Enteritidis phage type 1, 4, or 6 to phage type 7 with loss of lipopolysaccharide. J Clin Microbiol 1997; **35**: 330–3.
  23. Threlfall EJ, Chart H, Ward LR, de Sa JDH, Rowe B. Interrelationship between strains of *Salmonella enteritidis* belonging to phage types 4, 7, 7a, 8, 13a, 23, 24 and 30. J Appl Bacteriol 1993; **75**: 43–8.
  24. Brown DJ, Baggesen DL, Platt DJ, Olsen JE. Phage type conversion in *Salmonella enterica* serotype Enteritidis caused by the introduction of a resistance plasmid of incompatibility group X (IncX). Epidemiol Infect 1999; **122**: 19–22.
  25. Boyd EF, Hartl DL. Salmonella virulence plasmid: modular acquisition of the *spv* virulence region by an F-plasmid in *Salmonella enterica* subspecies I and insertion into the chromosome of subspecies II, IIIa, IV and VII isolates. Genet 1998; **149**: 1183–90.
  26. Bäumler AJ, Tsois RM, Ficht TA, Adams LG. Evolution of host adaptation in *Salmonella enterica*. Infect Immun 1998; **66**: 4579–87.
  27. Mills DM, Bajaj V, Lee CA. A 40 kb chromosomal fragment encoding *Salmonella typhimurium* invasion genes is absent from the corresponding region of the *Escherichia coli* K-12 chromosome. Mol Microbiol 1995; **15**: 749–59.
  28. Ochman H, Soncini FC, Solomon F, Groisman EA. Identification of a pathogenicity island required for *Salmonella* survival in host cells. Proc Natl Acad Sci USA 1996; **93**: 7800–4.
  29. Brown DJ, Threlfall EJ, Hampton MD, Rowe B. Molecular characterization of plasmids in *Salmonella enteritidis* phage types. Epidemiol Infect 1993; **110**: 209–16.
  30. Hovi M, Sukupolvi S, Edwards MF, Rhen M. Plasmid-associated virulence of *Salmonella enteritidis*. Microb Pathogen 1988; **4**: 385–91.
  31. Suzuki S, Ohmae K, Nakamura M, *et al.* Demonstration of the correlation of a 36-megadalton *Salmonella* serova enteritidis plasmid to virulence in mice by reintroduction of the plasmid. Japan J Vet Sci 1989; **51**: 203–5.
  32. Miller JF, Mekalanos JJ, Falkow S. Coordinate regulation and sensory transduction in the control of bacterial virulence. Science 1989; **243**: 916–21.
  33. Humphrey TJ, Williams A, McAlpine K, Jørgensen F, O'Byrne C. Pathogenicity in isolates of *Salmonella enterica* serotype Enteritidis PT4 which differ in RpoS expression: effects of growth phase and low temperature. Epidemiol Infect 1998; **121**: 295–301.
  34. Allen-Vercoe E, Collingham R, Woodward MJ. The variant *rpoS* allele of *S. enteritidis* strain 27655R does not affect virulence in a chick model nor constitutive curliation but does generate a cold-sensitive phenotype. FEMS Microbiol Lett 1998; **167**: 245–53.
  35. Ivanova A, Renshaw M, Guntaka RV, Eisenstark A. DNA base sequence variability in *katF* (putative sigma factor) gene of *Escherichia coli*. Nucl Acids Res 1992; **20**: 5479–80.
  36. Nevola JJ, Laux DC, Cohen PS. *In vivo* colonization of the mouse large intestine and *in vitro* penetration of intestinal mucus by an avirulent smooth strain of *Salmonella typhimurium* and its lipopolysaccharide-deficient mutant. Infect Immun 1987; **55**: 2884–90.
  37. Stone BJ, Garcia CM, Badger JL, Hassett T, Smith RIF, Miller V. Identification of novel loci affecting entry of *Salmonella enteritidis* into eucaryotic cells. J Bacteriol 1992; **174**: 3945–52.