

Published in final edited form as:

Tetrahedron Lett. 2009 November 25; 50(47): 6606.

Luminaolide, a novel metamorphosis-enhancing macrodiolide for scleractinian coral larvae from crustose coralline algae

Makoto Kitamura^a, Peter J. Schupp^{b,*}, Yoshikatsu Nakano^c, and Daisuke Uemura^{a,d,*}

^aDepartment of Biosciences and Informatics, Center for Biosciences and Informatics, Keio University, Hiyoshi, Yokohama 223-8522, Japan

^bUniversity of Guam Marine Laboratory, UOG Station, Mangilao, GUAM 96923, USA

^cSesoko Station, Tropical Biosphere Research Center, University of the Ryukyus, Motobu, Okinawa 905-0227, Japan

^dInstitute of Advanced Research Center, Nagoya University, Chikusa, Nagoya 464-8602, Japan

Abstract

A new metamorphosis-enhancing macrodiolide, luminaolide (**1**), was isolated from the crustose coralline algae (CCA) *Hydrolithon reinboldii*. Its structure was determined by spectroscopic analysis. A fraction (1.30 µg/mL) eluted with 80% aqueous MeOH by ODS gel column chromatography of the same CCA extract induced larval metamorphosis (25.9 ± 7.4%) against *Leptastrea purpurea*, and its metamorphosis-inducing activity was further enhanced to 92.6 ± 2.9% with the addition of **1** (25.6 ng/mL).

Keywords

Luminaolide; Enhancer; *Hydrolithon reinboldii*; *Leptastrea purpurea*; Coral larvae; Crustose coralline algae (CCA)

The settlement and metamorphosis of larvae of many marine invertebrates are known to be influenced by crustose coralline algae (CCA). In some species, recruitment is inhibited by CCA.¹ However, there are many more reports that coralline algae induce settlement and/or metamorphosis.² In scleractinian coral, various species of CCA have also been shown to be the primary sources of external inducers of metamorphosis in coral larvae.³ Larvae of the scleractinian coral *Agaricia humilis* settle and metamorphose when exposed to the extracts of CCA *Penyssonnelia* sp.⁴ and to the fragments of *Hydrolithon boergesenii*.⁵ These phenomena are thought to produce cell-wall-bound polysaccharides that are recognized by chemoreceptors on the planula larvae.⁴ While several CCA, such as *Lithophyllum inspidum*, *Hydrolithon onkodes*, *Neogoniolithon brassica-florida*, have been shown to induce metamorphosis of the coral larvae *Acropora millepora*, the bacteria *Pseudoalteromonas* sp. isolated from the surface of CCA have also been shown to induce the metamorphosis of *A. millepora* larvae.⁶ Moreover, the specific substratum preferences of CCA and bacteria have been reported in larvae of two species of scleractinian corals, *Goniastrea retiformis* and *Stylaraea punctata*.⁷ However, the chemical characterization of the external cues that act as natural inducers or enhancers^{8b} of larval metamorphosis has been very limited. In our continuing search for bioactive substances

in CCA,⁸ we found that fragments of coral rubble with the CCA *Hydrolithon reinboldii* induced larval metamorphosis in the scleractinian coral *Leptastrea purpurea* (88.9 ± 8.2 , $n = 6$). These biological phenomena were investigated by a simple bioassay⁹ using larvae of *L. purpurea* (Fig. 1). We describe here the isolation of a new macrodiolide, luminaolide (**1**) (Fig. 2), as a natural enhancer of larval metamorphosis, and report its structure and biological activity.

The CCA *H. reinboldii* (248 kg, wet weight, including coral rubble) which had overgrown a skeleton of dead Acroporidae (coral rubble) was collected by skin-diving at a depth of 0.5–1.5 m at Luminao Reef, Guam, USA. The coral rubble was extracted with methanol for 12 h. The extract was filtered, concentrated and partitioned between water and EtOAc. The EtOAc layer was further partitioned between 90% aqueous MeOH and hexane. The 90% aqueous MeOH layer was subjected to silica gel column chromatography (CHCl₃–MeOH) and then ODS gel column chromatography (40% aqueous MeOH–MeOH). The fractions eluted with 80% aqueous MeOH produced a lower percentage of larval metamorphosis compared to that in untreated live CCA. However, the combination of the fractions eluted with 80% aqueous MeOH and MeOH showed higher metamorphosis-inducing activity than the fraction eluted with 80% aqueous MeOH alone (data not shown). This effect guided isolation of the fraction eluted with MeOH by reversed-phase HPLC (MeOH) and preparative TLC (CHCl₃–MeOH, 20:1) to give luminaolide (**1**) [3.2 mg] as a white amorphous solid [α]_D²⁰ –49 (*c* 0.2, MeOH); IR (KBr) 3440 and 1650 cm⁻¹. The molecular weight of **1** was determined by ESIMS [1562.0 (M+Na)⁺, 792.5 (M+2Na)²⁺]. The molecular formula of **1** was found to be C₈₂H₁₄₂N₂O₂₄ [(M+Na)⁺, *m/z* 1561.9845, Δ –0.5 mmu, calcd for C₈₂H₁₄₂N₂O₂₄Na 1561.9850] by HR-FABMS.

¹H and ¹³C NMR (Table 1) and HSQC spectra of **1** in CD₃OD revealed the presence of four O-methyl, one N-methyl, and five secondary methyl groups, and nine methylenes, 10 oxygen-bearing methines, five C-methines, two disubstituted double bonds, one formamide, and two ketones. A terminal *N*-methyl-*N*-vinylformamide structure in **1** was deduced based on a comparison of the ¹H NMR data with those for aplyronines,¹⁰ scytophycins,¹¹ sphinxolide,¹² and macrocyclic trisoxazoles.¹³ Due to the limited rotation about the *N*-methyl-*N*-vinylformamide terminus, doubled NMR signals for some protons and carbons were observed in a ratio of approximately 2:1, as shown in Table 1. The Δ ^{29,30} double bond was assigned an *E* geometry based on a coupling constant of 14.2 Hz between the H29 and H30 signals. A detailed analysis of the ¹H–¹H COSY spectra of **1** allowed the construction of two partial structures: C2–C24 and C26–C30 (Fig. 3). The connection between C24 and C26 through a ketone carbonyl carbon (C25) was suggested from HMBC for H24/C25, H26/C25, and Me-26/C25. Similarly, an HMBC was observed for H2/C1 (δ_C 173.79), indicating that C2 was attached to an ester carbonyl carbon (C1). The existence of a dihydropyran ring at C7–C11 was implied by the HMBC for H7/C11. Furthermore, HMBC for OMe-5/C5, OMe-13/C13, OMe-21/C21, and OMe-27/C27 suggested the attachment of four methoxy groups at C5, C13, C21, and C27, respectively. The ester linkage between C1 and C19 was established based on a relatively lowerfield chemical shift for H19 (δ_H 5.62) and the HMBC for H19/C1. The remaining three oxymethine carbon signals showed a deuterium shift (C3; δ_C 66.04, C15; δ_C 72.68, and C17; δ_C 68.23, respectively) in the ¹³C NMR spectra in CD₃OH, which suggested that three hydroxyl groups were located at C3, C15, and C17, respectively. Overall, analysis of the NMR data indicated that **1** had a molecular formula of C₄₁H₇₁NO₁₂, which was exactly half of the molecular formula C₈₂H₁₄₂N₂O₂₄ determined by HR-FABMS. Thus, it was clear that **1** was a symmetrical macrodiolide dimer composed of two identical C₃₀ units containing two *N*-methyl-*N*-vinylformamide groups.

A fraction (1.30 μ g/mL) eluted with 80% aqueous MeOH by ODS gel column chromatography induced larval metamorphosis ($25.9 \pm 7.4\%$) of *L. purpurea*, as shown in Table 2. A higher concentration (6.50 μ g/mL) of this fraction showed malformed or dead individuals without an increase in the percentage of those undergoing metamorphosis (data not shown). These results

suggest that this fraction may contain at least one compound that acts as a natural inducer of larval metamorphosis, although no such inducer has yet been isolated from this fraction. On the other hand, when luminaolide (**1**) was offered to *L. purpurea* larvae at a concentration of 25.6 ng/mL together with this fraction by ODS gel column chromatography (1.30 µg/mL), metamorphosis significantly increased to $92.6 \pm 2.9\%$ compared to that with the fraction obtained by ODS gel column chromatography alone (Table 2). Furthermore, at different concentrations, **1** alone (12.8, 25.6, and 128 ng/mL, respectively) did not show any induction of larval metamorphosis in *L. purpurea* ($n = 4$). These results indicate that **1** serves to enhance larval metamorphosis in *L. purpurea* and does not act as an inducer. This kind of enhancing effect in larval metamorphosis has been generally reported for carotenoids in the scleractinian coral *Pseudosiderastrea tayamai*.^{8b}

Compounds with structures similar to that of **1**, such as tolytoxin,¹⁴ scytophycines,¹⁴ swinhoide A,¹⁵ and their analogs, have been isolated from cyanobacteria, and some related compounds are also thought to originate from cyanobacteria.¹⁶ Fenical and co-workers reported that lobophorolide was isolated from the seaweed *Lobophora variegata*. However, due to the similarity of its structure to related macrolides, they concluded that the true producer could be a microbial symbiont of *L. variegata*.¹⁷ Based on these previous reports, we can speculate that **1** is produced by epiphytic bacteria on the surface of the CCA *H. reinboldii* and not directly by the CCA itself, although no such symbiont has yet been identified.

In conclusion, a new metamorphosis-enhancing dimeric macrolide, luminaolide (**1**), was isolated from the CCA *H. reinboldii*. Compounds related to **1** which possess an *N*-methyl-*N*-vinylformamide terminus^{10–13,18} and macrodiolides¹⁹ with a wide range of activities, such as cytotoxic, antibiotic, antifungal, sea urchin egg cleavage-inhibitory, actin depolymerizing, and proteasome inhibitory activities, have been reported. However, this is the first report of a metamorphosis-enhancing macrolide for scleractinian coral larvae.

Acknowledgments

We thank Ciemon Caballes (University of Guam) for assisting in CCA collection and Chie Takase and Ingeborg Ipping Petterson (University of Guam) for the collection of larvae. This work was supported in part by Grants-in-Aid for Creative Scientific Research (16GS0206) and Scientific Research (C) (20611003) from the Japan Society for the Promotion of Science (JSPS) and the 21st century COE program (Establishment of COE on Materials Science) from the Ministry of Education, Science, Sports, and Culture (MEXT), Japan. We are indebted to Ono Pharmaceutical Co., Ltd for their financial support.

References and notes

1. Breitburg DL. Ecology 1984;65:1136.
2. Hadfield, MG.; Paul, VJ. Marine Chemical Ecology. McClintock, JB.; Baker, BJ., editors. CRC Press; Boca Raton: 2001. p. 443 (a) Hayakawa J, Kawamura T, Horii T, Watanabe Y. Fish. Sci 2007;73:371. (b) Williams EA, Craigie A, Yeates A, Degnan SM. Biol. Bull 2008;215:98. [PubMed: 18723641]
3. Morse ED, Hooker N, Morse ANC, Jensen RA. J. Exp. Mar. Biol. Ecol 1988;116:193.
4. Morse DE, Morse ANC. Biol. Bull 1991;181:104.
5. Morse ANC, Iwao K, Baba M, Shimoike K, Hayashibara T, Omori M. Biol. Bull 1996;191:149.
6. (a) Heyward AJ, Negri AP. Coral Reefs 1999;18:273. (b) Negri AP, Webster NS, Hill RT, Heyward AJ. Mar. Ecol. Prog. Ser 2001;223:121.
7. Golbuu Y, Richmond RH. Mar. Biol 2007;152:639.
8. (a) Kitamura M, Koyama T, Nakano Y, Uemura D. Chem. Lett 2005;34:1272. (b) Kitamura M, Koyama T, Nakano Y, Uemura D. J. Exp. Mar. Biol. Ecol 2007;340:96.
9. *Bioassay*: Assays for larval metamorphosis were performed in a glass dish at the same environmental conditions as stock larvae cultures. Samples of the test chemicals were dissolved in methanol and added into glass dishes. After the solvent was dried, 10 mL of filtered seawater and mobile planula

larvae (six individuals) were added into each dish, and the dishes were incubated in the dark for 24 h at 25–28 °C. The number of young polyps was then counted under an optical microscope. Our criteria were the major change from planula larva to developing primary polyp. For these experiments, we define metamorphosis to have occurred when the larvae changed into disk-shaped structures with septal mesenteries radiating from the central mouth region (Fig. 1B).

10. Yamada K, Ojika M, Ishigaki T, Yoshida Y. *J. Am. Chem. Soc.* 1993;115:11020.
11. (a) Moor RE, Patterson GML, Mynderse JS, Barchi J Jr. Barchi J Jr. Norton TR, Furusawa E, Furusawa S. *Pure & Appl. Chem* 1986;58:263. (b) Ishibashi M, Moor RE, Patterson GML, Xu C, Clardy J. *J. Org. Chem* 1986;51:5300.
12. Guella G, Mancini I, Chiasera G, Pietra F. *Helv. Chim. Acta* 1989;72:237.
13. (a) Roesener J, Scheuer PJ. *J. Am. Chem. Soc* 1986;108:846. (b) Matsumoto S, Fusetani N, Hashimoto K. *J. Am. Chem. Soc* 1986;108:847.
14. Carmeli S, Moore RE, Patterson GML. *J. Nat. Prod* 1990;53:1533. [PubMed: 2128517]
15. Andrianasolo EH, Gross H, Goeger D, Musafija-Girt M, McPhail K, Leal RM, Mooberry SL, Gerwick WH. *Org. Lett* 2005;7:1375. [PubMed: 15787510]
16. (a) Tsukamoto S, Ishibashi M, Sasaki T, Kobayashi J. *J. Chem. Soc., Perkin Trans* 1991;1:3185. (b) Todd JS, Alvi KA, Crews P. *Tetrahedron Lett* 1992;33:441.
17. Kubanek J, Jensen PR, Keifer PA, Sullards MC, Collins DO, Fenical W. *PNAS* 2003;100:6916. [PubMed: 12756301]
18. (a) Matsumoto S, Fusetani N, Hashimoto K, Noguchi H, Sankawa U. *J. Org. Chem* 1989;54:1360. (b) Saito S, Watabe S, Ozaki H, Fusetani N, Karaki H. *J. Biol. Chem* 1994;269:29710. [PubMed: 7961961] (c) Saito S, Watabe S, Ozaki H, Kigoshi H, Yamada K, Fusetani N, Karaki H. *J. Biochem* 1996;120:552. [PubMed: 8902620] (d) Tsukamoto S, Koimaru K, Ohta T. *Mar. Drugs* 2005;3:29.
19. (a) Nakamura H, Iitaka Y, Kitahara T, Okazaki T, Okami Y. *J. Antibiot* 1977;30:714. [PubMed: 924893] (b) Kaiser H, Keller-Schierlein W. *Helv. Chim. Acta* 1981;64:407. (c) Kobayashi M, Tanaka J, Katori T, Matsuura M, Kitagawa I. *Tetrahedron Lett* 1989;30:2963. (d) Tsuda M, Izui N, Shimbo K, Sato M, Fukushi E, Kawabata J, Katsumata K, Horiguchi T, Kobayashi J. *J. Org. Chem* 2003;68:5339. [PubMed: 12816496] (e) Kwon HC, Kauffman CA, Jensen PR, Fenical W. *J. Am. Chem. Soc* 2006;128:1622. [PubMed: 16448135]




Figure 1. (A) *Leptastrea purpurea* in Guam. Scale bar = 1.0 cm. (B) Early metamorphosis of *L. purpurea* larvae, 24 h after exposure to the extract of CCA *Hydrolithon reinboldii* in a glass dish. The central mouth and septal mesenteries are observed. Scale bar = 0.5 mm.




Figure 2.
Structure of luminaolide (**1**).




Figure 3.
Partial structure of luminaolide (**1**), based on 2D NMR correlations.

Table 1

NMR data for luminaolide (**1**) in CD³OD

No.	¹ H ^a	¹³ C ^b	HMBC
1		173.79	
2	2.50 dd (9.8, 16.2) ^c	44.00	C1, 3, 4
	2.59m		C1, 3
3	4.07m	65.95	
4	1.46m	42.71	C5
	1.85 ddd (3.7, 10.3, 14.1)		
5	3.73m	76.16	C5-OMe
5-OMe	3.40s	57.22	C5
6	1.35m	39.90	C5
	1.90 ddd (2.3, 11.5, 13.7)		C7
7	4.45 br d (10.6)	70.82	C6, 8, 9, 11
8	5.67 br d (10.1)	131.07	C7, 10
9	5.83 ddd (2.5, 5.5, 7.8)	125.11	C7
10	1.97m 2H	32.76	C9
11	3.55m	65.50	
12	1.64 ddd (1.3, 10.6, 14.2)	37.67	
	1.82m		C11
13	3.92 br d (9.7)	77.60	C11, 13-OMe
13-OMe	3.37 s	57.22	C13
14	1.51m	43.82	C14-Me, 15
14-Me	0.82 d (7.4)	9.64	C13, 14, 15
15	3.62m	72.62	C14, 17
16	1.74m	43.34	C17
	1.71m		
17	3.81m	68.14	
18	1.44m	42.29	C17
	1.95m		
19	5.62 br dd (1.4, 10.6)	71.94	C1
20	1.80m	41.69	C21
20-Me	0.98 d (7.8)	17.73	C19, 20
21	2.85 dd (3.2, 7.8)	89.40	C19, 20, 21-OMe, 23
21-OMe	3.44 s	61.89	C21
22	1.74m	35.89	
22-Me	0.97 d (7.3)	11.62	C21,22, 23
23	1.35m	25.31	
	1.74m		
24	2.54m	42.83	C22, 23, 25
	2.59m		
25		216.74	

No.	$^1\text{H}^a$	$^{13}\text{C}^b$	HMBC
25 ^e		216.64	
26	2.76 dq (6.9, 9.2)	50.28	C25, 26-Me, 27
26 ^e	2.74	50.31	
26-Me	0.95 d (6.9)	13.90	C25, 26, 27
26-Me ^e	0.94	14.44	
27	3.30 dd [2.6, 9.2] ^d	89.05	C27-OMe
27 ^e		88.97	
27-OMe	3.341 s	61.68	C27
27-OMe ^e	3.338 s	61.68	
28	2.46m	38.72	C28-Me, 29, 30
28 ^e	2.50	38.94	
28-Me	1.17 d (6.9)	19.71	C27, 28, 29
28-Me ^e		19.61	
29	5.19 dd (9.2, 14.2)	113.16	C30
29 ^e	5.26	115.32	
30	6.71 d (14.2)	130.50	C28, 29, 30-NMe, CHo
30 ^e	7.10	125.71	
30-NMe	3.03 s	27.68	C30, CHO
30-NMe ^e	3.12	33.58	
CHO	8.32 s	164.73	C30, 30-NMe
CHO ^e	8.09	163.36	

^a800 MHz.

^b150 MHz.

^cCoupling constants (Hz) are in parentheses.

^dCoupling constants (Hz) based on homo-*J*-resolved ^1H NMR spectral data are in parentheses.

^eSignal for minor conformer.

Table 2

Enhancing effects of luminaolide (**1**) on the induction of larval metamorphosis by a fraction eluted with 80% aqueous MeOH by ODS gel column chromatography

Additive (ng/mL)	% Metamorphosis \pm SE
Control	0 \pm 0.0 a
None	25.9 \pm 7.4 b
Lunimaolide (12.8)	27.8 \pm 6.8 b
Lunimaolide (25.6)	92.6 \pm 2.9 c

Metamorphosis was scored 24 h after the addition of *L. purpurea* larvae (1–3 days after larval release). Each value represents the mean (\pm SE) of nine replicates with six larvae in each replicate. Values with different letters were significantly different from each other ($P < 0.01$; Tukey's test).