



Published in final edited form as:

Biochemistry. 2008 February 12; 47(6): 1760. doi:10.1021/bi701755q.

DIFFERENTIAL REGULATED INTERACTIONS OF CALCIUM/ CALMODULIN-DEPENDENT PROTEIN KINASE II WITH ISOFORMS OF VOLTAGE GATED CALCIUM CHANNEL BETA SUBUNITS†

Chad E. Grueter¹, Sunday A. Abiria², Yunji Wu³, Mark E. Anderson³, and Roger J. Colbran^{1,2,4}

¹Department of Molecular Physiology and Biophysics, Vanderbilt University Medical Center, Nashville, Tennessee 37232.

²Center for Molecular Neuroscience, Vanderbilt University Medical Center, Nashville, Tennessee 37232.

³Department of Medicine, Vanderbilt University Medical Center, Nashville, Tennessee 37232.

⁴Kennedy Center for Research on Human Development, Vanderbilt University Medical Center, Nashville, Tennessee 37232.

Abstract

Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) phosphorylates the β_{2a} subunit of voltage-gated Ca²⁺ channels at Thr498 to facilitate cardiac L-type Ca²⁺ channels. CaMKII colocalizes with β_{2a} in cardiomyocytes and also binds to a domain in β_{2a} that contains Thr498 and exhibits amino acid sequence similarity to the CaMKII autoinhibitory domain and to a CaMKII binding domain in the NMDA receptor NR2B subunit (Grueter et al., 2006. Mol. Cell **23**:641). Here we explore the selectivity of the actions of CaMKII among Ca²⁺ channel β subunit isoforms. CaMKII phosphorylates the β_{1b} , β_{2a} , β_3 and β_4 isoforms with similar initial rates and final stoichiometries of 6–12 mole phosphate per mole protein. However, activated/autophosphorylated CaMKII binds to β_{1b} and β_{2a} with similar apparent affinity, but does not bind to β_3 or β_4 . Pre-phosphorylation of β_{1b} and β_{2a} by CaMKII substantially reduces the binding of autophosphorylated CaMKII. Residues surrounding Thr498 in β_{2a} are highly conserved in β_{1b} , but are different in β_3 and β_4 . Site-directed mutagenesis of this domain in β_{2a} showed that Thr498 phosphorylation promotes dissociation of CaMKII- β_{2a} complexes *in vitro* and reduces interactions of CaMKII with β_{2a} in cells. Mutagenesis of Leu493 to Ala substantially reduces CaMKII binding *in vitro* and in intact cells but does not interfere with β_{2a} phosphorylation at Thr498. In combination, these data show that phosphorylation dynamically regulates the interactions of specific isoforms of the VGCC β subunits with CaMKII.

†This work was supported by National Institutes of Health (NIH) research grants to R.J.C. (MH63232) and M.E.A. (HL062494, HL70250; HL46681). M.E.A. is an AHA Established Investigator. CEG was supported by an Institutional NIH training grant in cardiology (HL07411) and an AHA Fellowship. S.A.A. was partially supported by an AHA Fellowship.

Address correspondence to: RJC: Room 702 Light Hall, Vanderbilt University Medical Center, Nashville, TN 37232-0615. roger.colbran@vanderbilt.edu. Phone: 615-936-1630. FAX: 615-322-7236. MEA: Departments of Internal Medicine and Molecular Physiology, University of Iowa, Carver College of Medicine, Iowa City, IA 52242. mark-e-anderson@uiowa.edu. Phone: 319-356-2750. **Current addresses:** CEG: Department of Molecular Biology, 6000 Harry Hines Blvd., Room NA8. 142, Dallas, TX 75235-9148. MEA: Department of Internal Medicine, University of Iowa, Carver College of Medicine, Iowa City, IA 52242.

SUPPORTING INFORMATION AVAILABLE

Supplementary data (Supplementary Figure 1) showing that CaMKII binding to GST- β_{1b} and GST- β_{2a} in a glutathione agarose co-sedimentation assay is dependent on prior autophosphorylation of CaMKII is available free of charge via the Internet at <http://pubs.acs.org>.

Voltage-gated Ca²⁺ channel (VGCC) ion selectivity and responsiveness to pharmacological antagonist ligands are defined by the identity of the pore forming α_1 subunit. The biophysical properties are generally modified by differential association of auxiliary β , $\alpha_2\delta$ and γ subunits (1–3). Four genes encoding β isoforms have been identified (β_{1-4}) each having multiple mRNA splice variants, which differentially modulate the properties and cell surface expression of VGCC complexes (4–6). In addition, VGCC complexes are further modulated by a variety of posttranslational modifications.

The regulatory properties of Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) allow integration of signals conveyed by changes in the frequency, duration and amplitude of intracellular Ca²⁺ transients (7). This feature critically depends on Ca²⁺/calmodulin-dependent autophosphorylation of Thr286 in CaMKII α (or Thr287 in other CaMKII isoforms) in the autoregulatory domain, which confers subsequent autonomous kinase activity (reviewed in (8,9,10)). Thus, Ca²⁺ transients induce more prolonged kinase activation depending on specific parameters of Ca²⁺ signals in the local environment. Recent studies demonstrated that localization of CaMKII to specific subcellular microdomains confers distinct downstream phosphorylation events (11,12). These studies are consistent with the emerging concept that direct interactions of signaling molecules ensures accurate and timely responses to cell stimulation (13,14). Determining the mechanisms for CaMKII binding to target proteins, such as VGCCs, is thus an important goal for understanding the role of CaMKII in excitable cells.

CaMKII phosphorylates the α and/or β subunits of a variety of VGCCs to modulate Ca²⁺ entry. For example, CaMKII regulates T-type Ca²⁺ channels by binding to and phosphorylating the II-III intracellular loop in the Ca_v3.2 α subunit (15,16). The EF hand motif in the C-terminal domain of the L-type Ca²⁺ channel (LTCC) Ca_v1.3 α subunit is also phosphorylated by CaMKII, causing a negative voltage shift in LTCC current activation (17). CaMKII also phosphorylates multiple sites in the LTCC Ca_v1.2 α_{1c} subunit to promote both Ca²⁺- and voltage-dependent facilitation in heterologous cells (18–20). In addition to these roles in feedback regulation, CaMKII is involved in crosstalk between Ca²⁺ channels: for example, LTCC activation leads to depression of R-type Ca²⁺ channels in dendritic spines via a poorly defined CaMKII-dependent mechanism (21).

We recently showed that CaMKII-dependent facilitation of cardiac Ca_v1.2 LTCCs is mediated by phosphorylation of the β_{2a} subunit at Thr498 in cardiomyocytes (22). Moreover, β_{2a} acts as a CaMKII associated protein (CaMKAP) that directly interacts with CaMKII *in vitro* and in intact cells. Our findings showed that VGCC regulation may be strongly enhanced or modified by association of CaMKII with the β_{2a} subunit. In the current study, we demonstrate that CaMKII phosphorylates all of the β subunit isoforms, but interacts with β subunits in an isoform specific manner; this CaMKII- β subunit interaction is negatively modulated by phosphorylation of the β subunit.

MATERIALS AND METHODS

Generation of plasmid constructs

The open reading frames of the rat VGCC β_{1b} , β_{2a} , β_3 and β_4 subunits (Accession Numbers X61394, M80545, M88751 and L02315) (generous gifts from Dr. E. Perez-Reyes, University of Virginia), were amplified by PCR and ligated into pGEX-4T1 (Amersham Pharmacia Biotech). The β_{2a} subunit was also subcloned into pFLAG-CMV-2 (Sigma-Aldrich), pIRES (Clontech), and pLenti (Invitrogen). Murine CaMKII α and rat CaMKII δ coding sequences were inserted into pcDNA3. The cDNAs encoding β_{2a} were mutated essentially as described in the QuikChange kit (Stratagene). The pcDNA plasmid encoding a constitutively active T287D mutation of myc-CaMKII δ 2 was a generous gift from Dr. E. Olson (UTSW, Dallas).

GST fusion protein expression and purification

GST fusion proteins were expressed and purified as described in (22). Protein concentrations were determined by Bradford assay (BioRad) using bovine serum albumin as standard and confirmed by resolving proteins on SDS-polyacrylamide gels followed by Coomassie-Blue staining.

CaMKII purification and autophosphorylation

Recombinant rat CaMKII δ_2 or mouse CaMKII α purified from baculovirus-infected Sf9 insect cells were autophosphorylated at Thr287 or Thr286, respectively, using ATP or [γ - 32 P]ATP, essentially as described previously (23).

CaMKII plate binding assays

GST fusion proteins in 0.2 ml plate-binding buffer (50 mM Tris-HCl pH 7.5, 200 mM NaCl, 0.1 mM EDTA, 5 mM 2-mercaptoethanol, 0.1% (v/v) Tween-20, 5 mg/ml bovine serum albumin) were incubated for 18–24 hours at 4°C in glutathione-coated wells. After 3 washes with buffer, wells were incubated at 4°C with the indicated concentrations of 32 P-labeled, Thr287 autophosphorylated CaMKII δ (0.2 ml) for 2 hours then washed (8 times, 0.2 ml ice-cold buffer). The bound kinase was quantified using a scintillation counter.

In order to monitor dissociation of preformed CaMKII- β_{2a} complexes, GST- β_{2a} (WT or T498A: \approx 5 pmol) was immobilized in glutathione-coated multi-well plates (Pierce, Rockford, IL) and then incubated for 2 hours at 4°C with [32 P-T 286]CaMKII α (0.25 μ M) in binding buffer. Wells were rinsed 8 times in binding buffer and immobilized complexes were then incubated with 50 mM Tris-HCl pH 7.5, 0.1 M NaCl, 0.25 mg/ml bovine serum albumin, 0.1% Triton X-100, 1 mM dithiothreitol, 10 mM magnesium acetate, with or without 0.5mM ATP. Soluble/dissociated CaMKII was removed from the wells at the indicated times and quantified by scintillation counting.

CaMKII gel overlays

GST fusion proteins (50 pmoles) were resolved by SDS-polyacrylamide gel electrophoresis and transferred to nitrocellulose. Approximately equal protein loading was confirmed by staining membranes using Ponceau S. Membranes were blocked and then incubated for 2 hours at 4°C with 32 P-labeled, Thr287 autophosphorylated CaMKII δ_2 (100 nM), essentially as described (24). After washing, bound CaMKII was quantified using a phosphoimager.

CaMKII phosphorylation of GST- β subunits

Purified GST- β subunits (or GST alone as a blank) were incubated at either 30°C or 4°C in 50 mM HEPES, pH 7.5, 10 mM magnesium acetate, 1 mg/ml bovine serum albumin, 1 mM dithiothreitol, 0.4 mM [γ - 32]ATP (\approx 500 cpm/pmol) or 0.4 mM ATP containing purified CaMKII. At the indicated times aliquots were spotted on P81 phosphocellulose papers, washed in water and counted in a scintillation counter to measure the stoichiometry of phosphorylation. Parallel aliquots were resolved by SDS-polyacrylamide gel analysis, and dried gels were analyzed by autoradiography and/or a phosphoimager followed by densitometry, as described (22).

Immunoblotting

Samples were resolved on Tris-Glycine SDS-polyacrylamide gels and transferred to nitrocellulose membranes using standard methods. Membranes were blocked in 5% (w/v) milk in TTBS (50 mM Tris-HCl, pH 7.5; 0.2 M NaCl, 0.1% (v/v) Tween-20), then incubated with primary antibodies overnight at 4°C. After washing six times for >5 minutes each, membranes were incubated for one hour at room temperature with horseradish peroxidase conjugated

secondary antibody. The washed membranes were developed using enhanced chemiluminescence.

Co-immunoprecipitations from HEK293 cells

Experiments were performed as described in (22). Briefly, HEK293FT cells were transfected with FLAG- β_{2a} (WT, T498A, T498E or L493A), myc-CaMKII δ (T287D), and/or flag vector alone. Cell lysates were immunoprecipitated with FLAG-coated agarose beads (40 μ l; Sigma) and then immunoblotted (see above).

Statistics

Data are expressed as mean \pm sem. Paired comparisons were performed using the Student's t-test. Multiple group comparisons were performed using 1-way or 2-way ANOVA with Bonferroni post-hoc testing, unless otherwise noted. The null hypothesis was rejected if $p < 0.05$.

RESULTS

CaMKII efficiently phosphorylates β_{1-4} subunits

The highly efficient phosphorylation of the β_{2a} subunit at Thr498 by CaMKII is critical for CaMKII effects on cardiac LTCCs (22). In order to begin to analyze potential effects of CaMKII on the other β isoforms, we assessed the phosphorylation of full-length β subunit isoforms that had been expressed in bacteria as GST fusion proteins. First, incubations were performed at 4°C in order to provide an indication of the initial rate of phosphorylation by 10 nM CaMKII (Fig. 1A). Under these conditions CaMKII preferentially phosphorylates Thr498 in GST- β_{2a} (22). All four isoforms were phosphorylated, although β_{1b} and β_{2a} were phosphorylated at a relative rate \approx 2.5 fold faster than for β_3 or β_4 . We then assessed the extent of phosphorylation by various concentrations of CaMKII using a fixed concentration of the GST- β isoforms (1 μ M). We previously identified 6 CaMKII phosphorylation sites in β_{2a} that had been phosphorylated to a stoichiometry of \approx 3 mol/mol (22). In the present studies, each isoform was phosphorylated with a similar CaMKII concentration-dependence (Fig. 1B), with the β_{2a} , β_{1b} , β_3 and β_4 variants attaining stoichiometries of 13.1 ± 1.3 , 10.8 ± 0.4 , 7.4 ± 0.5 and 9.6 ± 1.7 moles of phosphate/mole β , respectively. While ANOVA detected significant variability in the final phosphorylation stoichiometries ($p = 0.038$), post-hoc testing revealed a single significant difference between β_3 and β_{2a} phosphorylation stoichiometries ($p < 0.05$). In combination, these data show that these four β subunit isoforms are rapidly phosphorylated by CaMKII to a high stoichiometry *in vitro* with relatively modest differences in the phosphorylation parameters that were measured.

Selective interactions of CaMKII with VGCC β subunits *in vitro*

Thr498 of the β_{2a} variant lies within a CaMKII-binding domain that is C-terminal to the SH3 and GK domains (Fig. 2A). Alignment of the amino acid sequence of the domain surrounding Thr498 in β_{2a} with other β subunit isoforms revealed variable conservation. A CaMKII consensus phosphorylation motif LXRXXS/T (25) is present in both β_{2a} and β_{1b} and there is additional amino acid sequence similarity outside this motif, but key residues from this motif are missing in β_3 and β_4 . Based on these alignments, we hypothesized that CaMKII would bind to β_{1b} in a similar manner to its interaction with β_{2a} , but not to β_3 or β_4 . To test this hypothesis we immobilized GST- β fusion proteins in glutathione coated 96-well plates and then incubated them with various concentrations of purified [32 P]-autophosphorylated CaMKII. In an initial experiment, the binding of CaMKII to GST- β_{1b} was indistinguishable from the binding to GST- β_{2a} , but binding to GST- β_3 and GST- β_4 was $< 1\%$ of the binding to GST- β_{2a} (Fig. 2B). Binding of Thr286-autophosphorylated CaMKII to both GST- β_{1b} and GST- β_{2a} was concentration dependent and saturable (Fig. 2C). GST- β_{1b} exhibited a significantly higher apparent affinity

for CaMKII than did GST- β_{2a} (apparent K_d 's 35 ± 12 and 120 ± 21 nM CaMKII subunit, respectively. $n=4$. $p<0.01$). GST- β_3 and GST- β_4 failed to bind significant amounts of the kinase even at a concentration of 1200 nM CaMKII subunit (data not shown). Autophosphorylated CaMKII also binds to GST- β_{1b} and GST- β_{2a} , but not to GST- β_3 and GST- β_4 in glutathione agarose cosedimentation assays, but there was no significant binding of nonphosphorylated CaMKII to any GST- β subunit isoforms (Supplementary Fig. 1). Thus, CaMKII does not appear to bind significantly to the β_3 and β_4 isoforms, but interacts with the β_{1b} isoform with an ≈ 3 -fold higher affinity than with β_{2a} following activation by Thr286/7 autophosphorylation.

Interaction with CaMKII is modulated by phosphorylation in vitro

Thr498 and several additional serine and threonine residues lie within the CaMKII binding domain of β_{2a} , suggesting that the interaction with CaMKII may be modulated by phosphorylation. Therefore, GST- β isoforms were preincubated with activated CaMKII in the presence or absence of ATP and then separated from reaction components by SDS-PAGE. The electrophoretic mobility of each GST- β isoform was reduced following preincubation with ATP and CaMKII (Fig. 3: protein stain), consistent with the relatively high phosphorylation stoichiometry under these conditions (c.f., Fig. 1B). An overlay assay was then used to assess the binding of ^{32}P -autophosphorylated CaMKII. Non-phosphorylated GST- β_{1b} and GST- β_{2a} proteins bound substantial, but comparable, amounts of CaMKII (Fig. 3). Pre-phosphorylation significantly reduced CaMKII binding to both proteins by 70–80%. No significant interactions were seen between CaMKII and either GST- β_3 or GST- β_4 , whether or not these protein were pre-phosphorylated. In combination, these data suggest that phosphorylation of the β subunit by CaMKII modulates the binding of CaMKII to the β_{1b} and β_{2a} isoforms.

The mechanism of CaMKII binding to β_{2a}

In order to dissect the mechanism for CaMKII binding we assessed the effect of mutating residues within the CaMKII binding domain of β_{2a} on the interaction with CaMKII. Mutation of Thr498 to Ala or Glu prevented or mimicked phosphorylation at this site, respectively (see above). In addition, residues homologous to Leu493 at the p-5 position relative to Thr498 in β_{2a} and β_{1b} are conserved in other high affinity CaMKII phosphorylation sites that form stable complexes with the CaMKII catalytic domain prior to phosphorylation (e.g., Ser1303 in NR2B and Thr286/7 in CaMKII α/δ). However, hydrophobic residues at the p-5 position are not conserved in β_3 or β_4 , and are not generally considered to be part of the minimal consensus phosphorylation site (25). Therefore, we also mutated Leu493 to Ala. Binding of ^{32}P -autophosphorylated CaMKII to GST- β_{2a} in glutathione coated multi-well plates was unaffected by the T498A mutation, whereas the T498E and L493A mutations significantly reduced CaMKII binding by >90% (Table 1). These data demonstrate that the region surrounding Thr498 is critical for stable binding of Thr286-autophosphorylated CaMKII to the full-length β_{2a} isoform.

Disruption of CaMKII binding does not affect phosphorylation of β_{2a} in vitro

We then explored the effects of the Thr498 and Leu493 mutations on CaMKII phosphorylation of the β_{2a} subunit. The T498A mutation substantially reduced the rate of β_{2a} phosphorylation at 4°C, as shown previously, and T498E mutation had a very similar effect. Interestingly, the rate of L493A- β_{2a} phosphorylation by CaMKII was indistinguishable from the rate of phosphorylation of the wild type protein (Fig. 4A), even though the L493A mutation interfered with CaMKII binding. When incubations were conducted at 30°C, CaMKII (10 nM) phosphorylated wild type β_{2a} to a stoichiometry of 2.2 mol/mol, consistent with stoichiometries reported in Fig. 1B using 10 nM CaMKII, but T498A mutation reduced phosphorylation to 0.3 mol/mol (Fig. 4B). In contrast, T498E and L493A mutations had no significant effect on the stoichiometry of CaMKII phosphorylation, although there was a trend for reduced

phosphorylation of T498E- β_{2a} . These data show that two mutations that severely compromise stable binding of activated CaMKII to β_{2a} have no significant effect on the overall stoichiometry of CaMKII phosphorylation *in vitro*.

As noted above, Leu493 is part of a conserved sequence surrounding high-affinity phosphorylation sites in multiple CaMKII-binding proteins. In order to determine whether L493A mutation altered the phosphorylation site specificity of CaMKII, we exploited similarities in protein sequences surrounding Thr498 in β_{2a} and Thr287 in CaMKII δ (Fig. 2A). Commercially available phospho-Thr287 antibodies detected the β_{2a} subunit following CaMKII phosphorylation, but not nonphosphorylated β_{2a} (Fig. 4B and data not shown). Phospho-Thr286/7 antibodies only weakly detected phosphorylated β_{1b} and failed to detect β_3 or β_4 before or after CaMKII phosphorylation (data not shown). Analysis of mutated β_{2a} proteins revealed that T498A- β_{2a} and T498E- β_{2a} could not be detected (Fig. 4B, middle blot), demonstrating that the antibody specifically recognized phospho-Thr498 in β_{2a} , in addition to the Thr286/287 autophosphorylation site in CaMKII. The lack of antibody cross-reactivity with T498E- β_{2a} is not surprising given differences in size and charge density between glutamate and phospho-threonine. Taken together, these findings validate the use of this antibody to report the phosphorylation status of Thr498 in the CaMKII binding domain of β_{2a} . Notably, Thr498 was phosphorylated to a similar extent in L493A- β_{2a} and wild type β_{2a} (Fig. 4B, inset). Thus, L493A mutation does not appear to significantly alter the ability of CaMKII to phosphorylate Thr498 *in vitro*.

Phosphorylation at Thr498 disrupts CaMKII binding

In order to explore the mechanism by which phosphorylation of β_{2a} by CaMKII interferes with subsequent binding of activated CaMKII, we pre-phosphorylated wild type and mutated β_{2a} proteins and analyzed CaMKII binding using overlay assays. As seen using the glutathione plate-binding assay, T498A- β_{2a} bound comparable amounts of activated CaMKII to wild type β_{2a} . However, pre-phosphorylation of T498A- β_{2a} had no significant effect on binding of activated CaMKII, whereas pre-phosphorylation of wild type β_{2a} significantly reduced binding by $\approx 80\%$ in these assays. The T498E mutation reduced CaMKII binding by $\approx 80\%$ and CaMKII phosphorylation had no significant additional effect on binding to T498E- β_{2a} . Interestingly, L493A mutation significantly reduced binding by $\approx 90\%$ in these assays, and pre-phosphorylation by CaMKII resulted in an additional significant decrease in binding. Taken together, these data suggest that pre-phosphorylation of β_{2a} at Thr498 substantially reduces the association of CaMKII.

We also investigated the role of Thr498 phosphorylation in the context of preformed CaMKII- β_{2a} complexes. Complexes of autonomously-active Thr286 autophosphorylated CaMKII bound to GST- β_{2a} (WT) were isolated in glutathione-coated multi-well plates and then incubated with or without addition of ATP. Addition of ATP to these complexes induced phosphorylation at Thr498 and several other sites, as reflected by immunoblotting with phospho-Thr286 CaMKII antibodies and by a substantial reduction in the electrophoretic mobility of GST- β_{2a} (data not shown). In the absence of ATP, CaMKII complexes with WT β_{2a} were remarkably stable ($<5\%$ dissociation over a 2 hour incubation), but addition of ATP induced dissociation of $\approx 50\%$ of bound CaMKII. T498A mutation of β_{2a} had little effect on the stability of complexes with CaMKII in the absence of ATP, but substantially reduced the rate and extent of dissociation following addition of ATP as compared to WT β_{2a} (Fig. 6). However, ATP still enhanced dissociation of CaMKII from preformed complexes with T498A- β_{2a} ($\approx 20\%$ of in 2 hours), suggesting that phosphorylation at other residues in β_{2a} and/or CaMKII may have a modest effect on the stability of these complexes, at least *in vitro*. In combination, these data suggest that phosphorylation of β_{2a} at Thr498 enhances the dissociation of CaMKII from preformed complexes with β_{2a} *in vitro*.

CaMKII interaction with β_{2a} is regulated in cells

Initially, we explored whether Thr498 in β_{2a} is phosphorylated in intact cells. Lysates of HEK293 cells expressing FLAG- β_{2a} (wild type or mutated) with a constitutively active CaMKII δ_2 mutant were immunoprecipitated using FLAG antibodies. Blotting the immune complexes with FLAG antibodies revealed a relatively consistent expression and immunoprecipitation of the β_{2a} proteins. The phospho-Thr286 CaMKII α antibodies detected the β_{2a} wild type and L493A mutant, but not T498A- or T498E- β_{2a} proteins (Fig. 6A). These data show that Thr498 can be phosphorylated to a similar extent in WT β_{2a} and L493A- β_{2a} in intact cells.

In order to investigate the regulation of CaMKII interaction with β_{2a} in intact cells by modification of Thr498 and the surrounding domain, lysates of HEK293 cells expressing constitutively active myc-tagged CaMKII δ without or with FLAG-tagged β_{2a} subunits (WT or mutated) were immunoprecipitated using FLAG antibodies. CaMKII was readily detected in immune complexes containing FLAG-tagged wild-type β_{2a} or T498A- β_{2a} , but could not be detected in immune complexes formed by T498E- β_{2a} or L493A- β_{2a} (Fig. 6B). These findings demonstrate that the domain surrounding Thr498 is critical for association of CaMKII with β_{2a} in cells, and suggest that phosphorylation at Thr498 diminishes CaMKII binding to β_{2a} in cells.

DISCUSSION

A diverse family of VGCCs regulates Ca^{2+} entry into excitable and non-excitable cells. The α_1 subunits confer core biophysical and pharmacological behavior of each type of VGCC. However, a cytosolic loop between the first and second major transmembrane domains in the α subunit is generally thought to constitutively interact with a β subunit. Alternative splicing of mRNAs from four mammalian genes generates >20 distinct β subunit proteins that have divergent roles in modulating the trafficking and biophysical properties of VGCCs (6). The β subunits share highly conserved SH3 and GK domains that form a compact structure, with a hydrophobic groove in the GK domain that interacts with the α subunit (26). However, some β subunits lack substantial parts of the SH3 and GK domains, yet still modulate LTCCs (27). These observations support recent findings showing that additional SH3-GK independent modulatory interactions between the α and β subunits are important for regulating VGCC activity (28–31). The variable domains presumably account for the unique effects of β subunit variants on the properties of α subunits.

Most studies have focused on the roles of protein interactions and post-translational modifications of the α subunit in modulating VGCCs. However, the importance of β subunits in regulating VGCCs has received increasing attention. Initial studies suggested that PKA phosphorylation of the β_2 subunit plays a role in facilitating LTCCs (32), although the importance of this modification in native cells has recently been questioned (33). In addition, β subunits have been shown to serve as scaffolding proteins that bind REM GTPases to inhibit VGCCs (34,35), and AHNAKs to link VGCCs to the actin cytoskeleton (36). Our recent work showed that CaMKII binds β_{2a} subunits and colocalizes with β_{2a} in adult cardiomyocytes. Moreover, phosphorylation of β_{2a} at Thr498 is required for Ca^{2+} and CaMKII-dependent facilitation of LTCCs in cardiomyocytes (22). While β subunit variants have unique direct effects on the biophysical properties and trafficking of LTCCs, the β isoform-selectivity of regulation by CaMKII and other modulators is poorly understood.

The β subunit variants tested here associate with multiple VGCC α subunits (reviewed in (2, 6)). Thus, our current results showing that activated/Thr286 autophosphorylated CaMKII forms stable complexes with β_{1b} and β_{2a} , but not with β_3 or β_4 , lead us to hypothesize that association of CaMKII with VGCC complexes will depend on the identity of the associated

β subunit. The β_{1b} and β_{2a} subunits contain an LXRXXS/T motif similar to sequences surrounding phosphorylation sites in NR2B and the CaMKII autoinhibitory domain that also form stable complexes with the CaMKII catalytic domain. Mutation of Leu493 to Ala within this motif reduced CaMKII binding to β_{2a} by >90%, but surprisingly had little effect on the initial rate of phosphorylation at Thr498, or on the overall phosphorylation stoichiometry at multiple sites. The lack of conservation of this motif accounts for the failure to detect binding to β_3 or β_4 .

Despite the apparent binding selectivity, CaMKII phosphorylates all four β subunit variants tested here with comparable relative rates and overall extent. The high maximal phosphorylation stoichiometries suggest that CaMKII efficiently phosphorylates several sites in each β isoform *in vitro*. Indeed, we previously identified 6 CaMKII phosphorylation sites in β_{2a} (22), and it will be important to identify phosphorylation sites in other β isoforms and determine their impact on the properties of VGCCs. Data presented here show that phosphorylation at Thr498 in β_{2a} negatively regulates CaMKII binding both *in vitro* and *in situ*. Presumably the conserved LXRXXS/T motif in β_{1b} is responsible for the regulated binding of CaMKII to this variant. Thus, interactions of CaMKII with β and/or β_1 variants are regulated by phosphorylation, likely playing an important role in modulating CaMKII targeting to LTCCs and/or other VGCCs.

Interestingly, β subunits are not generally thought to be important in the regulation of T-type VGCCs. However, recent studies show that CaMKII directly interacts with and phosphorylates the II-III linker of the $\text{Ca}_v3.2$ α subunit, shifting the current-voltage activation curve (15,16). In addition, recent data suggest that CaMKII may also interact with multiple cytoplasmic domains in $\text{Ca}_v1.2$ (19). Thus, association of CaMKII with VGCC α subunits may play additional important feature allowing localized Ca^{2+} concentrations to feedback and regulate Ca^{2+} influx.

Emerging data over the last couple of years suggest that mechanisms underlying CaMKII modulation of LTCCs are complex. Phosphorylation of multiple sites in the $\text{Ca}_v1.2$ α subunit by CaMKII appears to play distinct roles in voltage- and calcium-dependent modulation in heterologous cells (18,20). In addition, interactions of CaMKII with multiple intracellular domains of the $\text{Ca}_v1.2$ α subunit have been implicated in LTCC facilitation (19). In contrast, phosphorylation at Thr498 in β_{2a} is required for CaMKII to increase channel open probability at the single channel level and to facilitate whole cell Ca^{2+} currents in adult cardiomyocytes (22). The present observations that phosphorylation at Thr498 appears to be relatively independent of the stable binding of CaMKII to β_{2a} (Fig. 4), and also inhibits CaMKII binding (Fig. 3, Fig. 5, Fig. 6), provides insights that may reconcile these seemingly disparate observations. Interestingly, phosphorylation of β_{2a} in preformed complexes appears to promote dissociation of CaMKII (Fig. 6). CaMKII dissociation might be required to allow protein phosphatases to act on the phospho-Thr498 site to reset channels to their basal state. Presumably, such a mechanism would be most relevant if Thr498 phosphorylation directly modulates LTCC properties, as suggested by the importance of Thr498 in CaMKII-mediated increases in LTCC open probability and LTCC current facilitation in adult cardiomyocytes (22). Secondly, CaMKII may serve a structural role to dynamically assemble complexes containing additional, as yet unidentified, proteins associated with LTCCs. Thus, Thr498 phosphorylation in β_{2a} may promote re-organization of these protein complexes to enable facilitation. Structural roles for CaMKII have been postulated in neuronal postsynaptic densities (23). Previous studies showed that disruption of the actin- or tubulin-based cytoskeleton essentially abrogated CaMKII-dependent facilitation of cardiac LTCCs without affecting PKA-dependent facilitation, suggesting an important role for higher orders of molecular organization close to LTCCs in mediating the effects of CaMKII (37). A final possibility is that dissociation from β_{2a} induced by Thr498 phosphorylation is important in

allowing activated CaMKII to phosphorylate nearby regulatory sites in the α_{1c} or β subunits and/or other associated proteins. In other words, stable binding of activated CaMKII to β_{2a} may limit access of phosphorylation sites in other proteins to the CaMKII active site. Consistent with this hypothesis, dissociation of CaMKII from β_{2a} is substantially reduced by T498A mutation, perhaps explaining why CaMKII phosphorylates this mutant to a lower final stoichiometry than T498E- β_2 and L493A- β_2 , which have much weaker interactions with CaMKII (Fig. 4B). Further studies will be needed to clarify the mechanism of CaMKII actions at VGCC complexes and, perhaps, to identify new CaMKII targets.

In summary, findings reported here and in other recent papers suggest that feedback regulation of Ca^{2+} influx via VGCCs is precisely controlled in specific subcellular microdomains by multiple mechanisms that allow CaMKII and other Ca^{2+} -dependent signaling proteins to associate with channel subunits. The precise nature of the feedback regulation by CaMKII seems likely to depend on the identity of the β subunit associated with the complex. The regulated interaction of activated CaMKII with β_1 and β_2 variants seem likely to be important, although phosphorylation of β_3 and β_4 may also play a role in some cases. Our findings are in line with recent studies suggesting that subcellular targeting of CaMKII via its interactions with CaMKAPs modulates the specificity of its downstream actions (11,12). These complex biochemical mechanisms for feedback regulation of Ca^{2+} influx via VGCCs presumably provide great flexibility for modulating a variety of downstream signaling events such as cardiac excitation-contraction and excitation-transcription coupling and neuronal synaptic plasticity. Moreover, alterations in the association of β subunits with VGCCs might disrupt feedback regulation and downstream signaling in heart failure and other diseases (38).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ABBREVIATIONS

CaMKII	calcium/calmodulin-dependent protein kinase II
CaMKAP	CaMKII-associated protein
GK	guanylate kinase-like
GST	glutathione-S-transferase
LTCC	L-type VGCC
VGCC	voltage-gated calcium channel

Acknowledgments

We appreciate outstanding technical assistance from Martha A. Bass.

REFERENCES

1. Arikath J, Campbell KP. Auxiliary subunits: essential components of the voltage-gated calcium channel complex. *Curr Opin Neurobiol* 2003;13:298–307. [PubMed: 12850214]
2. Catterall WA. Structure and regulation of voltage-gated Ca^{2+} channels. *Annu Rev Cell Dev Biol* 2000;16:521–555. [PubMed: 11031246]
3. Hille, B. *Ion channels of excitable membranes*. 3rd ed.. Sunderland, Mass: Sinauer; 2001.
4. Perez-Reyes E, Castellano A, Kim HS, Bertrand P, Baggstrom E, Lacerda AE, Wei XY, Birnbaumer L. Cloning and expression of a cardiac/brain beta subunit of the L-type calcium channel. *The Journal of biological chemistry* 1992;267:1792–1797. [PubMed: 1370480]

5. Bichet D, Cornet V, Geib S, Carlier E, Volsen S, Hoshi T, Mori Y, De Waard M. The I-II loop of the Ca²⁺ channel alpha 1 subunit contains an endoplasmic reticulum retention signal antagonized by the beta subunit. *Neuron* 2000;25:177–190. [PubMed: 10707982]
6. Dolphin AC. Beta subunits of voltage-gated calcium channels. *J Bioenerg Biomembr* 2003;35:599–620. [PubMed: 15000522]
7. De Koninck P, Schulman H. Sensitivity of CaM kinase II to the frequency of Ca²⁺ oscillations. *Science* 1998;279:227–230. [PubMed: 9422695]
8. Lisman J, Schulman H, Cline H. The molecular basis of CaMKII function in synaptic and behavioural memory. *Nat Rev Neurosci* 2002;3:175–190. [PubMed: 11994750]
9. Hudmon A, Schulman H. Structure-function of the multifunctional Ca²⁺/calmodulin-dependent protein kinase II. *Biochem J* 2002;364:593–611. [PubMed: 11931644]
10. Colbran RJ. Targeting of calcium/calmodulin-dependent protein kinase II. *Biochem J* 2004;378:1–16. [PubMed: 14653781]
11. Tsui J, Inagaki M, Schulman H. Calcium/calmodulin-dependent protein kinase II (CaMKII) localization acts in concert with substrate targeting to create spatial restriction for phosphorylation. *The Journal of biological chemistry* 2005;280:9210–9216. [PubMed: 15582994]
12. Tsui J, Malenka RC. Substrate localization creates specificity in calcium/calmodulin-dependent protein kinase II signaling at synapses. *J Biol Chem* 2006;281:13794–13804. [PubMed: 16551613]
13. Scott JD, Pawson T. Cell communication: the inside story. *Scientific American* 2000;282:72–79. [PubMed: 10862426]
14. Pawson T, Nash P. Assembly of cell regulatory systems through protein interaction domains. *Science* 2003;300:445–452. [PubMed: 12702867]
15. Welsby PJ, Wang H, Wolfe JT, Colbran RJ, Johnson ML, Barrett PQ. A mechanism for the direct regulation of T-type calcium channels by Ca²⁺/calmodulin-dependent kinase II. *J Neurosci* 2003;23:10116–10121. [PubMed: 14602827]
16. Yao J, Davies LA, Howard JD, Adney SK, Welsby PJ, Howell N, Carey RM, Colbran RJ, Barrett PQ. Molecular basis for the modulation of native T-type Ca channels in vivo by Ca²⁺/calmodulin-dependent protein kinase II. *J Clin Invest* 2006;116:2403–2412. [PubMed: 16917542]
17. Gao L, Blair LA, Salinas GD, Needleman LA, Marshall J. Insulin-like growth factor-1 modulation of CaV1.3 calcium channels depends on Ca²⁺ release from IP3-sensitive stores and calcium/calmodulin kinase II phosphorylation of the alpha 1 subunit EF hand. *J Neurosci* 2006;26:6259–6268. [PubMed: 16763033]
18. Erxleben C, Liao Y, Gentile S, Chin D, Gomez-Alegria C, Mori Y, Birnbaumer L, Armstrong DL. Cyclosporin and Timothy syndrome increase mode 2 gating of CaV1.2 calcium channels through aberrant phosphorylation of S6 helices. *Proc Natl Acad Sci U S A* 2006;103:3932–3937. [PubMed: 16537462]
19. Hudmon A, Schulman H, Kim J, Maltez JM, Tsien RW, Pitt GS. CaMKII tethers to L-type Ca²⁺ channels, establishing a local and dedicated integrator of Ca²⁺ signals for facilitation. *J Cell Biol* 2005;171:537–547. [PubMed: 16275756]
20. Lee TS, Karl R, Moosmang S, Lenhardt P, Klugbauer N, Hofmann F, Kleppisch T, Welling A. Calmodulin Kinase II Is Involved in Voltage-dependent Facilitation of the L-type Cav1.2 Calcium Channel: IDENTIFICATION OF THE PHOSPHORYLATION SITES. *The Journal of biological chemistry* 2006;281:25560–25567. [PubMed: 16820363]
21. Yasuda R, Sabatini BL, Svoboda K. Plasticity of calcium channels in dendritic spines. *Nat Neurosci* 2003;6:948–955. [PubMed: 12937422]
22. Grueter CE, Abiria SA, Dzhura I, Wu Y, Ham AJ, Mohler PJ, Anderson ME, Colbran RJ. L-Type Ca(2+) Channel Facilitation Mediated by Phosphorylation of the beta Subunit by CaMKII. *Mol Cell* 2006;23:641–650. [PubMed: 16949361]
23. Robison AJ, Bass MA, Jiao Y, MacMillan LB, Carmody LC, Bartlett RK, Colbran RJ. Multivalent interactions of calcium/calmodulin-dependent protein kinase II with the postsynaptic density proteins NR2B, densin-180, and alpha-actinin-2. *The Journal of biological chemistry* 2005;280:35329–35336. [PubMed: 16120608]

24. McNeill RB, Colbran RJ. Interaction of autophosphorylated Ca²⁺/calmodulin-dependent protein kinase II with neuronal cytoskeletal proteins. Characterization of binding to a 190-kDa postsynaptic density protein. *The Journal of biological chemistry* 1995;270:10043–10049. [PubMed: 7730306]
25. White RR, Kwon YG, Taing M, Lawrence DS, Edelman AM. Definition of optimal substrate recognition motifs of Ca²⁺-calmodulin-dependent protein kinases IV and II reveals shared and distinctive features. *The Journal of biological chemistry* 1998;273:3166–3172. [PubMed: 9452427]
26. Van Petegem F, Clark KA, Chatelain FC, Minor DL Jr. Structure of a complex between a voltage-gated calcium channel beta-subunit and an alpha-subunit domain. *Nature* 2004;429:671–675. [PubMed: 15141227]
27. Harry JB, Kobrinsky E, Abernethy DR, Soldatov NM. New short splice variants of the human cardiac Cavbeta2 subunit: redefining the major functional motifs implemented in modulation of the Cav1.2 channel. *J Biol Chem* 2004;279:46367–46372. [PubMed: 15339916]
28. Butcher AJ, Leroy J, Richards MW, Pratt WS, Dolphin AC. The importance of occupancy rather than affinity of CaV(beta) subunits for the calcium channel I-II linker in relation to calcium channel function. *J Physiol* 2006;574:387–398. [PubMed: 16627564]
29. Leroy J, Richards MW, Butcher AJ, Nieto-Rostro M, Pratt WS, Davies A, Dolphin AC. Interaction via a key tryptophan in the I-II linker of N-type calcium channels is required for beta1 but not for palmitoylated beta2, implicating an additional binding site in the regulation of channel voltage-dependent properties. *J Neurosci* 2005;25:6984–6996. [PubMed: 16049174]
30. Zhang R, Dzhura I, Grueter CE, Thiel W, Colbran RJ, Anderson ME. A dynamic alpha-beta inter-subunit agonist signaling complex is a novel feedback mechanism for regulating L-type Ca²⁺ channel opening. *Faseb J* 2005;19:1573–1575. [PubMed: 15994413]
31. Maltez JM, Nunziato DA, Kim J, Pitt GS. Essential Ca(V)beta modulatory properties are AID-independent. *Nat Struct Mol Biol* 2005;12:372–377. [PubMed: 15750602]
32. Bunemann M, Gerhardstein BL, Gao T, Hosey MM. Functional regulation of L-type calcium channels via protein kinase A-mediated phosphorylation of the beta(2) subunit. *The Journal of biological chemistry* 1999;274:33851–33854. [PubMed: 10567342]
33. Ganesan AN, Maack C, Johns DC, Sidor A, O'Rourke B. {beta}-Adrenergic Stimulation of L-type Ca²⁺ Channels in Cardiac Myocytes Requires the Distal Carboxyl Terminus of {alpha}1C but Not Serine 1928. *Circ Res.* 2006
34. Beguin P, Mahalakshmi RN, Nagashima K, Cher DH, Ikeda H, Yamada Y, Seino Y, Hunziker W. Nuclear sequestration of beta-subunits by Rad and Rem is controlled by 14-3-3 and calmodulin and reveals a novel mechanism for Ca²⁺ channel regulation. *J Mol Biol* 2006;355:34–46. [PubMed: 16298391]
35. Finlin BS, Correll RN, Pang C, Crump SM, Satin J, Andres DA. Analysis of the complex between Ca²⁺ channel beta-subunit and the Rem GTPase. *The Journal of biological chemistry* 2006;281:23557–23566. [PubMed: 16790445]
36. Hohaus A, Person V, Behlke J, Schaper J, Morano I, Haase H. The carboxyl-terminal region of ahnak provides a link between cardiac L-type Ca²⁺ channels and the actin-based cytoskeleton. *Faseb J* 2002;16:1205–1216. [PubMed: 12153988]
37. Dzhura I, Wu Y, Colbran RJ, Corbin JD, Balsler JR, Anderson ME. Cytoskeletal disrupting agents prevent calmodulin kinase, IQ domain and voltage-dependent facilitation of L-type Ca²⁺ channels. *J Physiol* 2002;545:399–406. [PubMed: 12456820]
38. Bodi I, Mikala G, Koch SE, Akhter SA, Schwartz A. The L-type calcium channel in the heart: the beat goes on. *J Clin Invest* 2005;115:3306–3317. [PubMed: 16322774]




Figure 1. CaMKII efficiently phosphorylates all of the β subunit isoforms *in vitro*

A. Initial rate of phosphorylation of GST- β isoforms by CaMKII δ 2 at 4°C. Data (mean of 2 experiments) are plotted relative to the phosphorylation of GST- β_{2a} after 20 min. GST- β_{2a} , \circ . GST- β_{1b} , \bullet . GST- β_3 , \blacktriangledown . GST- β_4 , \triangle . **B.** Stoichiometry of phosphorylation of GST- β isoforms by increasing concentrations of CaMKII δ 2 after 20 minutes incubation at 30°C (mean \pm sem of 3 experiments). Symbols as in panel A. *: phosphorylation of GST- β_3 by 1000 nM CaMKII was significantly less than that of GST- β_{2a} ($p < 0.05$). Other differences were not statistically significant.




Figure 2. CaMKII association with VGCC β subunit isoforms

A. Schematic domain structure of the β_{2a} subunit showing SH3 and guanylate kinase-like (GK) domains, with the C-terminal CaMKII-binding domain containing the Thr498 phosphorylation site (indicated by the “P”). The amino acid sequence of the CaMKII binding domain in β_{2a} is aligned with similar sequences from other β isoforms and the CaMKIIδ autoregulatory domain (surrounding Thr287); identical residues are shown in gray boxes. **B.** GST-β subunit isoforms were immobilized in glutathione-coated multi-well plates (100 pmol/well) and incubated with ³²P-labeled Thr287 autophosphorylated CaMKIIδ2 (50 nM subunit). **C.** Concentration dependent binding of Thr287 autophosphorylated CaMKIIδ2 to GST-β_{2a} (○) and GST-β_{1b} (●)

in glutathione-coated multi-well plates. Both panels B and C plot binding as $\text{mean} \pm \text{sem}$ from 4 observations (β_{2a} and β_{1b}) or the mean of 2 observations (β_3 and β_4).




Figure 3. Prephosphorylation of GST- β_{2a} and GST- β_{1b} inhibits CaMKII binding

GST- β isoforms were preincubated with activated CaMKII δ_2 in the presence or absence of ATP, as indicated. Reactions were resolved by SDS-PAGE and transferred to nitrocellulose membrane for overlay with $[^{32}\text{P}]$ -labeled Thr287-autophosphorylated CaMKII δ_2 . The top panel shows a protein stain of the membrane prior to overlay and a representative autoradiograph to detect bound CaMKII. Binding was quantified using a phosphoimager and normalized to the binding detected using non-phosphorylated GST- β_{2a} ; the mean \pm sem from 3 experiments is plotted. Data were analyzed by 2-way ANOVA: *: $p < 0.001$ vs. binding to non-phosphorylated GST- β_{2a} . #: $p < 0.05$ vs. binding to the corresponding non-phosphorylated protein.




Figure 4. Phosphorylation of GST- β_{2a} by CaMKII is independent of stable binding
A. Initial rate for phosphorylation of GST- β_{2a} (wild type, \bullet : T498A, \blacktriangledown : T498E, \circ : L493A, \triangle) by CaMKII (10 nM) at 4°C. Data are the mean \pm sem of 3–4 observations normalized to the phosphorylation of the wild-type protein after 20 minutes. **B.** Wild type or mutated GST- β_{2a} proteins or GST alone were incubated for 20 min at 30°C with activated CaMKII (10 nM) and [γ -³²P]ATP. Aliquots of the reactions were resolved by SDS-PAGE and transferred to nitrocellulose membranes. Membranes were stained for total protein (bottom), and then immunoblotted using a phospho-specific antibody that was raised to phospho-Thr286 in CaMKII α , but also detects phospho-Thr498 in β_{2a} (middle). Total phosphorylation of β_{2a} was detected by autoradiography (top gel). The stoichiometry of total phosphorylation (bar graph)

was estimated after spotting parallel aliquots on phosphocellulose papers (see Methods). Data are plotted as mean \pm sem (n=4 experiments). *: p<0.01 vs. WT.




Figure 5. Pre-phosphorylation of β_{2a} at Thr498 inhibits the association of CaMKII
A. Wild type and mutated GST-β_{2a} proteins were preincubated with activated CaMKIIδ2 in the presence or absence of ATP. Samples were resolved by SDS-PAGE and transferred to nitrocellulose membranes to detect total protein loaded (top) and to probe with [³²P]CaMKIIδ2 by overlay assay and autoradiography (middle). Binding was quantified using a phosphoimager and normalized to binding to non-phosphorylated wild type protein: the mean ±sem from 3 experiments is plotted. #: p<0.001 vs. corresponding non-phosphorylated protein. *: p<0.001 vs. non-phosphorylated GST-β_{2a} wild type. Further post analyses using t tests showed that pre-phosphorylation significantly reduced CaMKII binding to the L493A mutant (†: p<0.05).




Figure 6. Phosphorylation of β_{2a} at Thr498 enhances dissociation of the CaMKII- β_{2a} complex
Complexes of GST- β_{2a} (WT or T498A) and [32 P-T 286]CaMKII α in glutathione-coated multi-well plates were incubated with or without ATP in a dissociation buffer (see Methods). At the indicated times, dissociated [32 P-T 286]CaMKII α was removed from wells and quantified by scintillation counting. Data points represent mean \pm sem (n=3) (n=2 for β_{2a} T498A in absence of ATP); error bars lie within symbols of some data points.




Figure 7. CaMKII interaction with β_{2a} is regulated by Thr498 phosphorylation *in situ*
 Myc-tagged T287D-CaMKII δ 2 was co-expressed with or without FLAG-tagged wild type or mutated β_{2a} proteins in HEK293 cells. Aliquots of the cell lysates (Inputs), FLAG immune supernatants (Super) and FLAG immune complexes (Pellets) were immunoblotted. **A:** Probing FLAG immune complexes using antibodies raised to phospho-Thr286 in CaMKII α showed that WT and L493A β_{2a} are partially phosphorylated at Thr498 in intact cells. **B:** Probing FLAG immune complexes using CaMKII δ antibodies revealed that CaMKII was associated with WT and T498A β_{2a} but could not be detected in the T498E and L493A β_{2a} immune complexes. The data are representative of >4 experiments.

Table 1
Disruption of CaMKII binding to GST- β_{2a} by site-directed mutagenesis

Purified GST- β_{2a} proteins (wild type or with the indicated point mutations) or GST alone were immobilized in a glutathione-coated 96-well plate (100 pmol/well) and then incubated with ^{32}P -labeled Thr287-autophosphorylated CaMKII δ_2 (100 nM subunit). After washing, bound CaMKII was quantified by scintillation counting. The data indicate the mean \pm sem (n = 3 experiments).

GST- β_{2a} Protein	CaMKII bound (<i>pmol subunit</i>)
Wild type	7.9 \pm 0.3
T498A	6.2 \pm 0.8
T498E	0.51 \pm 0.16
L493A	0.14 \pm 0.03