



Published in final edited form as:

Psychopharmacology (Berl). 2009 September ; 205(4): 529. doi:10.1007/s00213-009-1562-z.

The role of GABA_A receptors in the acute and chronic effects of ethanol: a decade of progress

Sandeep Kumar¹, Patrizia Porcu¹, David F. Werner¹, Douglas B. Matthews³, Jaime L. Diaz-Granados³, Rebecca S. Helfand³, and A. Leslie Morrow^{1,2}

Sandeep Kumar: ; Patrizia Porcu: ; David F. Werner: ; Douglas B. Matthews: ; Jaime L. Diaz-Granados: ; Rebecca S. Helfand: ; A. Leslie Morrow: morrow@med.unc.edu

¹ Department of Psychiatry, Bowles Center for Alcohol Studies, University of North Carolina School of Medicine, 3027 Thurston-Bowles Building, CB #7178, Chapel Hill, NC 27599-7178, USA

² Department of Pharmacology, Bowles Center for Alcohol Studies, University of North Carolina School of Medicine, 3027 Thurston-Bowles Building, CB #7178, Chapel Hill, NC 27599-7178, USA

³ Department of Psychology and Neuroscience, Baylor University, Waco, TX, USA

Abstract

The past decade has brought many advances in our understanding of GABA_A receptor-mediated ethanol action in the central nervous system. We now know that specific GABA_A receptor subtypes are sensitive to ethanol at doses attained during social drinking while other subtypes respond to ethanol at doses attained by severe intoxication. Furthermore, ethanol increases GABAergic neurotransmission through indirect effects, including the elevation of endogenous GABAergic neuroactive steroids, presynaptic release of GABA, and dephosphorylation of GABA_A receptors promoting increases in GABA sensitivity. Ethanol's effects on intracellular signaling also influence GABAergic transmission in multiple ways that vary across brain regions and cell types. The effects of chronic ethanol administration are influenced by adaptations in GABA_A receptor function, expression, trafficking, and subcellular localization that contribute to ethanol tolerance, dependence, and withdrawal hyperexcitability. Adolescents exhibit altered sensitivity to ethanol actions, the tendency for higher drinking and longer lasting GABAergic adaptations to chronic ethanol administration. The elucidation of the mechanisms that underlie adaptations to ethanol exposure are leading to a better understanding of the regulation of inhibitory transmission and new targets for therapies to support recovery from ethanol withdrawal and alcoholism.

Keywords

Alcohol; GABA_A receptor; Neuroactive steroids; Protein kinase C

Introduction

The past decade has brought many advances in our understanding of ethanol's acute and chronic actions mediated by gamma-aminobutyric acid type A (GABA_A) receptors in the central nervous system (CNS). The field has uncovered the GABAergic mechanisms that underlie many of the behavioral effects of ethanol, including its anxiolytic, anticonvulsant, sedative-hypnotic, cognitive-impairing, and motor incoordinating actions. These mechanisms include direct and indirect effects on GABA_A receptors as well as effects on GABA release and the

synthesis and availability of endogenous neuroactive steroids. The effects of chronic ethanol exposure have also been more clearly elucidated in rodents, monkeys, and humans, and there is a better understanding of the intracellular mechanisms that mediate adaptations of GABA_A receptors to ethanol. The purpose of this article is to highlight the advances of the past decade and to underscore new insights into the mechanisms of ethanol's actions and adaptations, particularly for their therapeutic relevance in the development of medications to support recovery from ethanol dependence and treatment of alcoholism.

GABA_A receptors are a family of chloride ion channels that predominately mediate rapid inhibitory neurotransmission throughout the CNS. The neurotransmitter GABA binds to these receptors, changing their conformation state and thereby opening the pore to allow chloride ions (Cl⁻) to pass down an electrochemical gradient. The flux of chloride ions hyperpolarizes the membrane leading to neuronal inhibition. However, it should be noted that, in developing animals, neurons have a higher intracellular chloride concentration at an early stage leading to an efflux of chloride and excitatory actions of GABA in immature neurons (Ben-Ari 2002).

GABA_A receptors are heteromeric protein complexes consisting of several homologous membrane-spanning glycoprotein subunits. Molecular cloning has revealed multiple GABA_A receptor subunits that can be divided by homology into subunit classes with several members: α (1–6), β (1–3), γ (1–3), δ , ϵ , θ , π , and ρ (1–3) (Olsen and Sieghart 2008; Sieghart and Sperk 2002). The large number of GABA_A receptor subunits generates the potential for various subunit compositions that may account for variable sensitivity to modulatory drugs such as benzodiazepines, barbiturates, neuroactive steroids, and possibly ethanol and general anesthetics. The GABA_A receptor α 1 subunit is the most abundant α subunit in adult brain, highly expressed throughout most brain regions and is a component of ~50% of all GABA_A receptors (Kralic et al. 2002a). The analysis of subunit composition by co-immunoprecipitation studies has demonstrated that 98% of α 1 subunit-containing receptors are assembled with a γ or δ subunit (Sieghart and Sperk 2002). In contrast, a significant fraction of α 4 subunit-containing receptors are comprised of α 4 and β subunits only (Bencsits et al. 1999). GABA_A receptors also associate with various other proteins that anchor the receptor, regulate phosphorylation state and trafficking, and modulate receptor function under various conditions. These protein interactions are not yet well understood but clearly influence GABA_A receptor assembly, expression on the membrane surface, endocytosis, and channel function that ultimately controls the inhibitory tone of the CNS.

GABA_A receptors can exist as either synaptic or extrasynaptic receptors that may facilitate rapid changes in inhibition. Most synaptic GABA_A receptors are composed of two α , two β , and one γ subunit, where the γ subunit is located between an α and β subunit (Sieghart et al. 1999; Tretter et al. 1997) and contribute to synaptically mediated (phasic) inhibition. In contrast, tonic inhibition is due to highly sensitive GABA_A receptors, activated by ambient extracellular GABA or from 'spillover' of GABA from synaptic signaling, thought to be in the range of 100 nM to 1 μ M (Santhakumar et al. 2006; Tossman et al. 1986). The subunit composition of GABA_A receptors at synaptic and extrasynaptic sites are thought to be different. Studies have shown that the α 1 subunit is usually expressed in synaptic GABA_A receptors while the α 4 subunit predominantly occurs in extrasynaptic receptors (Mody 2001; Olsen and Sieghart 2009). GABA_A receptor δ subunits are so far found to be exclusively present in extrasynaptic sites (Farrant and Nusser 2005). The α 4 subunit has relatively low expression in the CNS (Wisden et al. 1992) and α 4 β δ GABA_A receptors are exclusively extrasynaptic (Wei et al. 2003), whereas α 4 β γ 2 may be localized at the synapse, as well as extrasynaptically (Hsu et al. 2003; Liang et al. 2006). Given this distinction, the study of synaptic vs. extrasynaptic GABA_A receptors and the effect of ethanol on these receptors is essential to elucidate the mechanism involved in development of ethanol actions including tolerance and dependence. Additionally, the role of various subunits in GABA_A receptor-mediated actions has been

studied by recombinant expression studies *in vitro* and by genetically modified mouse models and pharmacologic modulators *in vivo*. In this review, we address the role of various subunits in ethanol actions by referring to various subunit-containing receptors. For example, the $\alpha 1$ subunit-containing receptors will be referred to as $\alpha 1$ -GABA_A receptors.

Acute effects of ethanol on GABA_A receptors

Behavioral evidence of GABAergic involvement in ethanol effects

Alcohol consumption produces many well-known behavioral effects including anxiolysis, impaired motor coordination, impaired cognitive function, sedation, hypnosis, anticonvulsant, and pro-aggressive action. Strikingly, many of these behavioral effects of ethanol overlap with the effects of GABA_A receptor agonists and can be altered by GABA_A receptor modifiers. GABA_A receptor agonists (benzodiazepines, muscimol) increase ethanol responses whereas inverse agonists and antagonists (Ro15-4513, picrotoxin, bicuculline) decrease ethanol responses (see Grobin et al. 1998 for review).

Genetically modified mouse models such as transgenics and global knockouts have increased our understanding of GABA_A receptor involvement in behavioral ethanol actions (see Boehm et al. 2006, 2004b; Crabbe et al. 2006 for more detailed reviews). For instance, the locomotor-stimulant effects of ethanol appear to be modulated by $\alpha 1$ -GABA_A receptors. The $\alpha 1$ -GABA_A receptor knockout mice have increased ethanol-induced activity (Blednov et al. 2003b; June et al. 2007; Kralic et al. 2003). Ethanol's sedative-hypnotic effects may also involve interactions with $\alpha 1$ -GABA_A receptors since male $\alpha 1$ -GABA_A receptor knockout mice have reduced loss of righting reflex duration (Blednov et al. 2003a). However, this effect was not observed in other studies (Kralic et al. 2003), possibly due to variation in genetic background of the strains used to create separate knockout mouse lines. Further, $\alpha 2$ -GABA_A receptors are implicated in the sedative-hypnotic effects of ethanol, as shown by reduced ethanol-induced loss of righting reflex in $\alpha 2$ -GABA_A receptor knockout animals (Boehm et al. 2004b). Studies using benzodiazepine-insensitive histidine101 to arginine (H101R) knockin mice have also indicated a role for $\alpha 2$ -GABA_A receptors in ethanol's sedative-hypnotic effects. The combined effect of subthreshold doses of diazepam and ethanol resulted in a loss of righting reflex in $\alpha 1$ -, $\alpha 3$ -, and $\alpha 5$ -, but not $\alpha 2$ - (H101R) GABA_A receptor knockin mice (Tauber et al. 2003). Additionally, $\beta 2$ -GABA_A receptors may also play a role in ethanol's sedative-hypnotic effects, as male $\beta 2$ -GABA_A receptor knockout mice displayed reduced loss of righting reflex duration compared to controls (Blednov et al. 2003a). Surprisingly, given the similarity of benzodiazepine and ethanol effects on anxiety, the $\alpha 1$ -, $\alpha 2$ -, $\alpha 4$ -, $\alpha 5$ -, $\gamma 2L$ -, and δ -GABA_A receptor knockout mice did not display changes in ethanol-induced anxiolysis (Boehm et al. 2004b; Chandra et al. 2008; Homanics et al. 1999a; Kralic et al. 2003; Mihalek et al. 2001). Further studies are clearly needed to elucidate whether specific GABAergic receptor subtypes mediate the anxiolytic effects of ethanol.

Several studies suggest that GABA_A receptors may be involved in ethanol preference. $\alpha 1$ -GABA_A receptor knockouts consumed less ethanol in a two-bottle choice drinking study, but also showed reduced saccharin preference compared to wildtype mice (Blednov et al. 2003b; June et al. 2007). In studies of ethanol reinforcement, $\alpha 1$ -GABA_A receptor knockouts exhibited reduced operant responding for ethanol, although this was also accompanied by a reduced responding for sucrose (June et al. 2007). In contrast, $\alpha 5$ - and δ -GABA_A receptor knockouts showed reduced ethanol preference with no changes in saccharin preference (Boehm et al. 2004b; Mihalek et al. 2001; Stephens et al. 2005). Additionally, a recent study utilizing viral vectors (Rewal et al. 2009) demonstrates that reduction of $\alpha 4$ -GABA_A receptor subunits in the nucleus accumbens shell by RNA interference decreases ethanol consumption and preference. Further studies in other brain areas as well as studies targeting other subunits will be of great interest. Remarkably, many other ethanol-induced behaviors remain unaltered in mice with

genetically modified GABA_A receptors. For instance, the δ -GABA_A receptors do not appear necessary for the discriminative stimulus effects of ethanol or substitution by barbiturates, benzodiazepines, and neuroactive steroids in knockout mice (Shannon et al. 2004). One caveat to these studies is that observed effects or lack of effects may be due to compensatory changes that are sometimes observed in knockout models and should be taken into consideration. Nonetheless, these studies implicate GABA_A receptor subtypes in some behavioral actions of ethanol.

Pharmacologic dissection of GABA_A receptors in ethanol action has also identified a potential role for GABA_A receptor subtypes in ethanol reinforcement. For instance, α 1-GABA_A receptors in the ventral pallidum are involved in the reinforcing properties of ethanol (Harvey et al. 2002). Systemic administration or infusion into the ventral pallidum of an α 1 selective mixed benzodiazepine agonist–antagonist, 3-propoxy- β -carboline hydrochloride, reduces ethanol-maintained responding in ethanol-preferring rats. α 1-GABA_A receptors have also been implicated in ethanol-related heightened aggressive behavior (de Almeida et al. 2004). Administration of the benzodiazepine antagonist flumazenil or the preferential α 1 antagonist, β -carboline-3-carboxylate-*t*-butyl ester, decreases ethanol-related aggression in mice trained to self-administer ethanol. The α 5-GABA_A receptors appear to play a role in the rewarding, motor-impairing, and sedative effects of ethanol in rats, since the selective inverse-agonist RY023 impacts these actions (Cook et al. 2005). Furthermore, interactions between ethanol and benzodiazepines increase spatial memory deficits in animals (Takiguchi et al. 2006) and humans (Simpson and Rush 2002) implicating γ 2-GABA_A receptors in spatial memory impairment. The development of more subtype specific/selective compounds will aid in further elucidating the role of specific GABA_A receptor subtypes in ethanol action. Newer, more selective compounds such as L-838,417 (which has selectivity for α 2-/ α 3-/ α 5-GABA_A receptors) will be particularly useful in identifying the role of specific receptor subtypes (see Atack 2003 for review).

The discriminative stimulus paradigm has been used as an *in vivo* assay of receptor-mediated activity and can be a useful tool to define the neurotransmitter systems that underlie the behavioral effects of a given dose and class of drugs (Holtzman 1990). Positive modulators of GABA_A receptors, like benzodiazepines, barbiturates, and certain neuroactive steroids, all produce ethanol-like discriminative stimulus effects in water vs. ethanol discrimination in pigeons, mice, rats, gerbils, and cynomolgus monkeys (Grant 1999; Grant et al. 1996, 2008a; Shelton and Grant 2002). Recent studies have aimed to understand the contribution of different GABA_A receptor subtypes to the discriminative stimulus properties of ethanol. Studies in non-human primates show that ethanol and zolpidem share similar discriminative stimulus effects suggesting that ethanol's sedative-hypnotic actions may involve α 1-GABA_A receptors (Helms et al. 2008). However, zolpidem antagonism by Ro15-4513 may involve additional zolpidem-sensitive α 2-/ α 3-/ α 5-GABA_A receptor subtypes (Helms et al. 2008). In contrast, Platt and collaborators (Platt et al. 2005) reported a prominent role for the α 5- but not the α 1-GABA_A receptors in the discriminative stimulus effects of ethanol and the ethanol-like effects of benzodiazepines in squirrel monkeys. However, the ethanol-like discriminative stimulus effects are complex. For instance, the discriminative stimulus effects of low doses of ethanol (1.0 g/kg) are predominantly mediated by GABA_A receptor positive modulation, while higher doses (2.0 g/kg) require other systems such as *N*-methyl-D-aspartate (NMDA) receptors (Grant 1999). Indeed, ethanol's discriminative stimulus effects are mediated by interactions of GABA_A, NMDA, and metabotropic glutamate receptor subtype 5 (mGluR5), located in specific limbic brain regions such as nucleus accumbens core and amygdala (Besheer et al. 2003; Besheer and Hodge 2005; Hodge and Cox 1998).

Direct effects on GABA_A receptor subtypes

Low dose effects of ethanol (≤30 mM) on tonic inhibition—While much work has shown that ethanol enhances GABA_A receptor potentiation, the exact mechanism of ethanol actions on these receptors remains unclear. Recent work has addressed the contribution of specific GABA_A receptor subtypes in distinct cellular localizations at ethanol concentrations typically found in social settings (3–30 mM). Several groups have shown that ethanol enhances the inhibition of $\alpha 4/6\beta\delta$ -GABA_A receptors (found extra-synaptically), whereas receptors containing a $\gamma 2$ subunit (found synaptically) are significantly less sensitive to the ethanol enhancement of GABAergic responses (Sundstrom-Poromaa et al. 2002; Wallner et al. 2003; Wei et al. 2004). Typically, δ -GABA_A receptors are found extrasynaptically and are commonly associated with $\alpha 4/\alpha 6$ subunits (Jones et al. 1997; Sur et al. 1999; Wei et al. 2003). Additionally, these receptors have a greater affinity for GABA (Brown et al. 2002; Wallner et al. 2003) and knockout models display substantially reduced/absent tonic inhibition (Brickley et al. 2001; Chandra et al. 2006; Stell et al. 2003). Importantly, δ -GABA_A receptor knockout mice also have reduced/ablated enhancement of tonic current by ethanol (Glykys et al. 2007). Therefore, it is likely that ethanol actions on $\alpha 4/6\beta\delta$ -GABA_A receptors involve the modulation of tonic inhibition *in vivo*.

In parallel, the ‘ethanol antagonist’ Ro15-4513 has been shown to bind with high affinity to δ -GABA_A receptors that are activated by low concentrations of ethanol (Hanchar et al. 2006). Ethanol-induced potentiation of $\alpha 4\beta\delta$ -GABA_A receptors by low ethanol concentrations is blocked by Ro15-4513 (Wallner et al. 2006). Changes in tonic inhibition may alter ethanol-related behavioral responses. Hanchar et al. (2005) noted that a naturally occurring mutation found in outbred Sprague-Dawley rats (arginine100 to glutamine, R100Q; also found in Sardinian non-preferring rats and ethanol non-tolerant rats) substantially increases the sensitivity of $\alpha 6\beta 3\delta$ -GABA_A receptors to ethanol. Importantly, rats with the naturally occurring mutation exhibit greater enhancement of tonic inhibition in cerebellar granule cells as well as greater motor impairment due to ethanol. Moreover, it now appears that tonic current is not solely dependent on the presence of $\alpha 4\delta$ - or $\alpha 6\delta$ -GABA_A receptors. $\alpha 1\delta$ -GABA_A receptors also mediate ethanol-induced increases in tonic inhibition in hippocampal interneurons in $\alpha 4$ -GABA_A receptor knockout mice (Glykys et al. 2007). Further studies of ethanol actions on these receptors are needed.

High dose effects of ethanol (>30 mM) on tonic inhibition—While the previously described studies have indicated a major role of ethanol potentiation of GABA inhibition at $\alpha 4/6\beta\delta$ extrasynaptic receptors, these observations are controversial. In a recent study by Borghese et al. (2006a), rat and human $\alpha 4\beta 3\delta$ -GABA_A subunits expressed in oocytes, $\alpha 4\beta 3\delta$ subunits stably transfected in fibroblasts, and slice recordings conducted in dentate granule cells all failed to exhibit enhanced GABA_A receptor potentiation by low doses of ethanol. Further, effects of lower doses of ethanol (30 mM or less) are not observed in cerebellar granule cells (Casagrande et al. 2007; Yamashita et al. 2006) or ventrobasal thalamic neurons (Jia et al. 2008a) and dentate gyrus granule cells that express $\alpha 4\beta 2\delta$ receptors (Liang et al. 2008). However, the latter study did observe some detectable effects at 10 mM ethanol. One possibility may be the orientation of receptor subunits within the GABA_A receptor pentamer. However, Kaur et al. (2009) have shown that 30 mM ethanol did not display potentiation in concatenated $\alpha 1\beta 3\delta$ receptors. In agreement with the lack of a low dose effect, autoradiographic experiments also suggest that Ro15-4513 binding is not altered by low concentrations of ethanol (Korpi et al. 2007). Additionally, the naturally occurring R100Q mutation in $\alpha 6$ -GABA_A receptors does not increase ethanol sensitivity at extrasynaptic receptors in outbred rats or in ethanol non-tolerant rats as described above (Botta et al. 2007a; Valenzuela et al. 2005). In agreement with the lack of a low dose effect, $\alpha 4$ - and δ -GABA_A receptor knockout mice do not show any major changes in acute ethanol-related behaviors (Chandra et al. 2008; Mihalek et al. 2001).

Nonetheless, electrophysiologic studies utilizing $\alpha 4$ -GABA_A receptor subunit knockout mice did not show ethanol-enhanced tonic inhibition compared to controls at 50 mM (Jia et al. 2008a; Liang et al. 2008). Many of these studies assessing low dose ethanol effects on extrasynaptic receptors and tonic inhibition are summarized in Table 1.

The discrepancies in ethanol actions observed across various laboratories may indicate very specific requirements for low dose effects of ethanol on extrasynaptic GABA_A receptors. It is also possible that presynaptic ethanol actions may play a role. Recent studies suggest that increased firing onto cerebellar granule cells increases extrasynaptically available GABA (Botta et al. 2007b; Carta et al. 2004; Valenzuela et al. 2005) that may contribute to changes in tonic inhibition. Other mechanisms may include increases in endogenous modulators that are particularly potent at extrasynaptic GABA_A receptors. Ethanol increases GABAergic neuroactive steroids in slice preparations (see 'Ethanol-induced elevation of neuroactive steroids'). Ethanol also increases taurine (De Witte et al. 1994), which has been shown to enhance tonic inhibition at low concentrations (Jia et al. 2008b). The possibility exists that other unknown chemicals are also released by ethanol. Alternatively, it is possible that the intracellular milieu of different preparations, including second messengers, kinases and phosphatases, may contribute to ethanol actions. Indeed, recent work by Choi et al. (2008) using knockout mice and stably transfected cells has shown that PKC δ activity may be required for ethanol enhancement of extrasynaptic GABA_A receptor responses and this requirement may explain the lack of consistent effects in prior studies. Irrespective of the low ethanol dose discrepancy, it is clear that ethanol does enhance tonic inhibition, and that low dose actions of ethanol may have multiple mechanisms. Further studies are needed to clarify these discrepancies.

High dose effects of ethanol (>30 mM) on phasic inhibition—GABA_A receptor subtypes associated with phasic inhibition are modulated by higher doses of ethanol. Using chimeric receptors, Mihic et al. (1997) identified a 45-amino-acid region spanning the second and third transmembrane domains required for the direct enhancement of glycine receptors by ethanol and volatile anesthetics. Moreover, the mutation of two residues (serine270 and alanine291) produces receptors insensitive to ethanol-induced potentiation. Interestingly, the volume of this putative ethanol 'binding pocket' dictates receptor sensitivity to ethanol (Wick et al. 1998; Ye et al. 1998). GABA_A receptors possessing the serine270 mutation have altered receptor gating and are hypersensitive to GABA (Ueno et al. 2000). Moreover, knockin mice with the serine270 to histidine (S270H) mutation show marked abnormal behaviors, premature mortality, and prolonged miniature inhibitory post-synaptic currents (mIPSCs) (Homanics et al. 2005). Nonetheless, the mutation of leucine277 to alanine (L277A) in combination with S270H restored GABA sensitivity to near normal affinity while retaining ethanol and volatile anesthetic insensitivity *in vitro* (Borghese et al. 2006b). Importantly, gene knockin mice with both mutations show decreased sensitivity to ethanol potentiation (80 mM) and recover more quickly from ethanol's motor ataxic effects, but do not differ on many other ethanol-related behaviors (Werner et al. 2006). These studies implicate a molecular site of action on GABA_A receptors found synaptically that contributes to high dose ethanol action.

A membrane-spanning region of the $\beta 3$ -GABA_A receptor subunit is also implicated in GABA_A receptor responses to higher amounts of ethanol. Work by Wallner et al. (2006) has indicated that while Ro15-4513 blocked ethanol-induced potentiation at low doses, higher concentrations of ethanol still enhanced GABA_A receptor function in the presence of Ro15-4513. Mutations in equivalent positions in $\beta 1$ -GABA_A receptors also modulate ethanol responses like intravenous anesthetic GABA_A receptor modulators etomidate and propofol in $\beta 3$ -GABA_A receptors (Jurd et al. 2003; Mihic et al. 1997). This evidence supports the notion that ethanol and Ro15-4513 may bind to the same site on GABA_A receptors at low ethanol

concentrations whereas a second ethanol-binding site involving α and β subunits may be responsible for ethanol responses at higher concentrations.

Overall, a considerable amount of evidence continues to accumulate that suggests GABA_A receptors may directly modulate some ethanol effects. More so, it is becoming apparent that different GABA_A receptor subtypes not only respond to different ethanol concentrations but their cellular localization may contribute to various ethanol-related responses.

Indirect ethanol effects on GABA_A receptor subtypes

Ethanol-mediated post-translational modification of GABA_A receptors—Protein phosphorylation regulates GABA_A receptor function under numerous physiologic conditions (Brandon et al. 2000; Kellenberger et al. 1992; Krishek et al. 1994; Kumar et al. 2005; Leidenheimer et al. 1992; Poisbeau et al. 1999), including alcoholism and epilepsy (Kumar et al. 2006; Niimura et al. 2005; Rakhade et al. 2008). Several protein kinases, including PKC and PKA, have consensus sites on GABA_A receptor subunits and can phosphorylate these subunits (Brandon et al. 2000; McDonald and Moss 1997). Phosphorylation of GABA_A receptor subunits can modify GABA binding to its receptor (Oh et al. 1999), channel conductance, and possibly internalization (Moss and Smart 2001). Additionally, studies using genetically modified mice clearly demonstrate that the pharmacological and behavioral effects of ethanol are partially mediated by PKC activity (Choi et al. 2008; Harris et al. 1995; Hodge et al. 1999; Proctor et al. 2003).

PKC phosphorylation: A large volume of recent work has focused on the role of PKC as a regulator of GABA_A receptor function following ethanol exposure. Various PKC isoforms associate with GABA_A receptors and therefore may alter receptor subunit phosphorylation and function. PKC β associates with GABA_A receptor β subunit and alters GABA_A receptor phosphorylation and function (Connolly et al. 1999). PKC γ co-immunoprecipitates with $\alpha 1$ - and $\alpha 4$ -GABA_A receptors in the cerebral cortex (Kumar et al. 2002) and PKC ϵ co-localizes with various GABA_A receptor subunits across the brain (Olive and Hodge 2000). Similarly, PKC δ alters the function of extrasynaptic GABA_A receptors (Choi et al. 2008). Despite the existence of several isoforms of PKC, genetic deletion of PKC γ , PKC ϵ , and PKC δ has clearly demonstrated that these isoforms are major players in ethanol-mediated modulation of GABA_A receptor function both *in vitro* and *in vivo*. For example, PKC γ knockout mice show reduced sensitivity to the anxiolytic effects of both intoxicating and sedative doses of ethanol (Bowers et al. 2001), while PKC δ knockout mice are insensitive to the ataxic effects of ethanol (Choi et al. 2008) and PKC ϵ knockout mice show increased ethanol sensitivity (Hodge et al. 1999). Furthermore, mice lacking the gene for PKC γ show a significant reduction in ethanol potentiation of GABA responses compared to responses in wildtype mice (Harris et al. 1995; Proctor et al. 2003). In contrast, ethanol and flunitrazepam potentiation of muscimol-stimulated chloride (Cl⁻) uptake is greater in microsacs from PKC ϵ knockout mice compared to wildtype controls (Hodge et al. 1999). Therefore, it appears that the ϵ and γ isoforms of PKC have opposing influences on ethanol enhancement of GABA_A receptor function. PKC δ plays a critical role in ethanol effects on extrasynaptic GABA_A receptors. PKC δ knockout mice display a reduction in ethanol-induced ataxia as well as ethanol-mediated tonic current in thalamic and dentate granule cell neurons (Choi et al. 2008). Taken together, it is likely that various PKC isoforms mediate phosphorylation of GABA_A receptors and/or associated proteins and modulate GABA_A receptor function following ethanol exposure.

Acute systemic ethanol administration directly alters PKC activity and expression in the brain (Kumar et al. 2006). However, ethanol administration to rats (2 g/kg) differentially alters PKC β , γ , and ϵ expression and translocation to the P2 fraction of rat cerebral cortex (Kumar et al. 2006). PKC β and PKC ϵ expression in subcellular fractions exhibited responses to ethanol

that likely represent increased synthesis and translocation, since increased levels of these kinases in the P2 fraction would require translocation of the enzyme. On the other hand, PKC γ expression and translocation to the P2 fraction of cortex are reduced by ethanol. These effects were associated with alterations in phosphorylation of both GABA $_A$ and NMDA receptors and/or associated proteins. Hence, ethanol differentially alters PKC expression in an isoform-specific manner and these effects are likely to contribute to GABAergic (and glutamatergic) actions of ethanol.

Phosphorylation of various GABA $_A$ receptor subunits can alter receptor function without altering receptor expression. We have shown that acute ethanol administration (2 g/kg) decreased serine phosphorylation of GABA $_A$ receptor β subunits without altering the expression of these subunits in the P2 fraction of rat cerebral cortex (Kumar et al. 2006). This suggests that ethanol may alter the phosphorylation state of receptors to alter GABA $_A$ receptor function and these effects may contribute to ethanol actions. This is consistent with the observation of increased GABA $_A$ receptor function following ethanol exposure. Likewise, reduced phosphorylation of the GABA $_A$ receptor $\gamma 2$ subunit enhances the action of ethanol (Qi et al. 2007). PKC ϵ phosphorylates the GABA $_A$ receptor $\gamma 2$ subunit at serine327 *in vitro* and $\gamma 2$ serine327 phosphorylation is reduced in mice lacking PKC ϵ . In addition, exposure to ethanol (4 g/kg) for 1 h increases $\gamma 2$ serine327 phosphorylation in the cerebellum of wildtype mice, but not PKC ϵ knockout mice and this leads to reduced GABA $_A$ receptor function in cerebellar microsacs (Qi et al. 2007). Finally, phosphorylation at serine327 on GABA $_A$ receptor $\gamma 2$ subunits is required for PKC ϵ -mediated modulation of GABA $_A$ receptor function (Qi et al. 2007). Further studies investigating post-translational modification of these receptors at PKC phosphorylation sites by specific PKC isoforms will lead to a better understanding of PKC-mediated effects of ethanol on GABA $_A$ receptors.

PKA phosphorylation: Accumulating evidence indicates that the cyclic adenosine monophosphate (cAMP)-dependent kinase PKA is involved in the neurobiological responses to ethanol. PKA is a tetramer composed of a regulatory homodimer and two catalytic subunits. PKA activation occurs when cAMP binds to the regulatory subunits of the PKA complex, liberating the catalytically active subunits, which diffuse throughout the cell and phosphorylate nearby proteins. Ethanol stimulates cAMP signaling through activation of adenosine A2a receptors. Since A2a receptors are coupled to stimulatory G proteins, this increases levels of intracellular cAMP followed by stimulation of PKA (see Diamond and Gordon 1994 for review). Recently, we have found that acute ethanol exposure increases the expression of PKA regulatory RII α and RII β subunits in the membrane fraction of rat cerebral cortex and also alters GABA $_A$ receptor expression (Kumar et al. 2008). Therefore, it appears that ethanol alters PKA expression in a subunit specific manner and may affect GABA $_A$ receptor expression.

Recent pharmacological and genetic studies have provided additional evidence for the importance of the cAMP/PKA system in regulating the neurobiological responses to ethanol. PKA inhibition (using micro-infusions of the PKA inhibitor Rp diastereomer of adenosine 3', 5'-cyclic mono-phosphorothioate, Rp-cAMPS) in the central amygdala prevents the development of anxiety in rats during ethanol withdrawal (Pandey et al. 2003). Other studies have shown that the PKA signaling pathway plays an important role in the modulation of several ethanol-induced behavioral actions such as loss of righting reflex (Maas et al. 2005; Thiele et al. 2000). For example, PKA regulatory RII β subunit mutant mice, which exhibit lower total cAMP-stimulated PKA activity, are less sensitive to the sedative-hypnotic effects of ethanol as assessed by the loss of righting reflex test (Lai et al. 2007). Furthermore, *in vivo* studies show that mutant *Drosophila* lacking production of the RII subunit of PKA are resistant to the intoxicating effects of ethanol (Park et al. 2000). In addition, PKA has been shown to alter GABA $_A$ receptor function. For example, PKA activation causes a decrease in GABA-activated currents in cultured cerebellar granule cells, hippocampal pyramidal cells,

and spinal cord neurons (Moss et al. 1992; Poisbeau et al. 1999; Porter et al. 1990; Robello et al. 1993). Taken together, it is clear that PKA also plays a role in ethanol-mediated behavior and may influence GABA_A receptor function.

Phosphorylation by fyn and tyrosine kinase: Recent studies have shown that *fyn* kinase is involved in ethanol-induced changes in GABA_A receptor function. For example, over-expression of *fyn* kinase reduces sedative-hypnotic sensitivity to ethanol (Boehm et al. 2004a). Furthermore, *fyn* kinase knockout mice are less sensitive to the sedative-hypnotic effects of etomidate (an intravenous anesthetic, selective for β 2- and/or β 3-GABA_A receptors; see Jurd et al. 2003) suggesting that *fyn* kinase alters GABA_A receptor function by acting on GABA_A receptor β 2 and β 3 subunits (Boehm et al. 2004c). Similarly, *in vitro* and *in vivo* studies have implicated tyrosine kinase in ethanol-mediated alterations of GABA_A receptor function. Ethanol exposure increases tyrosine kinase phosphorylation of the GABA_A receptor subunits α 1, β 2, and γ 2 in mouse cultured cortical neurons (Marutha Ravindran and Ticku 2006) and rat cerebral cortex (Marutha Ravindran et al. 2007). In addition, tyrosine kinase phosphorylation of GABA_A receptor γ 2 subunit regulates GABA_A receptor surface expression and function (Kittler et al. 2008). Hence, it appears that multiple kinases are involved in ethanol-mediated alteration in GABA_A receptor function. Therefore, studies of the interactions of GABA_A receptors with these kinases will lead to a better understanding of ethanol actions on GABA_A receptors.

Dephosphorylation: The phosphorylation state of a given protein is governed by the balance between protein kinases that transfer phosphate from adenosine triphosphate (ATP) to the protein (phosphorylation) and protein phosphatase that catalyzes the reverse reaction (dephosphorylation). It is now widely acknowledged that the regulation of protein phosphorylation requires coordinated control of both kinases and phosphatases. The effect of ethanol exposure on protein phosphatase expression and activity is still not known. However, phosphatase inhibitors alter the function of GABA_A receptors by reducing GABA_A receptor rundown (Huang and Dillon 1998). In addition, blockade of phosphatase activity during preparation of cerebral cortical synaptoneurosomes decreases muscimol-induced Cl⁻ uptake (Kumar et al. 2005). Hence, the balance between phosphatase and kinase activity is important for GABA_A receptor function. Further studies on the effects of ethanol on protein phosphatase activity are warranted.

Ethanol-induced elevation of neuroactive steroids—Neuroactive steroids are endogenous neuromodulators, synthesized *de novo* in the brain as well as adrenal glands, ovaries, and testes (see Biggio and Purdy 2001 for review). Among these compounds, the 3 α , 5 α - and 3 α ,5 β -reduced metabolites of progesterone (Majewska et al. 1986; Morrow et al. 1987), deoxycorticosterone (Majewska et al. 1986; Morrow et al. 1987), dihydroepiandrosterone (DHEA) (Frye et al. 1996; Kaminski et al. 2005; Park-Chung et al. 1999), and testosterone (Kaminski et al. 2005, 2006) enhance GABAergic neurotransmission. Their systemic administration induces anxiolytic, anticonvulsant, sedative-hypnotic, and cognitive-impairing effects, similar to other GABA_A receptor positive modulators and ethanol (see Morrow et al. 2006 for review).

Systemic administration of moderate doses of ethanol (1–2.5 g/kg) increases brain and plasma levels of (3 α ,5 α)-3-hydroxypregnan-20-one (3 α ,5 α -THP), (3 α ,5 α)-3,21-dihydroxypregnan-20-one (3 α ,5 α -THDOC), and their precursors in rodents (Barbaccia et al. 1999; Gabriel et al. 2004; Khisti et al. 2005; Korneyev et al. 1993; Morrow et al. 1998; 1999; O'Dell et al. 2004; Serra et al. 2003; VanDoren et al. 2000). The ethanol-induced increase in neuroactive steroids is mediated by the hypothalamic–pituitary–adrenal (HPA) axis, since it is no longer observed immediately following adrenalectomy/gonadectomy in rats (Khisti et al. 2003; O'Dell et al. 2004; Porcu et al. 2004). However, ethanol can increase neuroactive steroids

in hippocampal slices from both intact (Sanna et al. 2004) and adrenalectomized/gonadectomized rats (Follesa et al. 2006) and this finding is associated with an enhancement of GABAergic inhibition that can be blocked by the neuroactive steroid biosynthesis inhibitor finasteride (Sanna et al. 2004).

Ethanol-induced elevations in neuroactive steroids reach physiologically relevant concentrations that are capable of enhancing GABAergic transmission. A large body of evidence from multiple laboratories suggests that ethanol-induced elevations of GABAergic neuroactive steroids contribute to several behavioral effects of ethanol in rodents. Neuroactive steroids have been shown to modulate ethanol's anticonvulsant effects (VanDoren et al. 2000), sedation (Khisti et al. 2003), impairment of spatial memory (Matthews et al. 2002; Morrow et al. 2001), anxiolytic-like (Hirani et al. 2005), antidepressant-like (Hirani et al. 2002), and pro-aggressive (Fish et al. 2001) actions. Most of these behavioral responses are prevented by pretreatment with finasteride and/or by prior adrenalectomy (Hirani et al. 2002; 2005; VanDoren et al. 2000). The sedative-hypnotic effect of ethanol is partially blocked by adrenalectomy (Khisti et al. 2003). Importantly, administration of 5α -dihydroprogesterone, the immediate precursor of $3\alpha,5\alpha$ -THP, to adrenalectomized rats restores effects of ethanol, showing that brain synthesis of neuroactive steroids can modulate effects of ethanol (Khisti et al. 2003). However, neuroactive steroids do not appear to influence the motor incoordinating effects of ethanol, since neither finasteride administration nor adrenalectomy diminish these actions (Khisti et al. 2004). Taken together, these studies suggest that elevations in neuroactive steroids influence many of the GABAergic effects of ethanol *in vivo* and contribute to sensitivity to the behavioral effects of ethanol.

Acute application of ethanol (50 and 100 mM) to freshly isolated rat hippocampal slices increases tissue levels of $3\alpha,5\alpha$ -THP (Sanna et al. 2004). In addition, application of progesterone (the $3\alpha,5\alpha$ -THP precursor), CB34 (a peripheral benzodiazepine receptor agonist), or γ -hydroxybutyric acid, all of which promote steroidogenesis, also increases tissue levels of $3\alpha,5\alpha$ -THP. The increase in $3\alpha,5\alpha$ -THP induced by ethanol, progesterone, CB34, or γ -hydroxybutyric acid is prevented by pretreatment with the neuroactive steroid inhibitor finasteride. Furthermore, this increase in $3\alpha,5\alpha$ -THP levels is capable of potentiating GABA_A receptor function, measured by recordings of spontaneous mIPSCs. Ethanol increases mIPSC amplitude and frequency in a time- and concentration-dependent manner: the effect is observed within the first 3 min, then it is reduced at 10 min and reappears after 30 min. In contrast, the decay time is increased 30 min after ethanol exposure. However, pretreatment with finasteride inhibits the increase in mIPSC amplitude and decay time constant at 30 min, but has no effect on mIPSC frequency, suggesting that the presynaptic action of ethanol is not mediated by an increase in neuroactive steroid content (Sanna et al. 2004).

Presynaptic GABA release—Many GABAergic effects of ethanol likely result from the ability of ethanol to enhance presynaptic GABA release in spinal cord (Ziskind-Conhaim et al. 2003), amygdala (Roberto et al. 2003; 2004), hippocampus (Ariwodola and Weiner 2004), and cerebellum (Criswell and Breese 2005). An ethanol-induced increase in GABA release would be expected to modulate all subtypes of GABA_A receptors and could clearly contribute to the behavioral effects of ethanol mediated by GABA_A receptors in brain. This effect of ethanol exhibits regional and neuronal specificity (Criswell et al. 2008), suggesting that it may contribute to heterogeneity of ethanol interactions. Moreover, chronic exposure to GABA alters GABA_A receptor $\alpha 1$ subunit expression (Calkin and Barnes 1994; Montpied et al. 1991a), internalization (Tehrani and Barnes 1997), and decreases GABA_A receptor-mediated inhibition (Roca et al. 1990), similar to the effects of ethanol on GABA_A receptor $\alpha 1$ subunit expression (Kumar et al. 2003). Thus, ethanol-induced increases in GABA release likely contribute to ethanol-induced changes in GABA_A receptor surface expression that appear to be involved in ethanol tolerance, dependence, and withdrawal.

Chronic administration of ethanol

Behavioral effects

Chronic ethanol exposure results in CNS hyperexcitability during and following withdrawal from ethanol exposure. These effects include a heightened risk for seizures, increased anxiety, hyperalgesia, and disruptions in sleep states. While the exact molecular mechanisms attributable to each of these effects are not fully understood, an overwhelming amount of evidence points to GABA_A receptor involvement (see Grobin et al. 1998 for review).

Genetically modified mouse models have contributed to our understanding of the relationship between GABA_A receptor subtype expression and CNS hyperexcitability. Recent work in the ethanol-insensitive (S270H, L277A) α 1-GABA_A receptor knockin animals indicates that α 1 subunits play a significant role in CNS excitability. Male knockin mice display enhanced ethanol withdrawal-related handling-induced convulsions (HICs) (Werner et al. 2009). Additionally, a number of mortalities were observed in the knockin mice that were attributable to seizure severity, whereas mortalities were not observed following ethanol exposure in control mice. However, knockin mice exhibited spontaneous seizures that may have kindled ethanol-withdrawal seizure susceptibility. Interestingly, α 1-GABA_A receptor knockout mice had increased seizure susceptibility to bicuculline as well as more severe seizures and death (Kralic et al. 2002a). In other studies, α 1-GABA_A receptor knockout mice did not differ in withdrawal-related HICs, but these mice consumed less ethanol prior to withdrawal than wildtype control mice, possibly masking a difference between mouse lines (Blednov et al. 2003b). Studies have also assessed the role of β subunits in ethanol withdrawal seizure severity. β 2-GABA_A receptor knockout mice had increased HICs after chronic ethanol (Blednov et al. 2003b); however, knockouts had higher ethanol intake than controls. In other studies, β 3 asparagine265 to methionine (N265M)-GABA_A receptor knockin mice had enhanced withdrawal following chronic ethanol exposure (Sanchis-Segura et al. 2007). Withdrawal from acute high doses of ethanol has also been used to assess the role of specific GABA_A receptor subtypes. The expression of γ 2 subunits, which are involved in phasic inhibition, has also been thought to play a role. Work has shown that allelic variation in *gabrg2*, the gene encoding for the γ 2 subunit, was correlated with withdrawal-related behavior (Hood and Buck 2000). However, γ 2L-GABA_A receptor knockouts had similar withdrawal-related HICs (Homanics et al. 1999a). In studies of subunits involved in tonic inhibition, δ -GABA_A receptor knockout mice displayed decreased ethanol withdrawal HICs than controls; however, this effect was dependent on background strain of the mice (Mihalek et al. 2001). Interestingly, studies in α 6-GABA_A receptor knockout mice (which pairs with δ in the cerebellum) did not differ in ethanol withdrawal-related HICs (Homanics et al. 1997). Finally, withdrawal from acute high doses of ethanol has also been used to assess the role of specific GABA_A receptor subtypes. For instance, γ 2-GABA_A receptor transgenic mice did not differ from controls (Wick et al. 2000) in sensitivity to acute withdrawal HICs (Boehm et al. 2004b).

Apart from genetically modified mouse models, other studies have also implicated GABA_A receptors in ethanol dependence. For instance, pentylenetetrazole-kindled animals displayed ethanol withdrawal behavior similar to animals exposed to repeated cycles of ethanol withdrawal (Davidson et al. 1999). Moreover, animals subjected to repeated withdrawals exhibit faster pentylenetetrazole-kindled seizure activity compared to controls and animals experiencing only a single ethanol withdrawal (Ripley et al. 2002). Additionally, while lorazepam could decrease HIC severity in repeated withdrawals, this intervention led to an exacerbated final withdrawal compared to animals that did not receive lorazepam during multiple withdrawals from ethanol (Becker and Veatch 2002). These results hint that this interactive effect between benzodiazepines and ethanol may involve GABA_A receptors. Studies with benzodiazepines have also implicated GABA_A receptors in ethanol withdrawal-related hyperalgesic effects. Diazepam reversed ethanol withdrawal hyperalgesia, but this

effect could not be reversed by flumazenil (Gatch 1999). Studies have also shown that flumazenil reduces seizure severity (Buck et al. 1991b) and reverses the anxiogenic-like effects (Moy et al. 1997) associated with ethanol withdrawal. While known as a benzodiazepine antagonist, flumazenil has also been shown to have some partial agonist activity (see File and Pellow 1986 for review) and the effects of partial agonists are enhanced following chronic ethanol exposure (Liang et al. 2004). These studies, as well as those involving genetically modified mice, suggest that specific GABA_A receptor subtypes and/or regulation of these receptors contribute to ethanol dependence and further implicate GABA_A receptors as potential therapeutic targets for treating alcoholism.

Functional adaptations

Prolonged ethanol consumption and repeated ethanol withdrawals produce many adaptations of GABA_A receptor function (Table 2). For instance, ethanol exposures result in the development of tolerance to many of the GABAergic effects of ethanol including the sedative, motor incoordinating, and acute cognitive-impairing effects of ethanol (Le et al. 1986; Silvers et al. 2003c). Withdrawal from ethanol produces marked increases in CNS excitability that form a criterion for ethanol dependence (Becker et al. 1997; Devaud et al. 1996; McCown and Breese 1990; Moy et al. 2000). Substantial evidence suggests that these behavioral and neural adaptations involve marked alterations in the pharmacological properties of GABA_A receptors (Table 2).

Alterations in GABA_A receptor subunit expression, localization, and trafficking

The development of ethanol dependence is associated with alterations in many of the functional properties of GABA_A receptors throughout the brain which is attributed to altered expression or composition of GABA_A receptors on the cell surface. Chronic ethanol administration differentially alters the expression of distinct GABA_A receptor subunit mRNA and peptide levels in various brain regions (Table 3). The levels of GABA_A receptor $\alpha 1$, $\alpha 2$, and $\alpha 3$ subunit mRNA/peptides are reduced in the cerebral cortex while $\alpha 4$, $\beta 1$, $\beta 2$, $\beta 3$, $\gamma 1$, and $\gamma 2$ subunit mRNA/peptide levels are increased in cerebral cortex following chronic ethanol exposure. Since GABA_A receptor $\alpha 1$ subunits are the most abundant subunit in the cerebral cortex, the reduction in the expression of this subunit is likely to have significant functional consequences. GABA_A receptor $\alpha 4$ subunit expression measured by immunohistochemical labeling in brain slices is sparse across most brain regions except the thalamus, dentate gyrus, and striatum (Bencsits et al. 1999; Chandra et al. 2006; Pirker et al. 2000). However, selective modulation of $\alpha 4$ subunit expression in the hippocampus, using antisense oligonucleotides, has been shown to significantly alter the functional properties of GABA_A receptors and modulate steroid withdrawal excitability (Moran et al. 1998; Smith et al. 1998). Therefore, despite the relatively low expression of $\alpha 4$ subunit peptide, modulation of $\alpha 4$ subunit expression clearly influences GABA_A receptor function and GABA-mediated behaviors *in vivo*. Recombinant expression and homologous gene deletion studies have shown that functional properties are regulated by the subunit composition of GABA_A receptors. For example, recombinant GABA_A receptors with $\alpha 4\beta 2\gamma 2$ subunits respond to GABA and benzodiazepine agonists with lower efficacy than $\alpha 1\beta 2\gamma 2$ receptors (Whittemore et al. 1996). Moreover, homologous genetic deletion of $\alpha 1$ subunits dramatically alters the pharmacological properties of GABA_A receptors, reducing the potency and efficacy of GABA and of the benzodiazepine diazepam (Kralic et al. 2002a,b). Hence, alterations in GABA_A receptor subunit expression appear critically important in regulation of GABA_A receptor function following chronic ethanol administration.

The regulation of various GABA_A receptor subunits by ethanol differs across brain regions (Grobin et al. 2000). For example, chronic ethanol consumption for 14 days increases GABA_A receptor $\alpha 4$ subunit peptide expression in the cerebral cortex and the hypothalamus (Devaud et al. 1997), decreases $\alpha 4$ subunit peptide levels in the amygdala and nucleus

accumbens (Papadeas et al. 2001), and does not alter $\alpha 4$ subunit peptide levels in the hippocampus or ventral tegmental area (Matthews et al. 1998; Papadeas et al. 2001). However, repeated ethanol withdrawals or longer ethanol exposure increases $\alpha 4$ subunit peptide expression in the hippocampus (Cagetti et al. 2003; Matthews et al. 1998) and these effects are associated with alterations in the pharmacological responses of GABA_A receptors to benzodiazepine agonists and inverse agonists (Cagetti et al. 2003). Furthermore, in cerebellum, $\alpha 6$ subunit mRNA and peptide levels are increased while $\alpha 1$ subunit peptide and mRNA are decreased following chronic ethanol administration (Mhatre et al. 1993; Morrow et al. 1992). Therefore, it is clear that adaptation in GABA_A receptor subunits expression caused by chronic ethanol administration is not universal across brain regions. The mechanisms that underlie regional differences in the regulation of GABA_A receptors by ethanol are unclear but may involve mechanisms to be described in the following sections.

Alterations in synaptic localization of receptors—Given the important role of synaptic vs. extrasynaptic GABA_A receptors in acute ethanol effects, it is important to understand the role of these receptor populations following chronic ethanol exposure. Cagetti et al. (2003) found changes in synaptic GABA_A receptor function in the hippocampus of rats that had undergone chronic intermittent ethanol (CIE) treatment followed by 2 days of withdrawal. CIE treatment decreases the frequency, rise time, amplitude, and decay time of mIPSCs in pyramidal neurons from CA1, an overall change in GABA_A receptor function that is consistent with increased neuronal excitability. Interestingly, CIE eliminated the effects of diazepam on the mIPSCs and increased mIPSC sensitivity to the $\alpha 4$ subunit-selective benzodiazepine Ro15-4513. These pharmacological changes are consistent with a decreased expression of $\alpha 1$ -GABA_A receptors and an increased expression of $\alpha 4$ -GABA_A receptors in the synapse. Brief exposure (48–72 h) to the neuroactive steroid $3\alpha, 5\alpha$ -THP may cause a similar effect since there is a decrease in the decay time of mIPSCs recorded in hippocampal CA1 pyramidal cells (Hsu et al. 2003). The decrease in mIPSC decay time was blocked by infusion of $\alpha 4$ subunit antisense oligonucleotides into the hippocampus, suggesting that this change in synaptic GABA_A receptor kinetics is due to an increase in synaptic $\alpha 4$ subunit expression. Additionally, tonic current is also modified after chronic ethanol exposure. Specifically, the effects of various GABA_A receptor modulators on tonic currents are reduced or ablated after CIE treatment (Liang et al. 2004). Indeed, GABA_A receptor ligands with selectivity for $\alpha 4$ -GABA_A receptors exerted decreased effects on extrasynaptic currents as well as increased effects on synaptic currents. Electron microscopy revealed an increase in synaptic localization of $\alpha 4$, but not δ subunits on the dentate granule cells of CIE rats (Liang et al. 2006). Taken together, these studies show that translocation of $\alpha 4$ -GABA_A receptors from an extrasynaptic to synaptic localization may represent a mechanism of GABA_A receptor adaptation to ethanol exposure.

Endocytosis and trafficking—Several mechanisms for GABA_A receptor adaptation following chronic ethanol exposure have been proposed that include alterations in gene expression, post-translational modification, synaptic localization, intracellular signaling, and neuroactive steroid responses to ethanol. The expression of GABA_A receptors involves a highly regulated process of synthesis, assembly, endocytosis, and recycling or degradation (see Fig. 1). Golgi-derived vesicles provide newly synthesized receptors to the cell surface, whereas clathrin-coated vesicles mediate endocytosis of surface receptors that are ultimately degraded or recycled back to the cell surface (Lodish et al. 1996). Alterations in the expression and composition of various GABA_A receptors could result from selective endocytosis, recycling, and/or trafficking of newly synthesized receptors to the cell surface. Recent studies have shown the importance of GABA_A receptor trafficking on the surface expression of receptors following ethanol exposure (Kumar et al. 2004). Altered GABA_A receptor subunit composition and expression on the cell surface following ethanol exposure is thought to contribute to the development of ethanol dependence. Therefore, investigation of the mechanisms for altered

GABA_A receptor trafficking following ethanol exposure may provide novel targets for drug development.

Chronic ethanol exposure selectively increases the internalization of $\alpha 1$ -GABA_A receptors into clathrin-coated vesicles of the cerebral cortex with a corresponding decrease in these receptors in the synaptic fraction (Kumar et al. 2003). In contrast, there is no change in the internalization of $\alpha 4$ -GABA_A receptors into clathrin-coated vesicles although there is a significant increase in $\alpha 4$ subunit peptide in the synaptic fraction following chronic ethanol exposure. Hence, the regulation of intracellular trafficking following chronic ethanol exposure appears to alter the subtypes of GABA_A receptors on the cell surface. This may account for alterations in the pharmacological properties of GABA_A receptors that have been observed. Internalization of GABA_A receptors following chronic ethanol administration is mediated by clathrin and the adaptor complex (AP) since the association of adaptin- α and clathrin with $\alpha 1$ -GABA_A receptors in the intracellular fraction was increased following chronic ethanol administration (Kumar et al. 2003). GABA_A receptor $\beta 2$ and/or $\gamma 2$ subunits are required for recognition of the receptor by the AP-2 that precedes clathrin-dependent endocytosis (Herring et al. 2003; Kittler et al. 2005, 2008). Therefore, it is possible that PKC-mediated phosphorylation of GABA_A receptor subunits, receptor-associated proteins, and/or AP-2 following chronic ethanol administration alters the recognition and endocytosis of GABA_A receptors. Indeed, chronic ethanol consumption results in increased expression of $\alpha 4$ -, $\beta 2$ -, and $\beta 3$ -GABA_A receptor subunits in the cerebral cortex and all of these subunits contain consensus phosphorylation sites for PKC (Macdonald 1995; Mohler et al. 1996; Wisden et al. 1991). In contrast, $\alpha 1$ -, $\alpha 2$ -, and $\alpha 3$ -GABA_A receptor subunits are decreased in the cortex and these subunits do not contain consensus phosphorylation sites for PKC. Hence, it is likely that phosphorylation of GABA_A receptor subunits can prevent internalization by blocking AP-2 binding. A recent study has shown that a single dose of ethanol can increase the internalization of GABA_A receptor $\alpha 4$ and δ subunits (Liang et al. 2007). However, this study did not investigate the mechanism of GABA_A receptor internalization. Therefore, investigation of GABA_A receptor endocytosis and trafficking following ethanol exposure is essential for understanding the mechanisms of ethanol adaptations.

Trafficking of GABA_A receptors can be regulated by many protein kinases, including PKC. Chronic ethanol consumption decreases association of PKC γ with $\alpha 1$ -GABA_A receptors and decreases expression of the $\alpha 1$ subunit at the cell surface. In contrast, chronic ethanol exposure increases association of PKC γ with $\alpha 4$ -GABA_A receptors and increases expression of $\alpha 4$ at the cell surface (Kumar et al. 2002). However, in the hippocampus, the association of PKC γ with GABA_A receptors is not altered and there is no alteration of $\alpha 1$ subunit expression following chronic ethanol exposure (Kumar et al. 2004). Therefore, it appears that association of PKC γ with GABA_A receptors and phosphorylation of subunits may influence the trafficking of receptors *in vivo* following chronic ethanol administration. The increased association of PKC γ with $\alpha 4$ -GABA_A receptors may phosphorylate GABA_A receptor subunits and prevent recognition of the receptor by AP-2, thus preventing its internalization. Indeed, it has been shown that phosphorylation of GABA_A receptor subunits reduces the binding of receptors with AP-2 and subsequent internalization (Kittler et al. 2005, 2008). In addition, reduced PKC-dependent GABA_A receptor phosphorylation enhances receptor binding to the AP-2 and promotes their endocytosis (Terunuma et al. 2008). Taken together, it appears that PKC-mediated phosphorylation plays an important role in GABA_A receptor cell surface expression following chronic ethanol exposure.

GABA_A receptor subunits contain phosphorylation sites for many protein kinases including PKC, PKA, and *fyn* kinase. Phosphorylation can alter receptor function directly by changing its conformation and/or indirectly by altering receptor expression. Since PKA and *fyn* kinase alter GABA_A receptor function, it is possible that these protein kinases are also involved in

trafficking of GABA_A receptors. For example, chronic activation of PKA in cerebellar granule cells increases cell surface expression of GABA_A receptor $\alpha 1$ subunit (Ives et al. 2002). In addition, ethanol exposure alters expression and translocation of PKA (Diamond and Gordon 1994; Newton and Messing 2006; Pandey 1998). Hence, it is likely that PKA is also involved in trafficking of GABA_A receptors following ethanol exposure. However, to date, the role of these protein kinases has not yet been studied in trafficking of GABA_A receptors, especially following ethanol exposure. Future studies will determine the specific role of various protein kinases in GABA_A receptor trafficking and resolve the precise mechanisms that account for alterations in GABA_A receptor function and expression following chronic ethanol administration.

Alterations in ethanol-induced neuroactive steroid responses

Chronic ethanol consumption in rats results in blunted elevation of cerebral cortical $3\alpha,5\alpha$ -THP (Morrow et al. 2001) and plasma and brain deoxycorticosterone levels following acute ethanol challenge (Khisti et al. 2005) compared to pair-fed control rats. These findings suggest that there is tolerance to ethanol-induced increases in neuroactive steroid levels. The loss of ethanol-induced increases in neuroactive steroids may contribute to ethanol tolerance since the elevations of neuroactive steroids are required for anxiolytic, sedative, anticonvulsant, and cognitive-impairing effects of ethanol. Furthermore, since decreases in brain neuroactive steroid levels are concomitant with decreases in plasma neuroactive steroid levels, it is likely that the observed decreases in $3\alpha,5\alpha$ -THP and deoxycorticosterone levels are dependent on blunted HPA axis activity. It is well known that chronic stress results in adaptation of the HPA axis leading to decreases in stress-induced changes in the levels of corticosterone in rats (Spencer and McEwen 1990). This blunting of the HPA axis is associated with a reduction in corticotrophin releasing factor and adrenocorticotropin releasing hormone elevations following ethanol challenge (Lee et al. 2001). Furthermore, repeated ethanol withdrawal also leads to blunting of withdrawal-induced elevations in corticosterone levels (Borlikova et al. 2006). Thus, chronic ethanol administration induces adaptations of the HPA axis to promote loss of stress-induced increases in GABAergic neuroactive steroids that contribute to the regulation of homeostasis as well as ethanol sensitivity.

Chronic ethanol administration to rodents and humans produces tolerance to ethanol and cross-tolerance to benzodiazepines and barbiturates. In contrast, ethanol-dependent rats are sensitized to the anticonvulsant effects of $3\alpha,5\alpha$ -THP and $3\alpha,5\alpha$ -THDOC (Cagetti et al. 2004; Devaud et al. 1996; 1998). GABA_A receptor sensitivity to $3\alpha,5\alpha$ -THP and $3\alpha,5\alpha$ -THDOC is enhanced in the cortex, but sensitivity to alphasalone is reduced in the hippocampus (Cagetti et al. 2004) of ethanol-dependent rats, likely due to adaptations in GABA_A receptor expression described above. Since ethanol-dependent rats are sensitized to anticonvulsant actions of neuroactive steroids, this class of compounds may be useful during ethanol withdrawal. Indeed, neuroactive steroid therapy may have advantages over benzodiazepine therapy since benzodiazepines exhibit cross-tolerance with ethanol. Further studies are needed to explore this possibility.

Cross-talk with other neurotransmitter systems

As our understanding of cellular signaling pathways has advanced, it has become clear that intracellular communication between receptors is critical for the coordinated function of cells. There are multiple receptors expressed in every cell and their signals pass through a common set of downstream effectors distinguished by multiple isoforms with slightly different specificities and activities. The coupling among these pathways causes interactions among the signals sent by the different classes of receptors. Recent molecular pharmacology studies have demonstrated that ethanol acts on several neurotransmitter systems in the brain such as GABA, NMDA, glycine, 5-hydroxytryptamine (5-HT; serotonin) 3, and nicotinic acetylcholine

receptors as well as L-type Ca^{2+} channels and G protein-coupled receptors. Following the direct actions of ethanol on receptors in the brain, a second wave of indirect effects is initiated that subsequently leads to the typical acute behavioral effects of ethanol, ranging from disinhibition to sedation and hypnosis, with increasing concentrations of ethanol. Therefore, a clear understanding of the interactions between neurotransmitter receptors and their signaling pathways is critical for understanding the mechanisms that underlie ethanol tolerance and dependence.

Chronic ethanol exposure up-regulates NMDA receptor function and down-regulates GABA_A receptor function (see Crews et al. 1996 for review). Since many neurons contain both NMDA and GABA_A receptors and ethanol modulates both NMDA and GABA_A receptors, interactions between these receptors may play an important role in GABA_A receptor adaptation following ethanol administration. For example, administration of dizocilpine (a potent non-competitive antagonist of the NMDA receptor) to rats, like ethanol, increases membrane expression of GABA_A receptor $\alpha 4$ subunits in hippocampus (Matthews et al. 2000b). The effect on GABA_A receptors following NMDA receptor activation is mediated by intracellular Ca^{2+} , since the Ca^{2+} chelator 1,2-bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetra-acetic acid (BAPTA) blocks the down-regulation of GABA_A receptors following NMDA receptor activation (Robello et al. 1997). In addition, the application of glutamate in hippocampal cultures leads to translocation of $\text{PKC}\alpha$ and $\text{PKC}\gamma$ to the plasma membrane and cytoplasmic organelles, respectively, while $\text{PKC}\epsilon$ localization remains unaltered (Buchner et al. 1999; Etoh et al. 1991). Therefore, interactions of NMDA and GABA_A receptors via intracellular signaling pathways may play a vital role in GABA_A receptor adaptation following chronic ethanol administration. In addition, both NMDA and GABA_A receptor activation, as well as ethanol administration, alter brain-derived neurotrophic factor (BDNF) expression in neurons (MacLennan et al. 1995; Obrietan et al. 2002; Tapia-Arancibia et al. 2001; Zafra et al. 1991). BDNF has been shown to alter GABA_A receptor function and expression. For example, addition of BDNF in cortical cell cultures increases internalization of GABA_A receptor $\alpha 2$, $\beta 2$, $\beta 3$, and $\gamma 2$ subunits (Brünig et al. 2001). In contrast, chronic application of BDNF increases expression of GABA_A receptor $\alpha 6$ subunit in cerebellar granular cells that is blocked by concomitant application of ethanol (Ericson et al. 2003). Therefore, it is possible that ethanol-induced altered trafficking and expression of GABA_A receptors are partially mediated by BDNF. Future studies on interactions of BDNF and GABA_A receptor following chronic ethanol exposure are needed to clarify the role of BDNF on ethanol-induced adaptations of GABA_A receptors.

Likewise, serotonin receptors, a primary target of ethanol action, can modulate GABA_A receptor function and expression via PKC-mediated pathways. Activation of 5-HT₂ receptors results in inhibition of GABA_A receptor-mediated currents that can be blocked by the Ca^{2+} chelator BAPTA and the receptor for activated C-kinase (RACK) inhibitory peptide (RACK1-rVI) (Feng et al. 2001). In addition, even though 5-HT₃ is considered to be an excitatory receptor, it is often expressed on inhibitory GABA_A ergic interneurons, thus activation of 5-HT₃ receptors by ethanol may contribute to some of the inhibitory actions of ethanol via increased release of GABA (Lovinger 1999). Furthermore, activation of 5-HT₃ receptors is also known to increase the release of dopamine and glutamate (Lovinger 1999). Hence, it is clear that activation of serotonin receptors can alter GABA_A receptor function and expression. In addition, recent studies have concluded that dopamine and β -adrenergic receptors can also modulate GABA_A receptors via PKA-mediated pathways (Flores-Hernandez et al. 2000). Taken together, it is obvious that receptor cross-talk plays a vital role in modulation of GABA_A receptor function and expression following ethanol exposure.

Recent evidence indicates that many other neurotransmitter and cytokines may be involved in ethanol-mediated alteration in GABA_A receptor function and expression. For example, the

mGluR5 antagonist 2-methyl-6-(2-phenyl-ethynyl)pyridine (MPEP) modulates the GABAergic effects of ethanol (Besheer and Hodge 2005). Furthermore, the mGluR5 receptor selective agonist (R,S)-2-chloro-5-hydroxyphenylglycine (CHPG) enhances GABA-gated currents via post-synaptic effects in amacrine cells (Hoffpauir and Gleason 2002). Likewise, cytokines may be involved in ethanol-mediated alterations in GABA_A receptor function and expression. Tumor necrosis factor α (TNF- α) concentrations in the brain are elevated by ethanol administration (Crews et al. 2006; Kiefer et al. 2002) and this may contribute to a decrease in GABA_A receptor α 1 subunit expression (Stellwagen et al. 2005) following chronic ethanol exposure. Understanding of the mechanisms of ethanol dependence has been complicated by the multiple actions of ethanol at various neurotransmitters and/or intracellular signaling pathways. Hence, understanding of these interactions between neurotransmitter systems is essential to the development of novel therapeutic targets for alcoholism.

Studies in non-human primates

In the past 10 years, there have been new studies on the effects of ethanol administration to non-human primates that have also investigated effects on GABA_A receptor function and plasticity. Cynomolgus macaques (*Macaca fascicularis*) are a powerful model to study the effects of long-term ethanol exposure as they will freely self-administer intoxicating quantities of ethanol with drinking patterns that mimic human alcoholics (Grant et al. 2008b; Vivian et al. 2001). Long-term ethanol self-administration by cynomolgus macaques alters the pharmacological and functional properties of GABA_A receptors by decreasing GABA potency, but not efficacy, in the basolateral amygdala neurons. These alterations are associated with decreased α 2 and α 3 subunit mRNA expression and with a trend for a decrease in α 1 subunit mRNA expression. No significant effect of gender or significant interaction between gender and ethanol exposure is observed for any of the subunits examined. Furthermore, the mean expression levels for both the α 2 and the α 3 subunits were significantly correlated with the total amount of ethanol consumed (Floyd et al. 2004). In the same cohort of cynomolgus macaques, Hemby et al. (2006) found significant decreases in mRNA expression for: α 2, α 4, β 1, β 3, γ 1, γ 2, and γ 3 in orbitofrontal cortex; β 1, β 2, γ 1, and δ in dorsolateral prefrontal cortex; but no changes in anterior cingulate cortex. In addition, Anderson et al. (2007) showed that long-term ethanol self-administration reduces the sensitivity of amygdala GABA_A receptors to the benzodiazepine flunitrazepam and decreases mRNA expression of β 1 and γ 2 subunits. Furthermore, there are significant effects of ethanol drinking on the γ 2 subunit gene in males but not females; however, there was no correlation between subunit mRNA expression and total ethanol intake, suggesting that this gender adaptation is not related to drinking. Interestingly, GABA_A receptor density, measured by [³H]Ro15-4513 binding, does not differ in the parietal and temporal cortex, cerebellum, or hippocampus after long-term ethanol self-administration in these cynomolgus macaques (Sullivan et al. 2005). Overall, the effects of ethanol in cynomolgus macaques differ from rat models in several aspects and suggest that studies in non-human primates provide unique insights that may have more relevance for human alcoholism.

Alcohol tolerance mechanisms

Although many people consume ethanol, the fact that only a select number of individuals develop alcoholism remains enigmatic. Tolerance is thought to play a major role in the progression towards alcoholism and may therefore contribute to risk for the disease. Tolerance is defined as a reduced response to a constant amount of ethanol, such that a greater amount is needed to obtain the same effect. Tolerance is complex because it can involve multiple behaviors that respond to ethanol exposure in different ways. It can also be segregated as either innate or acquired (i.e., requiring a previous ethanol exposure), with the latter being further broken down into acute, rapid, and chronic tolerance, based on the temporal relationship

between ethanol exposures and their duration (Kalant et al. 1971). Acute tolerance is characterized by adaptations that occur immediately during exposure to ethanol. Rapid tolerance is typically defined as tolerance manifested after a single exposure has been eliminated from the system. Chronic tolerance involves various forms of multiple or continual ethanol exposure paradigms. GABA_A receptors are implicated in tolerance to ethanol's effects. For instance, in humans, sons of alcoholics exhibit decreased sensitivity to diazepam in eye-movement tasks (Cowley et al. 1994), indicative of reduced sensitivity to GABAergic stimulation. In animal studies, a considerable amount of work has been done to assess the role of GABA_A receptors in innate and acquired tolerance to ethanol, as described below.

Innate tolerance

Various studies using GABA_A receptor modulators in inbred, selectively bred, and recombinant inbred strains that differ in ethanol responses have correlated GABA_A receptor action to innate ethanol tolerance. For instance, mice selectively bred for differences in diazepam responses also differed in ethanol responses. Mice that were more resistant to diazepam were less ataxic to ethanol, compared to diazepam-sensitive mice (Gallaher and Gionet 1988). The GABA_A receptor agonist taurine may also be involved in different ethanol-related responses in Sardinian alcohol-preferring (sP) rats versus non-preferring (sNP) rats. Recent microdialysate studies indicate that sP rats had reduced ethanol-related taurine release compared to sNP rats (Quertemont et al. 2000). Notably, alcohol-preferring rats were less sensitive to the sedative-hypnotic effects of ethanol (Kurtz et al. 1996); therefore, it is possible that differences in ethanol-related behaviors in these animals may be related to differences in endogenous GABA_A receptor ligands like taurine or responses to other GABA_A receptor modulators. Indeed, extrasynaptic receptors involved in tonic inhibition have recently been shown to be sensitive to low concentrations of taurine (Jia et al. 2008a). However, sP and sNP rats do not differ in their sedative-hypnotic responses to other benzo-diazepines and barbiturates (Colombo et al. 2000), but sP rats are more sensitive to the sedative-hypnotic effects of γ -hydroxybutyric acid (Colombo et al. 1998). It is possible that neuroactive steroids may also be involved in innate responses in these animals. $3\alpha,5\alpha$ -THP and $3\alpha,5\alpha$ -THDOC are increased in sP vs. sNP rats (Barbaccia et al. 1999). In other selectively bred animals, withdrawal seizure-prone mice are not only found to be more sensitive to the anxiolytic effects of ethanol than withdrawal seizure-resistant mice but they are also more sensitive to the anxiolytic effects of pentobarbital (Atkins et al. 2000). In other studies, FAST & SLOW mice, selectively bred for differences in locomotor stimulatory effects of ethanol, also differ in responses to the convulsant effects of GABA_A receptor modulators, and to the locomotor-stimulant effects of benzodiazepines (Shen et al. 1998). Aside from selectively bred animals, inbred strains of mice also suggest a role for GABA_A receptors. C57BL/6J mice are not only less sensitive to the sedative-hypnotic effects of ethanol than 129/SvJ mice but are also less sensitive to the GABA agonist propofol (Homanics et al. 1999b). Interestingly, the same study indicated that C57BL/6J mice are more sensitive to zolpidem and midazolam, but not Ro15-4513 and pentobarbital. Besides pharmacologic modifiers, molecular biological techniques have been used to assess the role of specific GABA_A receptor subunits in innate tolerance. Using a coding polymorphism difference in C57BL/6J and DBA/2J mice, different lines of BXD recombinant inbred strains were used to determine whether variations in the gene for the GABA_A receptor γ 2 subunit correlated with ethanol-related behaviors. BXD lines with γ 2 of DBA/2J origin were less ataxic on the screen test, but had greater ethanol-induced hypothermia (Hood and Buck 2000). A trend was also noted for sedative-hypnotic effects as well as a second motor ataxia test.

Acquired tolerance

Apart from innate tolerance, various inbred, selectively bred, and recombinant inbred rodent strains, as well as genetically modified mouse models, have been used to assess acquired

ethanol tolerance. Acute functional tolerance (AFT) is a commonly used assay to assess acute tolerance. AFT is defined as greater sensitivity to ethanol effects during the rising phase of the blood ethanol curve than equivalent ethanol concentrations on the falling phase. However, because accurately assessing blood ethanol concentrations on the rising phase is difficult due to the immediate effects of ethanol, AFT is commonly assessed by measuring the ability of a subject to exhibit a specific ethanol-related behavior (e.g., motor coordination, loss of righting reflex) during successive ethanol exposures. Blood ethanol measurements are taken upon the appearance of each behavior and the difference in blood ethanol level on the successive measurements is indicative of acute tolerance.

AFT to the motor-impairing effects of ethanol is associated with a reduction in the ability of ethanol to enhance muscimol-stimulated Cl^- uptake in cerebellar microsacs (Wallace et al. 2007). Studies in GABA_A receptor genetically modified models have shown that AFT to motor incoordination is decreased in $\gamma 2\text{S}$ - and $\gamma 2\text{L}$ - GABA_A receptor transgenics (Wick et al. 2000), but does not differ in $\alpha 1$ -, $\alpha 5$ -, $\alpha 6$ -, $\gamma 2\text{L}$ -, or δ - GABA_A receptor knockout mice (Boehm et al. 2004b; Homanics et al. 1998, 1999a; Kralic et al. 2003; Mihalek et al. 2001). However, recent work in $\alpha 1$ - GABA_A receptor (S270H, L277A) knockin mice suggests that $\alpha 1$ - GABA_A receptors are involved in AFT to ethanol (Werner et al. 2009), since these mice display decreased AFT to motor coordination.

GABA_A receptor involvement in acute, rapid, and chronic ethanol tolerance has also been assessed in various animal models. For instance, selectively bred mice that differ in diazepam sensitivity also differ in rapid tolerance development. Mice that are more resistant to diazepam's effects also develop more rapid tolerance than diazepam-sensitive mice (Gallagher and Gionet 1988). Inbred long-and short-sleep mice display differences in rapid tolerance. Long-sleep mice that display a greater response to ethanol's sedative-hypnotic effects also develop greater rapid tolerance (Radcliffe et al. 2005). Indeed, these mice differ in GABA_A receptor inhibitory post-synaptic responses (Proctor et al. 2004), suggesting the involvement of different GABA_A receptor subunits in ethanol tolerance. BXD mice have also been used to assess tolerance. Assessment of the parental strains, C57BL/6J and DBA/2J, indicates that although DBA/2J mice were more sensitive to ethanol's motor ataxic effects, they also develop greater AFT (Gallagher et al. 1996). Subsequent tests show quantitative trait loci (QTL) for ethanol tolerance on chromosomes 9 and 11. While these studies did not identify a role for GABA_A receptors in tolerance, it is possible that subunits from the GABA_A receptor subunit gene cluster on chromosome 11 may be involved. Other studies show that mice selectively bred for ethanol withdrawal severity do not differ in tolerance to ethanol's hypothermic or sedative-hypnotic effects (Crabbe and Kosobud 1986), although these mice exhibit differences in GABA_A receptor mRNA subunit expression (Keir and Morrow 1994) as well as differential changes in subunit expression after chronic ethanol (Buck et al. 1991a). Together, these results support the notion that various genes contribute to ethanol-related responses and that tolerance to specific ethanol-related behaviors may be independent of withdrawal.

Genetically modified mice have also been used to assess the role of GABA_A receptors in chronic tolerance. Recent work suggests that $\alpha 1$ - GABA_A receptors are involved in chronic tolerance to ethanol's motor ataxic effects. GABA_A receptor $\alpha 1$ (S270H, L277A) knockin mice that are initially less sensitive to ethanol's motor ataxic effects display decreased tolerance after repeated ethanol exposure (Werner et al. 2009). Furthermore, $\beta 3$ - GABA_A receptors appear to play a role in chronic tolerance. $\beta 3$ (N265M) knockin mice display enhanced tolerance to ethanol's sedative-hypnotic effects compared to controls (Sanchis-Segura et al. 2007). However, $\alpha 6$ - and δ - GABA_A receptor knockout mice do not differ from controls in chronic tolerance to ethanol's sedative-hypnotic effects (Homanics et al. 1998; Mihalek et al. 2001). Studies have also been carried out in genetically modified mouse models of genes known to indirectly influence GABA_A receptor function. Work has shown that $\text{PKC}\gamma$ knockout mice

did not develop chronic tolerance to ethanol's sedative-hypnotic or hypothermic effects, but this response was strain dependent (Bowers et al. 1999). PKC ϵ knockout mice are resistant to tolerance to motor-impairing effects of ethanol (Wallace et al. 2007). Further studies are needed to establish the role of other GABA $_A$ receptor subunits and other GABA $_A$ receptor-associated proteins in AFT as well as acute and chronic tolerance to other behavioral measures.

Tolerance and altered GABA $_A$ receptor function and regulation

Recent work has suggested that alterations in GABA $_A$ receptor expression may underlie acquired ethanol tolerance. Studies indicate that a single high ethanol exposure results in alterations in GABA $_A$ receptor subunit expression. One hour following ethanol exposure, $\alpha 4$ and δ surface expression is decreased in the hippocampal CA1 region, but no difference in total subunit levels is found (Liang et al. 2007). This decrease in cell surface extrasynaptic receptors is associated with a decrease in ethanol-enhanced tonic current. Furthermore, a significant enhancement of surface expression of $\alpha 4$ and $\gamma 2$ subunit proteins, as well as a decrease in $\alpha 1$ and δ subunits, was noted 2 days following ethanol exposure. Additionally, the ethanol-induced increase in $\alpha 4$ subunit expression requires the binding of heat shock factor 1 at a downstream site (Pignataro et al. 2007). These adaptive changes are accompanied by an increase in mIPSC response to ethanol and decreased response to diazepam (Liang et al. 2007), suggesting that the observed changes in receptor expression might underlie altered sensitivity to ethanol. Furthermore, chronic ethanol exposure altered the recovery of baseline spontaneous neural activity of medial septal/diagonal band of Broca neurons following GABA microiontophoresis (Matthews et al. 2000a). These studies suggest that the observed changes in receptor expression following chronic ethanol exposure might underlie altered sensitivity to ethanol and GABA. Interestingly, the mIPSC effects were transient, and responses returned to normal after 2 weeks. It is possible that these transient changes contribute to ethanol tolerance. Tolerance may be restricted to selected brain regions or cell populations. Tolerance to ethanol's effects in the CA3 region of neonatal rats does not occur in slices exposed to ethanol for a prolonged period (Galindo and Valenzuela 2006). Finally, inhibition of synaptic GABA $_A$ receptors results in the incorporation of receptors diffusing from peri- and extrasynaptic sites (Thomas et al. 2005), independent of receptors arising from intracellular stores. Overall, while not pinpointing the exact role of GABA $_A$ receptors in ethanol action, these studies lend support to a possible role of changes in GABA $_A$ receptor subtype expression contributing to tolerance to various ethanol-related behaviors.

Adolescence

Adolescence (often defined as 12 to 20 years of age in humans) is a developmental period highlighted by continued neurobiological development accompanied by changes in personality. The neurobiological alterations include changes in neuronal networks, brain size, and receptor number and compositions, while changes in personality include increased risk choices and experimentation with drugs, including ethanol. In fact, increased drug use, particularly ethanol, is a commonly noted factor at this developmental stage in humans (Kim et al. 2008; Lim et al. 2007). The development of animal models to better understand the impact of ethanol exposure during adolescence has rapidly proceeded in the last decade. The need for animal models is obvious due to ethical and legal restrictions limiting ethanol research in human adolescence. The terms adolescence and puberty are often used interchangeably. While their timing overlaps, the terms are not synonymous (Sisk and Foster 2004; Spear 2000). Puberty refers to gonadal maturation, while adolescence refers to a gradual period of maturation of social and cognitive behaviors (Powell 2006; Sisk and Foster 2004). The precise timing of adolescence in rodents is difficult to determine as there is no specific event signaling the beginning and end of this developmental period and the boundaries vary depending on genetic differences, gender, nutrition, and other environmental periods (Spear 2007). In rats, a

conservative estimate of adolescence spans the period between postnatal days (PD) 28 and 42 (Spear 2007), but some harbingers of female adolescent development can begin as early as PD 20 and continue through PD 55 in males (Spear 2000, 2007). Studies investigating the effect of ethanol in adolescent rats have used the period between PD 28 and 60 (Smith 2003), while in mice, adolescence has been characterized as the period from PD 22 to 60 (Laviola et al. 2003).

Behavioral sensitivity

It has become clear that ethanol produces different behavioral responses in adolescence compared to adults. Specifically, age modulates ethanol effects on motor incoordination, anxiety, and cognitive function. Researchers have also shown that sensitivity to ethanol-induced ataxia increases during ontogeny (Land and Spear 2004; Smith 2003; Spear and Varlinskaya 2005). Specifically, adolescent rats exhibit less severe motor impairment following ethanol exposure (Spear and Varlinskaya 2005; White et al. 2002a). Throughout development, a natural increase in sensitivity to the motor-impairing effects of ethanol between adolescence and adulthood emerges (Silveri and Spear 1998).

Besides motor ataxia, anxiety has been shown to play a role in the recurrence and initiation of drinking behavior. Given the role of the GABA_A receptor system in anxiety, it is reasonable to surmise that the adolescent developmental period may be characterized by changes in the anxiolytic effects of ethanol. Recent studies report attenuated withdrawal-related anxiety and a lessened anxiolytic effect of ethanol in adolescent rats as measured in a social interaction test (Doremus et al. 2003; Varlinskaya and Spear 2002, 2004). Interestingly, early adolescent C57BL/6J mice show enhanced anxiety and greater anxiolytic response to ethanol on the elevated plus maze (Hefner and Holmes 2007). In addition to potential species differences, such contrary results are also likely confounded by different baseline levels of social interaction and plus maze behavior between adolescence and adults. However, it is possible that lessened anxiogenesis and increased anxiolysis during adolescence may also be a reflection of the developmental state of the GABA_A receptor system that underlies anxiety and may be a contributing factor to problem drinking. Salimov et al. (1996) reported that ethanol consumption during the adolescent period results in lower responsiveness to novelty-induced anxiety and stress. Again, these outcomes are predictive of a greater risk for problem drinking as a result of ethanol exposure during the adolescent developmental period.

Ethanol impairs cognition, executive function, and attentional processes that involve several brain regions. Among the constellation of cognitive impairments produced by ethanol, learning and memory, which is hippocampal dependent, is particularly vulnerable to ethanol-induced impairments (see Matthews and Silvers 2004 for review). However, the magnitude of these cognitive impairments is age dependent. For example, early studies reported that acute ethanol administration to adolescent rats produced greater learning and memory deficits compared to adults (Land and Spear 2004; Markwiese et al. 1998) and this differential effect was dose dependent (Acheson et al. 2001). However, the opposite effect (adolescents are less sensitive than adults to the effects of ethanol) has also been reported (Land and Spear 2004; Rajendran and Spear 2004). Regardless of whether adolescents are more or less sensitive to ethanol-induced cognitive impairments, it is clear that ethanol affects the hippocampus of adolescents differently than adults. Although the neurobiological mechanisms underlying this difference are not yet elucidated, it is known that ethanol more potently attenuates hippocampal long-term potentiation in brain slices from adolescent rats compared to adult rats (Pyapali et al. 1999). In addition, moderate ethanol doses (30 mM) enhance GABA_A receptor-mediated inhibitory tonic currents more potently in the dentate gyrus of adolescent rats compared to adult rats (Fleming et al. 2007). These results demonstrate that innate differences, including differences in GABA_A receptors, exist in the hippocampus between adolescent and adult

animals that likely underlie differential sensitivity to ethanol-induced impairments in cognition.

Developmental alterations in GABA_A receptor expression

Given that adolescent subjects differ from adults on many ethanol-related behaviors, it is possible that the spatial and temporal changes in GABA_A receptor subunit expression throughout the developmental process may contribute to their phenotypic differences. Each subunit exhibits unique developmental expression patterns based on regions and age, suggesting that the expression and function of GABA_A receptors during development may differ from those found in adults (Henschel et al. 2008). GABA_A receptor subunit expression of $\alpha 1$, $\alpha 2$, $\alpha 3$, $\alpha 5$, and $\gamma 2$ changes throughout development, reaching adult-like levels during the third month of life (Jenkins and Simmons 2006). Immunoreactivity of $\alpha 1$ subunit is low at birth and restricted to a few brain regions, with levels increasing during the first postnatal weeks (Fritschy et al. 1994). In contrast, $\alpha 2$ subunit distribution is widespread in the superficial layers (I–IV) of the neocortex (Yu et al. 2006) at birth. These levels decrease in some regions shortly after the appearance of the $\alpha 1$ subunit and co-expression occurs briefly, with $\alpha 1$ eventually replacing $\alpha 2$ subunit (Fritschy et al. 1994; Henschel et al. 2008). High levels of $\alpha 3$ subunit are found within the frontal and perirhinal cortex, with up-regulation occurring from PD 10 to 30; the $\alpha 5$ subunit has high expression throughout the brain, especially in the deep cortical layers (Yu et al. 2006). The initial increases in $\alpha 1$, $\alpha 2$, $\alpha 3$, and $\gamma 2$ found during the early postnatal period change during development, and decrease until PD 60, while $\alpha 5$ levels decrease from the high levels found in early development throughout the aging process (Yu et al. 2006). Prominent throughout development, $\beta 2$ and $\beta 3$ immunoreactivity shows consistent levels in both the neonatal and adult brain (Fritschy et al. 1994). In humans, a similar increase in $\alpha 1$ and $\gamma 2$ expression occurs with age in the temporal lobe, hippocampus, and basal ganglia (Kanaumi et al. 2006).

Consistent with developmental shifts in GABA_A receptor expression within specific brain regions, the cerebellum displays levels of $\alpha 6$ subunit expression that increase with age and expression of $\alpha 1$, $\gamma 2$, $\beta 2$, $\beta 3$, and δ subunits that decreases with age (Gutiérrez et al. 1997). Within the hippocampus, consistently high levels of $\gamma 2$ and $\alpha 5$ subunit expression are found across development, the $\alpha 1$ subunit shows decreasing levels through development and aging (Yu et al. 2006), and levels of the δ subunit mRNA increase throughout development, with the highest levels being found in adults (Laurie et al. 1992). Increasing levels of the δ subunit mRNA across development are interesting, given the proposed link of δ -GABA_A receptors to low-dose ethanol sensitivity (Choi et al. 2008). Considering that adolescents are less sensitive to ethanol's motor ataxic effects compared to adults, and δ subunit mRNA levels are lower in adolescents compared to adults, it is tantalizing to speculate that this subunit is a critical mediator underlying the changes in ethanol's effect on GABA_A receptors across development. Overall, because GABA_A receptor subunit expression differs across development, it is likely that changes in GABA_A receptor stoichiometry underlie many of the developmental specific effects of ethanol previously reviewed. Furthermore, chronic ethanol exposure alters GABA_A receptor mRNA and protein expression. As such, it is predicted that chronic ethanol exposure during adolescence, when GABA_A receptors are changing, should also alter GABA_A receptor protein expression in a manner that differs from the changes found following adult ethanol exposure.

Ethanol intake and chronic ethanol exposure

Adolescent rats consume more ethanol than adults (Doremus et al. 2005). Indeed, adolescents have been shown to consume two to three times more ethanol than adults relative to body weight (Lancaster et al. 1996). Some variation in adolescent ethanol self-administration has been found in consumption which may reflect methodological differences. When consumption

is examined in a 24-h continuous access paradigm, adolescents exhibit a greater increase in ethanol licking behavior over a 4-week period, compared to adults (Bell et al. 2006). However, when voluntary ethanol self-administration is examined, adolescents consumed significantly more ethanol when measured as g/kg body weight (Vetter et al. 2007). When the consumption off the floor paradigm is used, adolescents show high levels of consumption that decrease gradually into adulthood (Truxell et al. 2007). Overall, adolescents consume more ethanol than adults; this difference is particularly evident when consumption is calculated relative to body weight rather than proportion of overall consumption (Nance 1983). Complicating interpretation of these results, rates of ethanol metabolism vary with age. Adolescents have been observed to have faster rates of ethanol elimination than adults (Brasser and Spear 2002; Hollstedt et al. 1977); however, these results are not consistently found (Kelly et al. 1987; Silveri and Spear 2000). Additionally, these results have not been found to account for the attenuated ethanol sensitivity shown by adolescents to many of the acute effects of ethanol (Silveri and Spear 2000). The reported high levels of adolescent consumption may reflect age-related changes in both the rewarding and aversive properties of ethanol (Vetter et al. 2007).

The subjective properties of a drug play a large part in consumption, abuse, and addiction patterns. That is to say, the subjective experience of greater reward or less aversion as it relates to a drug will lead to a greater risk of abuse. Assessment of the motivational properties of ethanol has included the use of ethanol-induced conditioned place preference and ethanol-induced conditioned taste aversion (CTA) (Chester and Cunningham 1999; Chester et al. 1998). Interestingly, although not without inconsistencies, GABAergic involvement has been implicated in both of these tasks. For example, adult mice exposed to ethanol (via inhalation) during the adolescent period show an attenuated ethanol-induced CTA (Chester and Cunningham 1999; Smith 1989). Additionally, a chronic intermittent pattern of exposure, resulting in multiple withdrawals during adolescence, exacerbates this attenuated CTA (Diaz-Granados and Graham 2007; Graham and Diaz-Granados 2006). When subjects were administered diazepam during the withdrawal periods in order to assess the role of possible GABA_A receptor withdrawal-related perturbations, the attenuation of the ethanol-induced CTA was not dissimilar between diazepam-treated and control animals, suggesting that GABA_A receptor perturbations during withdrawal were not playing a role in the attenuation of CTA. However, adult mice that were exposed to only diazepam during the adolescent period showed the same attenuated CTA as adult mice exposed to only ethanol during adolescence. Thus, the rewarding and aversive properties of ethanol, and therefore ethanol self-administration, in adulthood are related to the interaction of these drugs with a developmentally dynamic GABAergic system.

Chronic exposure to ethanol during adolescence has also revealed long-lasting changes in hippocampal function and GABAergic neuroactive steroids. For example, CIE exposure to high dose ethanol during adolescence does not impair spatial learning during the exposure (Silvers et al. 2003a) but does produce cognitive tolerance to ethanol's memory impairing effects when animals are tested as young adults (Silvers et al. 2003b, 2006). However, the cognitive tolerance is not permanent (Silvers et al. 2006) and may in fact lead to sensitization of ethanol's effect on memory following a long delay (White et al. 2000). Mirroring the cognitive tolerance, CIE exposure during adolescence blunts ethanol-induced inhibition of hippocampal pyramidal neurons, and this effect lasts into young adulthood (Tokunaga et al. 2006). Furthermore, chronic intermittent ethanol during adolescence also reduces hippocampal 3 α ,5 α -THP levels in response to an ethanol challenge (Silvers et al. 2006) and this reduced neuroactive steroid response directly mirrors the reduced cognitive and neurophysiological effect following ethanol challenge. Therefore, it is likely that the prolonged cognitive and neurophysiological tolerance found following chronic ethanol exposure during adolescence is due to GABAergic mechanisms such as altered neuroactive steroid levels. Chronic ethanol exposure has also altered other ethanol-related behaviors in adolescents. Although sensitivity

to the motor-impairing effects of ethanol increases during development, repeated adolescent binge drinking episodes prevent the emergence of this normal developmental change (White et al. 2002b). Additional studies have also revealed a persistent tolerance to the sedative effects of ethanol, resulting in continued attenuation of the animal's loss of righting reflex (Silveri and Spear 1998; Silvers et al. 2003b).

However, few studies have investigated whether chronic ethanol administration during adolescence alters GABA_A receptor structure or function. In fact, only three studies directly compare the two age groups. Specifically, adolescent and adult rats do not differ on bicuculline-induced seizure threshold (Wills et al. 2008), but differ on pentylenetetrazole-induced seizures (Acheson et al. 1999) following chronic ethanol. However, chronic ethanol exposure for 1 month differentially altered GABA_A receptor function in adult vs. adolescents as measured by 3 α ,5 α -THDOC potentiated Cl⁻ flux (Grobin et al. 2001). These limited studies suggest that perhaps chronic ethanol might differentially alter GABA_A receptor function when animals are exposed during adolescence as compared to adulthood. However, additional work is clearly needed on this topic.

Human alcoholics

Do the effects of chronic ethanol exposure in animal studies translate to humans? Behavioral adaptations to ethanol in human beings mimic those observed in animal models. Further, GABAergic neurotransmission is important for many behavioral actions of ethanol and several studies in human alcoholics have highlighted changes in GABA_A receptor function and plasticity that resemble the adaptations found in animal models of ethanol dependence. For instance, human alcoholics exhibit ethanol tolerance and benzodiazepine and barbiturate cross-tolerance (Volkow et al. 1993; Woo and Greenblatt 1979). Early studies conducted in post-mortem human brains have suggested the involvement of GABA_A receptors: [³H]muscimol binding density is greater in alcoholic cerebral cortex (Tran et al. 1981) and superior frontal gyrus of non-cirrhotic alcoholics (Dodd et al. 1992) compared to healthy controls. Increases, decreases, or no changes in benzodiazepine binding have been reported in several studies of postmortem human alcoholics (Dodd 1995; Dodd et al. 1992; Freund and Ballinger 1988).

In vivo neuroimaging studies have shown that 1–6 months abstinent male ethanol-dependent subjects have reduced binding of tracers for the GABA/benzodiazepine receptor in the frontal cortex, measured by [¹¹C]-flumazenil positron emission tomography (PET, Gilman et al. 1996) or [¹²³I]-iomazenil single photon emission computed tomography (SPECT, Abi-Dargham et al. 1998; Lingford-Hughes et al. 1998). A trend for a reduced [¹²³I]-iomazenil binding was reported in ethanol-dependent women withdrawn from ethanol with benzodiazepines and abstinent from ethanol for at least 3 months (Lingford-Hughes et al. 2000). Using a [2-deoxy-2-[¹⁸F]-fluoro-D-glucose] PET study, Volkow et al. (1997) found that the inhibitory response to a lorazepam challenge was blunted in the orbitofrontal cortex and cingulate gyrus in 8–11 weeks (during detoxification) abstinent ethanol-dependent subjects, suggesting reduced GABA/benzodiazepine receptor function. However, this study did not report withdrawal intensity. In contrast, after 1 week of abstinence from ethanol dependence, higher [¹²³I]-iomazenil binding to GABA/benzodiazepine receptors was observed in the parietal, frontal, cingulate, temporal, insular, and occipital cortex of non-smoking (but not smoking) ethanol-dependent subjects. Furthermore, the GABA/benzodiazepine receptor availability in the cerebellum and occipital lobe at 1 week of abstinence was correlated with the severity of ethanol withdrawal determined in the subjects every 6 h immediately after admission to the clinic (Staley et al. 2005). Using [¹¹C]-flumazenil PET to measure the pharmacokinetics and pharmacodynamics of midazolam, Lingford-Hughes et al. (2005) showed that ethanol dependence was associated with decreased midazolam-induced time asleep in the absence of reduced benzodiazepine receptor occupancy in ethanol-dependent

patients abstinent for at least 6 weeks. The authors hypothesized that this reduced sensitivity in ethanol dependence may reflect a difference in the subunit profile of the GABA_A receptors, either as part of the vulnerability to, or as a consequence of their ethanol dependence. More recently, Taylor et al. (2008) found that GABA_A receptor function, assayed by midazolam-induced slow saccadic eye movement and sedation, was not affected in ethanol-dependent patients after 2 months of abstinence.

These changes in GABA_A receptor function are accompanied by changes in receptor plasticity. Several studies reported changes in GABA_A receptor subunit expression in post-mortem brains from human alcoholics, although results are often controversial. Lewohl et al. (1997) showed that GABA_A receptor α 1 subunit mRNA expression was elevated in human alcoholic superior frontal cortex, compared with non-alcoholic subjects, while α 2 and α 3 mRNA did not change. Further, α 1 protein expression was greater in the superior frontal compared to the motor cortex of cirrhotic alcoholics, but no such difference was observed in controls or uncomplicated alcoholics. In contrast, α 3 peptide expression was lower in superior frontal than in motor cortex of uncomplicated alcoholics (Lewohl et al. 2001). However, Mitsuyama et al. (1998) found no difference in α 1 and α 4 subunits mRNA expression in frontal cortex between controls and alcoholics, but β 3 mRNA expression was elevated in non-cirrhotic alcoholics. These changes were not accompanied by changes in α 1, α 4, or β 2/3 protein expression between alcoholics and controls. Others have reported no differences in mRNA expression for any β subunit between alcoholics and controls (Buckley and Dodd 2004).

With the development of more sophisticated fine-mapping techniques, studies on GABA_A receptor subunit expression have been associated to specific genotypes. Thus, Dodd and colleagues reported that the GABRB2 gene, which encodes for the β 2 subunit, did not influence the β 2 subunit expression; rather, expression of β 3 subunit was increased in controls with GABRB2-1,2 genotype, but not in alcoholics with the same genotype (Dodd et al. 2004). Furthermore, aldehyde dehydrogenase ADH1C, D2 dopamine receptor DRD2B, excitatory amino acid transporter 2 EAAT2, and apolipoprotein E APOE genotypes modulate GABA_A receptor β subunit expression in superior frontal cortex toward a less-effective form of the receptor (Dodd et al. 2004, 2006). Alleles associated with alcoholism also appear to reduce the β 3: β 2 ratios in alcoholic's superior frontal cortex. GABA_A receptors with the β 2 subunit are less efficacious at mediating chloride flux in response to any GABA concentration, so this subunit switching would have the effect of attenuating GABA_A receptor-mediated inhibition and may make this brain region more susceptible to the neuropathological consequences of chronic alcoholism (Buckley et al. 2006). More recently, Haughey et al. (2008) reported that levels of α 2 subunit mRNA and protein in prefrontal cortex differed between subjects with GABRA2 (the gene encoding the α 2 subunit) genotypes, with higher levels of α 2 subunit expression in prefrontal cortex of subjects with the AA genotype compared to subjects with the AG genotype (SNP rs279858); however, there was no difference between ethanol-dependent and control subjects.

Polymorphisms in the genes encoding the GABA_A receptor subunits have been linked to ethanol response and risk for alcoholism in humans. Genes encoding the GABA_A receptor subunits are clustered in several chromosomal regions including 4p13–q11 (α 2, α 4, β 1, and γ 1), 5q34–q35 (α 1, α 6, β 2, and γ 2), 15q11–q13 (α 5, β 3, and γ 3), and Xq28 (α 3, β 4, and ϵ 1) (Enoch 2008). Results from the Collaborative Study on the Genetics of Alcoholism (COGA) found an increased allele sharing among ethanol-dependent individuals in a region on chromosome 4p that includes a cluster of four genes: GABRA2, GABRA4, GABRB1, and GABRG1, encoding for α 2, α 4, β 1, and γ 1 subunits, respectively (Reich 1996; Reich et al. 1998). Since then, several studies have shown that GABRA2 is a key gene affecting the risk for alcoholism. Edenberg et al. (2004) reported that single nucleotide polymorphisms (SNPs) in the GABRA2 gene, but not other members in the gene cluster, were associated with both

ethanol dependence and changes in β -frequency electroencephalogram (EEG). Brain oscillations measured by EEG are a key phenotype in ethanol dependence: alcoholics and the offspring of male alcoholics have increased power in the β -frequency band (13–18 Hz) of the EEG (Costa and Bauer 1997). The association of GABRA2 markers with ethanol dependence was subsequently replicated in several independent studies in different populations (Agrawal et al. 2006; Bauer et al. 2007; Covault et al. 2004; Enoch et al. 2006; Fehr et al. 2006; Lappalainen et al. 2005; Soyka et al. 2008). Within these studies, strong association of the GABRA2 gene was found in European ancestry alcoholics without co-morbid drug dependence or major depression (Covault et al. 2004), although results from the COGA study reported the opposite (Agrawal et al. 2006). The association of GABRA2 with alcoholism was linked to anxiety scores (Enoch et al. 2006) and to severity of ethanol withdrawal (Fehr et al. 2006; Soyka et al. 2008), thus potentially influencing treatment outcome (Soyka et al. 2008). Indeed, Bauer et al. (2007) found that genetic vulnerability at the GABRA2 gene influenced the ability of different psychosocial treatments to reduce ethanol use.

Sensitivity to the acute effects of ethanol has also been linked to the GABRA2 gene (Haughey et al. 2008; Pierucci-Lagha et al. 2005). Interestingly, individuals homozygous for the A-allele at GABRA2 reported greater stimulant, sedative, and gastrointestinal subjective effects of acute ethanol, compared with subjects with one or more G-alleles (the G-allele is overrepresented among ethanol-dependent subjects; Covault et al. 2004). Administration of the neuroactive steroid biosynthesis inhibitor finasteride antagonized some of the behavioral effects of ethanol particularly among individuals homozygous for the A-allele, suggesting that neuroactive steroids may be mediating the subjective effects of ethanol, like in rodent studies (Pierucci-Lagha et al. 2005). The genetic association of ethanol dependence with GABRA2 is of significant functional interest as animal studies have identified the $\alpha 2$ subunit as the primary α subunit in limbic regions (McKernan and Whiting 1996). The $\alpha 2$ subunit mediates the anxiolytic effect of benzodiazepines and barbiturates (Dixon et al. 2008; Rudolph et al. 1999) as well as the sedative-hypnotic, but not the sedative, effects of combined exposure to ethanol and benzodiazepines in genetically modified mouse studies (Tauber et al. 2003).

More recent reports suggest that the association of the GABRA2 gene with alcoholism may be complicated by moderate linkage disequilibrium of GABRA2 markers with a functional genetic variation in the adjacent GABRG1 gene (Covault et al. 2008; Drgon et al. 2006; Ittiwut et al. 2008). Indeed, Covault et al. (2008) reported that markers in the GABRG1 region on chromosome 4 had the strongest association with ethanol dependence in the European American population. The $\gamma 1$ subunit-containing GABA_A receptors show increased sensitivity to neuroactive steroids compared to the $\gamma 2$ subunit-containing GABA_A receptors (Puia et al. 1993). This subunit is increased in ethanol-dependent rats (Devaud et al. 1995) that are sensitized to the anticonvulsant effects of GABAergic neuroactive steroids (Devaud et al. 1996). Altered sensitivity to neuroactive steroids may affect ethanol sensitivity and therefore modulate risk for ethanol dependence (Morrow et al. 2006). In addition to GABRA2, other genes that encode for the GABA_A receptor subunits have been associated with ethanol dependence. Thus, on chromosome 4, association of the GABRB1 gene with alcoholism was reported by several groups (Long et al. 1998; Parsian and Zhang 1999; Reich et al. 1998; Song et al. 2003).

Results for markers in the GABA_A receptor gene cluster on chromosome 5 are still controversial, with associations reported in some samples, but not others (Dick et al. 2005; Sander et al. 1999; Song et al. 2003). Thus, a positive association between ethanol dependence and SNPs at the GABRG2 gene, encoding for the $\gamma 2$ subunit, was observed in Scottish (Loh et al. 1999), Finnish (Radel et al. 2005), and Japanese populations (Loh et al. 2000). Association with the GABRB2 gene, encoding for the $\beta 2$ subunit, was reported in Scottish (Loh et al. 1999), Finnish, and Southwestern Native American (Radel et al. 2005) populations. The

GABRA1 gene, encoding for the $\alpha 1$ subunit, was associated with several drinking behavior phenotypes, including history of blackouts, age at first drunkenness, level of response to ethanol, and ethanol dependence (Dick et al. 2006). Associations with the GABRA6, encoding for the $\alpha 6$ subunit, have been found in Scottish (Loh et al. 1999) and Finnish (Radel et al. 2005) populations. Furthermore, sons of alcoholics with the proline385 to serine polymorphism at GABRA6 were less sensitive to the effects of ethanol and this polymorphism was a predictor for the development of alcoholism (Schuckit et al. 1999). In the same subjects, this polymorphism was also associated with reduced sensitivity to benzodiazepines (Iwata et al. 1999) and to the sedative effects of ethanol (Hu et al. 2005).

Finally, fine mapping of a GABA_A receptor gene cluster on chromosome 15 reported modest evidence of haplotypic association to ethanol dependence for the GABRG3 gene, encoding the $\gamma 3$ subunit (Dick et al. 2004). Moreover, Song et al. (2003) showed evidence for imprinting association, with paternal, but not maternal, transmission of GABRA5 and GABRB3 genes, encoding the $\alpha 5$ and $\beta 3$ subunits, being associated with alcoholism.

Therapeutic implications for alcoholism

The discovery of new mechanisms of ethanol actions on GABAergic transmission over the past 10 years opens new opportunities to intervene in the effects of ethanol on different aspects of brain function. It is now possible to inhibit effects of ethanol by blockade of $\alpha 4$ -GABA_A receptors or by inhibition of neuroactive steroid synthesis. The blockade of ethanol actions may prevent ethanol-induced toxicity and/or facilitate increased consumption due to the loss of ethanol sensitivity. Likewise, it is also possible to enhance ethanol sensitivity by targeting $\alpha 4$ -GABA_A receptors with selective agonists or promoting neuroactive steroid synthesis. While there are many advances in the molecular realm, it remains unclear whether it is advantageous or disadvantageous to manipulate ethanol actions in humans. Future studies are clearly needed to elucidate the therapeutic relevance of the advances of the last 10 years.

Future directions—unanswered questions

The advances of the past 10 years raise many new questions to explore. The identification of ethanol binding sites on various GABA_A receptor subtypes is yet unclear. The role of protein kinases and phosphatases in ethanol actions are limited to too few GABA_A receptor subtypes. The coordinated actions of ethanol on multiple non-GABAergic targets that indirectly influence GABAergic neurotransmission seem too complex to untangle with physiological approaches. There is a lot of work to do.

Following chronic ethanol administration, there are specific adaptations of GABA_A receptors that appear to mediate increased CNS excitability. These mechanisms could be targeted for therapeutic benefits. However, there are many unanswered questions in this realm as well. What is the significance of changes in $\alpha 2$ -, $\alpha 3$ -, $\beta 2$ -, $\beta 3$ -, and $\gamma 1$ -GABA_A receptors? Can we modulate PKC and PKA interactions with GABA_A receptors selectively? Would effects of these protein kinases on other ion channels interfere with therapeutic strategies for modulation of GABA_A receptors? Can we target specific phosphorylation sites on specific proteins? Would this interfere with the actions of other endogenous modulators?

Standing back to widen our perspective, must we ask if our animal models are adequate for understanding the effects of ethanol on humans? Can we begin to model the genetic diversity of humans, the neurobiology, endocrinology, and enhanced complexity of the human being? Can we apply the knowledge gained from laboratory animals to develop treatments for ethanol addiction, toxicity, and dependence? History would argue the answer is yes, so our challenge is to choose the most important questions and persevere in the complexity of the problem to make contributions to science and society.

References

- Abi-Dargham A, Krystal JH, Anjilvel S, Scanley BE, Zoghbi S, Baldwin RM, Rajeevan N, Ellis S, Petrakis IL, Seibyl JP, Charney DS, Laruelle M, Innis RB. Alterations of benzodiazepine receptors in type II alcoholic subjects measured with SPECT and [¹²³I]iomazenil. *Am J Psychiatry* 1998;155:1550–1555. [PubMed: 9812116]
- Acheson SK, Richardson R, Swartzwelder HS. Developmental changes in seizure susceptibility during ethanol withdrawal. *Alcohol* 1999;18:23–26. [PubMed: 10386661]
- Acheson SK, Ross EL, Swartzwelder HS. Age-independent and dose–response effects of ethanol on spatial memory in rats. *Alcohol* 2001;23:167–175. [PubMed: 11435027]
- Agrawal A, Edenberg HJ, Foroud T, Bierut LJ, Dunne G, Hinrichs AL, Nurnberger JI, Crowe R, Kuperman S, Schuckit MA, Begleiter H, Porjesz B, Dick DM. Association of GABRA2 with drug dependence in the collaborative study of the genetics of alcoholism sample. *Behav Genet* 2006;36:640–650. [PubMed: 16622805]
- Allan AM, Harris RA. Acute and chronic ethanol treatments alter GABA receptor-operated chloride channels. *Pharmacol Biochem Behav* 1987;27:665–670. [PubMed: 2443933]
- Anderson NJ, Daunais JB, Friedman DP, Grant KA, McCool BA. Long-term ethanol self-administration by the nonhuman primate, *Macaca fascicularis*, decreases the benzodiazepine sensitivity of amygdala GABA_A receptors. *Alcohol Clin Exp Res* 2007;31:1061–1070. [PubMed: 17428292]
- Ariwodola OJ, Weiner JL. Ethanol potentiation of GABAergic synaptic transmission may be self-limiting: role of presynaptic GABA_B receptors. *J Neurosci* 2004;24:10679–10686. [PubMed: 15564584]
- Atack JR. Anxiolytic compounds acting at the GABA_A receptor benzodiazepine binding site. *Curr Drug Targets CNS Neurol Disord* 2003;2:213–232. [PubMed: 12871032]
- Atkins AL, Rustay NR, Crabbe JC. Anxiety and sensitivity to ethanol and pentobarbital in alcohol withdrawal seizure-prone and withdrawal seizure-resistant mice. *Alcohol Clin Exp Res* 2000;24:1743–1749. [PubMed: 11141031]
- Barbaccia ML, Affricano D, Trabucchi M, Purdy RH, Colombo G, Agabio R, Gessa GL. Ethanol markedly increases “GABAergic” neurosteroids in alcohol-preferring rats. *Eur J Pharmacol* 1999;384:R1–R2. [PubMed: 10611449]
- Bauer LO, Covault J, Harel O, Das S, Gelernter J, Anton R, Kranzler HR. Variation in GABRA2 predicts drinking behavior in project MATCH subjects. *Alcohol Clin Exp Res* 2007;31:1780–1787. [PubMed: 17949392]
- Becker HC, Veatch LM. Effects of lorazepam treatment for multiple ethanol withdrawals in mice. *Alcohol Clin Exp Res* 2002;26:371–380. [PubMed: 11923591]
- Becker HC, Diaz-Granados JL, Weathersby RT. Repeated ethanol withdrawal experience increases the severity and duration of subsequent withdrawal seizures in mice. *Alcohol* 1997;14:319–326. [PubMed: 9209546]
- Bell RL, Rodd ZA, Sable HJ, Schultz JA, Hsu CC, Lumeng L, Murphy JM, McBride WJ. Daily patterns of ethanol drinking in peri-adolescent and adult alcohol-preferring (P) rats. *Pharmacol Biochem Behav* 2006;83:35–46. [PubMed: 16442608]
- Ben-Ari Y. Excitatory actions of gaba during development: the nature of the nurture. *Nat Rev Neurosci* 2002;3:728–739. [PubMed: 12209121]
- Bencsits E, Ebert V, Tretter V, Sieghart W. A significant part of native gamma-aminobutyric AcidA receptors containing alpha4 subunits do not contain gamma or delta subunits. *J Biol Chem* 1999;274:19613–19616. [PubMed: 10391897]
- Besheer J, Hodge CW. Pharmacological and anatomical evidence for an interaction between mGluR5- and GABA_A α1-containing receptors in the discriminative stimulus effects of ethanol. *Neuropsychopharmacology* 2005;30:747–757. [PubMed: 15549054]
- Besheer J, Cox AA, Hodge CW. Coregulation of ethanol discrimination by the nucleus accumbens and amygdala. *Alcohol Clin Exp Res* 2003;27:450–456. [PubMed: 12658110]
- Biggio, G.; Purdy, RH. *International Review of Neurobiology*. Vol. 46. Academic Press; New York: 2001. Neurosteroids and brain function.

- Blednov YA, Jung S, Alva H, Wallace D, Rosahl T, Whiting PJ, Harris RA. Deletion of the $\alpha 1$ or $\beta 2$ subunit of GABA_A receptors reduces actions of alcohol and other drugs. *J Pharmacol Exp Ther* 2003a;304:30–36. [PubMed: 12490572]
- Blednov YA, Walker D, Alva H, Creech K, Findlay G, Harris RA. GABA_A receptor $\alpha 1$ and $\beta 2$ subunit null mutant mice: behavioral responses to ethanol. *J Pharmacol Exp Ther* 2003b;305:854–863. [PubMed: 12626647]
- Boehm SL 2nd, Peden L, Jennings AW, Kojima N, Harris RA, Blednov YA. Over-expression of the *fyn*-kinase gene reduces hypnotic sensitivity to ethanol in mice. *Neurosci Lett* 2004a;372:6–11. [PubMed: 15531078]
- Boehm SL 2nd, Ponomarev I, Jennings AW, Whiting PJ, Rosahl TW, Garrett EM, Blednov YA, Harris RA. γ -Aminobutyric acid A receptor subunit mutant mice: new perspectives on alcohol actions. *Biochem Pharmacol* 2004b;68:1581–1602. [PubMed: 15451402]
- Boehm SL II, Peden L, Harris RA, Blednov Y. Deletion of the *fyn*-kinase gene alters sensitivity to GABAergic drugs: dependence on $\beta 2/\beta 3$ GABA_A receptor subunits. *J Pharmacol Exp Ther* 2004c;309:1154–1159. [PubMed: 14764659]
- Boehm SL 2nd, Ponomarev I, Blednov YA, Harris RA. From gene to behavior and back again: new perspectives on GABA_A receptor subunit selectivity of alcohol actions. *Adv Pharmacol* 2006;54:171–203. [PubMed: 17175815]
- Borghese CM, Sturustovu SI, Ebert B, Herd MB, Belelli D, Lambert JJ, Marshall G, Wafford KA, Harris RA. The δ subunit of γ -aminobutyric acid type A receptors does not confer sensitivity to low concentrations of ethanol. *J Pharmacol Exp Ther* 2006a;316:1360–1368. [PubMed: 16272217]
- Borghese CM, Werner DF, Topf N, Baron NV, Henderson LA, Boehm SL 2nd, Blednov YA, Saad A, Dai S, Pearce RA, Harris RA, Homanics GE, Harrison NL. An isoflurane- and alcohol-insensitive mutant GABA_A receptor $\alpha 1$ subunit with near-normal apparent affinity for GABA: characterization in heterologous systems and production of knockin mice. *J Pharmacol Exp Ther* 2006b;319:208–218. [PubMed: 16807363]
- Borlikova GG, Le Merrer J, Stephens DN. Previous experience of ethanol withdrawal increases withdrawal-induced *c-fos* expression in limbic areas, but not withdrawal-induced anxiety and prevents withdrawal-induced elevations in plasma corticosterone. *Psychopharmacology (Berl)* 2006;185:188–200. [PubMed: 16470400]
- Botta P, Mameli M, Floyd KL, Radcliffe RA, Valenzuela CF. Ethanol sensitivity of GABAergic currents in cerebellar granule neurons is not increased by a single amino acid change (R100Q) in the $\alpha 6$ GABA_A receptor subunit. *J Pharmacol Exp Ther* 2007a;323:684–691. [PubMed: 17704355]
- Botta P, Radcliffe RA, Carta M, Mameli M, Daly E, Floyd KL, Deitrich RA, Valenzuela CF. Modulation of GABA_A receptors in cerebellar granule neurons by ethanol: a review of genetic and electrophysiological studies. *Alcohol* 2007b;41:187–199. [PubMed: 17521847]
- Bowers BJ, Owen EH, Collins AC, Abeliovich A, Tonegawa S, Wehner JM. Decreased ethanol sensitivity and tolerance development in gamma-protein kinase C null mutant mice is dependent on genetic background. *Alcohol Clin Exp Res* 1999;23:387–397. [PubMed: 10195808]
- Bowers BJ, Elliott KJ, Wehner JM. Differential sensitivity to the anxiolytic effects of ethanol and flunitrazepam in PKC γ null mutant mice. *Pharmacol Biochem Behav* 2001;69:99–110. [PubMed: 11420074]
- Brandon NJ, Delmas P, Kittler JT, McDonald BJ, Sieghart W, Brown DA, Smart TG, Moss SJ. GABA_A receptor phosphorylation and functional modulation in cortical neurons by a protein kinase C-dependent pathway. *J Biol Chem* 2000;275:38856–38862. [PubMed: 10978327]
- Brasser SM, Spear NE. Physiological and behavioral effects of acute ethanol hangover in juvenile, adolescent, and adult rats. *Behav Neurosci* 2002;116:305–320. [PubMed: 11996316]
- Brickley SG, Revilla V, Cull-Candy SG, Wisden W, Farrant M. Adaptive regulation of neuronal excitability by a voltage-independent potassium conductance. *Nature* 2001;409:88–92. [PubMed: 11343119]
- Brown N, Kerby J, Bonnert TP, Whiting PJ, Wafford KA. Pharmacological characterization of a novel cell line expressing human $\alpha 4\beta 3\delta$ GABA_A receptors. *Br J Pharmacol* 2002;136:965–974. [PubMed: 12145096]

- Brüning I, Penschuck S, Berninger B, Benson J, Fritschy JM. BDNF reduces miniature inhibitory postsynaptic currents by rapid downregulation of GABA_A receptor surface expression. *Eur J Neurosci* 2001;13:1320–1328. [PubMed: 11298792]
- Buchner K, Adamec E, Beermann ML, Nixon RA. Isoform-specific translocation of protein kinase C following glutamate administration in primary hippocampal neurons. *Brain Res Mol Brain Res* 1999;64:222–235. [PubMed: 9931492]
- Buck KJ, Harris RA. Benzodiazepine agonist and inverse agonist actions on GABA_A receptor-operated chloride channels. II. Chronic effects of ethanol. *J Pharmacol Exp Ther* 1990;253:713–719. [PubMed: 2160008]
- Buck KJ, Hahner L, Sikela J, Harris RA. Chronic ethanol treatment alters brain levels of gamma-aminobutyric acid_A receptor subunit mRNAs: relationship to genetic differences in ethanol withdrawal seizure severity. *J Neurochem* 1991a;57:1452–1455. [PubMed: 1654394]
- Buck KJ, Heim H, Harris RA. Reversal of alcohol dependence and tolerance by a single administration of flumazenil. *J Pharmacol Exp Ther* 1991b;257:984–989. [PubMed: 1646333]
- Buckley ST, Dodd PR. GABA_A receptor β subunit mRNA expression in the human alcoholic brain. *Neurochem Int* 2004;45:1011–1020. [PubMed: 15337300]
- Buckley ST, Foley PF, Innes DJ, el Loh W, Shen Y, Williams SM, Harper CG, Tannenbergs AE, Dodd PR. GABA_A receptor β isoform protein expression in human alcoholic brain: interaction with genotype. *Neurochem Int* 2006;49:557–567. [PubMed: 16766085]
- Cagetti E, Liang J, Spigelman I, Olsen RW. Withdrawal from chronic intermittent ethanol treatment changes subunit composition, reduces synaptic function, and decreases behavioral responses to positive allosteric modulators of GABA_A receptors. *Mol Pharmacol* 2003;63:53–64. [PubMed: 12488536]
- Cagetti E, Pinna G, Guidotti A, Baicy K, Olsen RW. Chronic intermittent ethanol (CIE) administration in rats decreases levels of neurosteroids in hippocampus, accompanied by altered behavioral responses to neurosteroids and memory function. *Neuropharmacology* 2004;46:570–579. [PubMed: 14975681]
- Calkin PA, Barnes EM Jr. Gamma-aminobutyric acid-A (GABA_A) agonists down-regulate GABA_A/benzodiazepine receptor polypeptides from the surface of chick cortical neurons. *J Biol Chem* 1994;269:1548–1553. [PubMed: 8288621]
- Carta M, Mameli M, Valenzuela CF. Alcohol enhances GABAergic transmission to cerebellar granule cells via an increase in Golgi cell excitability. *J Neurosci* 2004;24:3746–3751. [PubMed: 15084654]
- Casagrande S, Cupello A, Pellistri F, Robello M. Only high concentrations of ethanol affect GABA_A receptors of rat cerebellum granule cells in culture. *Neurosci Lett* 2007;414:273–276. [PubMed: 17234340]
- Chandra D, Jia F, Liang J, Peng Z, Suryanarayanan A, Werner DF, Spigelman I, Houser CR, Olsen RW, Harrison NL, Homanics GE. GABA_A receptor α 4 subunits mediate extrasynaptic inhibition in thalamus and dentate gyrus and the action of gaboxadol. *Proc Natl Acad Sci U S A* 2006;103:15230–15235. [PubMed: 17005728]
- Chandra D, Werner DF, Liang J, Suryanarayanan A, Harrison NL, Spigelman I, Olsen RW, Homanics GE. Normal acute behavioral responses to moderate/high dose ethanol in GABA_A receptor alpha 4 subunit knockout mice. *Alcohol Clin Exp Res* 2008;32:10–18. [PubMed: 18076749]
- Charlton ME, Sweetnam PM, Fitzgerald LW, Terwilliger RZ, Nestler EJ, Duman RS. Chronic ethanol administration regulates the expression of GABA_A receptor α 1 and α 5 subunits in the ventral tegmental area and hippocampus. *J Neurochem* 1997;68:121–127. [PubMed: 8978717]
- Chen ZW, Olsen RW. GABA_A receptor associated proteins: a key factor regulating GABA_A receptor function. *J Neurochem* 2007;100:279–294. [PubMed: 17083446]
- Chester JA, Cunningham CL. GABA_A receptors modulate ethanol induced conditioned place preference and taste aversion in mice. *Psychopharmacology* 1999;144:363–372. [PubMed: 10435409]
- Chester JA, Risinger FO, Cunningham CL. Ethanol reward and aversion in mice bred for sensitivity to ethanol withdrawal. *Alcohol Clin Exp Res* 1998;22:468–473. [PubMed: 9581655]
- Choi DS, Wei W, Deitchman JK, Kharazia VN, Lesscher HM, McMahon T, Wang D, Qi ZH, Sieghart W, Zhang C, Shokat KM, Mody I, Messing RO. Protein kinase C δ regulates ethanol intoxication

- and enhancement of GABA-stimulated tonic current. *J Neurosci* 2008;28:11890–11899. [PubMed: 19005054]
- Colombo G, Agabio R, Lobina C, Loche A, Reali R, Gessa GL. High sensitivity to gamma-hydroxybutyric acid in ethanol-preferring sP rats. *Alcohol Alcohol* 1998;33:121–125. [PubMed: 9566473]
- Colombo G, Agabio R, Carai MA, Lobina C, Pani M, Reali R, Vacca G, Gessa GL. Different sensitivity to ethanol in alcohol-preferring sP and -nonpreferring sNP rats. *Alcohol Clin Exp Res* 2000;24:1603–1608. [PubMed: 11104106]
- Connolly CN, Kittler JT, Thomas P, Uren JM, Brandon NJ, Smart TG, Moss SJ. Cell surface stability of gamma-aminobutyric acid type A receptors: dependence on protein kinase C activity and subunit composition. *J Biol Chem* 1999;274:36565–36572. [PubMed: 10593956]
- Cook JB, Foster KL, Eiler WJ 2nd, McKay PF, Woods J 2nd, Harvey SC, Garcia M, Grey C, McCane S, Mason D, Cummings R, Li X, Cook JM, June HL. Selective GABA_A α 5 benzodiazepine inverse agonist antagonizes the neurobehavioral actions of alcohol. *Alcohol Clin Exp Res* 2005;29:1390–1401. [PubMed: 16131846]
- Costa L, Bauer L. Quantitative electroencephalographic differences associated with alcohol, cocaine, heroin and dual-substance dependence. *Drug Alcohol Depend* 1997;46:87–93. [PubMed: 9246556]
- Covault J, Gelernter J, Hesselbrock V, Nellisery M, Kranzler HR. Allelic and haplotypic association of GABRA2 with alcohol dependence. *Am J Med Genet B Neuropsychiatr Genet* 2004;129:104–109. [PubMed: 15274050]
- Covault J, Gelernter J, Jensen K, Anton R, Kranzler HR. Markers in the 5'-region of GABRG1 associate to alcohol dependence and are in linkage disequilibrium with markers in the adjacent GABRA2 gene. *Neuropsychopharmacology* 2008;33:837–848. [PubMed: 17507911]
- Cowley DS, Roy-Byrne PP, Radant A, Hommer DW, Greenblatt DJ, Vitaliano PP, Godon C. Eye movement effects of diazepam in sons of alcoholic fathers and male control subjects. *Alcohol Clin Exp Res* 1994;18:324–332. [PubMed: 8048734]
- Crabbe JC, Kosobud A. Sensitivity and tolerance to ethanol in mice bred to be genetically prone or resistant to ethanol withdrawal seizures. *J Pharmacol Exp Ther* 1986;239:327–333. [PubMed: 3772797]
- Crabbe JC, Phillips TJ, Harris RA, Arends MA, Koob GF. Alcohol-related genes: contributions from studies with genetically engineered mice. *Addict Biol* 2006;11:195–269. [PubMed: 16961758]
- Crews, F.; Morrow, AL.; Criswell, H.; Breese, G. Effects of ethanol on ion channels. In: Bradley, RJ.; Harris, RA., editors. *International review of neurobiology*. Vol. 39. Academic; New York: 1996. p. 283–367.
- Crews FT, Bechara R, Brown LA, Guidot DM, Mandrekar P, Oak S, Qin L, Szabo G, Wheeler M, Zou JY. Cytokines and alcohol. *Alcohol Clin Exp Res* 2006;30:720–730. [PubMed: 16573591]
- Criswell HE, Breese GR. A conceptualization of integrated actions of ethanol contributing to its GABA_A mimetic profile: a commentary. *Neuropsychopharmacology* 2005;30:1407–1425. [PubMed: 15856077]
- Criswell HE, Ming Z, Kelm MK, Breese GR. Brain regional differences in the effect of ethanol on GABA release from presynaptic terminals. *J Pharmacol Exp Ther* 2008;326:596–603. [PubMed: 18502983]
- Davidson M, Chen W, Wilce PA. Behavioral analysis of PTZ-kindled rats after acute and chronic ethanol treatments. *Pharmacol Biochem Behav* 1999;64:7–13. [PubMed: 10494991]
- de Almeida RM, Rowlett JK, Cook JM, Yin W, Miczek KA. GABA_A/ α 1 receptor agonists and antagonists: effects on species-typical and heightened aggressive behavior after alcohol self-administration in mice. *Psychopharmacology (Berl)* 2004;172:255–263. [PubMed: 14647970]
- De Witte P, Dahchour A, Quertemont E. Acute and chronic alcohol injections increase taurine in the nucleus accumbens. *Alcohol Alcohol Suppl* 1994;2:229–233. [PubMed: 8974341]
- Devaud LL, Smith FD, Grayson DR, Morrow AL. Chronic ethanol consumption differentially alters the expression of γ -aminobutyric acid type A receptor subunit mRNAs in rat cerebral cortex: competitive, quantitative reverse transcriptase-polymerase chain reaction analysis. *Mol Pharmacol* 1995;48:861–868. [PubMed: 7476917]

- Devaud LL, Purdy RH, Finn DA, Morrow AL. Sensitization of γ -aminobutyric acid_A receptors to neuroactive steroids in rats during ethanol withdrawal. *J Pharmacol Exp Ther* 1996;278:510–517. [PubMed: 8768698]
- Devaud LL, Fritschy J-M, Sieghart W, Morrow AL. Bidirectional alterations of GABA_A receptor subunit peptide levels in rat cortex during chronic ethanol consumption and withdrawal. *J Neurochem* 1997;69:126–130. [PubMed: 9202302]
- Devaud LL, Fritschy J-M, Morrow AL. Influence of gender on chronic ethanol-induced alternations of GABA_A receptors in rats. *Brain Res* 1998;796:222–230. [PubMed: 9689472]
- Diamond I, Gordon AS. The role of adenosine in mediating cellular and molecular responses to ethanol. *EXS* 1994;71:175–183. [PubMed: 8032148]
- Diaz-Granados JL, Graham DL. The effects of continuous and intermittent ethanol exposure in adolescence on the aversive properties of ethanol during adulthood. *Alcohol Clin Exp Res* 2007;12:2020–2027. [PubMed: 18034694]
- Dick DM, Edenberg HJ, Xuei X, Goate A, Kuperman S, Schuckit M, Crowe R, Smith TL, Porjesz B, Begleiter H, Foroud T. Association of GABRG3 with alcohol dependence. *Alcohol Clin Exp Res* 2004;28:4–9. [PubMed: 14745296]
- Dick DM, Edenberg HJ, Xuei X, Goate A, Hesselbrock V, Schuckit M, Crowe R, Foroud T. No association of the GABA_A receptor genes on chromosome 5 with alcoholism in the collaborative study on the genetics of alcoholism sample. *Am J Med Genet B Neuropsychiatr Genet* 2005;132B:24–28. [PubMed: 15690551]
- Dick DM, Plunkett J, Wetherill LF, Xuei X, Goate A, Hesselbrock V, Schuckit M, Crowe R, Edenberg HJ, Foroud T. Association between GABRA1 and drinking behaviors in the collaborative study on the genetics of alcoholism sample. *Alcohol Clin Exp Res* 2006;30:1101–1110. [PubMed: 16792556]
- Dixon CI, Rosahl TW, Stephens DN. Targeted deletion of the *GABRA2* gene encoding $\alpha 2$ -subunits of GABA_A receptors facilitates performance of a conditioned emotional response, and abolishes anxiolytic effects of benzodiazepines and barbiturates. *Pharmacol Biochem Behav* 2008;90:1–8. [PubMed: 18313124]
- Dodd PR, Thomas GJ, Harper CG, Kril JJ. Amino acid neurotransmitter receptor changes in cerebral cortex in alcoholism: effect of cirrhosis of the liver. *J Neurochem* 1992;59:1506–1515. [PubMed: 1328522]
- Dodd PR. Benzodiazepine binding sites in alcoholic cirrhotics: evidence for gender differences. *Metab Brain Dis* 1995;10:93–104. [PubMed: 7596333]
- Dodd PR, Foley PF, Buckley ST, Eckert AL, Innes DJ. Genes and gene expression in the brain of the alcoholic. *Addict Behav* 2004;29:1295–1309. [PubMed: 15345266]
- Dodd PR, Buckley ST, Eckert AL, Foley PF, Innes DJ. Genes and gene expression in the brains of human alcoholics. *Ann N Y Acad Sci* 2006;1074:104–115. [PubMed: 17105908]
- Doremus TL, Brunell SC, Varlinskaya EI, Spear LP. Anxio-genic effects during withdrawal from acute ethanol in adolescent and adult rats. *Pharmacol Biochem Behav* 2003;75:411–418. [PubMed: 12873633]
- Doremus TL, Brunell SC, Rajendran P, Spear LP. Factors influencing elevated ethanol consumption in adolescent relative to adult rats. *Alcohol Clin Exp Res* 2005;29:1796–1808. [PubMed: 16269909]
- Drgon T, D'Addario C, Uhl GR. Linkage disequilibrium, haplotype and association studies of a chromosome 4 GABA receptor gene cluster: candidate gene variants for addictions. *Am J Med Genet B Neuropsychiatr Genet* 2006;141B:854–860. [PubMed: 16894595]
- Edenberg HJ, Dick DM, Xuei X, Tian H, Almasy L, Bauer LO, Crowe RR, Goate A, Hesselbrock V, Jones K, Kwon J, Li TK, Nurnberger JI Jr, O'Connor SJ, Reich T, Rice J, Schuckit MA, Porjesz B, Foroud T, Begleiter H. Variations in *GABRA2*, encoding the $\alpha 2$ subunit of the GABA_A receptor, are associated with alcohol dependence and with brain oscillations. *Am J Hum Genet* 2004;74:705–714. [PubMed: 15024690]
- Enoch MA. The role of GABA_A receptors in the development of alcoholism. *Pharmacol Biochem Behav* 2008;90:95–104. [PubMed: 18440057]
- Enoch MA, Schwartz L, Albaugh B, Virkkunen M, Goldman D. Dimensional anxiety mediates linkage of *GABRA2* haplotypes with alcoholism. *Am J Med Genet B Neuropsychiatr Genet* 2006;141B:599–607. [PubMed: 16874763]

- Ericson M, Haythornthwaite AR, Yeh PW, Yeh HH. Brain-derived neurotrophic factor mitigates chronic ethanol-induced attenuation of gamma-aminobutyric acid responses in cultured cerebellar granule cells. *J Neurosci Res* 2003;73:722–730. [PubMed: 12929140]
- Etoh S, Baba A, Iwata H. NMDA induces protein kinase C translocation in hippocampal slices of immature rat brain. *Neurosci Lett* 1991;126:119–122. [PubMed: 1922921]
- Farrant M, Nusser Z. Variations on an inhibitory theme: phasic and tonic activation of GABA_A receptors. *Nat Rev Neurosci* 2005;6:215–229. [PubMed: 15738957]
- Fehr C, Sander T, Tadic A, Lenzen KP, Anghelescu I, Klawe C, Dahmen N, Schmidt LG, Szegedi A. Confirmation of association of the GABRA2 gene with alcohol dependence by subtype-specific analysis. *Psychiatr Genet* 2006;16:9–17. [PubMed: 16395124]
- Feng J, Cai X, Zhao J, Yan Z. Serotonin receptors modulate GABA_A receptor channels through activation of anchored protein kinase C in prefrontal cortical neurons. *J Neurosci* 2001;21:6502–6511. [PubMed: 11517239]
- File SE, Pellow S. Intrinsic actions of the benzodiazepine receptor antagonist Ro 15–1788. *Psychopharmacology (Berl)* 1986;88:1–11. [PubMed: 3003778]
- Fish EW, Faccidomo S, DeBold JF, Miczek KA. Alcohol, allopregnanolone and aggression in mice. *Psychopharmacology* 2001;153:473–483. [PubMed: 11243495]
- Fleming RL, Wilson WA, Swartzwelder HS. Magnitude and ethanol sensitivity of tonic GABA_A receptor-mediated inhibition in dentate gyrus changes from adolescence to adulthood. *J Neurophysiol* 2007;97:3806–3811. [PubMed: 17376852]
- Flores-Hernandez J, Hernandez S, Snyder GL, Yan Z, Fienberg AA, Moss SJ, Greengard P, Surmeier DJ. D(1) dopamine receptor activation reduces GABA_A receptor currents in neo-striatal neurons through a PKA/DARPP-32/PP1 signaling cascade. *J Neurophysiol* 2000;83:2996–3004. [PubMed: 10805695]
- Floyd DW, Friedman DP, Daunais JB, Pierre PJ, Grant KA, McCool BA. Long-term ethanol self-administration by cynomolgus macaques alters the pharmacology and expression of GABA_A receptors in basolateral amygdala. *J Pharmacol Exp Ther* 2004;311:1071–1079. [PubMed: 15280440]
- Follesa P, Biggio F, Talani G, Murru L, Serra M, Sanna E, Biggio G. Neurosteroids, GABA_A receptors, and ethanol dependence. *Psychopharmacology* 2006;186:267–280. [PubMed: 16163526]
- Freund G, Ballinger WEJ. Decrease of benzodiazepine receptors in frontal cortex of alcoholics. *Alcohol* 1988;5:275–282. [PubMed: 2852495]
- Fritschy J-M, Paysan J, Enna A, Möhler H. Switch in the expression of rat GABA_A receptor subtypes during postnatal development: an immunohistochemical study. *J Neurosci* 1994;14:5302–5324. [PubMed: 8083738]
- Frye CA, Van Keuren KR, Erskine MS. Behavioral effects of 3 α -androstenediol 1. Modulation of sexual receptivity and promotion of GABA-stimulated chloride flux. *Behav Brain Res* 1996;79:109–118. [PubMed: 8883822]
- Gabriel KI, Cunningham CL, Finn DA. Allopregnanolone does not influence ethanol-induced conditioned place preference in DBA/2J mice. *Psychopharmacology (Berl)* 2004;176:50–56. [PubMed: 15083256]
- Galindo R, Valenzuela CF. Immature hippocampal neuronal networks do not develop tolerance to the excitatory actions of ethanol. *Alcohol* 2006;40:111–118. [PubMed: 17307647]
- Gallagher EJ, Gionet SE. Initial sensitivity and tolerance to ethanol in mice genetically selected for diazepam sensitivity. *Alcohol Clin Exp Res* 1988;12:77–80. [PubMed: 2831751]
- Gallagher EJ, Jones GE, Belknap JK, Crabbe JC. Identification of genetic markers for initial sensitivity and rapid tolerance to ethanol-induced ataxia using quantitative trait locus analysis in BXD recombinant inbred mice. *J Pharmacol Exp Ther* 1996;277:604–612. [PubMed: 8627537]
- Gatch MB. Effects of benzodiazepines on acute and chronic ethanol-induced nociception in rats. *Alcohol Clin Exp Res* 1999;23:1736–1743. [PubMed: 10591589]
- Gilman S, Koeppe RA, Adams K, Johnson-Greene D, Junck L, Kluin KJ, Brunberg J, Martorello S, Lohman M. Positron emission tomographic studies of cerebral benzodiazepine-receptor binding in chronic alcoholics. *Ann Neurol* 1996;40:163–171. [PubMed: 8773597]

- Glykys J, Peng Z, Chandra D, Homanics GE, Houser CR, Mody I. A new naturally occurring GABA_A receptor subunit partnership with high sensitivity to ethanol. *Nat Neurosci* 2007;10:40–48. [PubMed: 17159992]
- Graham DL, Diaz-Granados JL. Periadolescent exposure to ethanol and diazepam alters the aversive properties of ethanol in adult mice. *Pharmacol Biochem Behav* 2006;84:406–414. [PubMed: 16844209]
- Grant KA. Strategies for understanding the pharmacological effects of ethanol with drug discrimination procedures. *Pharmacol Biochem Behav* 1999;64:261–267. [PubMed: 10515301]
- Grant KA, Azarov A, Bowen CA, Mirkis S, Purdy RH. Ethanol-like discriminative stimulus effects of the neurosteroid 3 α -hydroxy-5 α -pregnan-20-one in female *Macaca fascicularis* monkeys. *Psychopharmacology* 1996;124:340–346. [PubMed: 8739549]
- Grant KA, Helms CM, Rogers LS, Purdy RH. Neuroactive steroid stereospecificity of ethanol-like discriminative stimulus effects in monkeys. *J Pharmacol Exp Ther* 2008a;326:354–361. [PubMed: 18436788]
- Grant KA, Leng X, Green HL, Szeliga KT, Rogers LS, Gonzales SW. Drinking typography established by scheduled induction predicts chronic heavy drinking in a monkey model of ethanol self-administration. *Alcohol Clin Exp Res* 2008b;32:1824–1838. [PubMed: 18702645]
- Grobin AC, Matthews DB, Devaud LL, Morrow AL. The role of GABA_A receptors in the acute and chronic effects of ethanol. *Psychopharmacology* 1998;139:2–19. [PubMed: 9768538]
- Grobin AC, Fritschy J-M, Morrow AL. Chronic ethanol administration alters immunoreactivity for GABA_A receptor subunits in rat cortex in a region-specific manner. *Alcohol Clin Exp Res* 2000;24:1137–1144. [PubMed: 10968650]
- Grobin AC, Matthews DB, Montoya D, Wilson WA, Morrow AL, Swartzwelder HS. Age-related differences in neurosteroid potentiation of muscimol-stimulated ³⁶Cl-flux following chronic ethanol treatment. *Neuroscience* 2001;105:547–552. [PubMed: 11516822]
- Gutiérrez A, Khan ZU, Miralles CP, Mehta AK, Ruano D, Araujo F, Vitorica J, De Blas AL. GABA_A receptor subunit expression changes in the rat cerebellum and cerebral cortex during aging. *Brain Res* 1997;45:59–70.
- Hanchar HJ, Dodson PD, Olsen RW, Otis TS, Wallner M. Alcohol-induced motor impairment caused by increased extra-synaptic GABA_A receptor activity. *Nat Neurosci* 2005;8:339–345. [PubMed: 15696164]
- Hanchar HJ, Chutrinopkun P, Meera P, Supavilai P, Sieghart W, Wallner M, Olsen RW. Ethanol potently and competitively inhibits binding of the alcohol antagonist Ro15–4513 to alpha4/6beta3delta GABA_A receptors. *Proc Natl Acad Sci U S A* 2006;103:8546–8551. [PubMed: 16581914]
- Harris RA, McQuilkin SJ, Paylor R, Abeliovich A, Tonegawa S, Wehner JM. Mutant mice lacking the gamma isoform of protein kinase C show decreased behavioral actions of ethanol and altered function of gamma-aminobutyrate type A receptors. *Proc Natl Acad Sci U S A* 1995;92:3658–3662. [PubMed: 7731960]
- Harvey SC, Foster KL, McKay PF, Carroll MR, Seyoum R, Woods JE 2nd, Grey C, Jones CM, McCane S, Cummings R, Mason D, Ma C, Cook JM, June HL. The GABA_A receptor α 1 subtype in the ventral pallidum regulates alcohol-seeking behaviors. *J Neurosci* 2002;22:3765–3775. [PubMed: 11978852]
- Haughey HM, Ray LA, Finan P, Villanueva R, Niculescu M, Hutchison KE. Human γ -aminobutyric acid A receptor α 2 gene moderates the acute effects of alcohol and brain mRNA expression. *Genes Brain Behav* 2008;7:447–454. [PubMed: 18005236]
- Hefner K, Holmes A. Ontogeny of fear-, anxiety- and depression-related behavior across adolescence in C57BL/6J mice. *Behav Brain Res* 2007;176:210–215. [PubMed: 17098297]
- Helms CM, Rogers LS, Waters CA, Grant KA. Zolpidem generalization and antagonism in male and female cynomolgus monkeys trained to discriminate 1.0 or 2.0 g/kg ethanol. *Alcohol Clin Exp Res* 2008;32:1197–1206. [PubMed: 18482161]
- Hemby SE, O'Connor JA, Acosta G, Floyd D, Anderson N, McCool BA, Friedman D, Grant KA. Ethanol-induced regulation of GABA_A subunit mRNAs in prefrontal fields of cynomolgus monkeys. *Alcohol Clin Exp Res* 2006;30:1978–1985. [PubMed: 17117962]

- Henschel O, Gipson KE, Bordey A. GABA_A receptors, anesthetics and anticonvulsants in brain development. *CNS Neurol Disord Drug Targets* 2008;7:211–224. [PubMed: 18537647]
- Herring D, Huang R, Singh M, Robinson LC, Dillon GH, Leidenheimer NJ. Constitutive GABA_A receptor endocytosis is dynamin-mediated and dependent on a dileucine AP2 adaptin-binding motif within the β 2 subunit of the receptor. *J Biol Chem* 2003;278:24046–24052. [PubMed: 12707262]
- Hirani K, Khisti RT, Chopde CT. Behavioral action of ethanol in Porsolt's forced swim test: modulation by 3 α -hydroxy-5 α -pregnan-20-one. *Neuropharmacology* 2002;43:1339–1350. [PubMed: 12527484]
- Hirani K, Sharma AN, Jain NS, Ugale RR, Chopde CT. Evaluation of GABAergic neuroactive steroid 3 α -hydroxy-5 α -pregnan-20-one as a neurobiological substrate for the anti-anxiety effect of ethanol in rats. *Psychopharmacology* 2005;180:267–278. [PubMed: 15719223]
- Hodge CW, Cox AA. The discriminative stimulus effects of ethanol are mediated by NMDA and GABA_A receptors in specific limbic brain regions. *Psychopharmacology* 1998;139:95–107. [PubMed: 9768547]
- Hodge CW, Mehmert KK, Kelley SP, McMahon T, Haywood A, Olive MF, Wang D, Sanchez-Perez AM, Messing RO. Super-sensitivity to allosteric GABA_A receptor modulators and alcohol in mice lacking PKC ϵ . *Nat Neurosci* 1999;2:997–1002. [PubMed: 10526339]
- Hoffpauir B, Gleason E. Activation of mGluR5 modulates GABA_A receptor function in retinal amacrine cells. *J Neurophysiol* 2002;88:1766–1776. [PubMed: 12364505]
- Hollstedt C, Olsson O, Rydberg U. The effect of alcohol on the developing organism. Genetical, teratological and physiological aspects. *Med Biol* 1977;55:1–14. [PubMed: 321891]
- Holtzman, SG. Discriminative stimulus effects of drugs: relationship to potential for abuse. In: Adler, MW.; Cowan, A., editors. *Modern methods in pharmacology. Testing and evaluation of drugs of abuse*. Vol. 6. Wiley-Liss; New York: 1990. p. 193-210.
- Homanics GE, Ferguson C, Quinlan JJ, Daggett J, Snyder K, Lagenaur C, Mi ZP, Wang XH, Grayson DR, Firestone LL. Gene knockout of the α 6 subunit of the γ -aminobutyric acid type A receptor: lack of effect of responses to ethanol, pentobarbital, and general anesthetics. *Mol Pharmacol* 1997;51:588–596. [PubMed: 9106623]
- Homanics GE, Le NQ, Kist F, Mihalek R, Hart AR, Quinlan JJ. Ethanol tolerance and withdrawal responses in GABA_A receptor alpha 6 subunit null allele mice and in inbred C57BL/6J and strain 129/SvJ mice. *Alcohol Clin Exp Res* 1998;22:259–265. [PubMed: 9514316]
- Homanics GE, Harrison NL, Quinlan JJ, Krasowski MD, Rick CEM, de Blas AL, Mehta AK, Kist F, Mihalek RM, Aul JJ, Firestone LL. Normal electrophysiological and behavioral responses to ethanol in mice lacking the long splice variant of the γ 2 subunit of the γ -aminobutyrate type A receptor. *Neuropharmacology* 1999a;38:253–265. [PubMed: 10218866]
- Homanics GE, Quinlan JJ, Firestone LL. Pharmacologic and behavioral responses of inbred C57BL/6J and strain 129/SvJ mouse lines. *Pharmacol Biochem Behav* 1999b;63:21–26. [PubMed: 10340519]
- Homanics GE, Elsen FP, Ying S-W, Jenkins AE, Ferguson C, Sloat B, Yuditskaya S, Goldstein PA, Kralic JE, Morrow AL, Harrison NL. A gain-of-function mutation in the GABA_A receptor produces synaptic and behavioral abnormalities in the mouse. *Genes Brain Behav* 2005;4:10–19. [PubMed: 15660664]
- Hood HM, Buck KJ. Allelic variation in the GABA_A receptor γ 2 subunit is associated with genetic susceptibility to ethanol-induced motor incoordination and hypothermia, conditioned taste aversion, and withdrawal in BXD/Ty recombinant inbred mice. *Alcohol Clin Exp Res* 2000;24:1327–1334. [PubMed: 11003197]
- Hsu FC, Waldeck R, Faber DS, Smith SS. Neurosteroid effects on GABAergic synaptic plasticity in hippocampus. *J Neurophysiol* 2003;89:1929–1940. [PubMed: 12612040]
- Hu X, Oroszi G, Chun J, Smith TL, Goldman D, Schuckit MA. An expanded evaluation of the relationship of four alleles to the level of response to alcohol and the alcoholism risk. *Alcohol Clin Exp Res* 2005;29:8–16. [PubMed: 15654286]
- Huang RQ, Dillon GH. Maintenance of recombinant type A gamma-aminobutyric acid receptor function: role of protein tyrosine phosphorylation and calcineurin. *J Pharmacol Exp Ther* 1998;286:243–255. [PubMed: 9655866]

- Ittiwut C, Listman J, Mutirangura A, Malison R, Covault J, Kranzler HR, Sughondhabirom A, Thavichachart N, Gelernter J. Interpopulation linkage disequilibrium patterns of GABRA2 and GABRG1 genes at the GABA cluster locus on human chromosome 4. *Genomics* 2008;91:61–69. [PubMed: 17976953]
- Ives JH, Drewery DL, Thompson CL. Differential cell surface expression of GABA_A receptor α 1, α 6, β 2 and β 3 subunits in cultured mouse cerebellar granule cells influence of cAMP-activated signalling. *J Neurochem* 2002;80:317–327. [PubMed: 11902122]
- Iwata N, Cowley DS, Radel M, Roy-Byrne PP, Goldman D. Relationship between a GABA_A α 6 Pro385Ser substitution and benzodiazepine sensitivity. *Am J Psychiatry* 1999;156:1447–1449. [PubMed: 10484961]
- Jenkins SA, Simmons DD. GABAergic neurons in the lateral superior olive of the hamster are distinguished by differential expression of gad isoforms during development. *Brain Res* 2006;1111:12–25. [PubMed: 16919247]
- Jia F, Chandra D, Homanics GE, Harrison NL. Ethanol modulates synaptic and extrasynaptic GABA_A receptors in the thalamus. *J Pharmacol Exp Ther* 2008a;326:475–482. [PubMed: 18477766]
- Jia F, Yue M, Chandra D, Keramidas A, Goldstein PA, Homanics GE, Harrison NL. Taurine is a potent activator of extra-synaptic GABA_A receptors in the thalamus. *J Neurosci* 2008b;28:106–115. [PubMed: 18171928]
- Jones A, Korpi ER, McKernan RM, Pelz R, Nusser Z, Mäkelä R, Mellor JR, Pollard S, Bahn S, Stephenson FA, Randall AD, Sieghart W, Somogyi P, Smith AJH, Wisden W. Ligand-gated ion channel subunit partnerships: GABA_A receptor α 6 subunit gene inactivation inhibits δ subunit expression. *J Neurosci* 1997;17:1350–1362.
- June HL Sr, Foster KL, Eiler WJ 2nd, Goergen J, Cook JB, Johnson N, Mensah-Zoe B, Simmons JO, June HL Jr, Yin W, Cook JM, Homanics GE. Dopamine and benzodiazepine-dependent mechanisms regulate the EtOH-enhanced locomotor stimulation in the GABA_A alpha1 subunit null mutant mice. *Neuropsychopharmacology* 2007;32:137–152.
- Jurd R, Arras M, Lambert S, Drexler B, Sieghart R, Crestani F, Zaugg M, Vogt KE, Ledermann B, Antkowiak B, Rudolph U. General anesthetic actions in vivo strongly attenuated by a point mutation in the GABA_A receptor β 3 subunit. *FASEB J* 2003;17:250–252. [PubMed: 12475885]
- Kalant H, LeBlanc AE, Gibbins RJ. Tolerance to, and dependence on, some non-opiate psychotropic drugs. *Pharmacol Rev* 1971;23:135–191. [PubMed: 4398655]
- Kaminski RM, Marini H, Kim W, Rogawski MA. Anticonvulsant activity of androsterone and etiocholanolone. *Epilepsia* 2005;46:819–827. [PubMed: 15946323]
- Kaminski RM, Marini H, Ortinski PI, Vicini S, Rogawski MA. The pheromone androstenol (5 alpha-androst-16-en-3 alphaol) is a neurosteroid positive modulator of GABA_A receptors. *J Pharmacol Exp Ther* 2006;317:694–703. [PubMed: 16415088]
- Kanaumi T, Takashima S, Iwasaki H, Mitsudome A, Hirose S. Developmental changes in the expression of GABA_A receptor alpha 1 and gamma 2 subunits in human temporal lobe, hippocampus and basal ganglia: an implication for consideration on age-related epilepsy. *Epilepsy Res* 2006;71:47–53. [PubMed: 16829043]
- Kang M, Spigelman I, Sapp DW, Olsen RW. Persistent reduction of GABA_A receptor-mediated inhibition in rat hippocampus after chronic intermittent ethanol treatment. *Brain Res* 1996;709:221–228. [PubMed: 8833758]
- Kaur KH, Baur R, Sigel E. Unanticipated structural and functional properties of delta subunit containing GABA_A receptors. *J Biol Chem* 2009;284:7889–7896. [PubMed: 19141615]
- Keir WJ, Morrow AL. Differential expression of GABA_A receptor subunit mRNAs in ethanol-naive withdrawal seizure resistant (WSR) vs. withdrawal seizure prone (WSP) mouse brain. *Brain Res Mol Brain Res* 1994;25:200–208. [PubMed: 7808218]
- Kellenberger S, Malherbe P, Sigel E. Function of the α 1 β 2 γ 2s γ -aminobutyric acid type_A receptor is modulated by protein kinase C via multiple phosphorylation sites. *J Biol Chem* 1992;267:25660–25663. [PubMed: 1334482]
- Kelly SJ, Bonthius DJ, West JR. Developmental changes in alcohol pharmacokinetics in rats. *Alcohol Clin Exp Res* 1987;11:281–286. [PubMed: 3307494]

- Khisti RT, VanDoren MJ, O'Buckley TK, Morrow AL. Neuroactive steroid 3α -hydroxy- 5α -pregnan-20-one modulates ethanol-induced loss of righting reflex in rats. *Brain Res* 2003;980:255–265. [PubMed: 12867266]
- Khisti RT, VanDoren MJ, Matthews DB, Morrow AL. Ethanol-induced elevation of 3α -hydroxy- 5α -pregnan-20-one does not modulate motor incoordination in rats. *Alcohol Clin Exp Res* 2004;28:1249–1256. [PubMed: 15318125]
- Khisti RT, Boyd KN, Kumar S, Morrow AL. Systemic ethanol administration elevates deoxycorticosterone levels and chronic ethanol exposure attenuates this response. *Brain Res* 2005;1049:104–111. [PubMed: 15936001]
- Kiefer F, Jahn H, Schick M, Wiedemann K. Alcohol intake, tumour necrosis factor- α , leptin and craving: factors of a possibly vicious circle? *Alcohol* 2002;37:401–404. [PubMed: 12107045]
- Kim JH, Lee S, Chow J, Lau J, Tsang A, Choi J, Griffiths SM. Prevalence and the factors associated with binge drinking, alcohol abuse, and alcohol dependence: a population-based study of Chinese adults in Hong Kong. *Alcohol* 2008;43:360–370. [PubMed: 18230698]
- Kittler JT, Chen G, Honing S, Bogdanov Y, McAinsh K, Arancibia-Carcamo IL, Jovanovic JN, Pangalos MN, Haucke V, Yan Z, Moss SJ. Phospho-dependent binding of the clathrin AP2 adaptor complex to GABA_A receptors regulates the efficacy of inhibitory synaptic transmission. *Proc Natl Acad Sci U S A* 2005;102:14871–14876. [PubMed: 16192353]
- Kittler JT, Chen G, Kukhtina V, Vahedi-Faridi A, Gu Z, Tretter V, Smith KR, McAinsh K, Arancibia-Carcamo IL, Saenger W, Haucke V, Yan Z, Moss SJ. Regulation of synaptic inhibition by phospho-dependent binding of the AP2 complex to a YECL motif in the GABA_A receptor γ 2 subunit. *Proc Natl Acad Sci U S A* 2008;105:3616–3621. [PubMed: 18305175]
- Korneyev AY, Costa E, Guidotti A. During anesthetic-induced activation of hypothalamic pituitary adrenal axis, blood borne steroids fail to contribute to the anesthetic effect. *Neuroendocrinology* 1993;57:559–565. [PubMed: 8391666]
- Korpi ER, Debus F, Linden AM, Malecot C, Leppa E, Vekovischeva O, Rabe H, Bohme I, Aller MI, Wisden W, Luddens H. Does ethanol act preferentially via selected brain GABA_A receptor subtypes? The current evidence is ambiguous. *Alcohol* 2007;41:163–176. [PubMed: 17591542]
- Kralic JE, Korpi ER, O'Buckley TK, Homanics GE, Morrow AL. Molecular and pharmacological characterization of GABA_A receptor α 1 subunit knockout mice. *J Pharmacol Exp Ther* 2002a;302:1037–1045. [PubMed: 12183661]
- Kralic JE, O'Buckley TK, Khisti RT, Hodge CW, Homanics GE, Morrow AL. GABA_A receptor α 1 subunit deletion alters receptor subtype assembly, pharmacological and behavioral responses to benzodiazepines and zolpidem. *Neuropharmacology* 2002b;43:685–694. [PubMed: 12367614]
- Kralic JE, Wheeler M, Renzi K, Ferguson C, O'Buckley TK, Grobin AC, Morrow AL, Homanics GE. Deletion of GABA_A receptor α 1 subunit-containing receptors alters responses to ethanol and other anesthetics. *J Pharmacol Exp Ther* 2003;305:600–607. [PubMed: 12606632]
- Krishek BJ, Xie X, Blackstone C, Haganir RL, Moss SJ, Smart TG. Regulation of GABA_A receptor function by protein kinase C phosphorylation. *Neuron* 1994;12:1081–1095. [PubMed: 8185945]
- Kumar S, Sieghart W, Morrow AL. Association of protein kinase C with GABA_A receptors containing α 1 and α 4 subunits in the cerebral cortex: selective effects of chronic ethanol consumption. *J Neurochem* 2002;82:110–117. [PubMed: 12091471]
- Kumar S, Kralic JE, O'Buckley TK, Grobin AC, Morrow AL. Chronic ethanol consumption enhances internalization of α 1 subunit-containing GABA_A receptors in cerebral cortex. *J Neurochem* 2003;86:700–708. [PubMed: 12859683]
- Kumar S, Fleming R, Morrow AL. Ethanol regulation of γ -aminobutyric acid_A receptors: genomic and nongenomic mechanisms. *Pharmacol Ther* 2004;101:211–226. [PubMed: 15031000]
- Kumar S, Khisti RT, Morrow AL. Regulation of native GABA_A receptors by PKC and protein phosphatase activity. *Psychopharmacology* 2005;183:241–247. [PubMed: 16220337]
- Kumar S, Lane BM, Morrow AL. Differential effects of systemic ethanol administration on PKC ϵ , γ and β isoform expression, translocation to membranes and target phosphorylation: reversal by chronic ethanol exposure. *J Pharmacol Exp Ther* 2006;319:1366–1375. [PubMed: 16997974]

- Kumar S, Ren Q, Beckley JH, Morrow AL. Effects of ethanol on PKC and PKA expression in rat cerebral cortex contribute to the regulation of GABA_A α 1 subunit receptor surface expression. *Alcohol Clin Exp Res* 2008;32 21A Supplement to vol. 32.
- Kurtz DL, Stewart RB, Zweifel M, Li TK, Froehlich JC. Genetic differences in tolerance and sensitization to the sedative/hypnotic effects of alcohol. *Pharmacol Biochem Behav* 1996;53:585–591. [PubMed: 8866959]
- Lai CC, Kuo TI, Lin HH. The role of protein kinase A in acute ethanol-induced neurobehavioral actions in rats. *Anesth Analg* 2007;105:89–96. [PubMed: 17578962]
- Lancaster FE, Brown TD, Coker KL, Elliott JA, Wren SB. Sex differences in alcohol preference and drinking patterns emerge during early postpubertal period in Sprague-Dawley rats. *Alcohol Clin Exp Res* 1996;20:1043–1049. [PubMed: 8892526]
- Land C, Spear NE. Ethanol impairs memory of a single discrimination in adolescent rats at doses that leave adult memory unaffected. *Neurobiol Learn Mem* 2004;81:75–81. [PubMed: 14670361]
- Lappalainen J, Krupitsky E, Remizov M, Pchelina S, Taraskina A, Zvartau E, Somberg LK, Covault J, Kranzler HR, Krystal JH, Gelernter J. Association between alcoholism and γ -amino butyric acid α 2 receptor subtype in a Russian population. *Alcohol Clin Exp Res* 2005;29:493–498. [PubMed: 15834213]
- Laurie DJ, Wisden W, Seeburg PH. The distribution of thirteen GABA_A receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci* 1992;12:4151–4172. [PubMed: 1331359]
- Laviola G, Macri S, Morley-Fletcher S, Adriani W. Risk-taking behavior in adolescent mice: psychobiological determinants and early epigenetic influence. *Neurosci Biobehav Rev* 2003;27:19–31. [PubMed: 12732220]
- Le AD, Khanna JM, Kalant H, Grossi F. Tolerance to and cross-tolerance among ethanol, pentobarbital and chlordiazepoxide. *Pharmacol Biochem Behav* 1986;24:93–98. [PubMed: 3945670]
- Lee S, Schmidt ED, Tilders FJ, Rivier C. Effect of repeated exposure to alcohol on the response of the hypothalamic–pituitary–adrenal axis of the rat: I. Role of changes in hypothalamic neuronal activity. *Alcohol Clin Exp Res* 2001;25:98–105. [PubMed: 11198721]
- Leidenheimer NJ, McQuiklin SJ, Hahner LD, Whiting P, Harris RA. Activation of protein kinase C selectively inhibits the γ -aminobutyric acid_A receptor: role of desensitization. *Mol Pharmacol* 1992;41:1116–1123. [PubMed: 1319547]
- Lewohl JM, Crane DI, Dodd PR. Expression of the α 1, α 2, and α 3 isoforms of the GABA_A receptor in human alcoholic brain. *Brain Res* 1997;751:102–112. [PubMed: 9098573]
- Lewohl JM, Huygens F, Crane DI, Dodd PR. GABA_A receptor alpha-subunit proteins in human chronic alcoholics. *J Neurochem* 2001;78:424–434. [PubMed: 11483645]
- Liang J, Cagetti E, Olsen RW, Spigelman I. Altered pharmacology of synaptic and extrasynaptic GABA_A receptors on CA1 hippocampal neurons is consistent with subunit changes in a model of alcohol withdrawal and dependence. *J Pharmacol Exp Ther* 2004;310:1234–1245. [PubMed: 15126642]
- Liang J, Zhang N, Cagetti E, Houser CR, Olsen RW, Spigelman I. Chronic intermittent ethanol-induced switch of ethanol actions from extrasynaptic to synaptic hippocampal GABA_A receptors. *J Neurosci* 2006;26:1749–1758. [PubMed: 16467523]
- Liang J, Suryanarayanan A, Abriam A, Snyder B, Olsen RW, Spigelman I. Mechanisms of reversible GABA_A receptor plasticity after ethanol intoxication. *J Neurosci* 2007;27:12367–12377. [PubMed: 17989301]
- Liang J, Suryanarayanan A, Chandra D, Homanics GE, Olsen RW, Spigelman I. Functional consequences of GABA_A receptor α 4 subunit deletion on synaptic and extrasynaptic currents in mouse dentate granule cells. *Alcohol Clin Exp Res* 2008;32:19–26. [PubMed: 18070250]
- Lim WY, Fong CW, Chan JM, Heng D, Bhalla V, Chew SK. Trends in alcohol consumption in Singapore 1992–2004. *Alcohol Alcohol* 2007;42:354–361. [PubMed: 17496306]
- Lingford-Hughes AR, Acton PD, Gacinovic S, Suckling J, Busatto GF, Boddington SJ, Bullmore E, Woodruff PW, Costa DC, Pilowsky LS, Ell PJ, Marshall EJ, Kerwin RW. Reduced levels of GABA-benzodiazepine receptor in alcohol dependency in the absence of grey matter atrophy. *Br J Psychiatry* 1998;173:116–122. [PubMed: 9850223]

- Lingford-Hughes AR, Acton PD, Gacinovic S, Boddington SJ, Costa DC, Pilowsky LS, Ell PJ, Marshall EJ, Kerwin RW. Levels of gamma-aminobutyric acid-benzodiazepine receptors in abstinent, alcohol-dependent women: preliminary findings from an 123I-iomazenil single photon emission tomography study. *Alcohol Clin Exp Res* 2000;24:1449–1455. [PubMed: 11003213]
- Lingford-Hughes AR, Wilson SJ, Cunningham VJ, Feeney A, Stevenson B, Brooks DJ, Nutt DJ. GABA-benzodiazepine receptor function in alcohol dependence: a combined 11C-flumazenil PET and pharmacodynamic study. *Psychopharmacology (Berl)* 2005;180:595–606. [PubMed: 15864554]
- Lodish, H.; Baltimore, D.; Berk, A.; Zipursky, SL.; Matsui, P.; Darnell, J. *Molecular Cell Biology* (Scientific American Books). Freeman; New York: 1996. Synthesis and sorting of plasma membrane, secretory, and lysosomal proteins; p. 711-715.
- Loh EW, Smith I, Murray R, McLaughlin M, McNulty S, Ball D. Association between variants at the GABA_Aβ2, GABA_Aα6 and GABA_Aγ2 gene cluster and alcohol dependence in a Scottish population. *Mol Psychiatry* 1999;4:539–544. [PubMed: 10578235]
- Loh EW, Higuchi S, Matsushita S, Murray R, Chen CK, Ball D. Association analysis of the GABA_A receptor subunit genes cluster on 5q33–34 and alcohol dependence in a Japanese population. *Mol Psychiatry* 2000;5:301–307. [PubMed: 10889533]
- Long JC, Knowler WC, Hanson RL, Robin RW, Urbanek M, Moore E, Bennett PH, Goldman D. Evidence for genetic linkage to alcohol dependence on chromosomes 4 and 11 from an autosome-wide scan in an American Indian population. *Am J Med Genet* 1998;81:216–221. [PubMed: 9603607]
- Lovinger DM. 5-HT₃ receptors and the neural actions of alcohols: an increasingly exciting topic. *Neurochem Int* 1999;35:125–130. [PubMed: 10405996]
- Maas JW Jr, Vogt SK, Chan GC, Pineda VV, Storm DR, Muglia LJ. Calcium-stimulated adenylyl cyclases are critical modulators of neuronal ethanol sensitivity. *J Neurosci* 2005;25:4118–4126. [PubMed: 15843614]
- Macdonald RL. Ethanol, γ-aminobutyrate type A receptors, and protein kinase C phosphorylation. *Proc Natl Acad Sci U S A* 1995;92:3633–3635. [PubMed: 7731956]
- MacLennan AJ, Lee N, Walker DW. Chronic ethanol administration decreases brain-derived neurotrophic factor gene expression in the rat hippocampus. *Neurosci Lett* 1995;197:105–108. [PubMed: 8552271]
- Mahmoudi M, Kang MH, Tillakaratne N, Tobin AJ, Olsen RW. Chronic intermittent ethanol treatment in rats increases GABA_A receptor α4-subunit expression: possible relevance to alcohol dependence. *J Neurochem* 1997;68:2485–2492. [PubMed: 9166743]
- Majewska MD, Harrison NL, Schwartz RD, Barker JL, Paul SM. Steroid hormone metabolites are barbiturate-like modulators of the GABA receptor. *Science* 1986;232:1004–1007. [PubMed: 2422758]
- Markwiese BJ, Acheson SK, Levin ED, Wilson WA, Swartzwelder HS. Differential effects of ethanol on memory in adolescent and adult rats. *Alcohol Clin Exp Res* 1998;22:416–421. [PubMed: 9581648]
- Marutha Ravindran CR, Ticku MK. Tyrosine kinase phosphorylation of GABA_A receptor subunits following chronic ethanol exposure of cultured cortical neurons of mice. *Brain Res* 2006;1086:35–41. [PubMed: 16580644]
- Marutha Ravindran CR, Mehta AK, Ticku MK. Effect of chronic administration of ethanol on the regulation of tyrosine kinase phosphorylation of the GABA_A receptor subunits in the rat brain. *Neurochem Res* 2007;32:1179–1187. [PubMed: 17404840]
- Matthews DB, Silvers JR. The use of acute ethanol administration as a tool to investigate multiple memory systems. *Neurobiol Learn Mem* 2004;82:299–308. [PubMed: 15464411]
- Matthews DB, Devaud LL, Fritschy J-M, Sieghart W, Morrow AL. Differential regulation of GABA_A receptor gene expression by ethanol in the rat hippocampus vs. cerebral cortex. *J Neurochem* 1998;70:1160–1166. [PubMed: 9489737]
- Matthews DB, Criswell HE, Grobin AC, Morrow AL. Chronic ethanol consumption alters recovery of spontaneously active medial septal/diagonal band of Broca neurons from GABA-microiontophoresis. *Alcohol Clin Exp Res* 2000a;24:1427–1432. [PubMed: 11003210]

- Matthews DB, Kralic JE, Devaud LL, Fritschy J-M, Morrow AL. Chronic blockade of *N*-methyl-D-aspartate receptors alters γ -aminobutyric acid type A receptor gene expression and function in the rat. *J Neurochem* 2000b;74:1522–1528. [PubMed: 10737609]
- Matthews DB, Morrow AL, Tokunaga S, McDaniel JR. Acute ethanol administration and acute allopregnanolone administration impair spatial memory in the Morris water task. *Alcohol Clin Exp Res* 2002;26:1747–1751. [PubMed: 12436065]
- McCown TJ, Breese GR. Multiple withdrawals from chronic ethanol “kindle” inferior collicular seizure activity: evidence for kindling of seizures associated with alcoholism. *Alcohol Clin Exp Res* 1990;14:394–399. [PubMed: 2378423]
- McDonald BJ, Moss SJ. Conserved phosphorylation of the intracellular domains of GABA_A receptor β 2 and β 3 subunits by cAMP-dependent protein kinase, cGMP-dependent protein kinase, protein kinase C and Ca²⁺/calmodulin type II-dependent protein kinase. *Neuropharmacology* 1997;36:1377–1385. [PubMed: 9423925]
- McKernan RM, Whiting PJ. Which GABA_A-receptor subtypes really occur in the brain? *Trends Neurosci* 1996;19:139–143. [PubMed: 8658597]
- Mehta AK, Marutha Ravindran CR, Ticku MK. Low concentrations of ethanol do not affect radioligand binding to the delta-subunit-containing GABA_A receptors in the rat brain. *Brain Res* 2007;1165:15–20. [PubMed: 17662260]
- Mhatre MC, Mehta AK, Ticku MK. Chronic ethanol administration increases the binding of the benzodiazepine inverse agonist and alcohol antagonist [³H]Ro15–4513 in rat brain. *Eur J Pharmacol* 1988;153:141–145. [PubMed: 2850926]
- Mhatre MC, Ticku MK. Chronic ethanol administration alters γ -aminobutyric acid A receptor gene expression. *Mol Pharmacol* 1992;42:415–422. [PubMed: 1383684]
- Mhatre MC, Ticku MK. Chronic ethanol treatment upregulates GABA_A receptor β subunit expression. *Brain Res Mol Brain Res* 1994a;23:246–252. [PubMed: 8057781]
- Mhatre MC, Ticku MK. Chronic GABA treatment down-regulates the GABA_A receptor α 2 and α 3 subunit mRNAs as well as polypeptide expression in primary cultured cerebral cortical neurons. *Brain Res Mol Brain Res* 1994b;24:159–165. [PubMed: 7968353]
- Mhatre MC, Pena G, Sieghart W, Ticku MK. Antibodies specific for GABA_A receptor alpha subunits reveal that chronic alcohol treatment down-regulates alpha-subunit expression in rat brain regions. *J Neurochem* 1993;61:1620–1625. [PubMed: 8228981]
- Mihalek RM, Bowers BJ, Wehner JM, Kralic JE, VanDoren MJ, Morrow AL, Homanics GE. GABA_A-receptor δ subunit knockout mice have multiple defects in behavioral responses to ethanol. *Alcohol Clin Exp Res* 2001;25:1708–1718. [PubMed: 11781502]
- Mihic SJ, Ye Q, Wick MJ, Koltchine VV, Krasowski MA, Finn SE, Mascia MP, Valenzuela CF, Hanson KK, Greenblatt EP, Harris RA, Harrison NL. Sites of alcohol and volatile anaesthetic action on GABA_A and glycine receptors. *Nature* 1997;389:385–389. [PubMed: 9311780]
- Mitsuyama H, Little KY, Sieghart W, Devaud LL, Morrow AL. GABA_A receptor α 1, α 4 and β 3 subunit mRNA and protein expression in frontal cortex of human alcoholics. *Alcohol Clin Exp Res* 1998;22(4):815–822. [PubMed: 9660306]
- Mody I. Distinguishing between GABA_A receptors responsible for tonic and phasic conductances. *Neurochem Res* 2001;26:907–913. [PubMed: 11699942]
- Mohler, H.; Fritschy, J-M.; Lüscher, B.; Rudolph, U.; Benson, J.; Benke, D. The GABA_A receptors: from subunits to diverse functions. In: Narahashi, T., editor. *Ion channels*. Vol. 4. Plenum; New York: 1996. p. 89-111.
- Montpied P, Ginns EI, Martin BM, Roca D, Farb DH, Paul SM. Gamma-aminobutyric acid (GABA) induces a receptor-mediated reduction in GABA_A receptor α subunit messenger RNAs in embryonic chick neurons in culture. *J Biol Chem* 1991a;266:6011–6014. [PubMed: 1848843]
- Montpied P, Morrow AL, Karanian JW, Ginns EI, Martin BM, Paul SM. Prolonged ethanol inhalation decreases gamma-aminobutyric acid_A receptor α subunit mRNAs in the rat cerebral cortex. *Mol Pharmacol* 1991b;39:157–163. [PubMed: 1705000]
- Moran MH, Goldberg N, Smith SS. Progesterone withdrawal II: Insensitivity to the sedative effects of a benzodiazepine. *Brain Res* 1998;807:91–100. [PubMed: 9757006]

- Morrow AL, Suzdak PD, Paul SM. Steroid hormone metabolites potentiate GABA receptor-mediated chloride ion flux with nanomolar potency. *Eur J Pharmacol* 1987;142:483–485. [PubMed: 2828079]
- Morrow AL, Suzdak PD, Karanian JW, Paul SM. Chronic ethanol administration alters γ -aminobutyric acid, pentobarbital and ethanol-mediated $^{36}\text{Cl}^-$ uptake in cerebral cortical synapto-neurosomes. *J Pharmacol Exp Ther* 1988;246:158–164. [PubMed: 2839659]
- Morrow AL, Herbert JS, Montpied P. Differential effects of chronic ethanol administration on GABA_A receptor $\alpha 1$ and $\alpha 6$ subunit mRNA levels in rat cerebellum. *Mol Cell Neurosci* 1992;3:251–258. [PubMed: 19912867]
- Morrow AL, VanDoren MJ, Devaud LL. Effects of progester-one or neuroactive steroid? *Nature* 1998;395:652–653. [PubMed: 9790185]
- Morrow AL, Janis GC, VanDoren MJ, Matthews DB, Samson HH, Janak PH, Grant KA. Neurosteroids mediate pharmacological effects of ethanol: a new mechanism of ethanol action? *Alcohol Clin Exp Res* 1999;23:1933–1940. [PubMed: 10630613]
- Morrow AL, VanDoren MJ, Penland SN, Matthews DB. The role of GABAergic neuroactive steroids in ethanol action, tolerance and dependence. *Brain Res Rev* 2001;37:98–109. [PubMed: 11744078]
- Morrow AL, Porcu P, Boyd KN, Grant KA. Hypothalamic–pituitary–adrenal axis modulation of GABAergic neuroactive steroids influences ethanol sensitivity and drinking behavior. *Dialogues Clin Neurosci* 2006;8:463–477. [PubMed: 17290803]
- Moss SJ, Smart TG. Constructing inhibitory synapses. *Nat Rev Neurosci* 2001;2:240–250. [PubMed: 11283747]
- Moss SJ, Smart TG, Blackstone CD, Haganir RL. Functional modulation of GABA_A receptors by cAMP-dependent protein phosphorylation. *Science* 1992;257:661–665. [PubMed: 1323140]
- Moy SS, Knapp DJ, Criswell HE, Breese GR. Flumazenil blockade of anxiety following ethanol withdrawal in rats. *Psychopharmacology* 1997;131:354–360. [PubMed: 9226737]
- Moy SS, Knapp DJ, Duncan GE, Breese GR. Enhanced ultrasonic vocalization and Fos protein expression following ethanol withdrawal: effects of flumazenil. *Psychopharmacology* 2000;152:208–215. [PubMed: 11057525]
- Nance DM. The developmental and neural determinants of the effects of estrogen on feeding behavior in the rat: a theoretical perspective. *Neurosci Biobehav Rev* 1983;7:189–211. [PubMed: 6348605]
- Newton PM, Messing RO. Intracellular signaling pathways that regulate behavioral responses to ethanol. *Pharmacol Ther* 2006;109:227–237. [PubMed: 16102840]
- Niimura M, Moussa R, Bissoon N, Ikeda-Douglas C, Milgram NW, Gurd JW. Changes in phosphorylation of the NMDA receptor in the rat hippocampus induced by status epilepticus. *J Neurochem* 2005;92:1377–1385. [PubMed: 15748156]
- O'Dell LE, Alomary AA, Vallee M, Koob GF, Fitzgerald RL, Purdy RH. Ethanol-induced increases in neuroactive steroids in the rat brain and plasma are absent in adrenalectomized and gonadectomized rats. *Eur J Pharmacol* 2004;484:241–247. [PubMed: 14744609]
- Obrietan K, Gao XB, Van Den Pol AN. Excitatory actions of GABA increase BDNF expression via a MAPK-CREB-dependent mechanism—a positive feedback circuit in developing neurons. *J Neurophysiol* 2002;88:1005–1015. [PubMed: 12163549]
- Oh S, Jang CG, Ma T, Ho IK. Activation of protein kinase C by phorbol dibutyrate modulates GABA_A receptor binding in rat brain slices. *Brain Res* 1999;850:158–165. [PubMed: 10629760]
- Olive MF, Hodge CW. Co-localization of PKC ϵ with various GABA_A receptor subunits in the mouse limbic system. *Neuro-report* 2000;11:683–687.
- Olsen RW, Sieghart W. International Union of Pharmacology. LXX. Subtypes of gamma-aminobutyric acid(A) receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev* 2008;60:243–260. [PubMed: 18790874]
- Olsen RW, Sieghart W. GABA_A receptors: subtypes provide diversity of function and pharmacology. *Neuropharmacology* 2009;56:141–148. [PubMed: 18760291]
- Pandey SC. Neuronal signaling systems and ethanol dependence. *Mol Neurobiol* 1998;17:1–15. [PubMed: 9887443]
- Pandey SC, Roy A, Zhang H. The decreased phosphorylation of cyclic adenosine monophosphate (cAMP) response element binding (CREB) protein in the central amygdala acts as a molecular

- substrate for anxiety related to ethanol withdrawal in rats. *Alcohol Clin Exp Res* 2003;27:396–409. [PubMed: 12658105]
- Papadeas S, Grobin AC, Morrow AL. Chronic ethanol consumption differentially alters GABA_A receptor $\alpha 1$ and $\alpha 4$ subunit peptide expressions and GABA_A receptor-mediated ³⁶CL⁻ uptake in mesocorticolimbic regions of rat brain. *Alcohol Clin Exp Res* 2001;25:1270–1275. [PubMed: 11584145]
- Park-Chung M, Malayev A, Purdy RH, Gibbs TT, Farb DH. Sulfated and unsulfated steroids modulate gamma-aminobutyric acid_A receptor function through distinct sites. *Brain Research* 1999;830:72–87. [PubMed: 10350561]
- Park SK, Sedore SA, Cronmiller C, Hirsh J. Type II cAMP-dependent protein kinase-deficient *Drosophila* are viable but show developmental, circadian, and drug response phenotypes. *J Biol Chem* 2000;275:20588–20596. [PubMed: 10781603]
- Parsian A, Zhang ZH. Human chromosomes 11p15 and 4p12 and alcohol dependence: possible association with the GABRB1 gene. *Am J Med Genet* 1999;88:533–538. [PubMed: 10490712]
- Petrie J, Sapp DW, Tyndale RF, Park MK, Fanselow M, Olsen RW. Altered GABA_A receptor subunit and splice variant expression in rats treated with chronic intermittent ethanol. *Alcohol Clin Exp Res* 2001;25:819–828. [PubMed: 11410716]
- Pierucci-Lagha A, Covault J, Feinn R, Nellisery M, Hernandez-Avila C, Oncken C, Morrow AL, Kranzler HR. GABRA2 alleles moderate the subjective effects of alcohol, which are attenuated by finasteride. *Neuropsychopharmacology* 2005;30:1193–1203. [PubMed: 15702134]
- Pignataro L, Miller AN, Ma L, Midha S, Protiva P, Herrera DG, Harrison NL. Alcohol regulates gene expression in neurons via activation of heat shock factor 1. *J Neurosci* 2007;27:12957–12966. [PubMed: 18032669]
- Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G. GABA_A receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* 2000;101:815–850. [PubMed: 11113332]
- Platt DM, Duggan A, Spealman RD, Cook JM, Li X, Yin W, Rowlett JK. Contribution of $\alpha 1$ GABA_A and $\alpha 5$ GABA_A receptor subtypes to the discriminative stimulus effects of ethanol in squirrel monkeys. *J Pharmacol Exp Ther* 2005;313:658–667. [PubMed: 15650112]
- Poisbeau P, Cheney MC, Browning MD, Mody I. Modulation of synaptic GABA_A receptor function by PKA and PKC in adult hippocampal neurons. *J Neurosci* 1999;19:674–683. [PubMed: 9880588]
- Porcu P, Sogliano C, Ibbá C, Piredda M, Tocco S, Marra C, Purdy RH, Biggio G, Concas A. Failure of γ -hydroxybutyric acid both to increase neuroactive steroid concentrations in adrenalectomized-orchietomized rats and to induce tolerance to its steroidogenic effect in intact animals. *Brain Res* 2004;1012:160–168. [PubMed: 15158173]
- Porter NM, Twyman RE, Uhler MD, Macdonald RL. Cyclic AMP-dependent protein kinase decreases GABA_A receptor current in mouse spinal neurons. *Neuron* 1990;5:789–796. [PubMed: 2176510]
- Powell K. How does the teenage brain work? *Nature* 2006;442:865–867. [PubMed: 16929274]
- Proctor WR, Poelchen W, Bowers BJ, Wehner JM, Messing RO, Dunwiddie TV. Ethanol differentially enhances hippo-campal GABA A receptor-mediated responses in protein kinase C gamma (PKC gamma) and PKC epsilon null mice. *J Pharmacol Exp Ther* 2003;305:264–270. [PubMed: 12649378]
- Proctor WR, Wu PH, Bennett B, Johnson TE. Differential effects of ethanol on gamma-aminobutyric acid-A receptor-mediated synaptic currents in congenic strains of inbred long and short-sleep mice. *Alcohol Clin Exp Res* 2004;28:1277–1283. [PubMed: 15365296]
- Puia G, Ducic I, Vicini S, Costa E. Does neurosteroid modulatory efficacy depend on GABA_A receptor subunit composition? *Receptors Channels* 1993;1:135–142. [PubMed: 8081717]
- Pyapali GK, Turner DA, Wilson WA, Swartzwelder HS. Age and dose-dependent effects of ethanol on the induction of hippocampal long-term potentiation. *Alcohol* 1999;19:107–111. [PubMed: 10548153]
- Qi ZH, Song M, Wallace MJ, Wang D, Newton PM, McMahan T, Chou WH, Zhang C, Shokat KM, Messing RO. Protein kinase C ϵ regulates γ -aminobutyrate type A receptor sensitivity to ethanol and benzodiazepines through phosphorylation of $\gamma 2$ subunits. *J Biol Chem* 2007;282:33052–33063. [PubMed: 17875639]

- Quertemont E, Lallemand F, Colombo G, De Witte P. Taurine and ethanol preference: a microdialysis study using Sardinian alcohol-preferring and non-preferring rats. *Eur Neuropsychopharmacol* 2000;10:377–383.
- Radcliffe RA, Floyd KL, Drahnak JA, Deitrich RA. Genetic dissociation between ethanol sensitivity and rapid tolerance in mouse and rat strains selectively bred for differential ethanol sensitivity. *Alcohol Clin Exp Res* 2005;29:1580–1589. [PubMed: 16205358]
- Radel M, Vallejo RL, Iwata N, Aragon R, Long JC, Virkkunen M, Goldman D. Haplotype-based localization of an alcohol dependence gene to the 5q34 γ -aminobutyric acid type A gene cluster. *Arch Gen Psychiatry* 2005;62:47–55. [PubMed: 15630072]
- Rajendran P, Spear LP. The effects of ethanol on spatial and nonspatial memory in adolescent and adult rats studied using an appetitive paradigm. *Ann NY Acad Sci* 2004;1021:441–444. [PubMed: 15251925]
- Rakhade SN, Zhou C, Aujla PK, Fishman R, Sucher NJ, Jensen FE. Early alterations of AMPA receptors mediate synaptic potentiation induced by neonatal seizures. *J Neurosci* 2008;28:7979–7990. [PubMed: 18685023]
- Reich T. A genomic survey of alcohol dependence and related phenotypes: results from the Collaborative Study on the Genetics of Alcoholism (COGA). *Alcohol Clin Exp Res* 1996;20:133A–137A.
- Reich T, Edenberg HJ, Goate A, Williams JT, Rice JP, Van Eerdewegh P, Foroud T, Hesselbrock V, Schuckit MA, Bucholz K, Porjesz B, Li TK, Conneally PM, Nurnberger JI Jr, Tischfield JA, Crowe RR, Cloninger CR, Wu W, Shears S, Carr K, Crose C, Willig C, Begleiter H. Genome-wide search for genes affecting the risk for alcohol dependence. *Am J Med Genet* 1998;81:207–215. [PubMed: 9603606]
- Rewal M, Jurd R, Gill TM, He DY, Ron D, Janak PH. Alpha4-containing GABA_A receptors in the nucleus accumbens mediate moderate intake of alcohol. *J Neurosci* 2009;29:543–549. [PubMed: 19144854]
- Ripley TL, Dunworth SJ, Stephens DN. Consequences of amygdala kindling and repeated withdrawal from ethanol on amphetamine-induced behaviours. *Eur J Neurosci* 2002;16:1129–1138. [PubMed: 12383242]
- Robello M, Amico C, Cupello A. Regulation of GABA_A receptor in cerebellar granule cells in culture: differential involvement of kinase activities. *Neuroscience* 1993;53:131–138. [PubMed: 7682295]
- Robello M, Amico C, Cupello A. A dual mechanism for impairment of GABA_A receptor activity by NMDA receptor activation in rat cerebellum granule cells. *Eur Biophys J* 1997;25:181–187. [PubMed: 9037753]
- Roberto M, Madamba SG, Moore SD, Tallent MK, Siggins GR. Ethanol increases GABAergic transmission at both pre- and postsynaptic sites in rat central amygdala neurons. *Proc Natl Acad Sci U S A* 2003;100:2053–2058. [PubMed: 12566570]
- Roberto M, Madamba SG, Stouffer DG, Parsons LH, Siggins GR. Increased GABA release in the central amygdala of ethanol-dependent rats. *J Neurosci* 2004;24:10159–10166. [PubMed: 15537886]
- Roca DJ, Rozenberg I, Farrant M, Farb DH. Chronic agonist exposure induces down-regulation and allosteric uncoupling of the gamma-aminobutyric acid/benzodiazepine receptor complex. *Mol Pharmacol* 1990;37:37–43. [PubMed: 2153908]
- Rudolph U, Crestani F, Benke D, Brunig I, Benson JA, Fritschy JM, Martin JR, Bluethmann H, Mohler H. Benzodiazepine actions mediated by specific γ -aminobutyric acid_A receptor subtypes. *Nature* 1999;401:796–800. [PubMed: 10548105]
- Salimov RM, McBride WJ, McKinzie DL, Lumeng L, Li TK. Effects of ethanol consumption by adolescent alcohol-preferring P rats on subsequent behavioral performance in the cross-maze and slip funnel tests. *Alcohol* 1996;13:297–300. [PubMed: 8734846]
- Sanchis-Segura C, Cline B, Jurd R, Rudolph U, Spanagel R. Etomidate and propofol-hyposensitive GABA_A receptor β 3 (N265M) mice show little changes in acute alcohol sensitivity but enhanced tolerance and withdrawal. *Neurosci Lett* 2007;416:275–278. [PubMed: 17350761]
- Sander T, Ball D, Murray R, Patel J, Samochowiec J, Winterer G, Rommelspacher H, Schmidt LG, Loh EW. Association analysis of sequence variants of GABA_A α 6, β 2, and γ 2 gene cluster and alcohol dependence. *Alcohol Clin Exp Res* 1999;23:427–431. [PubMed: 10195814]

- Sanna E, Serra M, Cossu A, Colombo G, Follesa P, Cuccheddu T, Concas A, Biggio G. Chronic ethanol intoxication induces differential effects on GABA_A and NMDA receptor function in the rat brain. *Alcohol Clin Exp Res* 1993;17:115–123. [PubMed: 8383922]
- Sanna E, Talani G, Busonero F, Pisu MG, Purdy RH, Serra M, Biggio G. Brain steroidogenesis mediates ethanol modulation of GABA_A receptor activity in rat hippocampus. *J Neurosci* 2004;24:6521–6530. [PubMed: 15269263]
- Santhakumar V, Hancher HJ, Wallner M, Olsen RW, Otis TS. Contributions of the GABA_A receptor $\alpha 6$ subunit to phasic and tonic inhibition revealed by a naturally occurring polymorphism in the $\alpha 6$ gene. *J Neurosci* 2006;26:3357–3364. [PubMed: 16554486]
- Schuckit MA, Mazzanti C, Smith TL, Ahmed U, Radel M, Iwata N, Goldman D. Selective genotyping for the role of 5-HT_{2A}, 5-HT_{2C}, and GABA $\alpha 6$ receptors and the serotonin transporter in the level of response to alcohol: a pilot study. *Biol Psychiatry* 1999;45:647–651. [PubMed: 10088053]
- Serra M, Pisu MG, Floris I, Cara V, Purdy RH, Biggio G. Social isolation-induced increase in the sensitivity of rats to the steroidogenic effect of ethanol. *J Neurochem* 2003;85:257–263. [PubMed: 12641747]
- Shannon EE, Shelton KL, Vivian JA, Yount I, Morgan AR, Homanics GE, Grant KA. Discriminative stimulus effects of ethanol in mice lacking the γ -aminobutyric acid type A receptor δ subunit. *Alcohol Clin Exp Res* 2004;28:906–913. [PubMed: 15201633]
- Shelton KL, Grant KA. Discriminative stimulus effects of ethanol in C57BL/6J and DBA/2J inbred mice. *Alcohol Clin Exp Res* 2002;26:747–757. [PubMed: 12068241]
- Shen EH, Dorow J, Harland R, Burkhardt-Kasch S, Phillips TJ. Seizure sensitivity and GABAergic modulation of ethanol sensitivity in selectively bred FAST and SLOW mouse lines. *J Pharmacol Exp Ther* 1998;287:606–615. [PubMed: 9808687]
- Sieghart W, Sperk G. Subunit composition, distribution and function of GABA(A) receptor subtypes. *Curr Top Med Chem* 2002;2:795–816. [PubMed: 12171572]
- Sieghart W, Fuchs K, Tretter V, Ebert V, Jechlinger M, Hoyer H, Adamiker D. Structure and subunit composition of GABA_A receptors. *Neurochem Int* 1999;34:379–385. [PubMed: 10397365]
- Silveri MM, Spear LP. Decreased sensitivity to the hypnotic effects of ethanol early in ontogeny. *Alcohol Clin Exp Res* 1998;22 (3):670–676. [PubMed: 9622449]
- Silveri MM, Spear LP. Ontogeny of ethanol elimination and ethanol-induced hypothermia. *Alcohol* 2000;20:45–53. [PubMed: 10680716]
- Silvers JM, Tokunaga S, Berry RB, White AM, Matthews DB. Impairments in spatial learning and memory: ethanol, allopregnanolone, and the hippocampus. *Brain Res Rev* 2003a;43:275–284. [PubMed: 14629930]
- Silvers JM, Tokunaga S, Mittleman G, Matthews DB. Chronic intermittent injections of high-dose ethanol during adolescence produce metabolic, hypnotic, and cognitive tolerance in rats. *Alcohol Clin Exp Res* 2003b;27:1606–1612. [PubMed: 14574231]
- Silvers JM, Tokunaga S, Mittleman G, Matthews DB. Chronic intermittent injections of high-dose ethanol during adolescence produce metabolic, hypnotic, and cognitive tolerance in rats. *Alcohol Clin Exp Res* 2003c;27:1606–1612. [PubMed: 14574231]
- Silvers JM, Tokunaga S, Mittleman G, O'Buckley T, Morrow AL, Matthews DB. Chronic intermittent ethanol exposure during adolescence reduces the effect of ethanol challenge on hippocampal allopregnanolone levels and Morris water maze task performance. *Alcohol* 2006;39:151–158. [PubMed: 17127134]
- Simpson CA, Rush CR. Acute performance-impairing and subject-rated effects of triazolam and temazepam, alone and in combination with ethanol, in humans. *J Psychopharmacol* 2002;16:23–34. [PubMed: 11949768]
- Sisk CL, Foster DL. The neural basis of puberty and adolescence. *Nat Neurosci* 2004;7:1040–1047. [PubMed: 15452575]
- Smith SS. Progesterone enhances inhibitory responses of cerebellar Purkinje cells mediated by the GABA_A receptor subtype. *Brain Res Bull* 1989;23:317–322. [PubMed: 2556209]
- Smith RF. Animal models of periadolescent substance abuse. *Neurotoxicol Teratol* 2003;25:291–301. [PubMed: 12757826]

- Smith SS, Gong QH, Hsu F-C, Markowitz RS, Ffrench-Mullen JMH, Li X. GABA_A receptor α 4 subunit suppression prevents withdrawal properties of an endogenous steroid. *Nature* 1998;392:926–930. [PubMed: 9582073]
- Smith SS, Ruderman Y, Hua Gong Q, Gulinello M. Effects of a low dose of ethanol in an animal model of premenstrual anxiety. *Alcohol* 2004;33:41–49. [PubMed: 15353172]
- Song J, Koller DL, Foroud T, Carr K, Zhao J, Rice J, Nurnberger JI Jr, Begleiter H, Porjesz B, Smith TL, Schuckit MA, Edenberg HJ. Association of GABA_A receptors and alcohol dependence and the effects of genetic imprinting. *Am J Med Genet B Neuropsychiatr Genet* 2003;117B:39–45. [PubMed: 12555233]
- Soyka M, Preuss UW, Hesselbrock V, Zill P, Koller G, Bondy B. GABA-A2 receptor subunit gene (GABRA2) polymorphisms and risk for alcohol dependence. *J Psychiatr Res* 2008;42:184–191. [PubMed: 17207817]
- Spear LP. The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev* 2000;24:417–463. [PubMed: 10817843]
- Spear LP. Assessment of adolescent neurotoxicity: rationale and methodological considerations. *Neurotoxicol Teratol* 2007;29:1–9. [PubMed: 17222532]
- Spear LP, Varlinskaya EI. Adolescence. Alcohol sensitivity, tolerance, and intake. *Recent Dev Alcohol* 2005;17:143–159. [PubMed: 15789864]
- Spencer RL, McEwen BS. Adaptation of the hypothalamic–pituitary–adrenal axis to chronic ethanol stress. *Neuroendocrinology* 1990;52:481–489. [PubMed: 2126355]
- Staley JK, Gottschalk C, Petrakis IL, Gueorguieva R, O'Malley S, Baldwin R, Jatlow P, Verhoeff NP, Perry E, Weinzimmer D, Frohlich E, Ruff E, van Dyck CH, Seibyl JP, Innis RB, Krystal JH. Cortical gamma-aminobutyric acid type A-benzodiazepine receptors in recovery from alcohol dependence: relationship to features of alcohol dependence and cigarette smoking. *Arch Gen Psychiatry* 2005;62:877–888. [PubMed: 16061765]
- Stell BM, Brickley SG, Tang CY, Farrant M, Mody I. Neuroactive steroids reduce neuronal excitability by selectively enhancing tonic inhibition mediated by δ subunit-containing GABA_A receptors. *Proc Natl Acad Sci U S A* 2003;100:14439–14444. [PubMed: 14623958]
- Stellwagen D, Beattie EC, Seo JY, Malenka RC. Differential regulation of AMPA receptor and GABA receptor trafficking by tumor necrosis factor- α . *J Neurosci* 2005;25:3219–3228. [PubMed: 15788779]
- Stephens DN, Pistovcakova J, Worthing L, Atack JR, Dawson GR. Role of GABA_A α 5-containing receptors in ethanol reward: the effects of targeted gene deletion, and a selective inverse agonist. *Eur J Pharmacol* 2005;526:240–250. [PubMed: 16253225]
- Sullivan EV, Sable HJ, Strother WN, Friedman DP, Davenport A, Tillman-Smith H, Kraft RA, Wyatt C, Szeliga KT, Buchheimer NC, Daunais JB, Adalsteinsson E, Pfefferbaum A, Grant KA. Neuroimaging of rodent and primate models of alcoholism: initial reports from the integrative neuroscience initiative on alcoholism. *Alcohol Clin Exp Res* 2005;29:287–294. [PubMed: 15714052]
- Sundstrom-Poromaa I, Smith DH, Gong QH, Sabado TN, Li X, Light A, Wiedmann M, Williams K, Smith S. Hormonally regulated α ₄ β ₂ δ GABA_A receptors are a target for alcohol. *Nat Neurosci* 2002;5:721–722. [PubMed: 12118257]
- Sur C, Farrar SJ, Kerby J, Whiting PJ, Atack JR, McKernan RM. Preferential coassembly of α 4 and δ subunits of the γ -aminobutyric acid_A receptor in rat thalamus. *Mol Pharmacol* 1999;56:110–115. [PubMed: 10385690]
- Takiguchi A, Masuoka T, Yamamoto Y, Mikami A, Kamei C. Potentiation of ethanol in spatial memory deficits induced by some benzodiazepines. *J Pharmacol Sci* 2006;101:325–328. [PubMed: 16891765]
- Tapia-Arancibia L, Rage F, Givalois L, Digneon P, Arancibia S, Beauge F. Effects of alcohol on brain-derived neurotrophic factor mRNA expression in discrete regions of the rat hippocampus and hypothalamus. *J Neurosci Res* 2001;63:200–208. [PubMed: 11169630]
- Tauber M, Calame-Droz E, Prut L, Rudolph U, Crestani F. α 2- γ -Aminobutyric acid (GABA) A receptors are the molecular substrates mediating precipitation of narcosis but not of sedation by the combined use of diazepam and alcohol in vivo. *Eur J Neurosci* 2003;18:2599–2604. [PubMed: 14622161]

- Taylor C, Nash J, Rich A, Lingford-Hughes A, Nutt D, Potokar J. Assessment of GABA_A benzodiazepine receptor (GBzR) sensitivity in patients with alcohol dependence. *Alcohol Alcohol* 2008;43:614–618. [PubMed: 18586906]
- Tehrani MHJ, Barnes EM Jr. Sequestration of gamma-aminobutyric acid_A receptors on clathrin-coated vesicles during chronic benzodiazepine administration *in vivo*. *J Pharmacol Exp Ther* 1997;283:384–390. [PubMed: 9336347]
- Terunuma M, Xu J, Vitlani M, Sieghart W, Kittler J, Pangalos M, Haydon PG, Coulter DA, Moss SJ. Deficits in phosphorylation of GABA_A receptors by intimately associated protein kinase C activity underlie compromised synaptic inhibition during status epilepticus. *J Neurosci* 2008;28:376–384. [PubMed: 18184780]
- Thiele TE, Willis B, Stadler J, Reynolds JG, Bernstein IL, McKnight GS. High ethanol consumption and low sensitivity to ethanol-induced sedation in protein kinase A-mutant mice. *J Neurosci* 2000;20:RC75. [PubMed: 10783399]
- Thomas P, Mortensen M, Hosie AM, Smart TG. Dynamic mobility of functional GABA_A receptors at inhibitory synapses. *Nat Neurosci* 2005;8:889–897. [PubMed: 15951809]
- Ticku MK, Burch T. Alterations in γ -aminobutyric acid receptor sensitivity following acute and chronic ethanol treatments. *J Neurochem* 1980;34:417–423. [PubMed: 6251168]
- Tokunaga S, Silvers JM, Matthews DB. Chronic intermittent ethanol exposure during adolescence blocks ethanol-induced inhibition of spontaneously active hippocampal pyramidal neurons. *Alcohol Clin Exp Res* 2006;30:1–6. [PubMed: 16433726]
- Tossman U, Jonsson G, Ungerstedt U. Regional distribution and extracellular levels of amino acids in rat central nervous system. *Acta Physiol Scand* 1986;127:533–545. [PubMed: 2875604]
- Tran VT, Snyder SH, Major LF, Hawley RJ. GABA receptors are increased in brains of alcoholics. *Ann Neurol* 1981;9:289–292. [PubMed: 6261673]
- Tremwel MF, Hunter BE, Peris J. Chronic ethanol exposure enhances [³H]GABA release and does not affect GABA_A receptor mediated ³⁶Cl uptake. *Synapse* 1994;17:149–154. [PubMed: 7974196]
- Tretter V, Ehya N, Fuchs K, Sieghart W. Stoichiometry and assembly of a recombinant GABA_A receptor subtype. *J Neurosci* 1997;17:2728–2737. [PubMed: 9092594]
- Truxell E, Molina JC, Spear NE. Ethanol intake in the juvenile, adolescent, and adult rat: effects of age and prior exposure to ethanol. *Alcohol Clin Exp Res* 2007;31:755–765. [PubMed: 17386073]
- Ueno S, Lin A, Nikolaeva N, Trudell JR, Mihic SJ, Harris RA, Harrison NL. Tryptophan scanning mutagenesis in TM2 of the GABA_A receptor α subunit: effects on channel gating and regulation by ethanol. *Br J Pharmacol* 2000;131:296–302. [PubMed: 10991923]
- Valenzuela CF, Mameli M, Carta M. Single-amino-acid difference in the sequence of alpha6 subunit dramatically increases the ethanol sensitivity of recombinant GABA_A receptors. *Alcohol Clin Exp Res* 2005;29:1356–1357. Author reply 1358. [PubMed: 16046882]
- VanDoren MJ, Matthews DB, Janis GC, Grobin AC, Devaud LL, Morrow AL. Neuroactive steroid 3 α -hydroxy-5 α -pregnan-20-one modulates electrophysiological and behavioral actions of ethanol. *J Neurosci* 2000;20:1982–1989. [PubMed: 10684899]
- Varlinskaya EI, Spear LP. Acute effects of ethanol on social behavior of adolescent and adult rats: role of familiarity of the test situation. *Alcohol Clin Exp Res* 2002;26:1502–1511. [PubMed: 12394283]
- Varlinskaya EI, Spear LP. Acute ethanol withdrawal (hangover) and social behavior in adolescent and adult male and female Sprague-Dawley rats. *Alcohol Clin Exp Res* 2004;28:40–50. [PubMed: 14745301]
- Vetter CS, Doremus-Fitzwater TL, Spear LP. Time course of elevated ethanol intake in adolescent relative to adult rats under continuous, voluntary-access conditions. *Alcohol Clin Exp Res* 2007;31:1159–1168. [PubMed: 17511750]
- Vivian JA, Green HL, Young JE, Majerksy LS, Thomas BW, Shively CA, Tobin JR, Nader MA, Grant KA. Induction and maintenance of ethanol self-administration in cynomolgus monkeys (*Macaca fascicularis*): long-term characterization of sex and individual differences. *Alcohol Clin Exp Res* 2001;25:1087–1097. [PubMed: 11505038]
- Volkow ND, Wang GJ, Hitzemann R, Fowler JS, Wolf AP, Pappas N, Biegon A, Dewey SL. Decreased cerebral response to inhibitory neurotransmission in alcoholics. *Am J Psychiatr* 1993;150:417–422. [PubMed: 8382010]

- Volkow ND, Wang GJ, Overall JE, Hitzemann R, Fowler JS, Pappas N, Frecska E, Piscani K. Regional brain metabolic response to lorazepam in alcoholics during early and late alcohol detoxification. *Alcohol Clin Exp Res* 1997;21:1278–1284. [PubMed: 9347090]
- Wallner M, Hanchar HJ, Olsen RW. Ethanol enhances $\alpha 4\beta 3\delta$ and $\alpha 6\beta 3\delta$ γ -aminobutyric acid type A receptors at low concentrations known to affect humans. *Proc Natl Acad Sci U S A* 2003;100:15218–15223. [PubMed: 14625373]
- Wallace MJ, Newton PM, Oyasu M, McMahon T, Chou WH, Connolly J, Messing RO. Acute functional tolerance to ethanol mediated by protein kinase Cepsilon. *Neuropsychopharmacology* 2007;32:127–136. [PubMed: 16541084]
- Wallner M, Hanchar HJ, Olsen RW. Low-dose alcohol actions on $\alpha 4\beta 3\delta$ GABA_A receptors are reversed by the behavioral alcohol antagonist Ro15–4513. *Proc Natl Acad Sci USA* 2006;103:8540–8545. [PubMed: 16698930]
- Wei W, Zhang N, Peng Z, Houser CR, Mody I. Perisynaptic localization of δ subunit-containing GABA_A receptors and their activation by GABA spillover in the mouse dentate gyrus. *J Neurosci* 2003;23:10650–10661. [PubMed: 14627650]
- Wei W, Faria LC, Mody I. Low ethanol concentrations selectively augment the tonic inhibition mediated by delta subunit-containing GABA_A receptors in hippocampal neurons. *J Neurosci* 2004;24:8379–8382. [PubMed: 15385620]
- Werner DF, Blednov YA, Ariwodola OJ, Silberman Y, Logan E, Berry RB, Borghese CM, Matthews DB, Weiner JL, Harrison NL, Harris RA, Homanics GE. Knockin mice with ethanol-insensitive $\alpha 1$ -containing γ -aminobutyric acid type A receptors display selective alterations in behavioral responses to ethanol. *J Pharmacol Exp Ther* 2006;319:219–227. [PubMed: 16785315]
- Werner DF, Swihart AR, Ferguson C, Lariviere WR, Harrison NL, Homanics GE. Alcohol-induced tolerance and physical dependence in mice with ethanol insensitive alpha1 GABA_A receptors. *Alcohol Clin Exp Res* 2009;33:289–299. [PubMed: 19032579]
- White AM, Ghia AJ, Levin ED, Swartzwelder HS. Binge pattern ethanol exposure in adolescent and adult rats: differential impact on subsequent responsiveness to ethanol. *Alcohol Clin Exp Res* 2000;24:1251–1256. [PubMed: 10968665]
- White AM, Bae JG, Truesdale MC, Ahmad S, Wilson WA, Swartzwelder HS. Chronic-intermittent ethanol exposure during adolescence prevents normal developmental changes in sensitivity to ethanol-induced motor impairments. *Alcohol Clin Exp Res* 2002a;26:960–968. [PubMed: 12170104]
- White AM, Truesdale MC, Bae JG, Ahmad S, Wilson WA, Best PJ, Swartzwelder HS. Differential effects of ethanol on motor coordination in adolescent and adult rats. *Pharmacol Biochem Behav* 2002b;73:673–677. [PubMed: 12151043]
- Whittemore ER, Yang W, Drewe JA, Woodward RM. Pharmacology of the human γ -aminobutyric acid_A $\alpha 4$ subunit expressed in *Xenopus laevis* oocytes. *Mol Pharmacol* 1996;50:1365–1375.
- Wick MJ, Mihic SJ, Ueno S, Mascia MP, Trudell JR, Brozowski SJ, Ye Q, Harrison NL, Harris RA. Mutations of γ -aminobutyric acid and glycine receptors change alcohol cutoff: evidence for an alcohol receptor? *Proc Natl Acad Sci U S A* 1998;95:6504–6509. [PubMed: 9600996]
- Wick MJ, Radcliffe RA, Bowers BJ, Mascia MP, Luscher B, Harris RA, Wehner JM. Behavioural changes produced by transgenic overexpression of $\gamma 2L$ and $\gamma 2S$ subunits of the GABA_A receptor. *Eur J Neurosci* 2000;12:2634–2638. [PubMed: 10947837]
- Wills TA, Knapp DJ, Overstreet DH, Breese GR. Differential dietary ethanol intake and blood ethanol levels in adolescent and adult rats: effects on anxiety-like behavior and seizure thresholds. *Alcohol Clin Exp Res* 2008;32:1350–1360. [PubMed: 18540921]
- Wisden W, Herb A, Wieland H, Keinänen K, Lüddens H, Seeburg PH. Cloning, pharmacological characteristics and expression pattern of the rat GABA_A receptor $\alpha 4$ subunit. *FEBS Lett* 1991;289:227–230. [PubMed: 1655526]
- Wisden W, Laurie DH, Monyer H, Seeburg PH. The distribution of 13 GABA_A receptor subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon. *J Neurosci* 1992;12:1040–1062.
- Woo E, Greenblatt DJ. Massive benzodiazepine requirements during acute alcohol withdrawal. *Am J Psychiatry* 1979;136:821–823. [PubMed: 443469]

- Yamashita M, Marszalec W, Yeh JZ, Narahashi T. Effects of ethanol on tonic GABA currents in cerebellar granule cells and mammalian cells recombinantly expressing GABA_A receptors. *J Pharmacol Exp Ther* 2006;319:431–438. [PubMed: 16844844]
- Ye Q, Koltchine VV, Mihic SJ, Mascia MP, Wick MJ, Finn SE, Harrison NL, Harris RA. Enhancement of glycine receptor function by ethanol is inversely correlated with molecular volume at position $\alpha 267$. *J Biol Chem* 1998;273:3314–3319. [PubMed: 9452448]
- Yu Z-Y, Wang W, Fritschy J-M, Witte OW, Redecker C. Changes in neocortical and hippocampal GABA_A receptor subunit distribution during brain maturation and aging. *Brain Res* 2006;1099:73–81. [PubMed: 16781682]
- Zafra F, Castren E, Thoenen H, Lindholm D. Interplay between glutamate and gamma-aminobutyric acid transmitter systems in the physiological regulation of brain-derived neurotrophic factor and nerve growth factor synthesis in hippocampal neurons. *Proc Natl Acad Sci USA* 1991;88:10037–10041. [PubMed: 1658793]
- Ziskind-Conhaim L, Gao BX, Hinckley C. Ethanol dual modulatory actions on spontaneous postsynaptic currents in spinal motoneurons. *J Neurophysiol* 2003;89:806–813. [PubMed: 12574458]

Mechanisms of GABA_A Receptor Trafficking




Fig. 1. Mechanisms of GABA_A receptor endocytosis and recycling. Endocytosis by clathrin-coated vesicles requires adaptor complex-2 (AP-2) and clathrin binding to the receptor. AP-2 is specific for endocytosis of surface proteins including GABA_A receptors. First, AP-2 recognizes and binds to GABA_A receptors, then clathrin binds the complex and the receptor is internalized. Phosphorylation of GABA_A receptors can prevent AP-2 binding and subsequent internalization. Internalized receptors may be recycled back to the cell surface or degraded depending upon undefined intracellular signals. Protein kinases can alter many steps in the receptor trafficking process. For example, PKC phosphorylation can alter recycling of receptors, whereas PKA can increase surface expression of GABA_A receptors. Adaptor complex-1 (AP-1) transports newly synthesized receptors from the trans-Golgi reticulum to the cell surface. Many proteins are required for intracellular trafficking of GABA_A receptors. For example, GABA_A receptor-associated protein (GABARAP) is involved in trafficking of GABA_A receptors from the endoplasmic reticulum/Golgi complex towards the plasma membrane and possibly also in vesicle-mediated cycling between the plasma membrane and cytoplasmic pool of receptors. *N*-ethylmaleimide-sensitive factor (NSF) interacts with GABARAP and influences intracellular trafficking of GABA_A receptors. Gephyrin reduces the diffusion of GABA_A receptors from the membrane and facilitates their clustering at synapses. Protein linking integrin-associated protein to cytoskeleton-1 (Plic-1) is also involved in GABA_A receptor trafficking and facilitates surface expression of the receptors (see Chen and Olsen 2007 for review)

Table 1

Acute ethanol effects on extrasynaptic receptors and tonic inhibition

Experimental system	Brain region/cell type	Ethanol concentration ^a	Experimental considerations	Recombinant GABA _A receptor(s)	Reference
Recombinant recordings					
	<i>X. laevis</i> oocytes	1 mM		$\alpha 4\beta 2\delta$	(Sundstrom-Poromaa et al. 2002)
		3 mM		$\alpha 4/6\beta 3\delta$	(Wallner et al. 2003)
		3 mM		$\alpha 6\beta 3\delta$	(Hancher et al. 2006)
		3 mM		$\alpha 4\beta 3\delta$ (blocked by Ro15-4513)	(Wallner et al. 2006)
		30 mM		$\alpha 4/6\beta 2\delta$	(Wallner et al. 2003)
		100 mM		$\alpha 4\beta 3\delta$	(Borghese et al. 2006a, b)
		No effect (30 mM)		Concatenated $\alpha 1\beta 3\delta$	(Kaur et al. 2009)
	L(tk-) fibroblasts	3–100 mM		$\alpha 4\beta 3\delta$ (transfected with PKC δ)	(Choi et al. 2008)
	CHO cells	300 mM		$\alpha 4\beta 3\delta$	(Borghese et al. 2006a, b)
		No effect (10–100 mM)		$\alpha 4/6\beta 2\delta$, $\alpha 6\beta 3\delta$	(Yamashita et al. 2006)
Slice recordings					
	Hippocampal CA1 pyramidal cells	1–3 mM	Progesterone withdrawal		(Smith et al. 2004) (Sundstrom-Poromaa et al. 2002)
		50 mM			(Liang et al. 2007)
		No effect (30 mM)			(Wei et al. 2004)
	Dentate gyrus granule cells	30 mM			(Wei et al. 2004)
		30 mM			(Glykys et al. 2007)
		30 mM	\uparrow tonic current in adolescents; \uparrow basal tonic current in adults		(Fleming et al. 2007)
		30 mM	Not observed in PKC δ KOs		(Choi et al. 2008)
		50 mM			(Liang et al. 2008)
		No effect (30 mM)			(Borghese et al. 2006a, b)
	Molecular layer interneurons	20 mM	$\alpha 1$ co-localized with δ ; observed in $\alpha 4$ KOs		(Glykys et al. 2007)
		30 mM	Not observed in PKC δ KOs		(Choi et al. 2008)
Thalamus:					
	Ventrobasal neurons	50 mM			(Jia et al. 2008a)
	Relay neurons	30 mM	Not observed in PKC δ KOs		(Choi et al. 2008)

Experimental system	Brain region/cell type	Ethanol concentration ^a	Experimental considerations	Recombinant GABA _A receptor(s)	Reference
Ro15-4513 binding	Cerebellar granule cells	10 mM	↑ ethanol enhancement with R _{100Q} mutation in outbred rats		(Hanchar et al. 2005)
		20 mM	↑ spontaneous IPSCs		(Carta et al. 2004)
		25 mM	↑ spontaneous IPSCs; no ethanol enhancement with R _{100Q} mutation in outbred rats		(Botta et al. 2007a)
		40 mM	↑ spontaneous IPSCs; No ethanol enhancement with R _{100Q} mutation in outbred rats		(Botta et al. 2007b)
		50 mM	↑ spontaneous IPSCs; No ethanol enhancement with R _{100Q} mutation in inbred AT and ANT rats		(Valenzuela et al. 2005)
		100 mM			(Casagrande et al. 2007)
		No effect (30 mM)			(Yamashita et al. 2006)
		1–30 mM			(Hanchar et al. 2006)
		No effect (1–100 mM)			(Korpi et al. 2007)
		1–30 mM			(Hanchar et al. 2006)
		500 mM			(Mehta et al. 2007)
		No effect (1–50 mM)			(Korpi et al. 2007)
	No effect (1–100 mM)			(Korpi et al. 2007)	

↑, increased. Abbreviations: *ANT* alcohol non-tolerant; *AT* alcohol tolerant; *CHO* Chinese hamster ovary; *HEX* human embryonic kidney; *IPSCs* inhibitory post-synaptic currents; *KO* knockout; *PKC δ* protein kinase C δ

^aLowest effective ethanol concentration

Table 2Effects of chronic ethanol exposure on GABA_A receptor function in brain

GABA _A receptor activity	Alteration	Reference
Cl ⁻ channel function	↓ Cl ⁻ flux ^{a,b}	(Kang et al. 1996; Morrow et al. 1988; Sanna et al. 1993; Ticku and Burch 1980)
	↔ Cl ⁻ flux ^{c,d}	(Allan and Harris 1987; Tremwel et al. 1994)
	↓ mIPSC area ^c	(Cagetti et al. 2003; Liang et al. 2004; Liang et al. 2006, 2007)
	↓ Tonic current ^c	(Liang et al. 2007)
Ethanol-enhanced Cl ⁻ channel function	↓ Cl ⁻ flux ^d	(Allan and Harris 1987; Morrow et al. 1988)
	↑ mIPSC enhancement ^c	(Liang et al. 2006, 2007)
	↓ Tonic current enhancement ^c	(Liang et al. 2006, 2007)
TBPS binding	↑ Binding ^a	(Sanna et al. 1993)
Barbiturate modulation	↓ Pentobarbital-mediated Cl ⁻ flux ^a	(Morrow et al. 1988)
Benzodiazepine modulation	↓ Flunitrazepam Cl ⁻ Flux ^d	(Buck and Harris 1990; Sanna et al. 1993)
	↓ Diazepam-enhanced mIPSCs ^c	(Cagetti et al. 2003)
	↓ Diazepam-enhanced tonic current ^c	(Liang et al. 2004)
	↓ Zolpidem-enhanced mIPSCs ^c	(Liang et al. 2004)
	↓ Zolpidem-enhanced tonic current ^c	(Liang et al. 2004)
Ro15-4513 modulation	↑ Cl ⁻ flux inhibition ^a	(Buck and Harris 1990)
	↑ Binding ^{c,d}	(Mhatre et al. 1988; Cagetti et al. 2003)
	↑ mIPSCs enhancement ^c	(Cagetti et al. 2003)
	↓ Tonic current enhancement ^c	(Liang et al. 2004)
THIP modulation	↓ Enhancement of tonic current ^c	(Liang et al. 2004)
	↑ mIPSCs ^c	(Liang et al. 2004)
Neuroactive steroid modulation	↑ 3 α ,5 α -THP Cl ⁻ flux ^a	(Devaud et al. 1996)
	↑ 3 α ,5 α -THDOC Cl ⁻ flux ^a	(Devaud et al. 1996)
	↓ Alphaxalone mIPSC enhancement ^c	(Cagetti et al. 2003)

↑ Increased, ↓ decreased, ↓ abolished, ↔ no change

Abbreviations: 3 α ,5 α -THP (3 α ,5 α)-3-hydroxypregnan-20-one; 3 α ,5 α -THDOC (3 α ,5 α)-3,21-dihydroxypregnan-20-one; TBPS *t*-butylbicyclophosphorothionate; THIP 4,5,6,7-tetrahydroisothiazolo-[5,4-c]pyridine-3-ol

^aCerebral cortex

^bCerebellum

^cHippocampus

^dCerebral cortex, hippocampus, and cerebellum

Table 3

Effects of chronic ethanol administration on GABA_A receptor subunit expression in cerebral cortex, hippocampus, and cerebellum

Subunit	Alteration	Reference
$\alpha 1$	mRNA $\downarrow a,b,c$ $\leftrightarrow b$	(Charlton et al. 1997; Devaud et al. 1995; Liang et al. 2007; Mhatre and Ticku 1992; Montpied et al. 1991b; Morrow et al. 1992) (Petrie et al. 2001)
	Peptide $\downarrow a,b,c$ $\leftrightarrow c$	(Cagetti et al. 2003; Charlton et al. 1997; Devaud et al. 1997; Kumar et al. 2002; Marutha Ravindran et al. 2007; Mhatre et al. 1993; Werner et al. 2009) (Marutha Ravindran et al. 2007; Matthews et al. 1998)
$\alpha 2$	mRNA $\downarrow a$ $\leftrightarrow c$	(Mhatre and Ticku 1992, 1994b; Montpied et al. 1991b) (Cagetti et al. 2003)
	Peptide $\downarrow a,b$ $\leftrightarrow c$	(Marutha Ravindran et al. 2007; Mhatre et al. 1993; Mhatre and Ticku 1994b) (Marutha Ravindran et al. 2007; Matthews et al. 1998)
$\alpha 3$	mRNA $\downarrow a$ $\leftrightarrow a$	(Mhatre and Ticku 1994b) (Mhatre and Ticku 1992; Montpied et al. 1991b)
	Peptide $\downarrow a$ $\leftrightarrow c$	(Mhatre et al. 1993; Mhatre and Ticku 1994b) (Matthews et al. 1998)
$\alpha 4$	mRNA $\uparrow a,c$ $\leftrightarrow c$	(Devaud et al. 1995; Mahmoudi et al. 1997) (Petrie et al. 2001)
	Peptide $\uparrow a,c$ $\uparrow a$	(Cagetti et al. 2003; Devaud et al. 1997; Kumar et al. 2002; Liang et al. 2007; Marutha Ravindran et al. 2007; Matthews et al. 1998)
$\alpha 5$	mRNA $\uparrow a$ $\downarrow a$ $\leftrightarrow a,c$	(Charlton et al. 1997) (Mhatre and Ticku 1992) (Devaud et al. 1995; Mahmoudi et al. 1997; Petrie et al. 2001)
	Peptide $\downarrow a$ $\leftrightarrow c$	(Charlton et al. 1997) (Liang et al. 2007)
$\alpha 6$	mRNA $\uparrow b$ $\leftrightarrow b$	(Morrow et al. 1992) (Petrie et al. 2001)
	Peptide $\uparrow d$ $\uparrow a,b$ $\leftrightarrow a$	(Marutha Ravindran et al. 2007) (Mhatre and Ticku 1992, 1994a) (Devaud et al. 1995)

Subunit	Alteration	Reference
β2	mRNA	(Mhahre and Ticku 1994a)
	Peptide	(Devaud et al. 1995)
β3	mRNA	(Devaud et al. 1997; Marutha Ravindran et al. 2007)
	Peptide	(Marutha Ravindran et al. 2007; Matthews et al. 1998)
γ1	mRNA	(Mhahre and Ticku 1994a)
	Peptide	(Devaud et al. 1995)
γ2	mRNA	(Devaud et al. 1997)
	Peptide	(Matthews et al. 1998)
γ3	mRNA	(Cagetti et al. 2003; Devaud et al. 1995)
	Peptide	(Petrie et al. 2001)
δ	mRNA	(Devaud et al. 1997)
	Peptide	(Petrie et al. 2001)
γ2S	mRNA	(Cagetti et al. 2003; Liang et al. 2007; Marutha Ravindran et al. 2007)
	Peptide	(Devaud et al. 1997; Marutha Ravindran et al. 2007; Matthews et al. 1998)
γ2L	mRNA	(Cagetti et al. 2003; Devaud et al. 1995)
	Peptide	(Cagetti et al. 2003; Devaud et al. 1995)
γ3	mRNA	(Devaud et al. 1995)
	Peptide	(Devaud et al. 1995; Petrie et al. 2001)
δ	mRNA	(Cagetti et al. 2003; Liang et al. 2007)
	Peptide	(Cagetti et al. 2003; Liang et al. 2007)

↑ Increased; ↓ decreased; ↔ no change

^a Cerebral cortex

^b Cerebellum

^c Hippocampus