



Published in final edited form as:

Neurosci Lett. 2010 January 29; 469(3): 338. doi:10.1016/j.neulet.2009.12.023.

6-Hz Primed Low-Frequency rTMS to Contralesional M1 in Two Cases with Middle Cerebral Artery Stroke

James R. Carey, PhD, PT¹, David C. Anderson, MD², Bernadette T. Gillick, MS, PT [Doctoral Candidate]³, Maureen Whitford, MS, PT³, and Alvaro Pascual-Leone, MD, PhD⁴

¹)Program in Physical Therapy, University of Minnesota; Minneapolis, MN, USA

²)Department of Neurology, University of Minnesota

³)Graduate Program in Rehabilitative Science, University of Minnesota

⁴)Berenson-Allen Center for Noninvasive Brain Stimulation, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA; and Institut Gutmann de Neurorehabilitación, Universidad Autónoma, Barcelona, Spain

Abstract

This case study contrasted two subjects with stroke who received 6-Hz primed low-frequency repetitive transcranial magnetic stimulation (rTMS) to the contralesional primary motor area (M1) to disinhibit ipsilesional M1. Functional magnetic resonance imaging (fMRI) showed that the intervention disrupted cortical activation at contralesional M1. Subject 1 showed decreased intracortical inhibition and increased intracortical facilitation following intervention during paired-pulse TMS testing of ipsilesional M1. Subject 2, whose precentral knob was totally obliterated and who did not show an ipsilesional motor evoked potential at pretest, still did not show any at posttest; however, her fMRI did show a large increase in peri-infarct zone cortical activation. Behavioral results were mixed, indicating the need for accompanying behavioral training to capitalize on the brain organization changes induced with rTMS.

Keywords

fMRI; rTMS; stroke; hand

Introduction

Functional deficits following stroke are due not only to the ischemic loss of neurons but also to maladaptive brain reorganization associated with compensatory “learned non-use” [34] behaviors. Exaggerated interhemispheric inhibition of the ipsilesional primary motor area (M1) by contralesional M1 can lead to down-regulation of excitability in neurons that have survived the stroke and a worsening of the functional deficit [13,23,30]. Low-frequency repetitive transcranial magnetic stimulation (rTMS) applied to contralesional M1 has been shown to reduce its inhibition on ipsilesional M1 [14,31,33], leading to improved function in the paretic

© 2009 Elsevier Ireland Ltd. All rights reserved

carey007@umn.edu, phone: 612/626-2746; FAX: 612/625-4274.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

hand. Importantly, Iyer et al. [22] showed in healthy humans that the disruptive effect of low-frequency stimulation could be magnified if it was preceded (primed) with high-frequency stimulation. We recently have shown that 6-Hz primed low-frequency rTMS is safe in subjects with stroke [7]. The purpose of this case report was to contrast findings in two subjects who volunteered for a larger study examining the efficacy of 6-Hz primed low-frequency rTMS to contralesional M1.

Methods

Subjects

Both subjects had right middle cerebral artery strokes with dense paresis of the left upper extremity. For both, the stroke duration was 10 years and both were right-handed pre-stroke. Subject 1 was a 71-year-old male (upper extremity Fugl-Meyer [16] score = 36). Subject 2 was a 52-year-old woman (Fugl-Meyer = 35). The experiments were conducted in accordance with the Declaration of Helsinki and both subjects signed informed consent forms approved by our IRB. The study was conducted under an investigational device exemption (IDE# G050260) from the United States of America Food and Drug Administration.

Testing

For subject 1, testing occurred at pretest and two weeks later at posttest with five intervention sessions given every other day between the two tests. His posttest occurred 72 hours after his last session. Testers were blinded. In subject 2, we could not elicit a motor evoked potential (MEP) at pretest when applying TMS to ipsilesional M1 and so this person was not included in the larger. But as a case study, two pretests were given – one when she volunteered for the larger study and four months later after IRB approval as a case study. For subject 2, the five intervention sessions occurred daily after the second pretest and the posttest occurred 24 hours after the last session.

fMRI—Anatomical and functional images were acquired using a 3 Tesla magnet. The details of the data acquisition have been reported earlier [6]. Briefly, with electrogoniometers applied to the paretic and nonparetic index fingers, each subject attempted to track a target waveform displayed on a screen using finger flexion/extension movement. The task consisted of alternating 30-s blocks of rest (7 blocks), paretic (3 blocks) and nonparetic (3 blocks) finger tracking. We evaluated nonparetic finger tracking to screen for a possible adverse effect in the nonparetic hand. A total of 130 T₂-weighted scans of the blood-oxygen-level-dependent signal were taken over the brain volume divided into 36 slices. Voxel resolution was 3 × 3 × 3 mm. Finger tracking was quantified with an accuracy index [5] (maximum=100%).

Brain Voyager software was used for fMRI analysis as described previously [6]. The 3D functional volume was co-registered with the corresponding 3D anatomical volume, and both were normalized to standard Talairach space. A general linear model was run that created an activation map showing active voxels with significantly different signal intensity between paretic finger tracking and rest using a false discovery rate (FDR) [18] of $q(\text{FDR}) < 0.05$.

TMS—With EMG electrodes recording from the paretic extensor digitorum (ED) muscle, TMS was delivered to the scalp ipsilesionally using two Magstim 200² stimulators coupled by a Bistim module (Magstim, Dyfed, UK) and a figure-of-eight coil (wing diameter: 70-mm). Single-pulse stimulation was applied at 0.1 Hz starting at an intensity of 60% of machine maximum. The scalp location and stimulus intensity were adjusted until the optimal location was found for producing the resting motor threshold (RMT), defined as the lowest intensity that induced MEPs of at least 50 μV (peak-to-peak) on at least 5 of 10 trials in the target muscle [3].

Once the ipsilesional RMT was found (subject 1 only), 10 single pulses at 120% of his RMT were applied at 0.1 Hz. Then, paired pulses were applied with the conditioning pulse set at 80% of RMT and the test pulse at 120%. Short and long interpulse intervals between the conditioning and test pulses were used to assess intracortical inhibition (ICI) and facilitation (ICF), respectively [24]. We chose two short intervals (2 and 3 ms) and two long intervals (10 and 15 ms), consistent with Hamzei et al. [20], to ensure capturing of the suspected effects. Each MEP amplitude from 10 paired pulses applied randomly at each interval was normalized to the average of the 10 single-pulse MEPs.

Behavioral—Subjects performed three trials each of the Box and Block Test (BBT) [28], involving grasp and release of small cubes with the paretic hand over one minute, and a maximal finger extension force test against a calibrated load cell. The Motor Activity Log (MAL) [25] measured “real-world” function as judged by subject responses to structured interview questions on the amount of use and quality of movement (both rated 0–5) in using the paretic arm in 30 activities. The Test Évaluant la performance des Membres supérieurs des Personnes Âgées (TEMPA) [12] was used to assess manual skill, recorded as the total time to complete only those tasks that involved using both hands together (neither subject could perform the paretic-hand-only tasks).

Intervention

The RMT for the nonparetic extensor digitorum muscle was determined by stimulating the contralesional M1. rTMS intervention involved two phases, priming and low-rate stimulation. Priming consisted of 10 minutes of intermittent 6-Hz rTMS given in 5-s trains with 25-s intervals between trains (i.e. 30 pulses/train \times 2 trains/min \times 10 min. = 600 priming pulses) at an intensity of 90% RMT. Immediately following priming, 10 min. of continuous 1-Hz rTMS (i.e. 600 low-frequency pulses) was given at 90% RMT. We did not calculate a new RMT between the priming and low-frequency stimulation because we wanted to replicate the procedure used by Iyer et al [22] and not risk losing the priming effect that may depend on the immediacy of low-frequency following high-frequency stimulation.

Analysis

For fMRI, we restricted the location of our analysis to the active voxels located in the contralesional and ipsilesional sensorimotor cortex (SMC) at the level of the precentral knob because it is considered to be the focal area of M1 representing the hand [36]. We identified three brain slices in the transverse plane showing the precentral knob in both subjects ($Z = 52, 55, 58$). At these slices, we used Brain Voyager analysis features to record the active voxel count in the SMC and their average signal intensity (average t statistic reflecting difference between the paretic finger tracking and rest conditions).

For TMS paired-pulse testing (subject 1 only), following confirmation that the data were normally distributed, a one-tailed paired t test was applied to the paired-pulse/single-pulse ratios of MEP amplitude.

Results

fMRI

For all fMRI results, cortical activation corresponds to the paretic finger tracking condition. Subject 1 showed a decrease in active voxels in contralesional SMC following intervention (Fig. 1, Table 1) with little change in intensity, whereas ipsilesional SMC showed little change in active voxels but an increase in intensity. Subject 2 showed a relatively small increase in activation in contralesional SMC from pretest 1 to pretest 2 (Fig. 2, Table 1) followed by a more substantial decrease in active voxels and intensity after intervention. Ipsilesionally, the

small amount of available SMC substrate showed an increase in activation across the three tests; however, the more notable observation is the absence of change in peri-infarct zone activation from pretest 1 to pretest 2 but then a sizeable increase after intervention.

TMS

Subject 1's paired-pulse/single-pulse MEP ratios are shown in Fig. 3. For 2-ms and 3-ms interpulse intervals, Kujirai et al. [24] showed on healthy subjects that conditioned MEPs were reduced to roughly 20–30% of unconditional MEPs and our pretest values were close to that range. At posttest for the 3-ms interval, with the ratio increasing toward 1.0, there was a significant reduction ($p=0.029$) in ICI relative to pretest. For 10-ms and 15-ms intervals, Kujirai et al found that conditioned MEPs were facilitated to roughly 125–160% of unconditioned MEPs but our pretest values, being less than 1.0 on average, showed no facilitation. However, at posttest, both the 10-ms and 15-ms intervals showed increased ICF, relative to pretest, with the latter interval being significant ($p=0.017$). For subject 2, no single-pulse MEP could be elicited at pretest 1, pretest 2 or posttest even at intensities as high as 100% of machine maximum and with a broad search area. She then did not receive paired-pulse testing.

Behavioral

Subject 1's paretic hand tracking accuracy and finger extension force improved from pretest to at posttest but this did not translate into improved BBT or TEMPA scores (Table 1). His own rating of motor function (MAL) decreased in the amount of use but increased in the quality of movement. Subject 2's paretic hand tracking accuracy and TEMPA scores showed general decreases in skill across her three tests, whereas her BBT scores, finger extension force and MAL ratings showed general increases.

Safety

There were no adverse effects in either subject.

Discussion

The important finding of this study was that 6-Hz primed low-frequency rTMS to contralesional M1 was able to produce physiologic effects that were measurable 72 hours (subject 1) and 24 hours (subject 2) after their last intervention. However, the actual physiologic effects were different between the two subjects and this may relate to whether the precentral knob was damaged [20]. The fMRI results for both subjects suggest that the intervention disrupted the voluntary recruitment of contralesional M1, which was intended and is consistent with Nowak et al. [31]. Furthermore, in subject 1 only, there was an increase in BOLD signal intensity of the active voxels in ipsilesional SMC at the precentral knob, possibly as a consequence of the reduced voluntary recruitment of contralesional M1 leading to reduced interhemispheric inhibition from ipsilesional M1 [9,21]. Indeed, paired-pulse TMS testing to his ipsilesional M1 showed reduced ICI and increased ICF (Fig. 3), which is consistent with Wu et al. [35] in healthy subjects following 15 Hz rTMS.

Contrarily, whereas the ipsilesional precentral knob was largely spared in subject 1, it was obliterated in subject 2. Loss of this neural substrate likely accounts for the absence of any MEPs with single-pulse TMS to ipsilesional M1 at pretest 1 and pretest 2, which has also been reported by Bütefisch et al. [4]. At posttest, we were uncertain whether the small area of spared SMC at the medial edge of the infarct (Fig. 2) might, through vicariation of function [15], respond to the rTMS intervention and elicit MEPs during TMS testing. Although an increase in cortical activation of some of this spared SMC was observed (Fig. 2, $Z=58$), there still was no MEP. Instead, conceivably as an alternative brain reorganization strategy in the absence of

precentral knob substrate, there was an increase in peri-infarct zone activation extending into the ipsilesional premotor area [15].

The decreased tracking performance with her paretic hand at posttest may actually be a reflection of this large change in brain organization without sufficient time or training to capitalize on it. Other studies have reported on the rich potential for neuroplasticity in the peri-infarct zone, possibly leading to higher functional recovery if up-regulated [8,10,17,26]. Our speculation of different brain reorganization strategies between the two subjects being dependent on the presence or absence of the precentral knob is consistent with the findings of Hamzei et al. [20] in their use of constraint induced movement therapy in stroke.

Behavioral changes in our study were inconsistent, as some tests showed gains and others showed declines in both subjects. The upward trends in the BBT and finger force for subject 2 makes it unclear whether the improvement at posttest reflects an intervention effect or a learning effect from repeated exposure. With data from only two subjects, it is difficult to draw any conclusions on the functional effectiveness of the intervention. We postulate, however, that to fully capitalize on the rTMS-induced brain reorganization, it may be necessary to complement the rTMS intervention with behavioral training.

Studies on groups of subjects with stroke have found improved function in the paretic hand measured immediately after a single session of low-frequency rTMS to contralesional M1 [27,31,33]. But this raises the important therapeutic concern about the duration of the rTMS-induced after-effects, as Romero et al. [32] showed in healthy subjects that effects lasted only up to 15 minutes. Fregni et al. [14] addressed this concern in subjects with stroke by applying five sessions of low-frequency rTMS over two weeks. They found improvements in hand function two weeks after the last intervention. Although they found reduced excitability in contralesional M1 immediately after the last intervention, they did not do excitability testing at the two-week follow-up. Thus, to our knowledge, there are no studies in people with stroke that have determined the duration of cortical excitability changes beyond the day of intervention. In light of the transient effect found by Romero et al. [32] (albeit in healthy subjects), our findings in subject 1 of excitability effects 72 hours after the last day of intervention suggest that there may be value to the inclusion of priming, as shown by Iyer et al. [22] in healthy subjects. However, our design did not directly compare priming to no priming and so we are unable to conclude that priming was key to the observed duration of excitability after-effects. Future studies will need to explore this concern.

The exact mechanism of priming is not yet clear, but growing evidence [19,29] suggests that it operates through the Bienenstock-Cooper-Munro theory of bidirectional synaptic plasticity [1], which states that neuronal reactivity to conditioning stimuli depends on the recent history of activity. Restated, low levels of prior (i.e. prior to conditioning) neuronal activity bias the synapse toward long-term potentiation and high levels bias toward long-term depression [29], consistent with the underlying principle of preserving synaptic homeostasis. We believe that our results measured 72 hours (subject 1) and 24 hours (subject 2) after intervention suggest that the intended physiological effects were achieved but the mixed behavioral effects indicate that rehabilitative training should accompany the stimulation. Indeed, the extended duration of physiological after-effects gives the opportunity to apply other rehabilitative interventions during a critical time when the ipsilesional M1 is up-regulated. The duration of up-regulated excitability remains unknown for now in humans with stroke. Nonetheless, the theoretical principle remains strong in capitalizing on Hebbian-based rules for synaptic plasticity [2,11], which emphasize the importance of temporally correlated coactivation of presynaptic and postsynaptic activity (i.e. behavioral training amidst cortical disinhibition) to maximize excitability change. Ideally, surviving neurons in the ipsilesional SMC (and possibly other ipsilesional areas) could then emerge from their diaschisis and re-enter voluntary recruitment

leading to improved function, even in chronic stroke [20]. Further studies are needed to verify contralesional excitability changes and direct evidence of transcallosal inhibition.

Acknowledgments

This project was funded by the NIH (National Institute of Child Health and Human Development 1R01HD053153 and the National Center for Research Resources P41 RR008079 and M01-RR00400) . NIH grant K24 RR018875 supported in part Dr. Pascual-Leone's participation in the study.

References

- [1]. Bienenstock EL, Cooper LN, Munro PW. Theory for the development of neuron selectivity: orientation specificity and binocular interaction in visual cortex. *Journal of Neuroscience* 1982;2:32–48. [PubMed: 7054394]
- [2]. Buonomano DV, Merzenich MM. Cortical plasticity: from synapses to maps. *Annual Review of Neuroscience* 1998;21:149–186.
- [3]. Butefisch CM, Khurana V, Kopylev L, Cohen LG. Enhancing encoding of a motor memory in the primary motor cortex by cortical stimulation. *J. Neurophysiol* 2004;91:2110–2116. [PubMed: 14711974]
- [4]. Butefisch CM, Wessling M, Netz J, Seitz RJ, Homberg V. Relationship between interhemispheric inhibition and motor cortex excitability in subacute stroke patients. *Neurorehabilitation & Neural Repair* 2008;22:4–21. [PubMed: 17507644]
- [5]. Carey JR. Manual Stretch: Effect on Finger Movement Control and Force Control in Stroke Subjects with Spastic Extrinsic Finger Flexor Muscles. *Arch Phys Med Rehabil* 1990;71:888–894. [PubMed: 2222157]
- [6]. Carey JR, Durfee WK, Bhatt E, Nagpal A, Weinstein SA, Anderson KM, Lewis SM. Tracking vs. Movement Telerehabilitation Training to Change Hand Function and Brain Reorganization in Stroke. *Neurorehabilitation & Neural Repair* 2007;21:216–232. [PubMed: 17351083]
- [7]. Carey JR, Evans CD, Anderson DC, Bhatt E, Nagpal A, Kimberley TJ, Pascual-Leone A. Safety of 6-Hz Primed Low-Frequency rTMS in Stroke. *Neurorehabilitation & Neural Repair* 2008;22:185–192. [PubMed: 17876070]
- [8]. Carey JR, Kimberley TJ, Lewis SM, Auerbach E, Dorsey L, Rundquist P, Ugurbil K. Analysis of fMRI and Finger Tracking Training in Subjects with Chronic Stroke. *Brain* 2002;125:773–788. [PubMed: 11912111]
- [9]. Classen J, Schnitzler A, Binkofski F, Werhahn KJ, Kim YS, Kessler KR, Benecke R. The motor syndrome associated with exaggerated inhibition within the primary motor cortex of patients with hemiparetic. *Brain* 1997;120:605–619. [PubMed: 9153123]
- [10]. Cramer S, Nelles G, Benson R, Kaplan J, Parker R, Kwong K, Kennedy D, Finklestein S, Rosen B. A functional MRI study of subjects recovered from hemiparetic stroke. *Stroke* 1997;28:2518–2527. [PubMed: 9412643]
- [11]. Desai NS. Homeostatic plasticity in the CNS: synaptic and intrinsic forms. *Journal of Physiology-Paris* 2003;97:391–402.
- [12]. Desrosiers J, Hebert R, Bravo G, Dutil E. Upper extremity performance test for the elderly (TEMPA): normative data and correlates with sensorimotor parameters. *Test d'Evaluation des Membres Superieurs de Personnes Agees. Archives of Physical Medicine & Rehabilitation* 1995;76:1125–1129. [PubMed: 8540788]
- [13]. Duque J, Murase N, Celnik P, Hummel F, Harris-Love M, Mazzocchio R, Olivier E, Cohen LG. Intermanual Differences in movement-related interhemispheric inhibition. *Journal of Cognitive Neuroscience* 2007;19:204–213. [PubMed: 17280510]
- [14]. Fregni F, Boggio PS, Valle AC, Rocha RR, Duarte J, Ferreira MJ, Wagner T, Fecteau S, Rigonatti SP, Riberto M, Freedman SD, Pascual-Leone A. A sham-controlled trial of a 5-day course of repetitive transcranial magnetic stimulation of the unaffected hemisphere in stroke patients. *Stroke* 2006;37:2115–2122. [PubMed: 16809569]
- [15]. Fridman EA, Hanakawa T, Chung M, Hummel F, Leiguarda RC, Cohen LG. Reorganization of the human ipsilesional premotor cortex after stroke. *Brain* 2004;127:747–758. [PubMed: 14749291]

- [16]. Fugl-Meyer A, Jaasko L, Leyman I, Olsson S, Steglind S. The post-stroke hemiplegic patient: A method for evaluation of physical performance. *Scand J Rehab Med* 1975;7:13–31.
- [17]. Furlan M, Marchal G, Viader F, Derlon JM, Baron JC. Spontaneous neurological recovery after stroke and the fate of the ischemic penumbra. *Ann Neurol* 1996;40:216–226. [PubMed: 8773603]
- [18]. Genovese CR, Lazar NA, Nichols T. Thresholding of Statistical Maps in Functional Neuroimaging Using the False Discovery Rate. *NeuroImage* 2002;15:870–878. [PubMed: 11906227]
- [19]. Hamada M, Terao Y, Hanajima R, Shirota Y, Nakatani-Enomoto S, Furubayashi T, Matsumoto H, Ugawa Y. Bidirectional long-term motor cortical plasticity and metaplasticity induced by quadripulse transcranial magnetic stimulation. *Journal of Physiology* 2008;586:3927–3947. see comment. [PubMed: 18599542]
- [20]. Hamzei F, Liepert J, Dettmers C, Weiller C, Rijntjes M. Two different reorganization patterns after rehabilitative therapy: an exploratory study with fMRI and TMS. *NeuroImage* 2006;31:710–720. [PubMed: 16516499]
- [21]. Hummel FC, Cohen LG. Non-invasive brain stimulation: a new strategy to improve neurorehabilitation after stroke? *Lancet Neurology* 2006;5:708–712. [PubMed: 16857577]
- [22]. Iyer MB, Schleper N, Wassermann EM. Priming stimulation enhances the depressant effect of low-frequency repetitive transcranial magnetic stimulation. *Journal of Neuroscience* 2003;23:10867–10872. [PubMed: 14645480]
- [23]. Kirton A, Chen R, Friefeld S, Gunraj C, Pontigon AM, Deveber G. Contralesional repetitive transcranial magnetic stimulation for chronic hemiparesis in subcortical paediatric stroke: a randomised trial. *Lancet Neurology* 2008;7:507–513. see comment. [PubMed: 18455961]
- [24]. Kujirai T, Caramia MD, Rothwell JC, Day BL, Thompson PD, Ferbert A, Wroe S, Asselman P, Marsden CD. Corticocortical inhibition in human motor cortex. *Journal of Physiology* 1993;471:501–519. [PubMed: 8120818]
- [25]. Lin K-C, Huang Y-H, Hsieh Y-W, Wu C-Y. Potential predictors of motor and functional outcomes after distributed constraint-induced therapy for patients with stroke. *Neurorehabilitation & Neural Repair* 2009;23:336–342. [PubMed: 18984830]
- [26]. Liu Y, Rouiller E. Mechanisms of recovery of dexterity following unilateral lesion of the sensorimotor cortex in adult monkeys. *Exp Brain Res* 1999;128:149–159. [PubMed: 10473753]
- [27]. Mansur CG, Fregni F, Boggio PS, Riberto M, Gallucci-Neto J, Santos CM, Wagner T, Rigonatti SP, Marcolin MA, Pascual-Leone A. A sham stimulation-controlled trial of rTMS of the unaffected hemisphere in stroke patients. *Neurology* 2005;64:1802–1804. [PubMed: 15911819]
- [28]. Mathiowetz V, Volland G, Kashman N, Weber K. Adult norms for the Box and Block Test of manual dexterity. *Am J Occup Ther* 1985;39:386–391. [PubMed: 3160243]
- [29]. Muller JFM, Orekhov Y, Liu Y, Ziemann U. Homeostatic plasticity in human motor cortex demonstrated by two consecutive sessions of paired associative stimulation. *European Journal of Neuroscience* 2007;25:3461–3468. erratum appears in *Eur J Neurosci*. 2007 Aug;26(4):1077. [PubMed: 17553015]
- [30]. Murase N, Duque J, Mazzocchio R, Cohen LG. Influence of interhemispheric interactions on motor function in chronic stroke. *Ann. Neurol* 2004;55:400–409. [PubMed: 14991818]
- [31]. Nowak DA, Grefkes C, Dafotakis M, Eickhoff S, Kust J, Karbe H, Fink GR. Effects of low-frequency repetitive transcranial magnetic stimulation of the contralesional primary motor cortex on movement kinematics and neural activity in subcortical stroke. *Archives of Neurology* 2008;65:741–747. [PubMed: 18541794]
- [32]. Romero JR, Ansel D, Sparing R, Gangitano M, Pascual-Leone A. Subthreshold low frequency repetitive transcranial magnetic stimulation selectively decreases facilitation in the motor cortex. *Clinical Neurophysiology* 2002;113:101–107. [PubMed: 11801430]
- [33]. Takeuchi N, Chuma T, Matsuo Y, Watanabe I, Ikoma K. Repetitive Transcranial Magnetic Stimulation of Contralesional Primary Motor Cortex Improves Hand Function After Stroke. *Stroke* 2005;36:2681–2686. [PubMed: 16254224]
- [34]. Taub E, Crago J, Burgio L, Groomes T, Cook EI, DeLuca S. An operant approach to rehabilitation medicine: Overcoming learned nonuse by shaping. *J Exp Anal Behav* 1994;61:281–293. [PubMed: 8169577]

- [35]. Wu T, Sommer M, Tergau F, Paulus W. Lasting influence of transcranial magnetic stimulation on intracortical excitability in human subjects. *Neurosci Lett* 2000;287:37–40. [PubMed: 10841985]
- [36]. Yousry TA, Schmid UD, Alkadhi H, Schmidt D, Peraud A, Buettner A, Winkler P. Localization of the motor hand area to a knob on the precentral gyrus. A new landmark. *Brain* 1997;120:141–157. [PubMed: 9055804]




Figure 1.

Activation for subject 1 before (pretest) and after (posttest) intervention. Maps show a reduction in active voxels in left (contralateral) sensorimotor cortex (green circles) from pretest to posttest and an increase in active voxels and intensity in right (ipsilesional) sensorimotor cortex (yellow circles).




Figure 2.

Activation for subject 2 before (pretests 1 and 2) and after (posttest) intervention. Maps show a small increase in activation in the left (contralateral) sensorimotor cortex (green circles) from pretest 1 to pretest 2, followed by a reduction at posttest. In the ipsilesional hemisphere, an increase in peri-infarct zone activation is seen at posttest, including in the small amount of surviving sensorimotor cortex (yellow circles).




Figure 3. Excitability of ipsilesional primary motor area in patient 1 before (pretest) and after (posttest) intervention. Each bar represents mean \pm standard deviation of 10 measurements. (MEP= motor evoked potential, * $p=0.029$, ** $p=0.017$)

Table 1

fMRI and Behavioral Results Before and After Five Sessions of 6-Hz Primed Low-Frequency rTMS to Contralesional Primary Motor Area.

Measurement	Subject 1		Subject 2	
	Pretest	Posttest	Pretest 1	Posttest 1
fMRI				
Active Voxels				
Contra SMC	101	59	981	841
Ipsi SMC	322	314	2	85
Wt Avg Intensity				
Contra SMC	3.60	3.66	3.8	3.59
Ipsi SMC	3.52	4.28	2.65	3.59
Behavioral				
AI (%)	-53.0 ± 6.7	-18.5 ± 10.4	30.3 ± 11.9	-47.3 ± 23.1
BBT (blocks)				
	11.3 ± 0.6	9.0 ± 1.7	3.0 ± 1.0	7.0 ± 2.7
Force(N)				
	8.4 ± 1.1	to 10.7 ± 0.4	5.1 ± 1.6	8.3 ± 1.8
MAL				
Amount	0.6	0.4	0.9	1.2
Quality	0.4	0.7	0.9	1.6
TEMPA (s)				
	149.9	164.9	139.5	233.9

fMRI=functional magnetic resonance imaging, Contra=contralesional, Ipsi=Ipsilesional, Wt Avg=Weighted Average, BBT=Box and Block Test, N=Newtons, s=seconds, TEMPA = Test Évaluant la Performance des Membres Supérieurs des Personnes Âgées, MAL=Motor Activity Log