

Review

The endosymbiotic origin, diversification and fate of plastids

Patrick J. Keeling*

*Botany Department, Canadian Institute for Advanced Research, University of British Columbia,
3529-6270 University Boulevard, Vancouver, BC, Canada V6T 1Z4*

Plastids and mitochondria each arose from a single endosymbiotic event and share many similarities in how they were reduced and integrated with their host. However, the subsequent evolution of the two organelles could hardly be more different: mitochondria are a stable fixture of eukaryotic cells that are neither lost nor shuffled between lineages, whereas plastid evolution has been a complex mix of movement, loss and replacement. Molecular data from the past decade have substantially untangled this complex history, and we now know that plastids are derived from a single endosymbiotic event in the ancestor of glaucophytes, red algae and green algae (including plants). The plastids of both red algae and green algae were subsequently transferred to other lineages by secondary endosymbiosis. Green algal plastids were taken up by euglenids and chlorarachniophytes, as well as one small group of dinoflagellates. Red algae appear to have been taken up only once, giving rise to a diverse group called chromalveolates. Additional layers of complexity come from plastid loss, which has happened at least once and probably many times, and replacement. Plastid loss is difficult to prove, and cryptic, non-photosynthetic plastids are being found in many non-photosynthetic lineages. In other cases, photosynthetic lineages are now understood to have evolved from ancestors with a plastid of different origin, so an ancestral plastid has been replaced with a new one. Such replacement has taken place in several dinoflagellates (by tertiary endosymbiosis with other chromalveolates or serial secondary endosymbiosis with a green alga), and apparently also in two rhizarian lineages: chlorarachniophytes and *Paulinella* (which appear to have evolved from chromalveolate ancestors). The many twists and turns of plastid evolution each represent major evolutionary transitions, and each offers a glimpse into how genomes evolve and how cells integrate through gene transfers and protein trafficking.

Keywords: plastids; endosymbiosis; evolution; algae; protist; phylogeny

1. THE ORIGIN OF PLASTIDS: A SINGLE EVENT OF GLOBAL SIGNIFICANCE

Endosymbiosis has played many roles in the evolution of life, but the two most profound effects of this process were undoubtedly the origins of mitochondria and plastids in eukaryotic cells. There are many parallels in how these organelles originated and how their subsequent evolution played out, for example, the reduction of the bacterial symbiont genome and the development of a protein-targeting system (Whatley *et al.* 1979; Douglas 1998; Gray 1999; Gray *et al.* 1999; Gould *et al.* 2008). There are also many differences, however, and one of the more striking is the ultimate fate of the organelle once established: where mitochondria were integrated into the host and the two were seemingly never again separated (Williams & Keeling 2003; van der Giezen *et al.* 2005), plastid evolution has seen many more twists, turns and dead ends (for other reviews that cover various aspects of this history, see Delwiche 1999; McFadden 1999,

2001; Archibald & Keeling 2002; Stoebe & Maier 2002; Palmer 2003; Williams & Keeling 2003; Keeling 2004; Archibald 2005; Keeling *in press*).

Ultimately, mitochondria and plastids (with the small but interesting exception detailed in §2) have each been well established to have evolved from a single endosymbiotic event involving an alpha-proteobacterium and cyanobacterium, respectively (Gray 1999). This conclusion has not come without considerable debate, which has stemmed from several sources. First, the ancient nature of the event makes reconstructing its history difficult because a great deal of change has taken place since the origin of this system, and all of this change masks ancient history. In addition, however, the complexity of subsequent plastid evolution has made for special and less expected problems. Specifically, plastids were originally established in a subset of eukaryotes by a so-called 'primary' endosymbiosis with an ancient cyanobacterial lineage. Once established, primary plastids then spread from that lineage to other eukaryotes by additional rounds of endosymbiosis between two eukaryotes (secondary and tertiary endosymbioses, which are each discussed in detail within their own section below). This led to a very confusing

*pkeeling@interchange.ubc.ca

One contribution of 12 to a Theme Issue 'Evolution of organellar metabolism in unicellular eukaryotes'.

picture of plastid diversity and distribution, as the plastids and their hosts can have different evolutionary histories. Until this was realized, the tendency was, reasonably enough, to treat all photosynthetic lineages as close relatives, which created a long list of paradoxical observations where the plastid of some lineage seemed similar to that of one type of eukaryote, but the host component appeared more similar to another (Dougherty & Allen 1960; Leedale 1967; Brugerolle & Taylor 1977). For example, euglenids had plastids like those of green algae but their cytosolic features were like trypanosomes, which in fact proved to be exactly the case.

Once these added layers of complexity were resolved, or at least understood to exist (Gibbs 1978, 1981; Ludwig & Gibbs 1989), the multiple versus single origin of plastids boiled down to resolving the origin of primary plastids. Primary plastids are surrounded by two bounding membranes and are only found in three lineages. Glaucophytes are a small group of microbial algae with plastids that contain chlorophyll *a*, and are distinguished by the presence of a relict of the peptidoglycan wall that would have been between the two membranes of the cyanobacterial symbiont (Bhattacharya & Schmidt 1997; Steiner & Löffelhardt 2002). Red algae are considerably more diverse and conspicuous than glaucophytes, with 5000–6000 species ranging from tiny, non-flagellated coccoid cells in extreme environments to marine macroalgae that are known to anyone that has walked in the rocky intertidal zone (figure 1). Their plastids also contain chlorophyll *a* and phycobilisomes and are distinguished by the presences of phycoerythrin (Graham & Wilcox 2000). The green algae are also a diverse group and are abundant in both marine and freshwater environments (figure 1), and on dry land where a few green algal lineages have proliferated (e.g. Trentopohales and many Trebouxiophytes that interact with fungi in lichens), as well as giving rise to the land plants, a major lineage with global terrestrial impacts. Their plastids contain chlorophyll *a* and *b* and are also distinguished by the storage of starch within the plastid (Lewis & McCourt 2004).

The plastids in all three of these lineages share a good deal in common with one another, and genome sequences from all three lineages also share several features that are not known in any cyanobacterial lineage, for example, the presence of inverted repeats surrounding the rRNA operon (Palmer 1985; McFadden & Waller 1997). Molecular phylogenies of plastid-encoded genes have consistently shown them to be monophyletic to the exclusion of all currently known cyanobacterial lineages (Delwiche *et al.* 1995; Helmchen *et al.* 1995; Turner *et al.* 1999; Yoon *et al.* 2002; Hagopian *et al.* 2004; Khan *et al.* 2007). Further analysis of plastid-targeted proteins also showed that plastids contain a unique light harvesting complex protein that is not found in any known cyanobacteria (Wolfe *et al.* 1995; Durnford *et al.* 1999). All these observations are most consistent with the monophyly of plastids, but debate persisted, primarily because of analyses of nuclear sequences. In early molecular phylogenies, the three primary plastid lineages were not monophyletic (Bhattacharya *et al.* 1995; Bhattacharya &

Weber 1997), and in some cases gene trees seemed to show well-supported evidence against this monophyly (Stiller *et al.* 2001, 2003; Kim & Graham 2008). The early analyses have not held up, however, and more compellingly analysed large datasets of concatenated genes have most consistently demonstrated the monophyly of the nuclear lineages, and in those analyses with the most data this is recovered with strong support (Moreira *et al.* 2000; Rodriguez-Ezpeleta *et al.* 2005; Hackett *et al.* 2007; Burki *et al.* 2009).

The current consensus is that there is a single lineage, called Plantae or Archaeplastida (Adl *et al.* 2005), in some probably biflagellated heterotrophic ancestor of which the primarily endosymbiotic uptake of a cyanobacterium took place. The cyanobacterium was reduced by loss of genes and their corresponding functions, and also genetically integrated with its host. A complex mechanism for targeting nucleus-encoded proteins to the endosymbiont was progressively established, resulting in the outer and inner membrane complexes today known as translocon outer (TOC) and inner (TIC) chloroplast membranes (McFadden 1999; van Dooren *et al.* 2001; Wickner & Schekman 2005; Hormann *et al.* 2007; Gould *et al.* 2008). The targeted proteins mostly acquired amino terminal leaders called transit peptides, which are recognized by the TOC and used to drag the protein across the membranes, and which are subsequently cleaved in the plastid stroma by a specific peptidase (McFadden 1999; Wickner & Schekman 2005; Hormann *et al.* 2007; Patron & Waller 2007; Gould *et al.* 2008), a system remarkably similar to the protein-targeting system used by mitochondria (Lithgow 2000). This system probably coevolved with the transfer of a few genes to the nucleus, and once the system was established its presence would make it relatively easy for many genes to move to the host genome. Today, plastid genomes are a small fraction of the size of cyanobacterial genomes (Douglas 1998; Douglas & Raven 2003), and there is relatively little diversity in the size and overall structure of the genome, at least in comparison with mitochondrial genomes, and gene content is relatively stable, although many genes are known to have moved to the nucleus independently in multiple lineages (Martin *et al.* 1998; Gould *et al.* 2008). There has been some debate over whether the endosymbiont also donated a substantial number of genes whose products are not now targeted to the plastid (Martin *et al.* 1998, 2002; Martin 1999; Reyes-Prieto *et al.* 2006), but these will not be discussed here.

Once the endosymbiont was established and integrated with its host, the three major lineages of Archaeplastida diverged (figure 2). Again, there is considerable controversy about the branching order of these three groups, but current data lean towards the glaucophytes branching first. Glaucophyte plastids are unique in retaining the peptidoglycan wall between the two membranes, which would seem to support this (Bhattacharya & Schmidt 1997; Steiner & Löffelhardt 2002). However, all three groups have been proposed to be most basal at one time or another (Cavalier-Smith 1982; Kowallik 1997; Martin *et al.* 1998). Molecular phylogenies have not provided a particularly strong answer to this question, but many recent analyses based on large datasets of both nuclear




Figure 1. Diversity of phototrophic eukaryotes and their plastids. Primary plastids are found in a subset of photosynthetic eukaryotes, most conspicuously in green algae ((a) *Ulva*, or sea lettuce) and their close relatives the land plants ((b) *Typha*, or cattail), and in red algae ((c) *Chondracanthus*, or Turkish towel). Secondary plastids are known in many other lineages, including some large multicellular algae such as kelps and their relatives ((f) *Fucus*, a brown alga). In some secondary plastids, the nucleus of the endosymbiotic alga is retained and referred to as a nucleomorph ((d) the nucleomorph from *Partenskyella glossopodia*). In some dinoflagellates, an additional layer of symbiosis, tertiary symbiosis, has made cells of even greater complexity, for example, (e) *Durinskia*, where five different genetically distinct compartments have resulted from endosymbiosis: the host nucleus (red), the endosymbiont nucleus (blue), the endosymbiont plastid (yellow) and mitochondria from both host and endosymbiont (purple). (g) *Chromera velia* is a recently described alga that has shed a great deal of light on the evolution of plastids by secondary endosymbiosis. Image (a) is courtesy of K. Ishida, (e) is courtesy of K. Carpenter and all other images are by the author.

and plastid genes often show glaucophytes branching first, although some also show red algal branching earlier (Moreira *et al.* 2000; Rodriguez-Ezpeleta *et al.* 2005; Hackett *et al.* 2007). It has also been shown that glaucophytes alone retain the ancestral cyanobacterial fructose biphosphate aldolase, and that in both green and red algae the original gene has been replaced by a duplicate of the non-homologous (analogous) nuclear-encoded cytosolic enzyme (Gross *et al.* 1999; Nickol *et al.* 2000). Phylogenetic analysis of the red and green algal genes has shown that the plastid and cytosolic paralogues each form a (weak) clade

including both red and green algal genes (Rogers & Keeling 2003), which suggests that the gene replacement must have been ancestral to both red and green algae, and that the glaucophytes therefore diverged prior to this event, making them the first-diverging lineage of the Archaeplastida (figure 2).

2. PAULINELLA AND THE POSSIBILITY OF A SECOND ORIGIN OF PLASTIDS

All the data outlined above relate to the ultimate origin of plastids in nearly all major lineages of eukaryotes.




Figure 2. (Caption opposite.)

There is, however, a large number of endosymbiotic relationships seemingly based on photosynthesis that are less well understood and vary across the entire spectrum of integration, from passing associations to long term and seemingly well-developed partnerships (e.g. Rumpho *et al.* 2008). Indeed, the line between

what is an organelle and what is an endosymbiont is an arbitrary one. There are a few different, specific criteria that have been argued to distinguish the two, the most common being the genetic integration of the two partners, and the establishment of a protein-targeting system. Most photosynthetic endosymbionts probably

Figure 2. (*Opposite.*) Schematic view of plastid evolution in the history of eukaryotes. The various endosymbiotic events that gave rise to the current diversity and distribution of plastids involve divergences and reticulations whose complexity has come to resemble an electronic circuit diagram. Endosymbiosis events are boxed, and the lines are coloured to distinguish lineages with no plastid (grey), plastids from the green algal lineage (green) or the red algal lineage (red). At the bottom is the single primary endosymbiosis leading to three lineages (glaucophytes, red algae and green algae). On the lower right, a discrete secondary endosymbiotic event within the euglenids led to their plastid. On the lower left, a red alga was taken up in the ancestor of chromalveolates. From this ancestor, haptophytes and cryptomonads (as well as their non-photosynthetic relatives like katablepharids and telonemids) first diverged. After the divergence of the rhizarian lineage, the plastid appears to have been lost, but in two subgroups of Rhizaria, photosynthesis was regained: in the chlorarachniophytes by secondary endosymbiosis with a green alga, and in *Paulinella* by taking up a cyanobacterium (many other rhizarian lineages remain non-photosynthetic). At the top left, the stramenopiles diverged from alveolates, where plastids were lost in ciliates and predominantly became non-photosynthetic in the apicomplexan lineage. At the top right, four different events of plastid replacement are shown in dinoflagellates, involving a diatom, haptophyte, cryptomonad (three cases of tertiary endosymbiosis) and green alga (a serial secondary endosymbiosis). Most of the lineages shown have many members or relatives that are non-photosynthetic, but these have not all been shown for the sake of clarity.

are not integrated with their host at this level, but one case has attracted considerable recent attention for possibly having done so, and this is *P. chromatophora*.

Paulinella is a genus of euglyphid amoeba, which are members of the Cercozoa (Bhattacharya *et al.* 1995). These are testate amoebae with shells built from siliceous scales of great intricacy. Most euglyphids, including most members of the genus *Paulinella*, are non-photosynthetic heterotrophs that feed using granular filopods that emerge from an opening in their test (Johnson *et al.* 1988). However, one species, *P. chromatophora*, has lost its feeding apparatus and instead acquired a cyanobacterial endosymbiont that allows it to live without feeding: each *P. chromatophora* cell contains two kidney-bean-shaped cyanobacterial symbionts, called chromatophores (figure 2), and their division is synchronized with that of the host cell cycle so that each daughter amoeba retains the symbiont (Kies 1974; Kies & Kremer 1979). Early observations led to the suggestion that chromatophores were cyanobacteria related to *Synechococcus*, and molecular phylogenies later confirmed that they are related to the *Synechococcus/Prochlorococcus* lineage (Marin *et al.* 2005; Yoon *et al.* 2006). Early work also demonstrated that the symbiont transferred photosynthate to the amoeba host (Kies & Kremer 1979), altogether suggesting that the symbiont, called a chromatophore, was at least functionally equivalent to a plastid.

There has been a lot of debate over whether the chromatophore should be called an 'organelle' or a 'plastid', or an 'endosymbiont' (Theissen & Martin 2006; Yoon *et al.* 2006; Bhattacharya *et al.* 2007; Bodyl *et al.* 2007). To some extent, this is semantic, but in other ways the distinction in what we call it is important because it does affect the way we think about organelles. If we define 'organelles' in a narrow way, for instance restricted to cases in which a protein-targeting system has evolved, then we will inevitably come to the conclusion that 'organelles' can only originate or integrate in certain ways. All other cellular bodies will be given a different name and will have less impact on our thinking on organellogenesis. It could be beneficial to leave the definition a little more open, as two cells can become highly integrated in more ways than protein targeting. If, for example, an endosymbiont becomes dependent on its host for control over the division and segregation

of the endosymbiont, then one might consider the endosymbiont to be an organelle, which allows us to see the potential variation in the way that organellogenesis can take place, and two cells become integrated.

Returning to the chromatophore, the debate over whether or not it should be called an organelle has been stoked by the complete sequence of its genome (Nowack *et al.* 2008). As previously indicated by phylogenetic analysis of single genes, the genome clearly shows the chromatophore to be a member of the *Synechococcus/Prochlorococcus* lineage (Marin *et al.* 2005; Yoon *et al.* 2006), but rather than the roughly 3 Mbp and 3300 genes common to members of the genus *Synechococcus*, the chromatophore genome is a mere 1 Mbp in size and encodes only 867 genes (Nowack *et al.* 2008). Reduction has mostly taken place by the elimination of whole pathways and functional classes of genes. This reduction is different from that seen in plastids, because not only is it less severe (the chromatophore has more than four times the number of genes encoded by even the largest known plastid), but also in the tendency for whole pathways to be lost or kept: in plastids, most pathways that are retained are incomplete because many or most genes have moved to the nucleus (Nowack *et al.* 2008). This, at face value, suggested that there was no protein targeting or gene transfer (Keeling & Archibald 2008). However, analysis of expressed sequence tags from the *Paulinella* nuclear genome found a copy of *psaE* (Nakayama & Ishida 2009). The gene is phylogenetically related to the *Synechococcus/Prochlorococcus* lineage, and is missing from the chromatophore genome, so is most likely a case of transfer from the symbiont to the host (Nakayama & Ishida 2009). Whether the PsaE protein is targeted back to the chromatophore is not known, and intriguingly there is no evidence for an amino-terminal protein-targeting extension. It is possible that the *psaE* gene is non-functional, functions in the host (which seems unlikely given the gene), or that targeting has evolved by a completely new mechanism, which would not be surprising given it is an independent endosymbiosis. *Paulinella* is a fascinating system that will doubtless receive much more attention, and if it does prove to be a fully integrated 'organelle', determining how its evolution has paralleled that of canonical plastids and how it has differed will both provide valuable comparisons.

3. SECONDARY ENDOSYMBIOSIS AND THE RISE OF PLASTID DIVERSITY

As mentioned previously, primary plastids are found in glaucophytes, red algae and green algae (from which plants are derived). These groups represent a great deal of diversity and are collectively significant ecologically, but they only represent a fraction of eukaryotic phototrophs. Most algal lineages acquired their plastids through secondary endosymbiosis, which is the uptake and retention of a primary algal cell by another eukaryotic lineage (Delwiche 1999; McFadden 2001; Archibald & Keeling 2002; Stoebe & Maier 2002; Palmer 2003; Keeling 2004, *in press*; Archibald 2005; Gould *et al.* 2008). The plastids in most algal lineages can be attributed to this process, namely those of chlorarachniophytes and euglenids, which acquired plastids from green algae, and haptophytes, cryptomonads, heterokonts, dinoflagellates and apicomplexans, which acquired a plastid from a red alga (figure 2). Overall, this secondary spread of plastids had a major impact on eukaryotic diversity, evolution and global ecology (Falkowski *et al.* 2004). Many of the lineages with secondary plastids have grown to dominate primary production in their environment, and collectively they represent a significant fraction of known eukaryotic diversity: it has been estimated that the lineage encompassing all the red algal secondary plastids alone represents over 50 per cent of the presently described protist species (Cavalier-Smith 2004).

Our understanding of how such a process might have taken place is actually somewhat more clear than our understanding of how primary endosymbiosis might have played out, in part because the events were more recent, but more importantly because it happened more than once so parallels and differences can be compared. As was the case for the cyanobacterium in primary endosymbiosis, the secondary endosymbiotic algae progressively degenerated until all that remained in most cases was the plastid, and one or two additional membranes around it. Essentially, no trace of mitochondria, flagella, Golgi, endoplasmic reticulum (ER) or many other cellular features remain in these endosymbionts. In most lineages (see §4 for exceptions), the algal nucleus is completely absent as well, and any genes for plastid proteins that it once encoded have moved once again to the nuclear genome of the secondary host (it seems likely that many other genes also moved to that genome, but this has not been investigated very thoroughly). In general, it is thought that secondary endosymbiosis takes place through endocytosis, so the eukaryotic alga is taken up into a vacuole derived from the host endomembrane system (Delwiche 1999; McFadden 2001; Archibald & Keeling 2002; Keeling 2004; Archibald 2005; Gould *et al.* 2008). After reduction, such a plastid would be predicted to be surrounded by four membranes, corresponding to (from the outside inward) the host endomembrane, the plasma membrane of the engulfed alga and the two membranes of the primary plastid (McFadden 2001; Archibald & Keeling 2002; Keeling 2004; Gould *et al.* 2008). Most secondary plastids are indeed surrounded by four membranes, and there is

abundant evidence that the outermost membrane is derived from the host endomembrane system. Indeed, in cryptomonads, haptophytes and heterokonts, the outermost membrane is demonstrably contiguous with the host ER and nuclear envelope (Gibbs 1981). In other cases, the same situation is inferred from the way proteins are targeted (Foth *et al.* 2003; Gould *et al.* 2008; Bolte *et al.* 2009). Many plastid proteins are encoded in the new host nucleus, and these are targeted to the plastid post-translationally, but the process requires an additional step compared with the analogous process in primary plastids. Recall that targeting in primary plastids typically relies on the recognition of an N-terminal transit peptide by the TOC and TIC complexes on the outer and inner plastid membranes (McFadden 1999; Wickner & Schekman 2005; Hormann *et al.* 2007; Patron & Waller 2007; Gould *et al.* 2008). Secondary plastids are surrounded by additional membranes and are in effect situated within the endomembrane system of the secondary host (as opposed to the primary plastid, which is situated in the cytosol of its host), so any protein expressed in the cytosol is not directly exposed to the plastid outer membrane and could not be imported by the TIC–TOC system alone (McFadden 1999; Gould *et al.* 2008; Bolte *et al.* 2009; Kalanon *et al.* 2009). This has led to an additional leg in the journey from the cytosol to the plastid, and because these plastids are located in the endomembrane of their host, travel to the plastid is initially through the secretory system. As a result, proteins targeted to secondary plastids have a more complex, bipartite leader, which consists of a signal peptide followed by a transit peptide (Patron & Waller 2007). The signal peptide is recognized by the signal recognition particle, which stops translation and directs the protein to the rough ER (RER), where translation resumes and the nascent protein crosses the membrane as it is elongated. How the protein crosses the next membrane is only just emerging in some groups (Hempel *et al.* 2007; Sommer *et al.* 2007; Bolte *et al.* 2009), and it is not yet certain if the same mechanism is used by all secondary plastids, but once exposed to the inner pair of membranes, the transit peptide may be finally recognized by the TOC and TIC complexes and import completed.

The origin of membranes in four-membrane secondary plastids presents few mysteries and corresponds well with what is known about protein trafficking across those membranes. However, in euglenids and dinoflagellates, the secondary plastid is bounded by only three membranes (figure 2). This raised questions as to whether one membrane had been lost, and if so which one, or whether these plastids were derived by a different process. Specifically, it has been proposed that these plastids were acquired by myzocytosis, a feeding strategy whereby a predator attaches to a food cell and sucks its contents into a feeding vacuole, rather than engulfing the whole cell (Schneppf & Deichgraber 1984). Myzocytosis is known in dinoflagellates and euglenids, and such a process could indeed explain the membrane topology, but it remains unclear how an alga taken up by

myzocytosis could survive and divide without its plasma membrane. Moreover, it is now abundantly clear that the three-membrane dinoflagellate plastid is orthologous to the four-membrane plastid in apicomplexans (Moore *et al.* 2008; Keeling *in press*), so the three-membrane topology of dinoflagellate and euglenid plastids must be the result of some other convergent evolutionary pathway (Lukes *et al.* 2009). If the process of plastid origins is the same for three- and four-membrane secondary plastids (as must be the case in dinoflagellates), then which membrane was lost and why? There is no certain answer to this, but by a process of reduction it must have been the second membrane, corresponding to the plasma membrane of the engulfed alga. The rationale for this conclusion is that all other membranes are crossed by a known mechanism in protein targeting, and removing any one of them would have predictable and disastrous effects on trafficking. For example, as the outermost membrane is part of the host endomembrane system and protein trafficking uses the first steps in the secretion pathway to target plastid proteins to this compartment, the loss of this membrane would mean plastid-targeted proteins would be diverted to the secretory pathway. Similarly, the two inner membranes are involved in plastid function, and necessary to transit peptide recognition, so probably cannot be lost. In contrast, no satisfying explanation for why the second membrane is retained has been proposed (and we can therefore more readily imagine losing it), and the mechanism thought to be used to cross it in some groups is not clearly incompatible with loss (Bolte *et al.* 2009). This is not to say that the loss has not impacted targeting, because it has. Intriguingly, the plastid-targeted proteins of both euglenids and dinoflagellates have targeting peptides that are different in some respects from those of other secondary plastids (Sulli *et al.* 1999; Nassoury *et al.* 2003), and they share similar differences (Patron *et al.* 2005; Durnford & Gray 2006), despite having acquired their plastids independently from a green and red alga, respectively.

4. NUCLEOMORPHS

Another exception that has emerged more than once is the retention of a relict nucleus of the secondary endosymbiotic alga, structures called nucleomorphs (Gilson & McFadden 2002; Archibald 2007). In most cases, the secondary endosymbiont nucleus is completely lost, presumably owing to the movement of all genes necessary for the upkeep of the plastid to the nucleus of the new secondary host. In cryptomonads and chlorarachniophytes, however, this algal nucleus has persisted (figures 1 and 2) and has been the focus of much attention. When these tiny structures were first described using transmission electron microscopy (TEM) (Greenwood *et al.* 1977; Hibberd & Norris 1984), the process of secondary endosymbiosis was not appreciated and the distribution of plastids correspondingly difficult to understand. The demonstration that cryptomonad and chlorarachniophyte plastid compartments were associated with a eukaryotic nucleus and residual

cytoplasm with 80S ribosomes (Ludwig & Gibbs 1989; Douglas *et al.* 1991; McFadden *et al.* 1994) was a galvanizing discovery that ushered in widespread acceptance that eukaryote–eukaryote symbiosis was an important part of plastid evolution.

Not surprisingly, attention soon turned to the genomes retained in nucleomorphs. As the cryptomonad plastid is clearly derived from a red alga and the chlorarachniophyte plastid from a green alga, their nucleomorphs must have evolved independently from fully-fledged algal nuclei, but they were soon found to share a number of superficial similarities. In all species examined to date, the nucleomorph genome is composed of three small chromosomes, for a total genome size from as little as 373 kbp to over 650 kbp (Rensing *et al.* 1994; McFadden *et al.* 1997a; Gilson & McFadden 1999, 2002; Gilson *et al.* 2006; Archibald 2007; Silver *et al.* 2007). In nearly all cases, the rRNA operons are found as subtelomeric repeats on all six chromosome ends, although they face in opposite directions in some species (Gilson & McFadden 1996, 2002; Archibald 2007). Between these repeats, the chromosomes are ‘jam-packed’ with genes, the sequenced genomes have gene densities of about one gene per kilobase, the highest known density for a nuclear genome. This tight organization has apparently affected gene expression in nucleomorphs in both cryptomonads and chlorarachniophytes, so that there is now a high frequency of overlapping transcription: in *Guillardia theta*, nearly 100 per cent of characterized transcripts either begin in an upstream gene, terminate within or beyond a downstream gene, or both (Williams *et al.* 2005).

While these similarities certainly suggest that genomes in both lineages have been under similar pressures and constraints or been affected by similar modes of evolution, there are more differences the deeper one digs. Most importantly, there is no real pattern to the actual genes retained in the nucleomorphs. When nucleomorphs were first discovered, it was thought that they might harbour an extensive collection of genes for plastid-targeted proteins, but in the complete genomes sequenced to date, plastid-targeted protein genes are relatively scarce: only 17 in *Bigelowiella natans* (Gilson *et al.* 2006) and 30 in the cryptomonads (Douglas *et al.* 2001; Lane *et al.* 2007) have been identified. Moreover, there is no significant overlap in the identity of these genes, rather they seem to be two random subsets of possible plastid proteins (Gilson *et al.* 2006). Other interesting differences have been found in how these genomes have reacted to whatever process led to their severe reduction and compaction. For example, introns in cryptomonad nucleomorphs are not unusual in size or sequence, but they are extremely rare in number: the *G. theta* genome has only 18 introns (Douglas *et al.* 2001; Williams *et al.* 2005), and the nucleomorph of *Hemiselmis andersenii* has lost them altogether (Lane *et al.* 2007). In contrast, chlorarachniophyte nucleomorph genes are riddled with introns: the *B. natans* genome retains over 800 identified introns, and most seem to be ancient introns conserved with green algae and other chlorarachniophytes (Gilson *et al.* 2006). These introns, however, have dramatically

reduced in size so that all known cases in *B. natans* are between 18 and 21 bp, and the majority are 19 bp, a situation more or less conserved in other chlorarachniophytes (Slamovits & Keeling 2009).

The majority of genes in nucleomorph genomes are housekeeping genes responsible for the maintenance and expression of the genome itself, although in no case is the complement of genes in a nucleomorph genome sufficient for all necessary functions (Douglas *et al.* 2001; Gilson *et al.* 2006; Lane *et al.* 2007). Many genes are inferred to have moved to the host nucleus, and are presumably targeted back to the cytoplasm surrounding the nucleomorph (the periplastid compartment or PPC), or the nucleomorph itself. PPC-targeted proteins are interesting from a protein trafficking perspective, because in most secondary plastids there is little activity between the two pairs of membranes, and probably few genes targeted to this compartment. In cells with a nucleomorph, on the other hand, many genes appear to be targeted to this compartment and, as this is half way to the plastid, how they are targeted is an interesting question that could help resolve how proteins cross the second membrane (see above). The first such genes to be identified were in the cryptomonad, *G. theta*, and the leaders were shown to include a signal peptide and a transit peptide (Gould *et al.* 2006a,b). Intriguingly, the transit peptides share a single common feature, which was the lack of a phenylalanine residue immediately downstream of the signal peptide. This residue is part of a motif common to transit peptides of glaucophytes, red algae and all red-algal-derived plastids (Patron *et al.* 2005; Patron & Waller 2007). Adding a phenylalanine to this position of the PPC-targeted proteins led to their re-targeting to the plastid, suggesting that this position plays an important role in distinguishing PPC proteins from plastid proteins (Gould *et al.* 2006a,b), a function that seems to be directed by a derivative of the ERAD complex (Sommer *et al.* 2007; Bolte *et al.* 2009).

Interestingly, targeting to secondary green plastids does not rely on the F-residue (Patron *et al.* 2005; Patron & Waller 2007), so this information cannot participate even partially in the distinction between PPC- and plastid-targeting. Only two putative PPC-targeted proteins have been identified in chlorarachniophytes, for the translation factors EFL and eIF1 (Gile & Keeling 2008). Once again, the targeting information on these proteins appears to consist of a signal peptide followed by a sequence with all the characteristics of a transit peptide. Interestingly, the only difference between PPC-targeted and plastid-targeted protein leaders is the presence of an acid-rich domain at the C-terminus of the transit peptide (Gile & Keeling 2008). The *B. natans* genome is presently being sequenced, and will presumably yield a long list of PPC-targeted proteins whose targeting information, together with the ability to transform *Lotharella amoebiformis* (Hirakawa *et al.* 2008), will shed additional light on this problem.

The nucleomorphs of cryptomonads and chlorarachniophytes are relatively well-studied components of a eukaryotic endosymbiont, and they have evolved along remarkably similar lines. But this does not

mean such a path is inevitable. Indeed, in the tertiary endosymbiotic partnership between a diatom and dinoflagellates such as *Kryptoperidinium* and *Durinskia* (see below), the endosymbiont nucleus and its genome have not reduced at all, but grossly expanded (Kite *et al.* 1988). These endosymbionts are fully integrated in the host cell cycle, but are poorly studied at the molecular level and it is not known if any genetic exchange has taken place. Because of their enormous size, the nuclei are not typically referred to as nucleomorphs, but in some ways the situation is similar to the arguments about the *Paulinella* chromatophore and whether it should be called a plastid: the interesting point is not so much the name, but that the organelle has followed a different evolutionary path than have those in cryptomonad and chlorarachniophyte endosymbionts.

5. HOW MANY TIMES HAVE PLASTIDS MOVED BETWEEN EUKARYOTES?

We know secondary endosymbiosis has happened on multiple occasions, because both green and red algal endosymbionts are known, but exactly how many times secondary endosymbiosis has taken place has been a subject of ongoing debate.

On the green side, the question is more or less settled. Secondary green algal plastids are known in euglenids and chlorarachniophytes, and there is no strong similarity between the two. Euglenids have three-membrane plastids and store paramylon in the cytosol, whereas chlorarachniophytes have four-membrane plastids with a nucleomorph and store beta-1-3-glucans in the cytosol (McFadden *et al.* 1997b; Leedale & Vickerman 2000; Ishida *et al.* 2007). The hosts are also different, and although the chlorarachniophytes have only recently found a home in the tree of eukaryotes with Cercozoa (Bhattacharya *et al.* 1995; Cavalier-Smith & Chao 1997; Keeling 2001; Nikolaev *et al.* 2004; Burki *et al.* 2009), it was clear early on that euglenids were similar to trypanosomes (Leedale & Vickerman 2000), and not chlorarachniophytes. In keeping with this, early molecular phylogenies did not support a close relationship between the two groups. Nevertheless, it was proposed that their plastids shared a common ancestor, in the so-called Cabozoa hypothesis (Cavalier-Smith 1999). The rationale behind this hypothesis was that secondary endosymbiosis and the evolution of a protein-targeting system in particular is a complex process, and hypotheses for plastid evolution should severely limit the number of times this would have taken place. However, analysis of complete plastid genomes has refuted this proposal, because chlorarachniophytes and euglenid plastid genomes have now been shown to be specifically related to different green algal lineages (Rogers *et al.* 2007a; Turmel *et al.* 2009), and so could not have been derived from a single endosymbiosis (figure 2).

The evolution of secondary plastids from red algae is far more complex and a much greater number of lineages are involved. Secondary red algal plastids are found in cryptomonads, haptophytes, heterokonts, dinoflagellates and apicomplexans. Although these

organisms represent a great deal of plastid diversity (some have nucleomorphs, some are non-photosynthetic, some have three membranes, etc.), and although molecular phylogenies originally grouped neither nuclear nor plastid lineages together, again it was proposed that their plastids arose from a single common endosymbiosis, an idea known as the chromalveolate hypothesis (Cavalier-Smith 1999). In contrast to the Cabozoa hypothesis, data eventually emerged to support the chromalveolate hypothesis and, although still contentious, a variation on this hypothesis is gaining some general acceptance (figure 2). The evidence for this relationship and controversies surrounding it was recently reviewed elsewhere (Keeling *in press*), but as data are emerging quickly and because it does represent a major fraction of plastid diversity, the evidence will be summarized briefly here.

Plastid gene phylogenies have supported the relationship of heterokonts, haptophytes and cryptomonads to the exclusion of what few red algal lineages have been sampled (Yoon *et al.* 2002; Hagopian *et al.* 2004; Khan *et al.* 2007; Rogers *et al.* 2007a), but the strange plastid genomes of apicomplexans and dinoflagellates have essentially excluded them from such analyses. In addition, two nucleus-encoded genes for plastid-targeted proteins, glycerol-3-phosphate dehydrogenase and fructose biphosphate aldolase, have also supported the common origin of chromalveolate plastids because the chromalveolate genes have a unique evolutionary history that differs from homologues in other plastids (Fast *et al.* 2001; Harper & Keeling 2003; Patron *et al.* 2004). Nuclear gene phylogenies have typically not supported a monophyletic chromalveolate (e.g. Kim & Graham 2008), including analyses of some large multi-gene datasets (Patron *et al.* 2004; Burki *et al.* 2007, 2008). However, other large multi-gene analyses (e.g. Hackett *et al.* 2007; Hampl *et al.* 2009; Burki *et al.* 2009) have recovered a monophyletic chromalveolate, with one important provision: that it also includes Rhizaria. In the relatively short time since the monophyly of Rhizaria was discovered (Cavalier-Smith & Chao 1997; Keeling 2001; Archibald *et al.* 2002; Nikolaev *et al.* 2004), they have been considered a supergroup in their own right, but a string of large-scale analyses have consistently shown that they are closely related to alveolates and heterokonts (Burki *et al.* 2007, 2008, 2009; Hackett *et al.* 2007; Hampl *et al.* 2009), and were recently shown to share a novel class of Rab GTPase (Elias *et al.* 2009). If all of these observations are correct, it also means the ancestor of Rhizaria possessed the red algal plastid that is still present in many chromalveolates. This is interesting, not least because two rhizarian lineages are today photosynthetic: chlorarachniophytes and *Paulinella*, but they have acquired their plastids more recently and from different sources (see above). If Rhizaria are derived from an ancestrally photosynthetic chromalveolate, then these two lineages have reverted to phototrophy by entering into new symbioses (figure 2).

Although the single origin of all chromalveolate plastids remains in question, the ancestral state of two large and diverse subgroups has become much more clear in recent years. First, many large

multi-gene phylogenetic analyses, as well as the shared presence of a rare horizontal gene transfer in the plastid genome, have shown that cryptomonads and haptophytes are sisters (Rice & Palmer 2006; Burki *et al.* 2007, 2008, 2009; Hackett *et al.* 2007; Hampl *et al.* 2009; Okamoto *et al.* 2009). At the same time, they have been shown to be related to a number of more poorly studied lineages, including several that are non-photosynthetic (Okamoto & Inouye 2005; Not *et al.* 2007; Cuvelier *et al.* 2008; Burki *et al.* 2009; Okamoto *et al.* 2009). This fast-growing group of increasing importance and diversity has recently been called the Hacrobia (Okamoto *et al.* 2009). The other group where the ancestral state is now well established is the apicomplexans and dinoflagellates. Since the discovery of the apicomplexan plastid, there has been a long-running debate over the ancestry of plastids in these two lineages, and indeed whether the apicomplexan plastid is derived from a red or green alga (Williamson *et al.* 1994; Köhler *et al.* 1997; Roberts *et al.* 1998; Keeling *et al.* 1999; Fast *et al.* 2001; Funes *et al.* 2002; Waller *et al.* 2003). This debate is a difficult problem, because the plastid genomes of these two lineages are both highly derived and share almost no genes in common, so they are nearly impossible to compare directly (Keeling 2008). However, the discovery of *Chromera velia*, the photosynthetic sister to Apicomplexa (figure 1), changes this completely (Moore *et al.* 2008). *Chromera* forms a link between these two unusual lineages so that the initial characterization of molecular data (Moore *et al.* 2008) and now the complete sequence of its plastid genome (J. Janouskovec, A. Horak, M. Obernik, J. Lukes & P. J. Keeling 2009, unpublished data) completely eliminate any basis for the green origin of the apicomplexan plastid, and firmly supports the presence of a plastid in the common ancestor of apicomplexans and dinoflagellates.

6. PLASTID LOSS AND CRYPTIC PLASTIDS

One of the outcomes of the chromalveolate hypothesis has been an increased interest in the process of plastid loss and the prevalence of cryptic, non-photosynthetic plastids in heterotrophic lineages. This is because the early origin of the secondary red algal plastid central to the chromalveolate hypothesis requires that a number of currently non-photosynthetic lineages must have had photosynthetic ancestors. To understand the implications of this aspect of the hypothesis, it is necessary to consider several things carefully, in particular the difference between plastid loss and the loss of photosynthesis, and also how well we actually understand the distribution of cryptic plastids.

The difference between losing photosynthesis and losing a plastid may seem straightforward, but the distinction is surprisingly often ignored. This is problematic, because the available evidence suggests that the likelihoods of the two processes are quite different. Photosynthesis has been lost many times, and there is at least one case of this in nearly all photosynthetic groups (Williams & Hirt 2004; Krause 2008). Land plants have lost photosynthesis at least a dozen times (Nickrent *et al.* 1998), and dinoflagellates

are equally likely to dispense with the process (Saldarriaga *et al.* 2001; Hackett *et al.* 2004a). In many of these lineages, the plastid is readily detectable (e.g. Sepsenwol 1973; Siu *et al.* 1976; Nickrent *et al.* 1998; Williams & Keeling 2003; Krause 2008), whereas in others it has been harder to detect (see below). Plastid loss, on the other hand, could be viewed as an extreme subset of cases where photosynthesis has been lost, and it is an apparently rare subset that is very difficult to demonstrate: non-photosynthetic plastids can be challenging to detect, and showing they are absent is substantially harder again (Williams & Hirt 2004; Krause 2008). No plastid has unambiguously been demonstrated to exist without a genome (but see Nickrent *et al.* 1997), but it remains a possibility (mitochondria have lost their genome many times; Williams & Keeling 2003; van der Giezen *et al.* 2005; Hjort *et al.* 2010); so one could argue that a complete nuclear genome sequence is required to confidently conclude that the cell lacks a cryptic plastid, and all associated plastid-targeted proteins. Moreover, for plastid loss to have occurred, the ancestor of a lineage must have once *had* a plastid. While this may seem trivial, it is an important aspect of the argument in many cases. For example, within the chromalveolates, we can concretely state that there is no plastid in those ciliates and oomycetes with complete genomes (Aury *et al.* 2006; Eisen *et al.* 2006; Tyler *et al.* 2006), but whether their ancestors had a plastid is still open to debate, and we cannot be certain of plastid loss until this is demonstrated clearly (although a case has been made for both groups that relict, endosymbiont-derived genes have been retained; Tyler *et al.* 2006; Reyes-Prieto *et al.* 2008). Conversely, there is strong evidence that the ancestors of katablepharids contained a plastid (Patron *et al.* 2007), but we lack sufficient data from these non-photosynthetic heterotrophs to confidently conclude that a cryptic relict does not still exist in this lineage. Indeed, *Cryptosporidium* is arguably the only lineage of eukaryotes that can be concluded to have lost its plastid outright. It is presently unique among all eukaryotes in that the structural and genomic evidence is available to conclude that this organism does not contain a plastid (Abrahamsen *et al.* 2004; Xu *et al.* 2004), and at the same time the evolutionary evidence is also available to state that its ancestors did contain a plastid (Keeling 2008; Moore *et al.* 2008). This is not to say that this is the only lineage in which plastids have been lost: it is likely that many non-photosynthetic lineages have completely lost plastids, as well as many dinoflagellates with tertiary plastids (see below).

In all remaining non-photosynthetic lineages where some evolutionary argument for a plastid-bearing ancestry has been made, cryptic plastids have either now been found, or the presence or absence of a plastid is simply unknown. Indeed, evidence for cryptic plastids is now emerging in several lineages where phylogenetic relationships had suggested a possible plastid ancestry. The best characterized example is the so-called apicoplast of apicomplexan parasites, where the plastid may have been discovered relatively recently (McFadden *et al.* 1996; Wilson *et al.* 1996; Köhler *et al.* 1997), but is already among the best

studied plastids (for reviews, see Wilson 2002; Ralph *et al.* 2004; Lim & McFadden 2010). In another previously enigmatic lineage of parasites, the Helicosporidia, a similar story has unfolded: while these were previously believed to be related to parasites such as Apicomplexa, molecular phylogenetic analysis showed they are in fact green algae (Tartar *et al.* 2002). This led to the suggestion that they contain a cryptic plastid, the presence of which was confirmed by identifying the plastid genome (Tartar *et al.* 2003; de Koning & Keeling 2006), and several nuclear genes for plastid-targeted proteins (de Koning & Keeling 2004). Despite these advances, the organelle itself has yet to be identified.

A less complete picture is also forming for two non-photosynthetic sister lineages to dinoflagellates. In *Perkinsus*, genes for several plastid-derived proteins have been characterized and there is also evidence for the presence of an actual organelle (Grauvogel *et al.* 2007; Stelter *et al.* 2007; Teles-Grilo *et al.* 2007; Matsuzaki *et al.* 2008). In *Oxyrrhis marina*, another non-photosynthetic sister to dinoflagellates, several plastid-derived genes have also been found, and some of these have leaders which suggest that the proteins are targeted to an as-yet unidentified organelle, although interestingly other plastid-derived proteins no longer appear to be targeted (Slamovits & Keeling 2008).

The functions of these organelles are also known or partially known in many cases, and there are many parallels. Where the genome of a cryptic plastid is known (i.e. those of apicomplexans and helicosporidians), they give few clues as to the function of the organelle, and most of the functionally significant genes are nucleus-encoded genes whose products are targeted to the plastid. The cryptic plastid that is best characterized functionally is the apicoplast, which is involved in synthesis of fatty acids, isoprenoids and haem, although there is some variability between different species (Wilson 2002; Foth & McFadden 2003; Ralph *et al.* 2004; Sato *et al.* 2004; Goodman & McFadden 2008; Gould *et al.* 2008). In *Helicosporidium*, representative genes involved in all these pathways have also been found, as well as genes in several amino acid biosynthetic pathways and genes involved in controlling redox potential (de Koning & Keeling 2004), and a similar complement of putative plastid-targeted proteins has also been found in *Prototheca* (Borza *et al.* 2005), a non-photosynthetic green alga that is closely related to *Helicosporidium*. Smaller numbers of genes are known from *Oxyrrhis* and *Perkinsus*, but they too represent subsets of these pathways (Grauvogel *et al.* 2007; Stelter *et al.* 2007; Teles-Grilo *et al.* 2007; Matsuzaki *et al.* 2008; Slamovits & Keeling 2008), suggesting that these represent the core functions of diverse cryptic plastids.

The genomes of non-photosynthetic plastids are now known from a variety of lineages, and in many cases can be compared with close relatives that are photosynthetic. The genomes of the non-photosynthetic plastids are predictably reduced in size (so far, they are consistently smaller than those of their closest photosynthetic relatives: figures 2 and 3), but they generally retain some or all of the common features




Figure 3. Plastid genome structure variation in photosynthetic and non-photosynthetic lineages. Genomes from the green lineage are on the left, the red lineage on the right and the glaucophyte genome is shown in the centre in blue. Inverted repeats encoding rRNA operons are shown as thickened lines. Numbers with names indicate the per cent of the genome that encodes proteins. Where both are available, a genome from a non-photosynthetic species is shown within that of a photosynthetic relative to show the scale of reduction. In general, plastid genomes map as circles and have an inverted repeat that encodes the ribosomal RNA operon. The major exception to this is the plastid genome of dinoflagellates, which has been reduced in coding capacity and broken down to single gene mini-circles. In some rare cases, the repeat and/or operon has been lost (e.g. in *Helicosporidium* and *Chromera*), or the rRNA operon is encoded in tandem (e.g. in *Euglena*). Non-photosynthetic plastids are greatly reduced in size, but tend to retain the overall structure of their photosynthetic counterparts.

of a plastid genome, such as the inverted repeat or the ribosomal protein operons. As they have lost photosynthesis, one would expect all the genes related to this process to be gone as well, and in several cases they are (Wilson *et al.* 1996; de Koning & Keeling 2006). In other cases, however, some of the genes that have been retained are of interest given the absence of photosynthesis, such as ATP synthase genes in *Prototheca* or rubisco subunits in non-photosynthetic heterokonts and plants (Wolfe & dePamphilis 1997; Knauf & Hachtel 2002; Sekiguchi *et al.* 2002; McNeal *et al.* 2007; Barrett & Freudenstein 2008; Krause 2008).

7. TERTIARY ENDOSYMBIOSIS: A DINOFLAGELLATE ODDITY

In dinoflagellates, another layer of endosymbiotic complexity has been added to the evolutionary history

of plastids: tertiary endosymbiosis. The ancestor of dinoflagellates already had a secondary endosymbiotic plastid of red algal origin; however, in a few dinoflagellate lineages, this plastid is either gone or at least no longer photosynthetically active, and primary production is carried out by a new plastid that has been derived from another lineage with a secondary plastid of red algal origin. So far, dinoflagellates are known to have acquired such tertiary plastids from cryptomonads (Schnepf & Elbrächter 1988), haptophytes (Tengs *et al.* 2000) and diatoms (Dodge 1969; Chesnick *et al.* 1997), and they have also acquired a new secondary endosymbiont from the green algal lineage in one case called 'serial secondary endosymbiosis' (Watanabe *et al.* 1990).

Relatively little is known about the process of tertiary endosymbiosis, and the level of genetic integration appears to be variable. For example, the haptophyte endosymbiont of *Karlodinium* and *Karenia*

has been reduced to a similar extreme as most secondary plastids, so that all that remains is the plastid itself, and perhaps some additional membranes (Tengs *et al.* 2000). There is no evidence for the retention of the original dinoflagellate plastid, and the new plastid is serviced by many nucleus-encoded genes and a plastid-targeting system (Nosenko *et al.* 2006; Patron *et al.* 2006); so in this case the process appears to have completely substituted one fully integrated plastid for another. In contrast, the diatom endosymbiont found in another lineage of dinoflagellates (including the genera *Kryptoperidinium*, *Durinskia* and others) is far less reduced (figure 1): this endosymbiont retains a large nucleus with protein coding genes, a substantial amount of cytosol and even its original mitochondria—a condition unique among eukaryotes (Dodge 1969; Chesnick *et al.* 1997; McEwan & Keeling 2004; Imanian & Keeling 2007). It is unknown whether the dinoflagellate nucleus encodes genes for proteins targeted to the diatom plastid, but it may not be necessary given that the diatom nucleus is present and shows no signs of reduction. Moreover, in at least some of these genera, the original plastid is also thought to have been retained in the form of a three-membrane-bounded eyespot (Dodge 1969). Unlike *Karlodinium* and *Karenia*, where tertiary endosymbiosis led to an outright substitution of organelles, in this case the process has resulted in a considerable degree of redundancy and sub-functionalization (Imanian & Keeling 2007).

Tertiary endosymbiosis also presents another wrinkle in the evolution of plastid proteins because, in these events, a new plastid is introduced into a lineage that is already photosynthetic, or at least had photosynthetic ancestors and might have retained a cryptic plastid. During the integration of the new plastid, if and when a protein-targeting system was established and genes moved from the tertiary endosymbiont nucleus to the host nucleus, there might already be a number of plastid-derived genes in the host nucleus. What happens to the old genes? Are they 'overwritten' by the incoming genes that are better suited to the compartment where they have always functioned, or do some of the dinoflagellate plastid genes survive by re-compartmentalizing to the new plastid? Expressed sequence tag surveys of both *Karlodinium* and *Karenia* show that both kinds of events take place: most plastid-targeted proteins are derived from the haptophyte plastid lineage, and therefore probably came from the tertiary endosymbiont (although see below), but a significant fraction are derived from the dinoflagellate plastid lineage, and therefore have been re-directed to the new plastid (Nosenko *et al.* 2006; Patron *et al.* 2006). This also suggests that the old plastid coexisted with the new one, or plastid-targeted genes might have been lost before they could be targeted to the new plastid. Interestingly, the targeting peptides in these lineages bear little similarity to those of either dinoflagellates or haptophytes, so the integration of the tertiary plastid must have led to upheaval in the protein trafficking system (Patron *et al.* 2006). It is possible that this upheaval allowed dinoflagellate proteins to supplant the incoming haptophyte proteins that were presumably better adapted to the organelle

because both classes of proteins were equally likely to coadapt with the changing trafficking system.

Tertiary endosymbiosis is also important because the events are much less ancient than secondary and primary endosymbiotic plastid origins, and can offer a window into the process that shows not only diverse outcomes, but also perhaps evidence of transient states that have vanished in more ancient events. One of the most interesting of these relates to the shift between a transient host–symbiont association and the fixation of a permanent association. Although it is often characterized as a sudden event, endosymbiotic associations are more probably integrated gradually over long periods of time. It is likely that a host becomes adapted to associations with particular kinds of symbionts, and after a long period of transient associations, which may even include gene transfer and protein targeting, some event took place which made the association permanent (perhaps a shift in the control over cell division of the endosymbiont). The exact order of events was quite probably different in the fixation of different endosymbiotic organelles, but most cases probably involved an intermediate stage based on long-term but non-permanent associations. Although this seems likely, there is no direct evidence for this state in the highly integrated primary and secondary endosymbiotic organelles that remain today. In the case of tertiary endosymbionts, however, important evidence of this critical period does remain in at least two cases. The dinoflagellate hosts with haptophyte plastids are all closely related, as are the dinoflagellate hosts that took up diatoms. However, in both cases, the endosymbionts are derived from different lineages of haptophyte and diatom, respectively. At least two different haptophyte endosymbionts have been fixed, one in *Karlodinium* and a different one in *Karenia* (Gast *et al.* 2007). Similarly, at least three different diatom endosymbionts have been fixed, one centric and two distantly related pinnate diatoms (Horiguchi & Takano 2006; Takano *et al.* 2008). This implies that there was a period where these dinoflagellate lineages were forming transient relationships with particular kinds of algae, and that in multiple subgroups these associations were fixed with different symbionts. From these lineages, we ought to be able to detect events relating to integration that come from the illusive transient period (centric diatom-derived plastid-targeted genes in a lineage with a pennate diatom, for example), but currently there are few data of a comparative nature from these groups.

8. HORIZONTAL TRANSFER OF PLASTID GENES

Endosymbiosis and organelle evolution obviously involve a great deal of movement of genetic information, but the movement of genes between a host and symbiont is only one special variety of horizontal gene transfer. More generally, transfers can also take place between more transiently associated cells, food, vector transmission or perhaps just DNA from the environment (Gogarten *et al.* 2009). Whatever the mode of transfer, it is growing clear that eukaryotes can and have acquired genes from a variety of sources beyond their organelles (Keeling & Palmer 2008),

which raises an interesting question: have the organelles themselves been affected by horizontal gene transfer?

In the case of the mitochondrion of plants, a great deal of transfer has been described, making it perhaps the most promiscuous class of cellular genomes known (Bergthorsson *et al.* 2003, 2004). This means that, for plants at least, the mechanism exists to transfer genes between organelles from distinct species, so what about the plastid? Curiously, analysis of plant plastids not only failed to show the same high level of transfer as found in mitochondria, but revealed no evidence of transfer whatsoever (Rice & Palmer 2006). (One genus of parasitic plants has been found to encode a fragment of the plastid genome from another plant genus, but it is not known to reside in the plastid itself; Park *et al.* 2007.) This is thought to reflect the fact that plant mitochondria fuse when cells fuse, but plastids do not, a reasonable limitation of transfer in plants (Bergthorsson *et al.* 2003, 2004; Rice & Palmer 2006; Richardson & Palmer 2007). However, extending the search to all known plastids revealed a remarkable lack of evidence for transfer in general (Rice & Palmer 2006). Of all known plastid genes and genomes, a good case for horizontal acquisition of genes can only be made for a few genes and several introns. Intron transfers have been found to affect many plastid lineages, and involve several types of introns, including group I introns (Cho *et al.* 1998; Besendahl *et al.* 2000), group II introns (Sheveleva & Hallick 2004) and a subclass of group II introns called group III introns, in this case nested within a group II intron in a situation called 'twintrons'. Twintrons and group III introns were originally thought to be unique to euglenid plastids (Hallick *et al.* 1993), but have been found to have transferred to the cryptomonad *Rhodomonas salina* (Maier *et al.* 1995).

The first transfer of plastid gene sequences to be described was the large and small subunits of rubisco (*rbcS* and *rbcL*). Plastids in green algae, plants and glaucophytes use a cyanobacterium-derived type I rubisco that seems to be native to the plastid. In green plastids, only *rbcL* is encoded in the plastid, and *rbcS* is nucleus-encoded. In contrast, red algal plastid genomes encode both subunits, but they appear to be derived from proteobacteria by horizontal gene transfer (Valentin & Zetsche 1989; Delwiche & Palmer 1996). An early analysis of rubisco sequences revealed this transfer, and further analysis based on hundreds of organelle and bacterial genomes has lent additional support to the original interpretation (Rice & Palmer 2006). The second gene for which strong evidence for horizontal gene transfer exists is *rpl36*, where the copy found in both haptophyte and cryptomonad plastid genomes is clearly a paralogue of the copy found in all other plastids, and seems to be derived from an undefined bacterial lineage (Rice & Palmer 2006). Similarly, a bacterial *dnaX* gene was found in the plastid genome of *R. salina*, but not in any other plastid genome, including that of its close relative *G. theta* (Khan *et al.* 2007).

Plastid genomes may not be likely to acquire genes by horizontal transfer, but this does not mean the plastid proteome is not affected by the process, because

most plastid proteins are encoded in the nucleus. Indeed, one of the first and most famous cases of lateral gene transfer to a eukaryote was the plastid-targeted rubisco in dinoflagellates. Dinoflagellates would have originally used the proteobacterial rubisco acquired by horizontal transfer by the ancestor of red algae, but in dinoflagellates the proteobacterial gene has itself been replaced by another horizontal transfer of a single-subunit type II rubisco from another proteobacterium (Morse *et al.* 1995; Whitney *et al.* 1995; Palmer 1996). The possible impact of horizontal transfer on the plastid proteome has now been explored in large-scale analyses in a few lineages. Analysis of expressed sequence tags from the chlorarachniophyte *B. natans* showed that about 20 per cent of genes for which the phylogeny was resolvable were not evidently derived from a chlorophyte green alga, the source of the secondary plastid in this species (Archibald *et al.* 2003). Instead, several genes were related to red algae or red algal secondary plastids, streptophyte green algae (once again, the rubisco small subunit (SSU) was found to have been transferred), or non-cyanobacterial lineages of bacteria, although the history of transfer of some of these genes is more complex than originally envisioned (Rogers *et al.* 2007b). Similarly, genome-wide analyses of genes for plastid-targeted proteins in several dinoflagellates have found several cases of such transfer in this lineage (Bachvaroff *et al.* 2004; Hackett *et al.* 2004b; Waller *et al.* 2006a,b; Nosenko & Bhattacharya 2007), and a variety of individual gene transfer events have also been recorded in haptophytes and diatoms (Armbrust *et al.* 2004; Obornik & Green 2005; Patron *et al.* 2006). Overall, it appears that some lineages are prone to replacing genes for plastid-targeted proteins, and others are not, perhaps a reflection of mixotrophic or purely autotrophic ancestry, as eating other algae would provide an obvious ongoing source for potential new genes.

9. CONCLUDING REMARKS—THE CONTRASTING HISTORIES OF TWO ORGANELLES

Returning to the contrasts between plastid and mitochondrial evolution raised in the Introduction, the different paths travelled by these organelles through time and eukaryotic biodiversity are now becoming clearer than ever. Mitochondrial evolution has undeniably led to a rich diversity of highly derived organelles, but the evolution of this organelle appears to have been entirely vertical and without dead-ends. In contrast, plastid evolution has resulted in an equally rich diversity of organelles, but has added layers of complexity because plastids have moved between host lineages on a number of occasions—a conservative estimate based on current data would be seven times: two green secondary endosymbioses, one red secondary endosymbiosis, one green serial secondary endosymbiosis and three tertiary endosymbioses. In addition, there is mounting evidence of at least one other independent origin of a 'plastid' in *Paulinella*, and clear evidence for complete and outright plastid loss in at least one (*Cryptosporidium*) and probably many other

lineages. These deviations and dead-ends have no analogue in mitochondrial evolution, where secondary transfer and outright loss have never been observed. This is probably due to a variety of contributing factors: while it is certainly likely that the degree of metabolic integration of the two organelles differs and more highly restricts the evolution of mitochondria, it also seems likely that the original distribution of mitochondria and plastids sets the two organelles up for differing histories. Plastids originated within one of the eukaryotic supergroups, so most eukaryotes would not ancestrally have possessed a plastid. This would, at least intuitively, seem to support the likelihood of secondary endosymbiosis as it suggests that there is some point to the process (i.e. the spread of a useful organelle to new lineages). Mitochondria, in contrast, originated in the common ancestor of all eukaryotes; all eukaryotes ancestrally possessed a mitochondrion. This questions the point of and, by extension, the likelihood of secondary transfer because the only possible recipient cells would already have a homologous organelle. Such a transfer would either replace one organelle with another very similar organelle, or would require that the host lost or substantially reduced its mitochondrion, and then 'recovered' it through secondary endosymbiosis. Even if a mechanism for such an event were known, it vastly reduces the possible range of recipient lineages as reduced mitochondria are a relative rarity overall and tend to be restricted to habitats in which aerobic mitochondria are equally rare, and mitochondrial loss is completely unknown.

I wish to thank Kevin Carpenter and Ken Ishida for providing micrographs, Ken Ishida for sharing unpublished data and Geoff McFadden and Brian Leander for providing valuable comments on figure 2. Work in the Keeling laboratory on plastid evolution is supported by grants from the Natural Sciences and Engineering Research Council of Canada, the Canadian Institutes for Health Research and a grant to the Center for Microbial Diversity and Evolution from the Tula Foundation. P.J.K. is a fellow of the Canadian Institute for Advanced Research and a Senior Scholar of the Michael Smith Foundation for Health Research.

REFERENCES

- Abrahamsen, M. S. *et al.* 2004 Complete genome sequence of the apicomplexan, *Cryptosporidium parvum*. *Science* **304**, 441–445. (doi:10.1126/science.1094786)
- Adl, S. M. *et al.* 2005 The new higher level classification of eukaryotes with emphasis on the taxonomy of protists. *J. Eukaryot. Microbiol.* **52**, 399–451. (doi:10.1111/j.1550-7408.2005.00053.x)
- Archibald, J. M. 2005 Jumping genes and shrinking genomes—probing the evolution of eukaryotic photosynthesis with genomics. *IUBMB Life* **57**, 539–547. (doi:10.1080/15216540500167732)
- Archibald, J. M. 2007 Nucleomorph genomes: structure, function, origin and evolution. *Bioessays* **29**, 392–402. (doi:10.1002/bies.20551)
- Archibald, J. M. & Keeling, P. J. 2002 Recycled plastids: a green movement in eukaryotic evolution. *Trends Genet.* **18**, 577–584. (doi:10.1016/S0168-9525(02)02777-4)
- Archibald, J. M., Longet, D., Pawlowski, J. & Keeling, P. J. 2002 A novel polyubiquitin structure in Cercozoa and Foraminifera: evidence for a new eukaryotic supergroup. *Mol. Biol. Evol.* **20**, 62–66. (doi:10.1093/molbev/msg006)
- Archibald, J. M., Rogers, M. B., Toop, M., Ishida, K. & Keeling, P. J. 2003 Lateral gene transfer and the evolution of plastid-targeted proteins in the secondary plastid-containing alga *Bigeloviella natans*. *Proc. Natl Acad. Sci. USA* **100**, 7678–7683. (doi:10.1073/pnas.1230951100)
- Armbrust, E. V. *et al.* 2004 The genome of the diatom *Thalassiosira pseudonana*: ecology, evolution, and metabolism. *Science* **306**, 79–86. (doi:10.1126/science.1101156)
- Aury, J. M. *et al.* 2006 Global trends of whole-genome duplications revealed by the ciliate *Paramecium tetraurelia*. *Nature* **444**, 171–178. (doi:10.1038/nature05230)
- Bachvaroff, T. R., Concepcion, G. T., Rogers, C. R., Herman, E. M. & Delwiche, C. F. 2004 Dinoflagellate expressed sequence tags data indicate massive transfer of chloroplast genes to the nuclear genome. *Protist* **155**, 65–78. (doi:10.1078/1434461000165)
- Barrett, C. F. & Freudenstein, J. V. 2008 Molecular evolution of *rbcL* in the mycoheterotrophic coralroot orchids (*Corallorhiza* Gagnebin, Orchidaceae). *Mol. Phylogenet. Evol.* **47**, 665–679. (doi:10.1016/j.ympev.2008.02.014)
- Bergthorsson, U., Adams, K. L., Thomason, B. & Palmer, J. D. 2003 Widespread horizontal transfer of mitochondrial genes in flowering plants. *Nature* **424**, 197–201. (doi:10.1038/nature01743)
- Bergthorsson, U., Richardson, A. O., Young, G. J., Goertzen, L. R. & Palmer, J. D. 2004 Massive horizontal transfer of mitochondrial genes from diverse land plant donors to the basal angiosperm *Amborella*. *Proc. Natl Acad. Sci. USA* **101**, 17 747–17 752. (doi:10.1073/pnas.0408336102)
- Besendahl, A., Qiu, Y. L., Lee, J., Palmer, J. D. & Bhattacharya, D. 2000 The cyanobacterial origin and vertical transmission of the plastid tRNA(Leu) group-I intron. *Curr. Genet.* **37**, 12–23. (doi:10.1007/s002940050002)
- Bhattacharya, D. & Schmidt, H. A. 1997 Division Glaucocystophyta. In *Origin of algae and their plastids* (ed. D. Bhattacharya), pp. 139–148. Wien, New York: Springer-Verlag.
- Bhattacharya, D. & Weber, K. 1997 The actin gene of the glaucocystophyte *Cyanophora paradoxa*: analysis of the coding region and introns, and an actin phylogeny of eukaryotes. *Curr. Genet.* **31**, 439–446. (doi:10.1007/s002940050227)
- Bhattacharya, D., Helmchen, T. & Melkonian, M. 1995 Molecular evolutionary analyses of nuclear-encoded small subunit ribosomal RNA identify an independent rhizopod lineage containing the Euglyphidae and the Chlorarachniophyta. *J. Eukaryot. Microbiol.* **42**, 64–68. (doi:10.1111/j.1550-7408.1995.tb01541.x)
- Bhattacharya, D., Archibald, J. M., Weber, A. P. & Reyes-Prieto, A. 2007 How do endosymbionts become organelles? Understanding early events in plastid evolution. *Bioessays* **29**, 1239–1246. (doi:10.1002/bies.20671)
- Bodily, A., Mackiewicz, P. & Stiller, J. W. 2007 The intracellular cyanobacteria of *Paulinella chromatophora*: endosymbionts or organelles? *Trends Microbiol.* **15**, 295–296. (doi:10.1016/j.tim.2007.05.002)
- Bolte, K., Bullmann, L., Hempel, F., Bozarth, A., Zauner, S. & Maier, U. G. 2009 Protein targeting into secondary plastids. *J. Eukaryot. Microbiol.* **56**, 9–15. (doi:10.1111/j.1550-7408.2008.00370.x)
- Borza, T., Popescu, C. E. & Lee, R. W. 2005 Multiple metabolic roles for the nonphotosynthetic plastid of the green alga *Prototheca wickerhamii*. *Eukaryot. Cell* **4**, 253–261. (doi:10.1128/EC.4.2.253-261.2005)

- Brugerolle, G. & Taylor, F. J. R. 1977 Taxonomy, cytology and evolution of the Mastigophora. In *Proc. Fifth Int. Congress of Protozoology* (ed. S. H. Hutner), pp. 14–28. New York, NY: Pace University.
- Burki, F., Shalchian-Tabrizi, K., Minge, M., Skjæveland, A., Nikolaev, S. I., Jakobsen, K. S. & Pawlowski, J. 2007 Phylogenomics reshuffles the eukaryotic supergroups. *PLoS ONE* **2**, e790. (doi:10.1371/journal.pone.0000790)
- Burki, F., Shalchian-Tabrizi, K. & Pawlowski, J. 2008 Phylogenomics reveals a new ‘megagroup’ including most photosynthetic eukaryotes. *Biol. Lett.* **4**, 366–369. (doi:10.1098/rsbl.2008.0224)
- Burki, F. *et al.* 2009 Massive sequencing and phylogenomics of two enigmatic protist lineages reveal Telonemia and Centrohelizoa are related to photosynthetic chromalveolates. *Genom. Biol. Evol.* **1**, 231–238.
- Cavalier-Smith, T. 1982 The origins of plastids. *Biol. J. Linn. Soc.* **17**, 289–306. (doi:10.1111/j.1095-8312.1982.tb02023.x)
- Cavalier-Smith, T. 1999 Principles of protein and lipid targeting in secondary symbiogenesis: euglenoid, dinoflagellate, and sporozoan plastid origins and the eukaryote family tree. *J. Eukaryot. Microbiol.* **46**, 347–366. (doi:10.1111/j.1550-7408.1999.tb04614.x)
- Cavalier-Smith, T. 2004 Chromalveolate diversity and cell megaevolution: interplay of membranes, genomes and cytoskeleton. In *Organelles, genomes and eukaryotic evolution* (eds R. P. Hirt & D. Horner), pp. 71–103. London, UK: Taylor and Francis.
- Cavalier-Smith, T. & Chao, E. E. 1997 Sarcomonad ribosomal RNA sequences, Rhizopod phylogeny, and the origin of Euglyphid Amoeboae. *Arch. Protistenkd.* **147**, 227–236.
- Chesnick, J. M., Hooistra, W. H., Wellbrock, U. & Medlin, L. K. 1997 Ribosomal RNA analysis indicates a benthic pennate diatom ancestry for the endosymbionts of the dinoflagellates *Peridinium foliaceum* and *Peridinium balticum* (Pyrrophyta). *J. Eukaryot. Microbiol.* **44**, 314–320. (doi:10.1111/j.1550-7408.1997.tb05672.x)
- Cho, Y., Qiu, Y. L., Kuhlman, P. & Palmer, J. D. 1998 Explosive invasion of plant mitochondria by a group I intron. *Proc. Natl Acad. Sci. USA* **95**, 14 244–14 249. (doi:10.1073/pnas.95.24.14244)
- Cuvelier, M. L., Ortiz, A., Kim, E., Moehlig, H., Richardson, D. E., Heidelberg, J. F., Archibald, J. M. & Worden, A. Z. 2008 Widespread distribution of a unique marine protistan lineage. *Environ. Microbiol.* **10**, 1621–1634. (doi:10.1111/j.1462-2920.2008.01580.x)
- de Koning, A. P. & Keeling, P. J. 2004 Nucleus-encoded genes for plastid-targeted proteins in *Helicosporidium*: functional diversity of a cryptic plastid in a parasitic alga. *Eukaryot. Cell* **3**, 1198–1205. (doi:10.1128/EC.3.5.1198-1205.2004)
- de Koning, A. P. & Keeling, P. J. 2006 The complete plastid genome sequence of the parasitic green alga, *Helicosporidium* sp. is highly reduced and structured. *BMC Biol.* **4**, 12. (doi:10.1186/1741-7007-4-12)
- Delwiche, C. F. 1999 Tracing the thread of plastid diversity through the tapestry of life. *Am. Nat.* **154**(Suppl.), S164–S177. (doi:10.1086/303291)
- Delwiche, C. F. & Palmer, J. D. 1996 Rampant horizontal transfer and duplication of rubisco genes in eubacteria and plastids. *Mol. Biol. Evol.* **13**, 873–882.
- Delwiche, C. F., Kuhse, M. & Palmer, J. D. 1995 Phylogenetic analysis of *tufA* sequences indicates a cyanobacterial origin of all plastids. *Mol. Phylogenet. Evol.* **4**, 110–128. (doi:10.1006/mpev.1995.1012)
- Dodge, J. D. 1969 Observations on the fine structure of the eyespot and associated organelles in the dinoflagellate *Glenodinium foliaceum*. *J. Cell Sci.* **5**, 479–493.
- Dougherty, E. C. & Allen, M. B. 1960 Is pigmentation a clue to protistan phylogeny? In *Comparative biochemistry of photoreactive systems* (ed. M. B. Allen). New York, NY: Academic Press.
- Douglas, S. E. 1998 Plastid evolution: origins, diversity, trends. *Curr. Opin. Genet. Dev.* **8**, 655–661. (doi:10.1016/S0959-437X(98)80033-6)
- Douglas, A. E. & Raven, J. A. 2003 Genomes at the interface between bacteria and organelles. *Phil. Trans. R. Soc. Lond. B* **358**, 5–18. (doi:10.1098/rstb.2002.1188)
- Douglas, S. E., Murphy, C. A., Spencer, D. F. & Gray, M. W. 1991 Cryptomonad algae are evolutionary chimaeras of two phylogenetically distinct unicellular eukaryotes. *Nature* **350**, 148–151. (doi:10.1038/350148a0)
- Douglas, S. *et al.* 2001 The highly reduced genome of an enslaved algal nucleus. *Nature* **410**, 1091–1016. (doi:10.1038/35074092)
- Durnford, D. G. & Gray, M. W. 2006 Analysis of *Euglena gracilis* plastid-targeted proteins reveals different classes of transit sequences. *Eukaryot. Cell* **5**, 2079–2091. (doi:10.1128/EC.00222-06)
- Durnford, D. G., Deane, J. A., Tan, S., McFadden, G. I., Gantt, E. & Green, B. R. 1999 A phylogenetic assessment of the eukaryotic light-harvesting antenna proteins, with implications for plastid evolution. *J. Mol. Evol.* **48**, 59–68. (doi:10.1007/PL00006445)
- Eisen, J. A. *et al.* 2006 Macronuclear genome sequence of the ciliate *Tetrahymena thermophila*, a model eukaryote. *PLoS Biol.* **4**, e286. (doi:10.1371/journal.pbio.0040286)
- Elias, M., Patron, N. J. & Keeling, P. J. 2009 The RAB family GTPase Rab1A from *Plasmodium falciparum* defines a unique paralog shared by chromalveolates and rhizaria. *J. Eukaryot. Microbiol.* **56**, 348–356. (doi:10.1111/j.1550-7408.2009.00408.x)
- Falkowski, P. G., Katz, M. E., Knoll, A. H., Quigg, A., Raven, J. A., Schofield, O. & Taylor, F. J. 2004 The evolution of modern eukaryotic phytoplankton. *Science* **305**, 354–360. (doi:10.1126/science.1095964)
- Fast, N. M., Kissinger, J. C., Roos, D. S. & Keeling, P. J. 2001 Nuclear-encoded, plastid-targeted genes suggest a single common origin for apicomplexan and dinoflagellate plastids. *Mol. Biol. Evol.* **18**, 418–426.
- Foth, B. J. & McFadden, G. I. 2003 The apicoplast: a plastid in *Plasmodium falciparum* and other apicomplexan parasites. *Int. Rev. Cytol.* **224**, 57–110. (doi:10.1016/S0074-7696(05)24003-2)
- Foth, B. J., Ralph, S. A., Tonkin, C. J., Struck, N. S., Fraunholz, M., Roos, D. S., Cowman, A. F. & McFadden, G. I. 2003 Dissecting apicoplast targeting in the malaria parasite *Plasmodium falciparum*. *Science* **299**, 705–708. (doi:10.1126/science.1078599)
- Funes, S., Davidson, E., Reyes-Prieto, A., Magallón, S., Herion, P., King, M. P. & Gonzalez-Halphen, D. 2002 A green algal apicoplast ancestor. *Science* **298**, 2155. (doi:10.1126/science.1076003)
- Gast, R. J., Moran, D. M., Dennett, M. R. & Caron, D. A. 2007 Kleptoplasty in an Antarctic dinoflagellate: caught in evolutionary transition? *Environ. Microbiol.* **9**, 39–45. (doi:10.1111/j.1462-2920.2006.01109.x)
- Gibbs, S. P. 1978 The chloroplasts of *Euglena* may have evolved from symbiotic green algae. *Can. J. Bot.* **56**, 2883–2889. (doi:10.1139/b78-345)
- Gibbs, S. P. 1981 The chloroplast endoplasmic reticulum: structure, function, and evolutionary significance. *Int. Rev. Cytol.* **72**, 49–99. (doi:10.1016/S0074-7696(08)61194-8)
- Gile, G. H. & Keeling, P. J. 2008 Nucleus-encoded periplastid-targeted EFL in chlorarachniophytes. *Mol. Biol. Evol.* **25**, 1967–1977. (doi:10.1093/molbev/msn147)

- Gilson, P. R. & McFadden, G. I. 1996 The miniaturized nuclear genome of a eukaryotic endosymbiont contains genes that overlap, genes that are cotranscribed, and the smallest known spliceosomal introns. *Proc. Natl Acad. Sci. USA* **93**, 7737–7742. (doi:10.1073/pnas.93.15.7737)
- Gilson, P. R. & McFadden, G. I. 1999 Molecular, morphological and phylogenetic characterization of six chlorarachniophyte strains. *Phycol. Res.* **47**, 7–19. (doi:10.1111/j.1440-1835.1999.tb00278.x)
- Gilson, P. R. & McFadden, G. I. 2002 Jam packed genomes—a preliminary, comparative analysis of nucleomorphs. *Genetica* **115**, 13–28. (doi:10.1023/A:1016011812442)
- Gilson, P. R., Su, V., Slamovits, C. H., Reith, M. E., Keeling, P. J. & McFadden, G. I. 2006 Complete nucleotide sequence of the chlorarachniophyte nucleomorph: nature's smallest nucleus. *Proc. Natl Acad. Sci. USA* **103**, 9566–9571. (doi:10.1073/pnas.0600707103)
- Gogarten, M. B., Gogarten, J. P. & Olendzenski, L. (eds) 2009 *Horizontal gene transfer: genomes in flux*. New York, NY: Humana Press.
- Goodman, C. D. & McFadden, G. I. 2008 Fatty acid synthesis in protozoan parasites: unusual pathways and novel drug targets. *Curr. Pharm. Des.* **14**, 901–916. (doi:10.2174/138161208784041088)
- Gould, S. B., Sommer, M. S., Hadfi, K., Zauner, S., Kroth, P. G. & Maier, U. G. 2006a Protein targeting into the complex plastid of cryptophytes. *J. Mol. Evol.* **62**, 674–681. (doi:10.1007/s00239-005-0099-y)
- Gould, S. B., Sommer, M. S., Kroth, P. G., Gile, G. H., Keeling, P. J. & Maier, U. G. 2006b Nucleus-to-nucleus gene transfer and protein retargeting into a remnant cytoplasm of cryptophytes and diatoms. *Mol. Biol. Evol.* **23**, 2413–2422. (doi:10.1093/molbev/msl113)
- Gould, S. B., Waller, R. F. & McFadden, G. I. 2008 Plastid evolution. *Annu. Rev. Plant Biol.* **59**, 491–517. (doi:10.1146/annurev.arplant.59.032607.092915)
- Graham, L. E. & Wilcox, L. W. 2000 *Algae*. Upper Saddle River, NJ: Prentice-Hall.
- Grauvogel, C., Reece, K. S., Brinkmann, H. & Petersen, J. 2007 Plastid isoprenoid metabolism in the oyster parasite *Perkinsus marinus* connects dinoflagellates and malaria pathogens—new impetus for studying alveolates. *J. Mol. Evol.* **65**, 725–729. (doi:10.1007/s00239-007-9053-5)
- Gray, M. W. 1999 Evolution of organellar genomes. *Curr. Opin. Genet. Dev.* **9**, 678–687. (doi:10.1016/S0959-437X(99)00030-1)
- Gray, M. W., Burger, G. & Lang, B. F. 1999 Mitochondrial evolution. *Science* **283**, 1476–1481. (doi:10.1126/science.283.5407.1476)
- Greenwood, A. D., Griffiths, H. B. & Santore, U. J. 1977 Chloroplasts and cell compartments in Cryptophyceae. *Br. Phycol. J.* **12**, 119.
- Gross, W., Lenze, D., Nowitzki, U., Weiske, J. & Schnarrenberger, C. 1999 Characterization, cloning, and evolutionary history of the chloroplast and cytosolic class I aldolases of the red alga *Galdieria sulphuraria*. *Gene* **230**, 7–14. (doi:10.1016/S0378-1119(99)00059-1)
- Hackett, J. D., Anderson, D. M., Erdner, D. L. & Bhattacharya, D. 2004a Dinoflagellates: a remarkable evolutionary experiment. *Am. J. Bot.* **91**, 1523–1534. (doi:10.3732/ajb.91.10.1523)
- Hackett, J. D., Yoon, H. S., Soares, M. B., Bonaldo, M. F., Casavant, T. L., Scheetz, T. E., Nosenko, T. & Bhattacharya, D. 2004b Migration of the plastid genome to the nucleus in a peridinin dinoflagellate. *Curr. Biol.* **14**, 213–218.
- Hackett, J. D., Yoon, H. S., Li, S., Reyes-Prieto, A., Rummele, S. E. & Bhattacharya, D. 2007 Phylogenomic analysis supports the monophyly of cryptophytes and haptophytes and the association of rhizaria with chromalveolates. *Mol. Biol. Evol.* **24**, 1702–1713. (doi:10.1093/molbev/msm089)
- Hagopian, J. C., Reis, M., Kitajima, J. P., Bhattacharya, D. & de Oliveira, M. C. 2004 Comparative analysis of the complete plastid genome sequence of the red alga *Gracilaria tenuistipitata* var. *liui* provides insights into the evolution of rhodoplasts and their relationship to other plastids. *J. Mol. Evol.* **59**, 464–477. (doi:10.1007/s00239-004-2638-3)
- Hallick, R. B., Hong, L., Drager, R. G., Favreau, M. R., Monfort, A., Orsat, B., Spielmann, A. & Stutz, E. 1993 Complete sequence of *Euglena gracilis* chloroplast DNA. *Nucl. Acids Res.* **21**, 3537–3544. (doi:10.1093/nar/21.15.3537)
- Hapl, V., Hug, L., Leigh, J. W., Dacks, J. B., Lang, B. F., Simpson, A. G. & Roger, A. J. 2009 Phylogenomic analyses support the monophyly of Excavata and resolve relationships among eukaryotic 'supergroups'. *Proc. Natl Acad. Sci. USA* **106**, 3859–3864. (doi:10.1073/pnas.0807880106)
- Harper, J. T. & Keeling, P. J. 2003 Nucleus-encoded, plastid-targeted glyceraldehyde-3-phosphate dehydrogenase (GAPDH) indicates a single origin for chromalveolate plastids. *Mol. Biol. Evol.* **20**, 1730–1735. (doi:10.1093/molbev/msg195)
- Helmchen, T. A., Bhattacharya, D. & Melkonian, M. 1995 Analyses of ribosomal RNA sequences from glaucocystophyte cyanelles provide new insights into the evolutionary relationships of plastids. *J. Mol. Evol.* **41**, 203–210.
- Hempel, F., Bozarth, A., Sommer, M. S., Zauner, S., Przyborski, J. M. & Maier, U. G. 2007 Transport of nuclear-encoded proteins into secondarily evolved plastids. *Biol. Chem.* **388**, 899–906. (doi:10.1515/BC.2007.119)
- Hibberd, D. J. & Norris, R. E. 1984 Cytology and ultrastructure of *Chlorarachnion reptans* (Chlorarachniophyta division nova, Chlorarachniophyceae classis nova). *J. Phycol.* **20**, 310–330. (doi:10.1111/j.0022-3646.1984.00310.x)
- Hirakawa, Y., Kofuji, R. & Ishida, K.-I. 2008 Transient transformation of a chlorarachniophyte alga, *Lotharella amoebiformis* (Chlorarachniophyceae), with *uidA* and *egfp* reporter genes. *J. Phycol.* **44**, 814–820. (doi:10.1111/j.1529-8817.2008.00513.x)
- Hjort, K., Goldberg, A. V., Tsaousis, A. D., Hirt, R. P. & Embley, T. M. 2010 Diversity and reductive evolution of mitochondria among microbial eukaryotes. *Phil. Trans. R. Soc. B* **365**, 713–727. (doi:10.1098/rstb.2009.0224)
- Horiguchi, T. & Takano, Y. 2006 Serial replacement of a diatom endosymbiont in the marine dinoflagellate *Peridinium quinquecorne* (Peridinales, Dinophyceae). *Phycol. Res.* **54**, 193–200. (doi:10.1111/j.1440-1835.2006.00426.x)
- Hormann, F., Soll, J. & Bolter, B. 2007 The chloroplast protein import machinery: a review. *Meth. Mol. Biol.* **390**, 179–193. (doi:10.1007/978-1-59745-466-7_12)
- Imanian, B. & Keeling, P. J. 2007 The dinoflagellates *Durinskia baltica* and *Kryptoperidinium foliaceum* retain functionally overlapping mitochondria from two evolutionarily distinct lineages. *BMC Evol. Biol.* **7**, 172. (doi:10.1186/1471-2148-7-172)
- Ishida, K., Yabuki, A. & Ota, S. 2007 The chlorarachniophytes: evolution and classification. In *Unravelling the algae—the past, present, and future of algal systematics*, vol. 75 (eds J. Brodie & J. Lewis). Boca Raton, FL: CRC Press.
- Johnson, P. W., Hargraves, P. E. & Sieburth, J. M. 1988 Ultrastructure and ecology of *Calycomonas ovalis* Wulff, 1919, (Chrysoophyceae) and its redescription as a testate

- rhizopod, *Paulinella ovalis* n. comb. (Filosea: Euglyphina). *J. Protozool.* **35**, 618–626.
- Kalanon, M., Tonkin, C. J. & McFadden, G. I. 2009 Characterization of two putative protein translocation components in the apicoplast of *Plasmodium falciparum*. *Eukaryot. Cell* **8**, 1146–1154. (doi:10.1128/EC.00061-09)
- Keeling, P. J. 2001 Foraminifera and Cercozoa are related in actin phylogeny: two orphans find a home? *Mol. Biol. Evol.* **18**, 1551–1557.
- Keeling, P. J. 2004 The diversity and evolutionary history of plastids and their hosts. *Am. J. Bot.* **91**, 1481–1493. (doi:10.3732/ajb.91.10.1481)
- Keeling, P. J. 2008 Evolutionary biology: bridge over troublesome plastids. *Nature* **451**, 896–897. (doi:10.1038/451896a)
- Keeling, P. J. In press. Chromalveolates and the evolution of plastids by secondary endosymbiosis. *J. Eukaryot. Microbiol.*
- Keeling, P. J. & Archibald, J. M. 2008 Organelle evolution: what's in a name? *Curr. Biol.* **18**, R345–R347. (doi:10.1016/j.cub.2008.02.065)
- Keeling, P. J. & Palmer, J. D. 2008 Horizontal gene transfer in eukaryotic evolution. *Nat. Rev. Genet.* **9**, 605–618. (doi:10.1038/nrg2386)
- Keeling, P. J., Palmer, J. D., Donald, R. G., Roos, D. S., Waller, R. F. & McFadden, G. I. 1999 Shikimate pathway in apicomplexan parasites. *Nature* **397**, 219–220. (doi:10.1038/16618)
- Khan, H., Parks, N., Kozera, C., Curtis, B. A., Parsons, B. J., Bowman, S. & Archibald, J. M. 2007 Plastid genome sequence of the cryptophyte alga *Rhodomonas salina* CCMP1319: lateral transfer of putative DNA replication machinery and a test of chromist plastid phylogeny. *Mol. Biol. Evol.* **24**, 1832–1842. (doi:10.1093/molbev/msm101)
- Kies, L. 1974 Elektronenmikroskopische Untersuchungen an *Paulinella chromatophora* Lauterborn, einer Thekamöbe mit blau-grünen Endosymbionten (Cyanellen). *Protoplasma* **80**, 69–89. (doi:10.1007/BF01666352)
- Kies, L. & Kremer, B. P. 1979 Function of cyanelles in the Tecamoeba *Paulinella chromatophora*. *Naturewissenschaften* **66**, 578–579. (doi:10.1007/BF00368819)
- Kim, E. & Graham, L. E. 2008 EEF2 analysis challenges the monophyly of Archaeplastida and Chromalveolata. *PLoS ONE* **3**, e2621. (doi:10.1371/journal.pone.0002621)
- Kite, G. C., Rothschild, L. J. & Dodge, J. D. 1988 Nuclear and plastid DNAs from the binucleate dinoflagellates *Glenodinium foliaceum* and *Peridinium balticum*. *Biosystems* **21**, 151–164. (doi:10.1016/0303-2647(88)90008-1)
- Knauf, U. & Hachtel, W. 2002 The genes encoding subunits of ATP synthase are conserved in the reduced plastid genome of the heterotrophic alga *Prototheca wickerhamii*. *Mol. Genet. Genom.* **267**, 492–497. (doi:10.1007/s00438-002-0681-6)
- Köhler, S., Delwiche, C. F., Denny, P. W., Tilney, L. G., Webster, P., Wilson, R. J. M., Palmer, J. D. & Roos, D. S. 1997 A plastid of probable green algal origin in apicomplexan parasites. *Science* **275**, 1485–1489. (doi:10.1126/science.275.5305.1485)
- Kowallik, K. V. 1997 Origin and evolution of chloroplasts: current status and future perspectives. In *Eukaryotism and symbiosis: intertaxonic combination versus symbiotic adaptation* (eds H. E. Schenk, R. G. Herrmann, K. W. Jeon, N. E. Müller & W. Schwemmler), pp. 3–23. Berlin, Germany: Springer.
- Krause, K. 2008 From chloroplasts to 'cryptic' plastids: evolution of plastid genomes in parasitic plants. *Curr. Genet.* **54**, 111–121. (doi:10.1007/s00294-008-0208-8)
- Lane, C. E., van den Heuvel, K., Kozera, C., Curtis, B. A., Parsons, B. J., Bowman, S. & Archibald, J. M. 2007 Nucleomorph genome of *Hemiselmis andersenii* reveals complete intron loss and compaction as a driver of protein structure and function. *Proc. Natl Acad. Sci. USA* **104**, 19 908–19 913. (doi:10.1073/pnas.0707419104)
- Leedale, G. F. 1967 *Euglenoid flagellates*. Englewood Cliffs, NJ: Prentice-Hall.
- Leedale, G. F. & Vickerman, K. 2000 Phylum Euglenozoa. In *An illustrated guide to the Protozoa* (eds J. J. Lee, G. F. Leedale & P. Bradbury), pp. 1135–1185, 2nd edn. Lawrence, KA: Society of Protozoologists.
- Lewis, L. A. & McCourt, R. M. 2004 Green algae and the origin of land plants. *Am. J. Bot.* **91**, 1535–1556. (doi:10.3732/ajb.91.10.1535)
- Lim, L. & McFadden, G. 2010 The evolution, metabolism and functions of the apicoplast. *Phil. Trans. R. Soc. B* **365**, 749–763. (doi:10.1098/rstb.2009.0273)
- Lithgow, T. 2000 Targeting of proteins to mitochondria. *FEBS Lett.* **476**, 22–26. (doi:10.1016/S0014-5793(00)01663-X)
- Ludwig, M. & Gibbs, S. P. 1989 Evidence that nucleomorphs of *Chlorarachnion reptans* (Chlorarachniophyceae) are vestigial nuclei: morphology, division and DNA-DAPI fluorescence. *J. Phycol.* **25**, 385–394. (doi:10.1111/j.1529-8817.1989.tb00135.x)
- Lukes, J., Leander, B. S. & Keeling, P. J. 2009 Cascades of convergent evolution: the corresponding evolutionary histories of euglenozoans and dinoflagellates. *Proc. Natl Acad. Sci. USA* **106**, 9963–9970. (doi:10.1073/pnas.0901004106)
- Maier, U. G., Rensing, S. A., Igloi, G. L. & Maerz, M. 1995 Twintrons are not unique to the *Euglena* chloroplast genome: structure and evolution of a plastome cpn60 gene from a cryptomonad. *Mol. Gen. Genet.* **246**, 128–131. (doi:10.1007/BF00290141)
- Martin, W. 1999 A briefly argued case that mitochondria and plastids are descendants of endosymbionts, but that the nuclear compartment is not. *Proc. R. Soc. Lond. B* **266**, 1387–1395. (doi:10.1098/rspb.1999.0792)
- Martin, W., Stoebe, B., Goremykin, V., Hansmann, S., Hasegawa, M. & Kowallik, K. V. 1998 Gene transfer to the nucleus and the evolution of chloroplasts. *Nature* **393**, 162–165. (doi:10.1038/30234)
- Martin, W. *et al.* 2002 Evolutionary analysis of *Arabidopsis*, cyanobacterial, and chloroplast genomes reveals plastid phylogeny and thousands of cyanobacterial genes in the nucleus. *Proc. Natl Acad. Sci. USA* **99**, 12 246–12 251. (doi:10.1073/pnas.182432999)
- Marin, B., Nowack, E. C. & Melkonian, M. 2005 A plastid in the making: evidence for a second primary endosymbiosis. *Protist* **156**, 425–432. (doi:10.1016/j.protis.2005.09.001)
- Matsuzaki, M., Kuroiwa, H., Kuroiwa, T., Kita, K. & Nozaki, H. 2008 A cryptic algal group unveiled: a plastid biosynthesis pathway in the oyster parasite *Perkinsus marinus*. *Mol. Biol. Evol.* **25**, 1167–1179. (doi:10.1093/molbev/msn064)
- McEwan, M. L. & Keeling, P. J. 2004 HSP90, tubulin and actin are retained in the tertiary endosymbiont genome of *Kryptoperidinium foliaceum*. *J. Eukaryot. Microbiol.* **51**, 651–659. (doi:10.1111/j.1550-7408.2004.tb00604.x)
- McFadden, G. I. 1999 Plastids and protein targeting. *J. Eukaryot. Microbiol.* **46**, 339–346. (doi:10.1111/j.1550-7408.1999.tb04613.x)
- McFadden, G. I. 2001 Primary and secondary endosymbiosis and the origin of plastids. *J. Phycol.* **37**, 951–959. (doi:10.1046/j.1529-8817.2001.01126.x)
- McFadden, G. I. & Waller, R. F. 1997 Plastids in parasites of humans. *Bioessays* **19**, 1033–1040. (doi:10.1002/bies.950191114)

- McFadden, G. I., Gilson, P. R., Hofmann, C. J., Adcock, G. J. & Maier, U. G. 1994 Evidence that an amoeba acquired a chloroplast by retaining part of an engulfed eukaryotic alga. *Proc. Natl Acad. Sci. USA* **91**, 3690–3694. (doi:10.1073/pnas.91.9.3690)
- McFadden, G. I., Reith, M., Munholland, J. & Lang-Unnasch, N. 1996 Plastid in human parasites. *Nature* **381**, 482. (doi:10.1038/381482a0)
- McFadden, G. I., Gilson, P. R., Douglas, S. E., Cavalier-Smith, T., Hofmann, C. J. & Maier, U. G. 1997a Bonsai genomics: sequencing the smallest eukaryotic genomes. *Trends Genet.* **13**, 46–49. (doi:10.1016/S0168-9525(97)01010-X)
- McFadden, G. I., Gilson, P. R. & Hofmann, C. J. B. 1997b *Division Chlorarachniophyta. Origins of algae and their plastids*. Wien, New York: Springer-Verlag.
- McNeal, J. R., Kuehl, J. V., Boore, J. L. & de Pamphilis, C. W. 2007 Complete plastid genome sequences suggest strong selection for retention of photosynthetic genes in the parasitic plant genus. *Cuscuta*. *BMC Plant Biol.* **7**, 57. (doi:10.1186/1471-2229-7-57)
- Moore, R. B. *et al.* 2008 A photosynthetic alveolate closely related to apicomplexan parasites. *Nature* **451**, 959–963. (doi:10.1038/nature06635)
- Moreira, D., Le Guyader, H. & Phillippe, H. 2000 The origin of red algae and the evolution of chloroplasts. *Nature* **405**, 69–72. (doi:10.1038/35011054)
- Morse, D., Salois, P., Markovic, P. & Hastings, J. W. 1995 A nuclear-encoded form II RuBisCO in dinoflagellates. *Science* **268**, 1622–1624. (doi:10.1126/science.7777861)
- Nakayama, T. & Ishida, K. 2009 Another acquisition of a primary photosynthetic organelle is underway in *Paulinella chromatophora*. *Curr. Biol.* **19**, R284–R285. (doi:10.1016/j.cub.2009.02.043)
- Nassoury, N., Cappadocia, M. & Morse, D. 2003 Plastid ultrastructure defines the protein import pathway in dinoflagellates. *J. Cell Sci.* **116**, 2867–2874. (doi:10.1242/jcs.00517)
- Nickol, A. A., Muller, N. E., Bausenwein, U., Bayer, M. G., Maier, T. L. & Schenk, H. E. 2000 *Cyanophora paradoxa*: nucleotide sequence and phylogeny of the nucleus encoded muroplast fructose-1,6-bisphosphate aldolase. *Z. Naturforsch.* **55**, 991–1003.
- Nickrent, D. L., Ouyang, Y., Duff, R. J. & dePamphilis, C. W. 1997 Do nonasterid holoparasitic flowering plants have plastid genomes? *Plant Mol. Biol.* **34**, 717–729. (doi:10.1023/A:1005860632601)
- Nickrent, D. L., Duff, R. J., Colwell, A. E., Wolfe, A. D., Young, N. D., Steiner, K. E. & dePamphilis, C. W. 1998 Molecular phylogenetic and evolutionary studies of parasitic plants. In *Molecular systematics of plants II. DNA sequencing* (eds D. E. Soltis, P. S. Soltis & J. J. Doyle), pp. 211–241. Boston, MA: Kluwer.
- Nikolaev, S. I., Berney, C., Fahrni, J. F., Bolivar, I., Polet, S., Mylnikov, A. P., Aleshin, V. V., Petrov, N. B. & Pawlowski, J. 2004 The twilight of Heliozoa and rise of Rhizaria, an emerging supergroup of amoeboid eukaryotes. *Proc. Natl Acad. Sci. USA* **101**, 8066–8071. (doi:10.1073/pnas.0308602101)
- Nosenko, T. & Bhattacharya, D. 2007 Horizontal gene transfer in chromalveolates. *BMC Evol. Biol.* **7**, 173. (doi:10.1186/1471-2148-7-173)
- Nosenko, T., Lidie, K. L., Van Dolah, F. M., Lindquist, E., Cheng, J. F. & Bhattacharya, D. 2006 Chimeric plastid proteome in the Florida 'red tide' dinoflagellate *Karenia brevis*. *Mol. Biol. Evol.* **23**, 2026–2038. (doi:10.1093/molbev/msl074)
- Not, F., Valentin, K., Romari, K., Lovejoy, C., Massana, R., Tobe, K., Vaultot, D. & Medlin, L. K. 2007 Picobiliphytes: a marine picoplanktonic algal group with unknown affinities to other eukaryotes. *Science* **315**, 253–255. (doi:10.1126/science.1136264)
- Nowack, E. C., Melkonian, M. & Glockner, G. 2008 Chromatophore genome sequence of *Paulinella* sheds light on acquisition of photosynthesis by eukaryotes. *Curr. Biol.* **18**, 410–418. (doi:10.1016/j.cub.2008.02.051)
- Obornik, M. & Green, B. R. 2005 Mosaic origin of the heme biosynthesis pathway in photosynthetic eukaryotes. *Mol. Biol. Evol.* **22**, 2343–2353. (doi:10.1093/molbev/msi230)
- Okamoto, N. & Inouye, I. 2005 The katablepharids are a distant sister group of the Cryptophyta: a proposal for Katablepharidophyta *divisio nova*/Kathablepharida phylum novum based on SSU rDNA and beta-tubulin phylogeny. *Protist* **156**, 163–179. (doi:10.1016/j.protis.2004.12.003)
- Okamoto, N., Chantangsi, C., Horak, A., Leander, B. S. & Keeling, P. J. 2009 Molecular phylogeny and description of the novel katablepharid, *Roombia truncata* gen. et sp. nov., and establishment of the Hacrobia taxon nov. *PLoS One* e7080. (doi:10.1371/journal.pone.0007080)
- Palmer, J. D. 1985 Comparative organization of chloroplast genomes. *Annu. Rev. Genet.* **19**, 325–354. (doi:10.1146/annurev.ge.19.120185.001545)
- Palmer, J. D. 1996 Rubisco surprises in dinoflagellates. *Plant Cell* **8**, 343–345. (doi:10.1105/tpc.8.3.343)
- Palmer, J. D. 2003 The symbiotic birth and spread of plastids: how many times and whodunit? *J. Phycol.* **39**, 1–9. (doi:10.1046/j.1529-8817.2003.02185.x)
- Park, J. M., Manen, J. F. & Schneeweiss, G. M. 2007 Horizontal gene transfer of a plastid gene in the non-photosynthetic flowering plants *Orobanche* and *Phelipanche* (Orobanchaceae). *Mol. Phylogenet. Evol.* **43**, 974–985. (doi:10.1016/j.ympev.2006.10.011)
- Patron, N. J. & Waller, R. F. 2007 Transit peptide diversity and divergence: a global analysis of plastid targeting signals. *Bioessays* **29**, 1048–1058. (doi:10.1002/bies.20638)
- Patron, N. J., Rogers, M. B. & Keeling, P. J. 2004 Gene replacement of fructose-1,6-bisphosphate aldolase (FBA) supports a single photosynthetic ancestor of chromalveolates. *Eukaryot. Cell* **3**, 1169–1175. (doi:10.1128/EC.3.5.1169-1175.2004)
- Patron, N. J., Waller, R. F., Archibald, J. M. & Keeling, P. J. 2005 Complex protein targeting to dinoflagellate plastids. *J. Mol. Biol.* **348**, 1015–1024. (doi:10.1016/j.jmb.2005.03.030)
- Patron, N. J., Waller, R. F. & Keeling, P. J. 2006 A tertiary plastid uses genes from two endosymbionts. *J. Mol. Biol.* **357**, 1373–1382. (doi:10.1016/j.jmb.2006.01.084)
- Patron, N. J., Inagaki, Y. & Keeling, P. J. 2007 Multiple gene phylogenies support the monophyly of cryptomonad and haptophyte host lineages. *Curr. Biol.* **17**, 887–891. (doi:10.1016/j.cub.2007.03.069)
- Ralph, S. A., Van Dooren, G. G., Waller, R. F., Crawford, M. J., Fraunholz, M. J., Foth, B. J., Tonkin, C. J., Roos, D. S. & McFadden, G. I. 2004 Tropical infectious diseases: metabolic maps and functions of the *Plasmodium falciparum* apicoplast. *Nat. Rev. Microbiol.* **2**, 203–216. (doi:10.1038/nrmicro843)
- Rensing, S. A., Goddemeier, M., Hofmann, C. J. & Maier, U. G. 1994 The presence of a nucleomorph hsp70 gene is a common feature of Cryptophyta and Chlorarachniophyta. *Curr. Genet.* **26**, 451–455. (doi:10.1007/BF00309933)
- Reyes-Prieto, A., Hackett, J. D., Soares, M. B., Bonaldo, M. F. & Bhattacharya, D. 2006 Cyanobacterial contribution to algal nuclear genomes is primarily limited to plastid functions. *Curr. Biol.* **16**, 2320–2325. (doi:10.1016/j.cub.2006.09.063)

- Reyes-Prieto, A., Moustafa, A. & Bhattacharya, D. 2008 Multiple genes of apparent algal origin suggest ciliates may once have been photosynthetic. *Curr. Biol.* **18**, 956–962. (doi:10.1016/j.cub.2008.05.042)
- Rice, D. W. & Palmer, J. D. 2006 An exceptional horizontal gene transfer in plastids: gene replacement by a distant bacterial paralog and evidence that haptophyte and cryptophyte plastids are sisters. *BMC Biol.* **4**, 31. (doi:10.1186/1741-7007-4-31)
- Richardson, A. O. & Palmer, J. D. 2007 Horizontal gene transfer in plants. *J. Exp. Bot.* **58**, 1–9. (doi:10.1093/jxb/erl148)
- Roberts, F. *et al.* 1998 Evidence for the shikimate pathway in apicomplexan parasites. *Nature* **393**, 801–805. (doi:10.1038/30718)
- Rodriguez-Ezpeleta, N., Brinkmann, H., Burey, S. C., Roure, B., Burger, G., Löffelhardt, W., Bohnert, H. J., Philippe, H. & Lang, B. F. 2005 Monophyly of primary photosynthetic eukaryotes: green plants, red algae, and glaucophytes. *Curr. Biol.* **15**, 1325–1330. (doi:10.1016/j.cub.2005.06.040)
- Rogers, M. B. & Keeling, P. J. 2003 Lateral gene transfer and re-compartmentalisation of Calvin cycle enzymes in plants and algae. *J. Mol. Evol.* **58**, 367–375. (doi:10.1007/s00239-003-2558-7)
- Rogers, M. B., Gilson, P. R., Su, V., McFadden, G. I. & Keeling, P. J. 2007a The complete chloroplast genome of the chlorarachniophyte *Bigeloviella natans*: evidence for independent origins of chlorarachniophyte and euglenid secondary endosymbionts. *Mol. Biol. Evol.* **24**, 54–62. (doi:10.1093/molbev/msl129)
- Rogers, M. B., Watkins, R. F., Harper, J. T., Durnford, D. G., Gray, M. W. & Keeling, P. J. 2007b A complex and punctate distribution of three eukaryotic genes derived by lateral gene transfer. *BMC Evol. Biol.* **7**, 89. (doi:10.1186/1471-2148-7-89)
- Rumpho, M. E., Worful, J. M., Lee, J., Kannan, K., Tyler, M. S., Bhattacharya, D., Moustafa, A. & Manhart, J. R. 2008 Horizontal gene transfer of the algal nuclear gene *psbO* to the photosynthetic sea slug *Elysia chlorotica*. *Proc. Natl Acad. Sci. USA* **105**, 17 867–17 871. (doi:10.1073/pnas.0804968105)
- Saldarriaga, J. F., Taylor, F. J. R., Keeling, P. J. & Cavalier-Smith, T. 2001 Dinoflagellate nuclear SSU rRNA phylogeny suggests multiple plastid losses and replacements. *J. Mol. Evol.* **53**, 204–213. (doi:10.1007/s002390010210)
- Sato, S., Clough, B., Coates, L. & Wilson, R. J. 2004 Enzymes for heme biosynthesis are found in both the mitochondrion and plastid of the malaria parasite *Plasmodium falciparum*. *Protist* **155**, 117–125. (doi:10.1078/1434461000169)
- Schnepf, E. & Deichgraber, G. 1984 'Myzocytosis,' a kind of endocytosis with implications to compartmentalization in endosymbiosis. *Naturwissenschaften* **71**, 218–219. (doi:10.1007/BF00490442)
- Schnepf, E. & Elbrächter, M. 1988 Cryptophyte-like double membrane-bound chloroplast in the dinoflagellate, *Dinophysis* Ehrenb.: evolutionary, phylogenetic and toxicological implications. *Bot. Acta.* **101**, 196–203.
- Sekiguchi, H., Moriya, M., Nakayama, T. & Inouye, I. 2002 Vestigial chloroplasts in heterotrophic stramenopiles *Pteridomonas danica* and *Ciliophrys infusionum* (Dictyochophyceae). *Protist* **153**, 157–167. (doi:10.1078/1434-4610-00094)
- Sepsenwol, S. 1973 Leucoplast of the cryptomonad *Chilomonas paramecium*. Evidence for presence of a true plastid in a colorless flagellate. *Exp. Cell Res.* **76**, 395–409. (doi:10.1016/0014-4827(73)90392-3)
- Sheveleva, E. V. & Hallick, R. B. 2004 Recent horizontal intron transfer to a chloroplast genome. *Nucl. Acids Res.* **32**, 803–810. (doi:10.1093/nar/gkh225)
- Silver, T. D., Koike, S., Yabuki, A., Kofuji, R., Archibald, J. M. & Ishida, K. 2007 Phylogeny and nucleomorph karyotype diversity of chlorarachniophyte algae. *J. Eukaryot. Microbiol.* **54**, 403–410. (doi:10.1111/j.1550-7408.2007.00279.x)
- Siu, C., Swift, H. & Chiang, K. 1976 Characterization of cytoplasmic and nuclear genomes in the colorless alga *Polytoma*. I. Ultrastructural analysis of organelles. *J. Cell. Biol.* **69**, 352–370. (doi:10.1083/jcb.69.2.352)
- Slamovits, C. H. & Keeling, P. J. 2008 Plastid-derived genes in the nonphotosynthetic alveolate *Oxyrrhis marina*. *Mol. Biol. Evol.* **25**, 1297–1306. (doi:10.1093/molbev/msn075)
- Slamovits, C. H. & Keeling, P. J. 2009 Evolution of ultra-small spliceosomal introns in highly reduced nuclear genomes. *Mol. Biol. Evol.* **26**, 1699–1705. (doi:10.1093/molbev/msp081)
- Sommer, M. S., Gould, S. B., Lehmann, P., Gruber, A., Przyborski, J. M. & Maier, U. G. 2007 Der1-mediated preprotein import into the periplastid compartment of chromalveolates? *Mol. Biol. Evol.* **24**, 918–928. (doi:10.1093/molbev/msm008)
- Steiner, J. M. & Löffelhardt, W. 2002 Protein import into cyanelles. *Trends Plant Sci.* **7**, 72–77. (doi:10.1016/S1360-1385(01)02179-3)
- Stelter, K., El-Sayed, N. M. & Seeber, F. 2007 The expression of a plant-type ferredoxin redox system provides molecular evidence for a plastid in the early dinoflagellate *Perkinsus marinus*. *Protist* **158**, 119–130. (doi:10.1016/j.protis.2006.09.003)
- Stiller, J. W., Riley, J. & Hall, B. D. 2001 Are red algae plants? A critical evaluation of three key molecular data sets. *J. Mol. Evol.* **52**, 527–539.
- Stiller, J. W., Reel, D. C. & Johnson, J. C. 2003 A single origin of plastids revisited: convergent evolution in organellar genome content. *J. Phycol.* **39**, 95–105. (doi:10.1046/j.1529-8817.2003.02070.x)
- Stoebe, B. & Maier, U. G. 2002 One, two, three: nature's tool box for building plastids. *Protoplasma* **219**, 123–130. (doi:10.1007/s007090200013)
- Sulli, C., Fang, Z., Muchhal, U. & Schwartzbach, S. D. 1999 Topology of *Euglena* chloroplast protein precursors within endoplasmic reticulum to Golgi to chloroplast transport vesicles. *J. Biol. Chem.* **274**, 457–463. (doi:10.1074/jbc.274.1.457)
- Takano, Y., Hansen, G., Fujita, D. & Horiguchi, T. 2008 Serial replacement of diatom endosymbiont in two freshwater dinoflagellates, *Peridiniopsis* spp. (Peridinales, Dinophyceae). *Phycologia* **47**, 41–53. (doi:10.2216/07-36.1)
- Tartar, A., Boucias, D. G., Adams, B. J. & Becnel, J. J. 2002 Phylogenetic analysis identifies the invertebrate pathogen *Helicosporidium* sp. as a green alga (Chlorophyta). *Int. J. Syst. Evol. Microbiol.* **52**, 273–279.
- Tartar, A., Boucias, D. G., Adams, B. J. & Becnel, J. J. 2003 Comparison of plastid 16S rDNA (*rrn16*) from *Helicosporidium* spp.: evidence supporting the reclassification of Helicosporidia as green algae (Chlorophyta). *Int. J. Syst. Evol. Microbiol.* **53**, 1719–1723. (doi:10.1099/ijs.0.02559-0)
- Teles-Grilo, M. L., Tato-Costa, J., Duarte, S. M., Maia, A., Casal, G. & Azevedo, C. 2007 Is there a plastid in *Perkinsus atlanticus* (Phylum Perkinsozoa)? *Eur. J. Protistol.* **43**, 163–167. (doi:10.1016/j.ejop.2007.02.002)
- Tengs, T., Dahlberg, O. J., Shalchian-Tabrizi, K., Klaveness, D., Rudi, K., Delwiche, C. F. & Jakobsen, K. S. 2000

- Phylogenetic analyses indicate that the 19'hexanoyloxy-fucoanthin-containing dinoflagellates have tertiary plastids of haptophyte origin. *Mol. Biol. Evol.* **17**, 718–729.
- Theissen, U. & Martin, W. 2006 The difference between organelles and endosymbionts. *Curr. Biol.* **16**, R1016–R1017. Author reply R1017–8. (doi:10.1016/j.cub.2006.11.020)
- Turmel, M., Gagnon, M. C., O'Kelly, C. J., Otis, C. & Lemieux, C. 2009 The chloroplast genomes of the green algae *Pyramimonas*, *Monomastix*, and *Pycnococcus* shed new light on the evolutionary history of prasino-phytes and the origin of the secondary chloroplasts of euglenids. *Mol. Biol. Evol.* **26**, 631–648. (doi:10.1093/molbev/msn285)
- Turner, S., Pryer, K. M., Miao, V. P. & Palmer, J. D. 1999 Investigating deep phylogenetic relationships among cyanobacteria and plastids by small subunit rRNA sequence analysis. *J. Eukaryot. Microbiol.* **46**, 327–338. (doi:10.1111/j.1550-7408.1999.tb04612.x)
- Tyler, B. M. *et al.* 2006 *Phytophthora* genome sequences uncover evolutionary origins and mechanisms of pathogenesis. *Science* **313**, 1261–1266. (doi:10.1126/science.1128796)
- Valentin, K. & Zetsche, K. 1989 The genes of both subunits of ribulose-1,5-bisphosphate carboxylase constitute an operon on the plastome of a red alga. *Curr. Genet.* **16**, 203–209. (doi:10.1007/BF00391478)
- van der Giezen, M., Tovar, J. & Clark, C. G. 2005 Mitochondrion-derived organelles in protists and fungi. *Int. Rev. Cytol.* **244**, 175–225.
- van Dooren, G. G., Schwartzbach, S. D., Osafune, T. & McFadden, G. I. 2001 Translocation of proteins across the multiple membranes of complex plastids. *Biochim. Biophys. Acta* **1541**, 34–53. (doi:10.1016/S0167-4889(01)00154-9)
- Waller, R. F., Keeling, P. J., van Dooren, G. G. & McFadden, G. I. 2003 Comment on 'A green algal apicoplast ancestor'. *Science* **301**, 49. (doi:10.1126/science.1083647)
- Waller, R. F., Patron, N. J. & Keeling, P. J. 2006a Phylogenetic history of plastid-targeted proteins in the peridinin-containing dinoflagellate *Heterocapsa triquetra*. *Int. J. Syst. Evol. Microbiol.* **56**, 1439–1447. (doi:10.1099/ijs.0.64061-0)
- Waller, R. F., Slamovits, C. H. & Keeling, P. J. 2006b Lateral gene transfer of a multigene region from cyanobacteria to dinoflagellates resulting in a novel plastid-targeted fusion protein. *Mol. Biol. Evol.* **23**, 1437–1443. (doi:10.1093/molbev/msl008)
- Watanabe, M. M., Suda, S., Inouye, I., Sawaguchi, I. & Chihara, M. 1990 *Lepidodinium viride* gen et sp. nov. (Gymnodiniales, Dinophyta), a green dinoflagellate with a chlorophyll *a*- and *b*-containing endosymbiont. *J. Phycol.* **26**, 741–751. (doi:10.1111/j.0022-3646.1990.00741.x)
- Whatley, J. M., John, P. & Whatley, F. R. 1979 From extracellular to intracellular: the establishment of mitochondria and chloroplasts. *Proc. R. Soc. Lond. B* **204**, 165–187. (doi:10.1098/rspb.1979.0020)
- Whitney, S. M., Shaw, D. C. & Yellowlees, D. 1995 Evidence that some dinoflagellates contain a ribulose-1,5-bisphosphate carboxylase/oxygenase related to that of the alpha-proteobacteria. *Proc. R. Soc. Lond. B* **259**, 271–275. (doi:10.1098/rspb.1995.0040)
- Wickner, W. & Schekman, R. 2005 Protein translocation across biological membranes. *Science* **310**, 1452–1456. (doi:10.1126/science.1113752)
- Williams, B. A. & Hirt, R. P. 2004 RACE and RAGE cloning in parasitic microbial eukaryotes. *Meth. Mol. Biol.* **270**, 151–172.
- Williams, B. A. P. & Keeling, P. J. 2003 Cryptic organelles in parasitic protists and fungi. *Adv. Parasitol.* **54**, 9–67. (doi:10.1016/S0065-308X(03)54001-5)
- Williams, B. A. P., Slamovits, C. H., Patron, N. J., Fast, N. M. & Keeling, P. J. 2005 A high frequency of overlapping gene expression in compacted eukaryotic genomes. *Proc. Natl Acad. Sci. USA* **102**, 10 936–10 941. (doi:10.1073/pnas.0501321102)
- Williamson, D. H., Gardner, M. J., Preiser, P., Moore, D. J., Rangachari, K. & Wilson, R. J. 1994 The evolutionary origin of the 35 kb circular DNA of *Plasmodium falciparum*: new evidence supports a possible rhodophyte ancestry. *Mol. Gen. Genet.* **243**, 249–252.
- Wilson, R. J. 2002 Progress with parasite plastids. *J. Mol. Biol.* **319**, 257–274. (doi:10.1016/S0022-2836(02)00303-0)
- Wilson, R. J. M. I. *et al.* 1996 Complete gene map of the plastid-like DNA of the malaria parasite *Plasmodium falciparum*. *J. Mol. Biol.* **261**, 155–172. (doi:10.1006/jmbi.1996.0449)
- Wolfe, A. D. & dePamphilis, C. W. 1997 Alternate paths of evolution for the photosynthetic gene *rbcL* in four non-photosynthetic species of *Orobanche*. *Plant Mol. Biol.* **33**, 965–977. (doi:10.1023/A:1005739223993)
- Wolfe, G. R., Cunningham, F. X., Durnford, D. G., Green, B. R. & Gantt, E. 1995 Evidence for a common origin of chloroplasts with light-harvesting complexes of different pigmentation. *Nature* **367**, 566–568. (doi:10.1038/367566a0)
- Xu, P. *et al.* 2004 The genome of *Cryptosporidium hominis*. *Nature* **431**, 1107–1112. (doi:10.1038/nature02977)
- Yoon, H. S., Hackett, J. D., Pinto, G. & Bhattacharya, D. 2002 A single, ancient origin of the plastid in the Chromista. *Proc. Natl Acad. Sci. USA* **99**, 15 507–15 512. (doi:10.1073/pnas.242379899)
- Yoon, H. S., Reyes-Prieto, A., Melkonian, M. & Bhattacharya, D. 2006 Minimal plastid genome evolution in the *Paulinella* endosymbiont. *Curr. Biol.* **16**, R670–R672. (doi:10.1016/j.cub.2006.08.018)