



Published in final edited form as:

Bioorg Med Chem. 2010 January 15; 18(2): 508. doi:10.1016/j.bmc.2009.12.018.

2-Dialkynyl Derivatives of (N)-Methanocarba Nucleosides: “Clickable” A₃ Adenosine Receptor-Selective Agonists

Dilip K. Tosh^a, Moshe Chinn^a, Lena S. Yoo^a, Dong Wook Kang^b, Hans Luecke^b, Zhan-Guo Gao^a, and Kenneth A. Jacobson^{a,*}

^aMolecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland 20892

^bGene Regulation Group, Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland 20892

Abstract

We modified a series of (N)-methanocarba nucleoside 5-uronamides to contain dialkyne groups on an extended adenine C2 substituent, as synthetic intermediates leading to potent and selective A₃ adenosine receptor (AR) agonists. The proximal alkyne was intended to promote receptor recognition, and the distal alkyne reacted with azides to form triazole derivatives (click cycloaddition). Click chemistry was utilized to couple an octadiynyl A₃AR agonist to azido-containing fluorescent, chemically reactive, biotinylated, and other moieties with retention of selective binding to the A₃AR. A bifunctional thiol-reactive crosslinking reagent was introduced. The most potent and selective novel compound was a 1-adamantyl derivative (K_i 6.5 nM), although some of the click products had K_i values in the range of 200–400 nM. Other potent, selective derivatives (K_i at A₃AR in nM) were intended as possible receptor affinity labels: 3-nitro-4-fluorophenyl (10.6), α -bromophenacyl (9.6), thiol-reactive isothiazolone (102), and arylisothiocyanate (37.5) derivatives. The maximal functional effects in inhibition of forskolin-stimulated cAMP were measured, indicating that this class of click adducts varied from partial to full A₃AR agonist compared to other widely used agonists. Thus, this strategy provides a general chemical approach to linking potent and selective A₃AR agonists to reporter groups of diverse structure and to carrier moieties.

Keywords

G protein-coupled receptor; purines; azide; structure activity relationship; radioligand binding

Introduction

The A₃ adenosine receptor (AR), a G protein-coupled receptor (GPCR), is found in myocytes, astrocytes, neurons, neutrophils, eosinophils, and other cell types.¹ Both A₃AR agonists and antagonists are proposed for the treatment of cancer and inflammatory diseases.^{2–6} An

*Corresponding author: kajacobs@helix.nih.gov, Molecular Recognition Section, Bldg. 8A, Rm. B1A-19, NIH, NIDDK, LBC, Bethesda, MD 20892-0810, Tel.: 301-496-9024, Fax: 301-480-8422.

Supporting information available: An alternate synthetic procedure the synthesis of compound **32** and NMR spectral characterization of compounds **9**, **11**, and **15** are available.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

antiischemic effect of A₃AR agonists also suggests their use in protection of skeletal muscle and cardiac muscle.⁷

The crystallographic structure of the human (h) A_{2A}AR has been determined, but a directly determined structure is lacking for the A₃AR.⁸ We recently used the coordinates of the A_{2A}AR structure as a template for homology modeling of the A₃AR and have predicted specific interactions of the ligand with the receptor protein.⁹ Hypotheses for agonist binding at the A₃AR have been supported using site-directed mutagenesis and reengineering of the putative binding site to recognize tailored nucleoside ligands (neoreceptors).¹⁰ Thus, structural insights into recognition in the A₃AR binding site have been gained even in the absence of an X-ray structure. The 2 and 6 positions of the adenine ring are most amenable to chain derivatization without sterically interfering in the binding process.

Molecular probes for the A₃AR containing a rigid bicyclo[3.1.0]hexane ring system of the North conformation ((N)-methanocarba) in place of the freely twisting ribose moiety were recently reported.^{9,11,12} Amide-linked fluorescent **1** and biotin-containing probes **2** and **3** of high A₃AR affinity and selectivity were designed using a functionalized congener approach based on stepwise chain extension designed to preserve or enhance receptor affinity (Chart 1).¹³

The present study introduces 2-dialkynyl groups into selective A₃AR ligands of the structural class of the (N)-methanocarba-5'*N*-methyluronamides. The proximal alkyne was intended to promote receptor recognition.⁹ Earlier studies established the enhancing effect in recognition at both A_{2A} and A₃ARs of introducing an alkynyl group adjacent to the purine C2.¹⁴ The distal alkyne was shown to react selectively with alkyl or aryl azides by click cycloaddition to form triazole derivatives of the general structure **4**.¹⁵ Furthermore, an optimized *N*⁶-(3-chlorobenzyl) group, which favors A₃AR selectivity in human, mouse, and rat, was included in the nucleoside structures.¹² Fluorescent, chemically reactive, biotinylated, and other moieties were incorporated by this chemical route, leading to selective A₃AR agonists.

Results

Chemical Synthesis

We have introduced terminal alkynyl groups on the C2 position substituents of (N)-methanocarba nucleoside derivatives to serve as cross-linking sites on the A₃AR agonists, such as for the introduction of reporter groups (R¹ in general structure **4** of Chart 1). Prior to introduction of a terminal alkyne for coupling of the nucleosides to reporter groups, the structure activity relationship (SAR) of this series of *N*⁶-(3-chlorobenzyl)-(N)-methanocarba-5'*N*-methyluronamide derivatives was explored in detail.^{9,11} Thus, here we started with a nucleoside pharmacophore that was already optimized for activation of the A₃AR. Only the adenine C2 position was structurally modified, and all modifications were intended for click chemistry.

The synthetic route to the small molecular 5'*N*-methyluronamide (N)-methanocarba 2-alkynyl triazole-containing derivatives **10** – **23** is shown in Scheme 1. The synthesis of the 2',3'-protected dialkynyl intermediates **6** and **7** was performed using a Sonogashira coupling¹⁶ on the corresponding 2-iodo intermediate **5**. After deprotection of the 2',3'-hydroxyl groups to provide nucleosides **8** and **9**, compounds **10** – **23** were obtained using the Cu(I)-catalyzed 2 + 3 cyclization reaction of the terminal acetylene group with an appropriate azide.¹⁵ The reactions were generally selective for the terminal alkyne, but reaction with 4-isothiocyanatophenylazide **24** initially produced the disubstituted product **25** as the major product isolated (Scheme 2). It is well known that copper-catalyzed click reactions direct the formation of only one regioisomer, i.e. the 1,4-regioisomer is formed (out of the two possibilities).^{17,18} In the case

of compound **25**, although there is a possibility of two regioisomers, we obtained only one isomer. Based on the regiochemistry of a single click reaction, the reaction on the internal alkyne moiety was assumed to occur from its less hindered face. Use of the Cu(I)-stabilizing catalyst tris[(1-benzyl-1H-1,2,3-triazol-4-yl)methyl] amine (TBTA)¹⁹ in this reaction provided the desired monosubstituted analogue **26**.

Compounds **15** and **17** were intended as model compounds for the amide coupling of amine functionalized congeners such as **16** to alkyl carboxylic carriers. Compound **18** was included in order to probe steric tolerance at the distal position.

The fluorescent squaraine-rotaxane derivative **22** was prepared from a commercial reactive azide, of undisclosed structure, which fluoresces strongly at 680 nm.^{20,21} Such rotaxane derivatives have the following advantages over other small fluorophores: improved chemical and photochemical stability, sharp absorption and emission bands, and stability over the pH range from 2–12. They are typically much brighter in fluorescence than the Alexa dyes.²² The elemental composition of this proprietary dye moiety, but not the full chemical structure, was revealed by the supplier, and the integrity of the synthetic product **22** was demonstrated by high resolution mass spectroscopy. The expected molecular weight of the click product of the 1:1 reaction with dialkyne **9** was obtained.

An isothiazolone azide derivative **32** designed as a bifunctional linker for thiol reactivity was synthesized as shown in Scheme 3. This served as the precursor for the adenosine derivative **19**. The synthesis of isothiazolone derivatives from 3-benzoylpropionic acid was reported by Tsolomitis *et al.*²³ An amide coupling reaction between **27** and *p*-toluidine formed 4-oxo-4-phenyl-*N-p*-tolyl-butylamide **28**, which was treated with an excess of thionyl chloride to give 5-benzoyl-2-*p*-tolylisothiazol-3-one **29**. Debenzylation of **29** afforded 2-*p*-tolylisothiazol-3-one **30** in the presence of 10% aqueous sodium hydroxide in benzene, along with dithietane derivatives as by-products. Compound **30** was transformed with *N*-bromosuccinimide in the presence of a catalytic amount of benzoyl peroxide in carbon tetrachloride to the bromomethyl derivative **31**, which upon treatment with sodium azide afforded 2-(4-azidomethylphenyl) isothiazol-3-one **32**.

Quantification of Pharmacological Activity

Binding assays at three hAR subtypes were carried out on the alkyne and triazole derivatives using standard radioligands^{24–26} and membrane preparations from Chinese hamster ovary (CHO) cells (A₁ and A₃) or human embryonic kidney (HEK293) cells (A_{2A}) stably expressing a hAR subtype (Table 1).^{27,28} The previously reported molecular probes (**1** – **3**) were used for comparison in the biological assays.

The two homologous dialkyne intermediates **8** and **9** were equipotent with K_i values in A₃AR binding of 24–29 nM. The selectivity of both intermediates in comparison to the A₁ and A_{2A}ARs was >400 and roughly 300-fold, respectively. Following the click reaction with either aryl or alkyl azides, considerable affinity and selectivity at the A₃AR were preserved. Thus, the main pharmacophore maintained its receptor recognition function in the triazole-extended series. The 3-nitro-4-fluorophenyl adducts **10** and **11** were roughly equipotent to the dialkyne precursors. The A₃AR affinity of the *p*-substituted phenyl adducts **12**, **13**, **14**, and **26** varied depending on the aryl substituent. The order of potency depending on the 4 position substitution was: bromoacetyl **14** > isothiocyanate **26** > amino **12** > carboxy **13**. Compound **14** displayed a K_i value of 9.6 nM. The disubstituted click product **25** was highly potent in binding to the A₃AR, with a K_i value of 8.8 nM. Thus, in spite of the added steric bulk, it was surprisingly 4-fold more potent than the monosubstituted isothiocyanate derivative **26**.

Among adducts of alkyl azides, a short chain acetamidoalkyl derivative **15** was moderately potent with a K_i value of 22 nM, and this ethyl derivative was 4-fold more potent than a higher butyl homologue **17**. Compound **15** was 109-fold selective for the A_3AR in comparison to the $A_{2A}AR$ (Figure 1A). The free aminobutyl precursor derivative **16** displayed only moderate affinity at the A_3AR . The sterically bulky 1-adamantyl adduct **18** was highly potent in binding with a K_i value of 6.5 nM at the A_3AR and 500-fold selectivity in comparison to the $A_{2A}AR$ (Figure 1B).

A thiol-reactive derivative **19** displayed only moderate affinity at the A_3AR . The two derivatives of biotin **20** and **21** having unbranched chains and a fluorescent derivative of Alexa Fluor 488 **23** were relatively weak in binding to the A_3AR with K_i values in the range of 200 – 400 nM. The longer biotin derivative **21** contained a tetraethylene glycol spacer, which apparently did not enhance affinity. The rotaxane derivative **22** bound to the A_3AR receptor with a K_i value of 239 nM.

The click products were tested in a functional assay at the A_3AR (inhibition of forskolin-stimulated cAMP production^{29,30} in A_3AR -expressing CHO cells) as shown in Table 2. The (N)-methanocarba derivatives displayed A_3AR agonist properties with varying degrees of maximal inhibition of cAMP production at 10 μ M. In general, the degree of inhibition was similar to or less than that of 5'-*N*-ethylcarboxamidoadenosine (NECA), taken as a reference standard. Therefore, some of these derivatives, e.g., 4-fluoro-3-nitrophenyl **11**, bromophenacyl **14**, acetamidoethyl **15**, and rotaxane **22** derivatives, were highly efficacious agonists. Compounds that displayed intermediate (50–90%) efficacies were: **9**, **13**, **16**, **17**, **19**, and **25**. Agonists of lower efficacy (<50%) were the short dialkyne derivative **8** and the aminophenyl **12**, isothiocyanate **26**, and 1-adamantyl **18** triazole derivatives. The affinity of compounds **20**, **21**, and **23** was so low that a 10 μ M test concentration in the cAMP assay was not sufficient to ensure full receptor occupancy. The corresponding efficacy values for two A_3AR agonists, *N*⁶-(3-iodobenzyl)-5'-*N*-methylcarboxamidoadenosine (IB-MECA) and its 2-chloro analogue Cl-IB-MECA, which are currently in Phase II clinical trials, are included for comparison.¹ Both are full agonists in this assay.

This series of (N)-methanocarba nucleosides is known to be much weaker in interaction with the $A_{2B}AR$ than with other subtypes.¹² We have tested selected nucleosides at 10 μ M for agonist activity in the stimulation of cAMP accumulation in CHO cells stably expressing the human $A_{2B}AR$. The percent stimulation was: 100% (NECA), 30.2% (**11**), 19.4%, (**15**), and 36.5% (**18**).

Discussion

Click chemistry as a means of assembling complex ligands and for introducing structural diversity is finding increasing application in biological systems.¹⁵ With respect to the ARs, a series of triazole derivatives of adenosine was previously prepared using click chemistry, resulting in A_3AR selective agonists, partial agonists, and antagonists.³¹ The triazole ring was directly attached at the 2 position of adenine, which caused wide variation of the receptor affinity and relative efficacy of the analogues. However, in the present study, the triazole was incorporated at a more distal position of an elongated and flexible 2-adenine substituent. This resulted in a greater degree of preservation of the agonist properties in this series than in the previous study, in which some derivatives became antagonists.³¹ Thus, we have used the azide/alkyne cycloaddition reaction to easily synthesize a wide range of biologically active molecules. The click reaction was intended for linking adenosine functionalized congeners to other moieties and carriers, such as reporter groups, chemically reactive groups, and macromolecular carriers. Other studies of biologically-active small molecules have used click cycloaddition reactions to incorporate reporter groups, such as fluorescent dyes and biotin.^{32,33}

The small molecule adducts in the present series were not uniformly potent as A₃AR selective agonists. Rather, only certain derivatives displayed exceptionally high affinity at this subtype (K_i <20 nM): **14**, **18**, and **25**. Two different lengths of dialkynyl chains were compared in binding at the A₃AR. The octadiynyl derivative **9** appeared to be better suited than the shorter homologue **8** for binding to this receptor when coupled to an aryl azide (cf. **10** and **11**). Therefore, the octadiynyl chain was used in subsequent derivatives. A study by Seela and coworkers also utilized octadiynyl-derivatized nucleosides that were incorporated into oligonucleotides for click reactions that took place exclusively at the distal alkynyl group.³⁴ Compounds **10**, **14**, **19**, and **26** were designed as potential affinity labels of the A₃AR, but the chemical irreversibility of binding was not tested in the present study. The use of isothiocyanates, 3-isothiazolones, and other electrophilic derivatives to affinity label biopolymers has been described.^{13,35}

In conclusion, the most potent and selective novel compound was a 1-adamantyl derivative (K_i 6.5 nM), which suggested the existence of a hydrophobic binding pocket in this region of the receptor. Curiously, this compound proved to be a partial agonist of the A₃AR with only 27.8% of the maximal efficacy in comparison to NECA. Various other click products were in the K_i range of 200–400 nM. Other potent, selective derivatives (K_i at A₃AR in nM) were intended as possible receptor affinity labels: 3-nitro-4-fluorophenyl derivative **11** (10.6), α -bromophenacyl **14** (9.6), thiol-reactive isothiazolone **19** (102), and arylisothiocyanate **26** (37.5). The maximal functional effects in inhibition of forskolin-stimulated cAMP were measured, indicating that this class of click adducts varied from partial to full A₃AR agonist compared to other widely used agonists. Thus, this strategy provides a general chemical approach to linking potent and selective A₃AR agonists to reporter groups of diverse structure and to carrier moieties.

One disadvantage of the standard cycloaddition reaction is that cuprous ions are required. This reactant might not be compatible with the full range of GPCR ligands desired to be coupled to carriers and is certainly not useful in cell systems. To overcome this drawback, Bertozzi and colleagues have explored cyclooctyne derivatives that are substrates for “copper-free” click chemistry.³⁶ These groups provide greater compatibility with living systems and allow covalent coupling to azido groups on biopolymers and other biomolecules. We are currently extending these results to application of cyclooctynyl derivatives of A₃AR agonists to conjugation using copper-free click chemistry.

Experimental Section

Chemical Synthesis

¹H NMR spectra were obtained with a Varian Gemini 300 spectrometer. When using D₂O was used as a solvent, the chemical shifts are expressed as relative ppm from HOD (4.80 ppm).

The purity of the final nucleotide derivatives were determined using a Hewlett–Packard 1100 HPLC equipped with a Zorbax Eclipse 5 mm XDB-C18 analytical column (250 × 4.6 mm; Agilent Technologies Inc, Palo Alto, CA), using a linear gradient solvent system: 5 mM TBAP (tetrabutylammonium dihydrogenphosphate)–CH₃CN from 80:20 to 40:60 in 20 min with a flow rate of 1 mL/min. Peaks were detected by UV absorption (254 nm) using a diode array detector. All derivatives tested for biological activity were shown to be at least 97% pure using this analytical HPLC system.

High-resolution mass measurements were performed on a Micromass/Waters LCT Premier Electrospray Time of Flight (TOF) mass spectrometer coupled with a Waters HPLC system. Unless noted otherwise, reagents and solvents were purchased from Sigma-Aldrich (St. Louis,

MO). Solutions of the nucleoside analogues in DMSO (5 mM) were prepared for biological testing and stored at -20°C .

The squaraine-rotaxane azide derivative SRfluor® 680 Azide was obtained from Molecular Targeting Technologies, Inc. (West Chester, PA). Alexa Fluor 488 azide and biotin(PEG)4 azide were purchased from Invitrogen Corp. (Carlsbad, CA). DMEM/F12 medium and 1 M Tris-HCl (pH 7.5) were purchased from Mediatech, Inc. (Herndon, VA). Unless noted otherwise, reagents and solvents were purchased from Sigma-Aldrich (St. Louis, MO).

(1'S, 2'R, 3'S, 4'S, 5'S)-4'-[6-(3-Chlorobenzylamino)-2-(1,6-heptadiynyl)-9H-purin-9-yl]-(1'S, 2'R, 3'S, 4'S, 5'S)-4'-[6-(3-Chlorobenzylamino)-2-(1,7-octadiynyl)-9H-purin-9-yl]-2',3'-O-isopropylidenebicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (7)

To a solution of compound **5** (440 mg, 0.73 mmol) in anhydrous DMF (12 mL), Pd(PPh₃)₄ (92 mg, 0.08 mmol), CuI (30.5 mg, 0.16 mmol), 1,7-octadiyne (1.0 mL, 8.01 mmol) and then triethylamine (0.22 mL, 1.6 mmol) was added. The reaction mixture was heated at 60 °C for overnight. Solvent was evaporated under vacuum and the residue was purified on flash silica gel column chromatography (CH₂Cl₂:MeOH = 70:1) to give the compound **7** (352 mg, 83%) as foamy syrup. ¹H NMR (CD₃OD, 300 MHz) δ 8.11 (s, 1H), 7.43 (s, 1H), 7.26–7.33 (m, 3H), 5.74 (d, *J* = 7.2 Hz, 1H), 5.01 (s, 1H), 4.83 (m, 1H), 2.87 (s, 3H), 2.53 (t, *J* = 6.9 Hz, 2H), 2.34–2.31 (m, 3H), 2.10–2.15 (m, 1H), 1.71–1.84 (m, 4H), 1.54–1.57 (m, 4H), 1.40 (t, *J* = 5.4 Hz, 1H), 1.29 (s, 3H). HRMS calculated for C₃₁H₃₄ClN₆O₃ (M + H)⁺: 573.2381; found 573.2397.

2',3'-O-isopropylidenebicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (6)

Compound **6** (81%) was synthesized from **5** following same procedure as for compound **7**. ¹H NMR (CD₃OD, 300 MHz) δ 8.12 (s, 1H), 7.45 (s, 1H), 7.28–7.34 (m, 3H), 5.76 (d, *J* = 6.9 Hz, 1H), 5.02 (s, 1H), 4.85–4.87 (m, 1H), 2.88 (s, 3H), 2.64 (t, *J* = 6.9 Hz, 2H), 2.41–2.46 (m, 2H), 2.31 (t, *J* = 2.7, 1H), 2.12–2.17 (m, 1H), 1.85–1.94 (m, 2H), 1.56 (m, 4H), 1.40–1.44 (m, 1H), 1.30 (s, 3H). HRMS calculated for C₃₀H₃₂ClN₆O₃ (M + H)⁺: 559.3046; found 559.3085.

(1'S, 2'R, 3'S, 4'S, 5'S)-4'-[6-(3-Chlorobenzylamino)-2-(1,6-heptadiynyl)-9H-purin-9-yl]-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (8)

Compound **8** (86%) was synthesized from **6** following same procedure as for compound **9**. ¹H NMR (CD₃OD, 300 MHz) δ 8.07 (s, 1H), 7.42 (s, 1H), 7.25–7.42 (m, 3H), 5.01 (d, *J* = 6.9 Hz, 1H), 4.84–4.87 (m, 1H), 3.98 (d, *J* = 6.6 Hz, 1H), 2.86 (s, 3H), 2.59 (t, *J* = 7.2 Hz, 2H), 2.36–2.44 (m, 2H), 2.28 (t, *J* = 2.4 Hz, 1H), 2.06–2.10 (m, 1H), 1.79–1.88 (m, 3H), 1.34–1.39 (m, 1H). HRMS calculated for C₂₇H₂₈ClN₆O₃ (M + H)⁺: 519.1911; found 519.1912.

(1'S, 2'R, 3'S, 4'S, 5'S)-4'-[6-(3-Chlorobenzylamino)-2-(1,7-octadiynyl)-9H-purin-9-yl]-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (9)

To a solution of compound **7** (350 mg, 0.61 mmol) in methanol (7 mL), 10% trifluoromethane sulfonic acid was added and heated at 70 °C for 6 h. Solvent was evaporated and the residue was purified on flash silica gel chromatography (CH₂Cl₂:MeOH = 40:1) to give the compound **9** (295 mg, 91%) as a syrup. ¹H NMR (CD₃OD, 300 MHz) δ 8.08 (s, 1H), 7.43 (s, 1H), 7.28–7.33 (m, 3H), 5.02 (d, *J* = 6.6 Hz, 1H), 4.80–4.82 (m, 1H), 4.00 (dd, *J*₁ = 1.2 Hz, *J*₂ = 5.7 Hz, 1H), 2.87 (s, 3H), 2.51 (t, *J* = 6.9 Hz, 1H), 2.22–2.30 (m, 3H), 2.07–2.10 (m, 1H), 1.86 (t, *J* = 5.1 Hz, 1H), 1.70–1.81 (m, 4H), 1.35–1.40 (m, 1H). HRMS calculated for C₂₈H₃₀ClN₆O₃ (M + H)⁺: 533.2068; found 533.2082.

(1S,2R,3S,4R,5S)-4'-(6-(3-Chlorobenzylamino)-2-(6-(1-(4-fluoro-3-nitrophenyl)-1H-1,2,3-triazol-4-yl)pent-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (10)

Compound **10** (89%) was synthesized from **8** following same procedure as for compound **11**. ¹H NMR (CD₃OD, 300 MHz) δ 8.52–8.56 (m, 2H), 8.18–8.22 (m, 1H), 8.08 (s, 1H), 7.59–7.65 (m, 1H), 7.41 (s, 1H), 7.25–7.32 (m, 3H), 5.04 (d, *J* = 5.4 Hz, 1H), 4.80–4.83 (m, 1H), 3.99 (dd, *J*₁ = 0.9 Hz, *J*₂ = 5.7 Hz, 1H), 3.04 (t, *J* = 7.5 Hz, 2H), 2.87 (s, 3H), 2.60 (t, *J* = 6.6 Hz, 2H), 2.07–2.17 (m, 3H), 1.86 (t, *J* = 5.1 Hz, 1H), 1.38–1.45 (m, 2H), 0.89–0.96 (m, 1H). HRMS calculated for C₃₃H₃₁ClFN₁₀O₅ (M + H)⁺: 701.2151; found 701.2172.

(1S,2R,3S,4R,5S)-4'-(6-(3-Chlorobenzylamino)-2-(6-(1-(4-fluoro-3-nitrophenyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (11)

To a mixture of compound **9** (34 mg, 0.063 mmol) and 4-fluoro-3-nitro-phenyl azide (16.2 mg, 0.088 mmol) in in THF/H₂O 3:1 (2 mL), was added freshly prepared 1M sodium ascorbate (51 μL, 0.05 mmol) followed by 7.5% aqueous copper sulfate pentahydrate solution (42 μL, 0.012 mmol) and stirred for over night at room temperature. Solvent was evaporated and the residue was purified on flash silica gel column chromatography (CH₂Cl₂:MeOH = 45:1) to give the clicked product **11** (42 mg, 94%) as a syrup. ¹H NMR (CD₃OD, 300 MHz) δ 8.57–8.58 (m, 1H), 8.49 (s, 1H), 8.20–8.25 (m, 1H), 8.09 (s, 1H), 7.60–7.67 (m, 1H), 7.40 (s, 1H), 7.20–7.32 (m, 3H), 5.03 (d, *J* = 6.6 Hz, 1H), 4.84–4.87 (m, 1H), 3.99 (d, *J* = 6.3 Hz, 1H), 2.91 (t, *J* = 7.5 Hz, 2H), 2.87 (s, 3H), 2.56 (t, *J* = 6.9 Hz, 2H), 1.96–2.11 (m, 3H), 1.75–1.87 (m, 3H), 1.31–1.40 (m, 2H). HRMS calculated for C₃₄H₃₃ClFN₁₀O₅ (M + H)⁺: 715.2308; found 715.2347.

(1S,2R,3S,4R,5S)-4'-(6-(3-Chlorobenzylamino)-2-(6-(1-(4-amino-phenyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (12)

To a mixture of compound **9** (4.46 mg, 0.008 mmol) and 4-amino-phenylazide (2 mg, 0.011 mmol) in a mixture of *t*-butanol (0.5 mL) and water (0.5 mL), was added TBTA (1 mg, 0.001 mmol) and freshly prepared sodium ascorbate (8.3 μL, 0.008 mmol) followed by copper sulfate (8.3 μL, 0.003 mmol). The reaction mixture was stirred at room temperature for overnight, solvent was evaporated and the residue was purified on flash silica gel column chromatography (CH₂Cl₂:MeOH = 30:1) to give the compound **12** (4 mg, 72%) as a syrup. ¹H NMR (CD₃OD, 300 MHz) δ 8.08 (s, 1H), 8.07 (s, 1H), 7.23–7.41 (m, 6H), 6.74 (d, *J* = 8.7 Hz, 2H), 5.01 (d, *J* = 6.9 Hz, 1H), 4.78–4.82 (m, 1H), 3.96 (d, *J* = 6.0 Hz, 1H), 2.82–2.86 (m, 5H), 2.53 (t, *J* = 6.9 Hz, 2H), 2.06–2.16 (m, 2H), 1.93–1.98 (m, 2H), 1.71–1.87 (m, 4H), 1.29–1.39 (m, 1H). HRMS calculated for C₃₄H₃₅ClN₁₀O₃Na (M + Na)⁺: 689.2480; found 689.2465.

(1S,2R,3S,4R,5S)-4'-(6-(3-Chlorobenzylamino)-2-(6-(1-(4-carboxyl-phenyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (13)

Compound **13** (81%) was synthesized from **9** following same procedure as for compound **12**. ¹H NMR (CD₃OD, 300 MHz) δ 8.39 (s, 1H), 8.09–8.14 (m, 3H), 7.81–7.84 (m, 2H), 7.22–7.44 (m, 4H), 5.04 (d, *J* = 6.6 Hz, 1H), 4.83–4.85 (m, 1H), 3.99 (d, *J* = 6.6 Hz, 1H), 2.89–2.94 (m, 5H), 2.57 (t, *J* = 6.6 Hz, 2H), 1.98–2.14 (m, 4H), 1.77–1.90 (m, 4H), 1.38–1.43 (m, 1H). HRMS calculated for C₃₅H₃₃ClN₉O₅ (M - H)⁺: 694.2293; found 694.2311.

(1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-4-(2-(6-(4-(2-bromoacetyl)phenyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (14)

Compound **14** (79%) was synthesized from **9** following same procedure as for compound **11**. ¹H NMR (CD₃OD, 300 MHz) δ 8.43 (s, 1H), 8.33 (s, 1H), 8.07–8.15 (m, 4H), 7.21–7.42 (m, 4H), 5.01 (d, *J* = 6.6 Hz, 1H), 4.77–4.82 (m, 1H), 4.70 (s, 2H), 3.96 (d, *J* = 6.9 Hz, 1H), 2.86–2.92 (m, 5H), 2.41 (t, *J* = 5.1 Hz, 2H), 1.93–2.10 (m, 4H), 1.74–1.87 (m, 4H), 1.35–1.40 (m, 1H). HRMS calculated for C₃₆H₃₅ClBrN₉O₄Na (M + Na)⁺: 796.1738; found 796.1713.

(1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-(2-(6-(1-(4-acetamidoethyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (15)

Compound **15** (86%) was synthesized from **9** following same procedure as for compound **11**. ¹H NMR (CD₃OD, 300 MHz) δ 8.09 (s, 1H), 7.78 (s, 1H), 7.43 (s, 1H), 7.27–7.33 (m, 3H), 5.03 (d, *J* = 5.1 Hz, 1H), 4.83–86 (m, 1H), 4.47 (t, *J* = 5.7 Hz, 2H), 4.01 (d, *J* = 6.6 Hz, 1H), 3.62 (t, *J* = 5.7 Hz, 2H), 2.87 (s, 3H), 2.79 (t, *J* = 7.5 Hz, 2H), 2.53 (t, *J* = 7.2 Hz, 2H), 2.07–2.10 (m, 1H), 1.85–1.93 (m, 5H), 1.69–1.74 (m, 2H), 1.36–1.41 (m, 1H). HRMS calculated for C₃₂H₃₈ClN₁₀O₄ (M + H)⁺: 661.2766; found 661.2751.

(1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-(2-(6-(1-(4-aminobutyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (16)

Compound **16** (73%) was synthesized from **9** following same procedure as for compound **12**. ¹H NMR (CD₃OD, 300 MHz) δ 8.11 (s, 1H), 7.81 (s, 1H), 7.45 (s, 1H), 7.28–7.34 (m, 3H), 5.06 (d, *J* = 6.3 Hz, 1H), 4.80–4.83 (m, 1H), 4.42 (t, *J* = 6.6 Hz, 2H), 4.01 (d, *J* = 6.6 Hz, 1H), 2.85–2.88 (m, 5H), 2.79–2.84 (m, 4H), 2.54 (t, *J* = 7.2 Hz, 2H), 2.08–2.12 (m, 1H), 1.86–1.99 (m, 5H), 1.66–1.78 (m, 2H), 1.54–1.62 (m, 2H), 1.38–1.43 (m, 1H). HRMS calculated for C₃₂H₄₀ClN₁₀O₃ (M + H)⁺: 647.2973; found 647.2968.

(1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-(2-(6-(1-(4-acetamidobutyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (17)

To a solution of compound **16** (1.79 mg, 0.002 mmol) in anhydrous DMF (0.5 mL), acetic acid *N*-hydroxysuccinimide ester (1 mg, 0.006 mmol) was added and the mixture stirred at room temperature for overnight. Solvent was evaporated and the residue was purified in preparative TLC (CH₂Cl₂:MeOH = 25:1) to give the compound **17** as a syrup (1.26 mg, 66 %). ¹H NMR (CD₃OD, 300 MHz) δ 8.11 (s, 1H), 7.80 (s, 1H), 7.44 (s, 1H), 7.27–7.33 (m, 3H), 5.04 (d, *J* = 6.3 Hz, 1H), 4.83–4.85 (m, 1H), 4.39 (t, *J* = 6.9 Hz, 2H), 3.70 (t, *J* = 3.9 Hz, 4H), 3.59 (m, 4H), 2.87 (s, 3H), 2.78–2.86 (m, 2H), 2.48–2.57 (m, 2H), 1.93 (s, 3H), 1.85–1.91 (m, 2H), 1.68–1.79 (m, 2H), 1.39–1.52 (m, 2H), 1.23–1.43 (m, 2H). HRMS calculated for C₃₄H₄₁ClN₁₀O₄Na (M + Na)⁺: 711.2898; found 711.2917.

(1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-(2-(6-(1-(adamantyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (18)

Compound **18** (76%) was synthesized from **9** following same procedure as for compound **11**. ¹H NMR (CD₃OD, 300 MHz) δ 8.09 (s, 1H), 7.81 (s, 1H), 7.44 (s, 1H), 7.25–7.33 (m, 3H), 5.03 (d, *J* = 6.6 Hz, 1H), 4.83–4.85 (m, 1H), 3.98 (d, *J* = 6.0 Hz, 1H), 2.86 (s, 3H), 2.79 (t, *J* = 7.2 Hz, 2H), 2.51 (t, *J* = 6.9 Hz, 2H), 2.16 (s, 9H), 2.06–2.10 (m, 1H), 1.68–1.95 (m, 13H), 1.36–1.40 (m, 1H). HRMS calculated for C₃₈H₄₅ClN₉O₃ (M + H)⁺: 710.3334; found 710.3352.

(1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-(2-(1-(4-3-oxoisothiazol-2(3H)-yl)benzyl-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (19)

Compound **19** (81%) was synthesized from **9** following same procedure as for compound **11**. ¹H NMR (CD₃OD, 300 MHz) δ 8.56 (d, *J* = 6.3 Hz, 1H), 8.08 (s, 1H), 7.83(s, 1H), 7.53–7.56 (m, 2H), 7.39–7.42 (m, 2H), 7.25–7.30 (m, 4H), 6.29 (d, *J* = 6.3 Hz, 1H), 5.6 (s, 2H), 5.02 (d, *J* = 6.9 Hz, 1H), 4.80–4.83 (m, 1H), 3.97 (d, *J* = 6.0 Hz, 1H), 2.86 (s, 3H), 2.80 (t, *J* = 7.2 Hz, 2H), 2.50 (t, *J* = 7.2 Hz, 2H), 2.06–2.10 (m, 1H), 1.84–1.92 (m, 4H), 1.67–1.74 (m, 2H), 1.35–1.40 (m, 1H). HRMS calculated for C₃₈H₃₈ClN₁₀O₄ (M + H)⁺: 765.2487; found 765.2461.

(1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-(2-(6-(1-(6-(5-(3aS,4S,6aR)-oxohexahydro-1H-thieno[3,4-d]imidazol-4-yl)pentanamido)hexyl-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (20)

Compound **20** (73%) was synthesized from **9** and the appropriate azide [ref. 38] following same procedure as for compound **12**. ¹H NMR (CD₃OD, 300 MHz) δ 8.08 (s, 1H), 7.77 (s, 1H), 7.42 (s, 1H), 7.23–7.31 (m, 3H), 5.02 (d, *J* = 6.3 Hz, 1H), 4.83–4.86 (m, 1H), 4.45–4.49 (m, 1H), 4.26–4.35 (m, 3H), 3.98 (d, *J* = 7.2 Hz, 1H), 3.65–3.69 (m, 3H), 3.54–3.57 (m, 3H), 3.03–3.24 (m, 4H), 2.85 (s, 3H), 2.78 (t, *J* = 7.5 Hz, 2H), 2.51 (t, *J* = 7.2 Hz, 2H), 2.06–2.28 (m, 6H), 1.81–1.89 (m, 6H), 1.60–1.72 (m, 6H), 1.34–1.46 (m, 2H). HRMS calculated for C₄₄H₅₇ClN₁₂O₅SNa (M + Na)⁺: 923.3882; found 923.3887.

1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-(2-(6-(1-(7-oxo-7-(6-(5-(3aS,4S,6aR)-oxohexahydro-1H-thieno[3,4-d]imidazol-4-yl)pentanamido)hexylamino)heptyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (21)

Compound **21** (82%) was synthesized from **9** following same procedure as for compound **12**. ¹H NMR (CD₃OD, 300 MHz) δ 8.03 (s, 1H), 7.81 (s, 1H), 4.46 (s, 1H), 7.29–7.35 (m, 3H), 5.05 (d, *J* = 6.5 Hz, 1H), 4.80–4.85 (m, 1H), 4.42–4.46 (m, 4H), 3.99 (d, *J* = 6.9 Hz, 1H), 3.79–3.85 (m, 8H), 3.56–3.62 (m, 8H), 3.15–3.26 (m, 4H), 2.89 (s, 3H), 2.23–2.57 (m, 6H), 1.03–1.91 (bm, 28H). HRMS calculated for C₅₅H₇₈ClN₁₃O₁₀SNa (M + Na)⁺: 1170.5302; found 1170.5288.

2-(6-Amino-3-imino-4,5-disulfonato-3H-xanthen-9-yl)-5-(6-(4-(6-(6-3-chlorobenzylamino)-9-((1S,2R,3S,4R,5S)-3,4-dihydroxy-5-(methylcarbamoyl)bicyclo[3.1.0]hexane-2-yl)-9H-purin-2-yl)hex-5-ynyl)-1H-1,2,3-triazol-1-yl)hexylcarbamoyl)benzoate (23)

Compound **23** (76%) was synthesized from **9** following same procedure as for compound **12**. ¹H NMR (CD₃OD, 300 MHz) δ 8.56 (s, 1H), 8.07(s, 1H), 7.83 (s, 1H), 7.26–7.43 (m, 4H), 7.09–7.12 (m, 2H), 6.90 (d, *J* = 6.9 Hz, 2H), 5.75 (d, *J* = 6.7 Hz, 1H), 5.57 (d, *J* = 6.8 Hz, 1H), 4.98 (d, *J* = 6.6 Hz, 1H), 4.80–4.84 (m, 1H), 4.13 (t, *J* = 3.6 Hz, 2H), 3.98 (d, *J* = 7.2 Hz, 1H), 3.69–3.72 (m, 4H), 3.59–3.64 (m, 4H), 2.89 (s, 3H), 2.50–2.55 (m, 4H), 2.19–2.32 (m, 4H), 1.21–1.46 (m, 7H). HRMS calculated for C₅₅H₅₄ClN₁₂O₁₃S₂ (M + H)⁺: 1189.3063; found 1189.3038. Fluorescence (aq. pH 7.4): λ_{ex} 490 nm, λ_{em} 520 nm.

(1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-(2-(1,6-bis(4-isothiocyanate-phenyl)-1H-bis(1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (25)

To a mixture of compound **9** (17 mg, 0.031 mmol) and 4-azidophenyl isothiocyanate (7.8 mg, 0.044 mmol) in THF/H₂O 3:1 (1.2 mL), was added freshly prepared 1M sodium ascorbate (26 μL, 0.025 mmol) followed by 7.5% aqueous copper sulfate pentahydrate solution (21 μL, 0.006 mmol) and stirred for 2 d at room temperature. Solvent was evaporated and the residue

was purified on flash silica gel column chromatography (CH₂Cl₂:MeOH = 20:1) to give an unusual clicked product **25** (21 mg, 75%) as a syrup. ¹H NMR (CD₃OD, 300 MHz) δ 8.29 (s, 1H), 8.12 (s, 1H), 7.71 (d, *J* = 8.7 Hz, 2H), 7.63 (d, *J* = 9.0 Hz, 2H), 7.50 (d, *J* = 6.6 Hz, 2H), 7.41 (s, 1H), 7.23–7.30 (m, 3H), 7.08 (d, *J* = 6.6 Hz, 2H), 5.02 (d, *J* = 5.2 Hz, 1H), 4.79–4.81 (m, 1H), 3.97 (d, *J* = 6.6 Hz, 1H), 2.83–2.91 (m, 5H), 2.55 (t, *J* = 6.9 Hz, 2H), 1.97–2.09 (m, 4H), 1.74–1.86 (m, 4H), 1.39–1.47 (m, 1H). HRMS calculated for C₄₁H₄₀ClN₁₄O₃S (M – CS)⁺: 843.2817; found 843.2812.

(1S,2R,3S,4R,5S)-6-(3-Chlorobenzylamino)-(2-(6-(1-(4-isothiocyanate-phenyl)-1H-1,2,3-triazol-4-yl)hex-1-ynyl)-9H-purin-9-yl)-2',3'-dihydroxybicyclo[3.1.0]hexane-1'-carboxylic acid N-methylamide (26)

To a mixture of compound **9** (4.22 mg, 0.007 mmol) and 4-isothiocyanate-phenylazide (2 mg, 0.011 mmol) in a mixture of *t*-butanol (0.3 mL) and water (0.3 mL), was added TBTA (1 mg, 0.001 mmol) and freshly prepared sodium ascorbate (7.9 μL, 0.007 mmol) followed by copper sulfate (7.9 μL, 0.002 mmol). The reaction mixture was stirred at room temperature for overnight. The starting material and product came in same R_f value in TLC. Solvent was evaporated and the residue was purified on flash silica gel column chromatography (CH₂Cl₂:MeOH = 25:1) to give the compound **26** (3.8 mg, 69%) as a syrup, no diclicked product was detected. ¹H NMR (CD₃OD, 300 MHz) δ 8.35 (s, 1H), 8.09 (s, 1H), 7.83 (d, *J* = 6.9 Hz, 2H), 7.40–7.43 (m, 3H), 7.24–7.32 (m, 3H), 5.02 (d, *J* = 6.3 Hz, 1H), 4.83–4.85 (m, 1H), 3.97 (d, *J* = 6.3 Hz, 1H), 2.87–2.92 (m, 5H), 2.55 (t, *J* = 6.9 Hz, 2H), 1.97–2.13 (m, 4H), 1.87 (t, *J* = 4.5 Hz, 2H), 1.74–1.83 (m, 2H), 1.38–1.41 (m, 1H). HRMS calculated for C₃₅H₃₄ClN₁₀O₃S (M + H)⁺: 709.2225; found 709.2236.

4-Oxo-4-phenyl-N-*p*-tolyl-butylamide(28)

EDC (347 mg, 1.81 mmol) and HOBt (254 mg, 1.88 mmol) were added to mixture of *p*-toluidine (129 mg, 1.20 mmol) and 3-benzoylpropionic acid **27** (218 mg, 1.21 mmol) in 1,4-dioxane (3 mL) at room temperature. It was stirred for 20 min and then added triethylamine (0.5 mL, 3.59 mmol). The reaction mixture was stirred for 20 h at room temperature and the solvent was evaporated under vacuum. The residue was purified by flash silica gel column chromatography (hexane:ethyl acetate = 3:1 to 2:1) yielded the compound **28** (252 mg, 78%). ¹H-NMR (CDCl₃, 300MHz) δ 2.30 (s, 3H), 2.80 (t, *J* = 6.3 Hz, 2H), 3.45 (t, *J* = 6.3 Hz, 2H), 7.10 (d, *J* = 8.4 Hz, 2H), 7.39 (d, *J* = 8.4 Hz, 2H), 7.47 (m, 2H), 7.58 (m, 1H), 7.99 (m, 1H).

5-Benzoyl-2-*p*-tolylisothiazol-3-one (29)

A solution of 4-oxo-4-phenyl-N-*p*-tolyl-butylamide **28** (252 mg, 0.944 mmol) in thionyl chloride (1 mL) was stirred for 16 h at room temperature. Excess thionyl chloride was removed under vacuum and the residue was purified by flash silica gel column chromatography (hexane:ethyl acetate = 10:1 to 4:1) to give the compound **29** (216 mg, 77%). ¹H-NMR (CDCl₃, 300 MHz) δ 2.40 (s, 3H), 6.82 (s, 1H), 7.28 (m, 2H), 7.54 (m, 4H), 7.71 (m, 1H), 7.96 (m, 2H). HRMS: calculated for C₁₇H₁₄NO₂S (M+H)⁺ 296.0745; found 296.0762.

2-*p*-Tolylisothiazol-3-one (30)

Aqueous sodium hydroxide (4 mL, 10%) was added to the solution of 5-benzoyl-2-*p*-tolylisothiazol-3-one **29** (203 mg, 0.678 mmol) in benzene (9 mL), and the mixture was stirred for 3 d at room temperature. The reaction mixture was added water (20 mL) and extracted with ethyl acetate. The combined organic layer was dried (MgSO₄), filtered and evaporated. The residue was purified by flash silica gel column chromatography (hexane:ethyl acetate = 2:1 to 1:1) and gave the compound **30** (77 mg, 59%). ¹H-NMR (CDCl₃, 300MHz) δ 2.38 (s, 3H),

6.32 (d, $J = 6.6$ Hz, 1H), 7.25 (d, $J = 8.1$ Hz, 2H), 7.44 (d, $J = 8.1$ Hz, 2H), 8.14 (d, $J = 6.6$ Hz, 1H). HRMS: calculated for $C_{10}H_{10}NOS$ (M+H)⁺ 192.0483; found 192.0494.

2-(4-Bromomethylphenyl)isothiazol-3-one (31)

Catalytic amount of benzoyl peroxide (4 mg) was added to a solution of 2-*p*-tolylisothiazol-3-one **30** (123 mg, 0.643 mmol), and *N*-bromosuccinimide (121 mg, 0.673 mmol) in carbon tetrachloride (8 mL), which was refluxed for 2 h and then cooled to room temperature. Solvent was evaporated under low pressure and the residue was purified by flash silica gel column chromatography (methylene chloride:methanol = 50:1 to 20:1), afforded the compound **31** (182 mg, 93%). ¹H-NMR (CDCl₃, 300MHz) δ 4.50 (s, 2H), 6.33 (d, $J = 6.3$ Hz, 1H), 7.47 (d, $J = 8.4$ Hz, 2H), 7.59 (d, $J = 8.4$ Hz, 2H), 8.16 (d, $J = 6.3$ Hz, 1H). HRMS: calculated for $C_{10}H_9NOSBr$ (M+H)⁺ 269.9588; found 269.9593.

2-(4-Azidomethylphenyl)isothiazol-3-one (32)

Sodium azide (6.0 mg, 0.0923 mmol) was added to a solution of 2-(4-bromomethylphenyl)isothiazol-3-one **31** (18.2 mg, 0.0674 mmol) in DMF (1 mL) and stirred for 20 h at room temperature. Water (10 mL) was added and the mixture was extracted with diethyl ether. The combined organic layer was dried (MgSO₄), filtered and evaporated. The residue was purified on flash silica gel column chromatography (hexane:ethyl acetate = 4:1 to 1:1), afforded the desired compound **32** (11.7 mg, 75%). ¹H-NMR (CDCl₃, 300 MHz) δ 4.38 (s, 2H), 6.34 (d, $J = 6.3$ Hz, 1H), 7.41 (d, $J = 8.4$ Hz, 2H), 7.64 (d, $J = 8.4$ Hz, 2H), 8.17 (d, $J = 6.3$ Hz, 1H). HRMS: found 233.0496; calcd for $C_{10}H_9N_4OS$ (M+H)⁺ 233.0497.

Receptor binding and functional assays

[³H]Adenosine-5'-*N*-methyluronamide (**36**, [³H]NECA, 42.6 Ci/mmol) was obtained from Perkin Elmer. [³H](2-[*p*-(2-Carboxyethyl)phenyl-ethylamino]-5'-*N*-ethylcarboxamidoadenosine) (**37**, [³H]CGS21680, 40.5 Ci/mmol) and [¹²⁵I]*N*⁶-(4-amino-3-iodobenzyl)adenosine-5'-*N*-methyluronamide (**38**, [¹²⁵I]I-AB-MECA, 2200 Ci/mmol) were purchased from Perkin-Elmer Life and Analytical Science (Boston, MA). Test compounds were prepared as 5 mM stock solutions in DMSO and stored frozen at -20°C.

Cell Culture and Membrane Preparation—CHO cells stably expressing the recombinant hA₁ and hA₃Rs, and HEK-293 cells stably expressing the hA_{2A}AR were cultured in Dulbecco's modified Eagle medium (DMEM) and F12 (1:1) supplemented with 10% fetal bovine serum, 100 units/mL penicillin, 100 µg/mL streptomycin, and 2 µmol/mL glutamine. In addition, 800 µg/mL geneticin was added to the A_{2A} media, while 500 µg/mL hygromycin was added to the A₁ and A₃ media. After harvesting, cells were homogenized and suspended in PBS. Cells were then centrifuged at 240 *g* for 5 min, and the pellet was resuspended in 50 mM Tris-HCl buffer (pH 7.5) containing 10 mM MgCl₂. The suspension was homogenized and was then ultracentrifuged at 14,330 *g* for 30 min at 4 °C. The resultant pellets were resuspended in Tris buffer, incubated with adenosine deaminase (3 units/mL) for 30 min at 37 °C. The suspension was homogenized with an electric homogenizer for 10 sec, pipetted into 1 mL vials and then stored at -80 °C until the binding experiments were conducted. The protein concentration was measured using the BCA Protein Assay Kit from Pierce Biotechnology, Inc. (Rockford, IL).

Binding assays—Into each tube in the binding assay was added 50 µL of increasing concentrations of the test ligand in Tris-HCl buffer (50 mM, pH 7.5) containing 10 mM MgCl₂, 50 µL of the appropriate agonist radioligand, and finally 100 µL of membrane suspension. For the A₁AR (22 µg of protein/tube) the radioligand used was [³H]**36** (final concentration of 3.5 nM). For the A_{2A}AR (20 µg/tube) the radioligand used was [³H]**37** (10

nM). For the A₃AR (21 µg/tube) the radioligand used was [¹²⁵I]**38** (0.34 nM). Nonspecific binding was determined using a final concentration of 10 µM unlabeled **36** diluted with the buffer. The mixtures were incubated at 25 °C for 60 min in a shaking water bath. Binding reactions were terminated by filtration through Brandel GF/B filters under a reduced pressure using a M-24 cell harvester (Brandel, Gaithersburg, MD). Filters were washed three times with 3 mL of 50 mM ice-cold Tris-HCl buffer (pH 7.5). Filters for A₁ and A_{2A}AR binding were placed in scintillation vials containing 5 mL of Hydrofluor scintillation buffer and counted using a Perkin Elmer Liquid Scintillation Analyzer (Tri-Carb 2810TR). Filters for A₃AR binding were counted using a Packard Cobra II γ-counter. The K_i values were determined using GraphPad Prism for all assays.

Cyclic AMP accumulation assay—Intracellular cyclic AMP levels were measured with a competitive protein binding method.^{29,30} CHO cells that expressed the recombinant human A₃AR were harvested by trypsinization. After centrifugation and resuspended in medium, cells were planted in 24-well plates in 1.0 mL medium. After 24 h, the medium was removed and cells were washed three times with 1 mL DMEM, containing 50 mM HEPES, pH 7.4. Cells were then treated with the agonist NECA and/or test compound in the presence of rolipram (10 µM) and adenosine deaminase (3 units/mL). After 45 min forskolin (10 µM) was added to the medium, and incubation was continued for an additional 15 min. The reaction was terminated by removing the supernatant, and cells were lysed upon the addition of 200 µL of 0.1 M ice-cold HCl. The cell lysate was resuspended and stored at -20°C. For determination of cyclic AMP production, protein kinase A (PKA) was incubated with [³H]cyclic AMP (2 nM) in K₂HPO₄/EDTA buffer (K₂HPO₄, 150 mM; EDTA, 10 mM), 20 µL of the cell lysate, and 30 µL 0.1 M HCl or 50 µL of cyclic AMP solution (0–16 pmol/200 µL for standard curve). Bound radioactivity was separated by rapid filtration through Whatman GF/C filters and washed once with cold buffer. Bound radioactivity was measured by liquid scintillation spectrometry.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Dr. John Lloyd and Dr. Noel Whittaker (NIDDK) for mass spectral determinations. This research was supported by the Intramural Research Program of the NIH, National Institute of Diabetes and Digestive and Kidney Diseases. We thank Arunkumar Easwaran, Ph.D., Molecular Targeting Technologies, Inc. West Chester, PA for helpful discussions.

Abbreviations

AR	adenosine receptor
cAMP	adenosine 3',5'-cyclic phosphate
CHO	Chinese hamster ovary
CI-IB-MECA	2-chloro- <i>N</i> ⁶ -(3-iodobenzyl)-5'- <i>N</i> -methylcarboxamidoadenosine
DMEM	Dulbecco's modified Eagle's medium
EDC	1-ethyl-3-(3-dimethylaminopropyl) carbodiimide
EDTA	ethylenediaminetetraacetic acid
GPCR	G protein-coupled receptor

HATU	2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium
HEK	human embryonic kidney
HOBt	1-hydroxybenzotriazole
I-AB-MECA	<i>N</i> ⁶ -(3-iodo-4-aminobenzyl)-5'- <i>N</i> -methylcarboxamidoadenosine
IB-MECA	<i>N</i> ⁶ -(3-iodobenzyl)-5'- <i>N</i> -methylcarboxamido-adenosine
NECA	5'- <i>N</i> -ethylcarboxamidoadenosine
DMF	<i>N,N</i> -dimethylformamide
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
HRMS	high resolution mass spectroscopy
TBTA	tris[(1-benzyl-1H-1,2,3-triazol-4-yl)methyl] amine
TEA	triethylamine
TLC	thin layer chromatography

References

- Jacobson, KA.; Klutz, AM.; Tosh, DK.; Ivanov, AA.; Preti, D.; Baraldi, PG. Handbook of Experimental Pharmacology. Vol. 193. Springer; 2009. p. 123
- Gessi S, Merighi S, Varani K, Cattabriga E, Benini A, Mirandola P, Leung E, Ma c Lennan S, Feo C, Baraldi S, Borea PA. J. Cell. Physiol 2007;211:826. [PubMed: 17348028]
- Yang H, Avila MY, Peterson-Yantorno K, Coca-Prados M, Stone RA, Jacobson KA, Civan MM. Current Eye Res 2005;30:747.
- Hua X, Chason KD, Fredholm BB, Deshpande DA, Penn RB, Tilley SL. J Allergy Clin Immunol 2008;122:107.
- Fishman P, Jacobson KA, Ochaion A, Cohen S, Bar-Yehuda S. Immun. Endoc. Metab. Agents in Med. Chem 2007;7:298.
- Silverman MH, Strand V, Markovits D, Nahir M, Reitblat T, Molad Y, Rosner I, Rozenbaum M, Mader R, Adawi M, Caspi D, Tishler M, Langevitz P, Rubinow A, Friedman J, Green L, Tanay A, Ochaion A, Cohen S, Kerns WD, Cohn I, Fishman-Furman S, Farbstein M, Bar-Yehuda S, Fishman P. J. Rheumatol 2008;35:41. [PubMed: 18050382]
- Zheng J, Wang R, Zambraski E, Wu D, Jacobson KA, Liang BT. Am. J. Physiol.; Heart and Circ. Physiol 2007;293:3685.
- Jaakola V-P, Griffith MT, Hanson MA, Cherezov V, Chien EYT, Lane JR, IJzerman AP, Stevens RC. Science 2008;322:1211. [PubMed: 18832607]
- Tosh DK, Chinn M, Ivanov AA, Klutz AM, Gao ZG, Jacobson KA. J. Med. Chem 2009;52:7580. DOI: 10.1021/jm900426g. [PubMed: 19499950]
- Jacobson KA, Gao ZG, Liang BT. Trends Pharmacol. Sci 2007;28:111. [PubMed: 17280720]
- Melman A, Wang B, Joshi BV, Gao ZG, de Castro S, Heller CL, Kim SK, Jeong LS, Jacobson KA. Bioorg. Med. Chem 2008;16:8546. [PubMed: 18752961]
- Melman A, Gao ZG, Kumar D, Wan TC, Gizewski E, Auchampach JA, Jacobson KA. Bioorg. Med. Chem. Lett 2008;18:2813. [PubMed: 18424135]
- Jacobson KA. Bioconjugate Chem 2009;20:1816.
- a) Matsuda A, Shinozaki M, Yamaguchi T, Homma H, Nomoto R, Miyasaka T, Watanabe Y, Abiru T. J. Med. Chem 1992;35:241. [PubMed: 1732541] b) Cristalli G, Volpini R, Vittori S, Camaioni E, Monopoli A, Conti A, Dionisotti S, Zocchi C, Ongini E. J. Med. Chem 1994;37:1720. [PubMed: 8201607] c) Rieger JM, Brown ML, Sullivan GW, Lind en J, Macdonald TL. J. Med. Chem

- 2001;44:531. [PubMed: 11170643] d) Volpini R, Costanzi S, Lambertucci C, Vittori S, Klotz KN, Cristalli G. *J. Med. Chem* 2002;45:3271. [PubMed: 12109910]
15. Kolb HC, Sharpless KB. *Drug Discov. Today* 2003;8:1128. [PubMed: 14678739]
 16. Chinchilla R, Nájera C. *Chem. Rev* 2007;107:874. [PubMed: 17305399]
 17. Kolb HC, Finn MG, Sharpless KB. *Angew. Chem. Int. Ed* 2001;40:2004.
 18. Bock VD, Hiemstra H, van Maarseveen JH. *Eur. J. Org. Chem* 2006;51:68.
 19. Chan TR, Hilgraf R, Sharpless KB, Fokin VV. *Org. Lett* 2004;6:2853. [PubMed: 15330631]
 20. Gassensmith JJ, Arunkumar E, Barr L, Baumes JM, DiVittorio KM, Johnson JR, Noll BC, Smith BD. *J. Am. Chem. Soc* 2007;129:15054. [PubMed: 17994746]
 21. Johnson JR, Fu N, Arunkumar E, Leevy WM, Gammon ST, Piwinica-Worms D, Smith BD. *Angew. Chem. Int. Ed* 2007;46:5528.
 22. Berlier JE, Rothe A, Buller G, Bradford J, Gray DR, Filanoski BJ, Telford WG, Yue S, Liu J, Cheung C-Y, Chang W, Hirsch JD, Beechem JM, Haugland RP, Haugland RP. *J. Histochem. Cytochem* 2003;51:1699. [PubMed: 14623938]
 23. Tsolomitis A, Sandris C. *Heterocycles* 1987;25:569.
 24. Klotz KN, Lohse MJ, Schwabe U, Cristalli G, Vittori S, Grifantini M. *Naunyn Schmiedebergs Arch. Pharmacol* 1989;340:679. [PubMed: 2615857]
 25. Jarvis MF, Schutz R, Hutchison AJ, Do E, Sills MA, Williams M. *J. Pharmacol. Exp. Ther* 1989;251:888. [PubMed: 2600819]
 26. Olah ME, Gallo-Rodriguez C, Jacobson KA, Stiles GL. *Mol. Pharmacol* 1994;45:978. [PubMed: 8190112]
 27. Englert M, Qwitterer U, Klotz KN. *Biochem. Pharmacol* 2002;64:61. [PubMed: 12106606]
 28. Jacobson KA, Park KS, Jiang J.-I. Kim YC, Olah ME, Stiles GL, Ji X.d. *Neuropharmacology* 1997;36:1157. [PubMed: 9364471]
 29. Nordstedt C, Fredholm BB. *Anal. Biochem* 1990;189:231. [PubMed: 2177960]
 30. Post SR, Ostrom RS, Insel PA. *Methods Mol. Biol* 2000;126:363. [PubMed: 10685423]
 31. Cosyn L, Palaniappan KK, Kim SK, Duong HT, Gao ZG, Jacobson KA, Van Calenbergh S. *J. Med. Chem* 2006;49:7373. [PubMed: 17149867]
 32. Hsu T-L, Hanson SR, Kishikawa K, Wang S-K, Sawa M, Wong C-H. *Proc. Natl. Acad. Sci USA* 2007;104:2614. [PubMed: 17296930]
 33. Salic A, Mitchison TJ. *Proc. Natl. Acad. Sci USA* 2008;105:2415. [PubMed: 18272492]
 34. Ming X, Leonard P, Heindl D, Seela F. *Nucleic Acids Symposium Series* 2008;52:471. [PubMed: 18776458]
 35. Morley JO, Kapur AJO, Charlton MH. *Int. J. Chem. Kinetics* 2007;39:254.
 36. Codelli JA, Baskin JM, Agard NJ, Bertozzi CR. *J. Am. Chem. Soc* 2008;130:11486. [PubMed: 18680289]
 37. Bradford MM. *Anal. Biochem* 1976;72:248. [PubMed: 942051]
 38. Inverarity IA, Viguier RFH, Cohen P, Hulme AN. *Bioconjugate Chem* 2007;18:1593.







Figure 1. Inhibition of radioligand binding by (N)-methanocarba nucleoside analogues in membranes of CHO cells expressing the human A₁ (■) and A_{2A} (▲) and A₃ (▼) ARs. Inhibition curves are shown for the agonist analogues containing a triazole group, the adamantyl derivative **18**, and the amide model compound for click linkage to carriers **15**. Both of the analogues shown were highly selective for the hA₃AR in comparison to the A₁ and A_{2A}ARs.

**Scheme 1.**


Use of click reactions to couple a terminal alkynyl groups present on a functionalized chain of an adenosine analogue to various azido moieties. Reagents: A. (i) $\text{HC}\equiv\text{C}(\text{CH}_2)_n\text{C}\equiv\text{CH}$ ($n = 3, 4$), $\text{Pd}(\text{PPh}_3)_4$, CuI , Et_3N , DMF , rt; (ii) 10% TFA , MeOH , 70°C ; (iii) appropriate azide, $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, sodium ascorbate, TBTA , $\text{THF}/\text{H}_2\text{O}$ or $t\text{-BuOH}/\text{H}_2\text{O}$, rt; (iv) acetic acid *N*-hydroxysuccinimide ester, DMF , rt.

**Scheme 2.**

The unusual example of click reaction occurring at both alkyne groups present on a functionalized chain of an adenosine analogue. When the catalyst TBTA was used, only the monosubstituted product **26** was isolated. Reagents: (i) $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, sodium ascorbate, THF/ H_2O , rt.; (ii) $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, sodium ascorbate, TBTA, *t*-BuOH/ H_2O , rt.

**Scheme 3.**


Synthesis of isothiazolone derivative **32** to serve as a thiol-reactive crosslinker. Reagents: (i) *p*-toluidine, EDC, HOBT, Et₃N, 1,4-dioxane, rt, 16 h; (ii) thionyl chloride, rt, 16 h; (iii) 10% aqueous NaOH, benzene, rt, 3 d; (iv) *N*-bromosuccinimide, benzoyl peroxide, carbon tetrachloride, reflux, 2 h; (v) sodium azide, DMF.

**Chart 1.**

Molecular probes in the (N)-methanocarba series of adenosine A₃AR agonists that were previously reported (**1 – 3**)⁹ and a general formula (**4**) for triazole derivatives prepared in this study.


Table 1

Potency of a series of (N)-methanocarba adenosine derivatives at three subtypes of human ARs.



Bioorg Med Chem. Author manuscript; available in PMC 2011 January 15.

Compd	Structure	Affinity (K _i , nM) or % inhibition ^d		
		A ₁	A _{2A}	A ₃
	R ¹			
<i>b,c</i>		(36±3%)	4730±1020	17.2±3.1
<i>b,c</i>		(1±1%)	(51±2%)	36.4±5.6
<i>b,c</i>		(12±4%)	(47±11%)	57.7±16.2
		(4±2%)	(40±3%)	8.8±1.3
8		(36±4%)	4330±500	23.6±3.9
9		(31±2%)	7040±1430	29.4±9.8

Compd	Structure	Affinity (K _i , nM) or % inhibition ^a		
		A ₁	A _{2A}	A ₃
n	R ²			
10	3-nitro-4-fluorophenyl	(10±3%)	(39±4%)	26.0±8.2
11 ^e	3-nitro-4-fluorophenyl	(8±4%)	6730±280	10.6±3.8
12	4-aminophenyl	(3±1%)	5490±1150	87.1±13.1
13	4-carboxyphenyl	(0±0%)	(40±1%)	180±23
16	4-isothiocyanatophenyl	(2±1%)	(28±4%)	37.5±16.0
14 ^e	4-(α-Br-phenacyl)	(12±3%)	5740±730	9.6±1.3
15	(CH ₂) ₂ NHCOCH ₃	(12±4%)	2440±320	22.3±1.6
16	(CH ₂) ₄ NH ₂	(13±3%)	1630±350	47.0±1.8
17	(CH ₂) ₄ NHCOCH ₃	(17±1%)	7240±510	89.5±12.6
18 ^e	1-adamantyl	(22±2%)	3280±700	6.5±0.5
19		(15±3%)	(49±1%)	102±25
20 ^d	-(CH ₂) ₆ NH-biotin	(0±0%)	(27±1%)	285±54
21 ^d	-(CH ₂) ₆ CONH(CH ₂) ₂ [O(CH ₂) ₄ NH-biotin	(0±0%)	(19±5%)	235±43
22	Rotaxane derivative	(0±0%)	(2±1%)	239±43
23 ^d	(CH ₂) ₆ NHCO-Alexa Fluor	(0±0%)	(23±5%)	416±45

^a All experiments were done on CHO or HEK293 (A_{2A} only) cells stably expressing one of four subtypes of human ARs. The binding affinity for A₁, A_{2A} and A₃ ARs was expressed as K_i values (n = 3–5) and was determined by using agonist radioligands ([³H]36; [³H]37; or [¹²⁵I]38; respectively), unless noted. A percent in parentheses refers to inhibition of radioligand binding at 10 μM.

^b Values from Tosh et al.⁹

^c Structure given in Chart 1.

^d Structure given in Scheme 1.

^e 11, MRS5223; 14, MRS5226; 15, MRS5233; 18, MRS5224; 25, MRS5225.

Table 2Maximal efficacy of (N)-methanocarba adenosine derivatives in a functional assay at the A₃AR.

Compound	% Inhibition of cAMP formation ^a at hA ₃ AR
<i>IB-MECA</i>	99±6
<i>CI-IB-MECA</i>	97
<i>36, NECA</i>	100
<i>1^b</i>	94.4±9.6
<i>2^b</i>	84.5±12.0
<i>3^b</i>	107±18
<i>25</i>	83.6±7.2
<i>8</i>	18.3±7.6
<i>9</i>	60.2±17.0
<i>10</i>	ND
<i>11</i>	116±22
<i>12</i>	44.3±2.4
<i>13</i>	83.4±13.1
<i>26</i>	19.5±14.5
<i>14</i>	109±12
<i>15</i>	93.6±17.7
<i>16</i>	64.4±12.4
<i>17</i>	59.7±19.2
<i>18</i>	27.8±17.4
<i>19</i>	75.9±16.5
<i>20</i>	55.6±12.5
<i>21</i>	41.8±11.3
<i>22</i>	111±18
<i>23</i>	37.8±14.6

^aThe efficacy at the human A₃AR was determined by inhibition of forskolin-stimulated cyclic AMP production in AR-transfected CHO cells, as described in the text. At a concentration of 10 μM, in comparison to the maximal effect of a full agonist NECA at 10 μM. Data are expressed as mean ± standard error (n = 3). ND, not determined.

^bValues from Tosh et al.⁹