

RESEARCH

Open Access

Effects of Parecoxib and Fentanyl on nociception-induced cortical activity

Yuan-Zhi Peng, Xiao-Xi Li, Ying-Wei Wang*

Abstract

Background: Analgesics, including opioids and non-steroid anti-inflammatory drugs reduce postoperative pain. However, little is known about the quantitative effects of these drugs on cortical activity induced by nociceptive stimulation. The aim of the present study was to determine the neural activity in response to a nociceptive stimulus and to investigate the effects of fentanyl (an opioid agonist) and parecoxib (a selective cyclooxygenase-2 inhibitor) on this nociception-induced cortical activity evoked by tail pinch. Extracellular recordings (electroencephalogram and multi-unit signals) were performed in the area of the anterior cingulate cortex while intracellular recordings were made in the primary somatosensory cortex. The effects of parecoxib and fentanyl on induced cortical activity were compared.

Results: Peripheral nociceptive stimulation in anesthetized rats produced an immediate electroencephalogram (EEG) desynchronization resembling the cortical arousal (low-amplitude, fast-wave activity), while the membrane potential switched into a persistent depolarization state. The induced cortical activity was abolished by fentanyl, and the fentanyl's effect was reversed by the opioid receptor antagonist, naloxone. Parecoxib, on the other hand, did not significantly affect the neural activity.

Conclusion: Cortical activity was modulated by nociceptive stimulation in anesthetized rats. Fentanyl showed a strong inhibitory effect on the nociceptive-stimulus induced cortical activity while parecoxib had no significant effect.

Background

Pain is a complex subjective sensory experience which is comprised of sensory discrimination, cognitive appreciation and affective motivation components [1]. The treatment of pain is a fundamental issue in the practice of anesthesia. However, the mechanisms of pain and analgesia are unclear. With the development of imaging technology (e.g., functional magnetic resonance imaging and positron emission tomography), cortical areas that are commonly activated by noxious stimuli have been illuminated. These areas include the primary and secondary somatosensory cortices (S1, S2), anterior cingulate cortex (ACC), insular cortex and motor cortex [2,3]. Electrophysiological studies show that cortical neuronal activity under anesthesia is associated with an Up-Down membrane potential fluctuation, while EEG signals are characterized by large-amplitude and slow-wave activity

[4-6]. It is well known that synapses are highly plastic in the central nervous system including ACC, and that long-term changes in synaptic transmission contribute to different functions of brain including chronic pain. Long-term potentiation (LTP) is believed to be the basis of learning and memory, and it can also be induced by different protocols or peripheral injury at the spinal cord dorsal horn [7,8]. Elucidation of the electrophysiological mechanisms of the neuron in encoding nociceptive information may provide new advances in the detection of pain under anesthesia. Despite the effectiveness of opioid drugs in postoperative pain management, there are limitations in their use, partly due to tolerance effects. In addition, opioid drugs can potentially cause respiratory depression, nausea and vomiting [9]. Non-steroid anti-inflammatory drugs (NSAIDs) have been used in combination with opioid analgesics to reduce postoperative opioid administration [10,11]. Parecoxib, a selective inhibitor of cyclooxygenase-2 (COX-2), can block prostaglandin biosynthesis associated with

* Correspondence: wangyingwei@yahoo.com
Department of Anesthesiology, Xinhua Hospital, Shanghai Jiaotong University School of Medicine, Shanghai 200092, China

inflammatory pain. Moreover, parecoxib lacks the potential adverse effects of COX-1 inhibition, including gastroduodenal ulceration and bleeding or impairment of platelet function [10,11]. However, it has been argued that parecoxib is lacking in adequate analgesic effects on postoperative pain [12].

The present study investigates the neuronal activity of ACC and S1 areas (two regions known for their involvement in pain processing) during noxious stimulation. In addition, the effect of parecoxib was compared with fentanyl on evoked cortical activity.

Results

Cortical response to nociceptive stimulation

We simultaneously measured extracellular activity in the ACC and intracellular activity in the S1 region in response to nociceptive stimulation. We found that both the ACC and S1 areas were activated by tail pinch in urethane anesthetized rats (Fig. 1A). To examine whether the neural activity of other cortical areas could be altered by nociceptive stimulation, we performed recordings at V1(primary visual cortex), PtA(parietal association cortex) and FrA(frontal association cortex), which presented similar activity to that seen in ACC and S1 (Fig. 1D and 1E).

The EEG recordings showed a shift from large-amplitude, low-frequency waves (commonly seen in anesthetized rats) towards low-amplitude, high-frequency waves (resembling that of awake rats) during tail pinch (Fig. 1A and 1C, upper panels). In addition, the mean firing rate obtained from multi-unit(MU) recordings was enhanced and the Down state disappeared (Fig. 1B). The power spectrum analysis revealed that during tail pinch the ratio of power within low frequency bands was markedly reduced (Fig. 1C), along with the decreased absolute power of each EEG band (Fig. 2A).

Intracellular recordings showed that the membrane potential (V_m) exhibited a bimodal distribution with Up-Down transitions in the resting condition. This pattern quickly switched to a persistent Up state following nociceptive stimulation (Fig. 1A, D and 1E, lower panels; Fig. 2B). Furthermore, waveform correlation analysis revealed that highly synchronized activity was observed between the ACC and S1 recordings in the resting state, while desynchronization was presented when nociceptive stimulation was applied (Fig. 2C and 2D). The desynchronization was observed when comparing extracellular recordings in both the ACC and SI (Fig. 2C right panel, $p < 0.01$), and when comparing extracellular recordings in the ACC with intracellular recordings in SI (Fig. 2D,





Figure 1 Electrophysiological recordings from cortical areas during nociceptive stimulation. A, upper panel: Trace of the extracellular signal from the ACC, showing that the amplitude was reduced while the frequency was increased when a nociceptive stimulus was applied; lower panel: Intracellular trace from S1, showing that the Down state disappeared and the membrane potential (V_m) switched to a persistent Up state during nociceptive stimulation. B: The Multi-Unit signal was filtered from the upper panel in A. The spike firing rate of the MU signal was increased, and the Down state disappeared under nociceptive stimulation. C, upper panel: the trace filtered from A (upper panel), note that EEG waves switched to low-amplitude, high-frequency pattern activity; lower panel: The power spectrum of the corresponding EEG (C, upper panel), showing that the low frequency band was markedly reduced. D: The traces recorded from the ACC (upper panel) and from V1 (lower panel). E: The traces recorded from PtA (upper panel) and FrA (lower panel). Triangles signify the start and end of nociceptive stimulation. The color bar represents the proportion of EEG bands.



$p < 0.001$). Together, these results provide direct electrophysiological evidence for extensive alterations in cortical activity by nociceptive stimulation, as suggested by previous EEG and imaging studies [2,3,13].

Effects of parecoxib and fentanyl on cortical activity


The effects of parecoxib and fentanyl on cortical activity in response to acute mechanical pain were compared. Extracellular recordings at ACC and intracellular recordings at S1 in urethane anesthetized rats were obtained. EEG recordings revealed that normalized total power (Fig. 3A) was decreased, while MF (medial frequency; see Fig. 3B), SEF (95% spectral edge frequency; see Fig. 3C) and distance of cumulative power distribution (Fig. 3D) were increased by nociceptive stimulation compared to the resting state. Fentanyl significantly inhibited these effects, and moreover, its action was reversed by naloxone (see Fig. 3). Parecoxib administration, on the other hand, did not inhibit the observed response in any of these measures (Fig. 3).

There was a significant decrease in δ band power and an increase in other band powers when nociceptive stimulation was applied (Fig. 4A and 4B, control group),

and a similar change was observed in the parecoxib group (Fig. 4A, right panel). However, there were no significant changes in any EEG band after fentanyl injected (Fig. 4B, middle panel). In addition, naloxone reversed the action of fentanyl (Fig. 4B, right panel).

Although the Down state was reduced or disappeared (Fig. 1), nociceptive stimulation did not influence the spike mean firing rate from MU recordings (Fig. 5A). However, the firing rate was reduced in some neurons (Fig. 1A and 1E; lower panel), and this inhibition was also observed in previous studies [14,15]. We therefore, further investigated spike firing rate by a Non-uniform index of spike frequency to evaluate effects of parecoxib and fentanyl. This measure showed that nociceptive stimulation decreased the “non-uniform index” in control (untreated) animals and that parecoxib did not inhibit this response; The response was partially inhibited, however by fentanyl and again, naloxone abolished the fentanyl effect (Fig. 5B).

Finally, we investigated the Up and Down states under different conditions in the S1 region. The ratio of time in the Down state to time in the Up state was reduced



and the firing rate of the Up state increased (Fig. 6A, B; control) during nociceptive stimulation. Administration of fentanyl reversed these responses while the addition of naloxone abolished the fentanyl's effects (Fig. 6A and 6B, right panels). In addition, the distance of membrane potential distribution was significantly increased during tail pinch (Fig. 6C; control) and administration of fentanyl significantly inhibited this response. Naloxone was again able to reverse the fentanyl effect (Fig. 6C, right panel). Parecoxib had no significant effect on any of these measures (Fig. 6, left panels).


Fentanyl (30 µg/kg, i.m.) did not significantly influence heart rate, while respiratory rate was lightly depressed (Fig. 7, n = 9). Further study showed that there was no significant change in the index of blood gas, glucose or hematocrit (see table 1).

Discussion

The present study investigated the neuronal response in the ACC and S1 area of urethane anesthetized rats during acute mechanical noxious stimulation. In addition, the effects of parecoxib and fentanyl on evoked neuronal responses were assessed. Acute mechanical stimulation led to neuronal activation in all cortical areas studied. In

a more detailed investigation using the ACC and S1 areas specifically, we found that peripheral nociceptive stimulation induced a transition from typical Up and Down state dynamics to a long-lasting EEG activated state. In addition, intracellular recordings showed that membrane potentials were locked into a persistent Up state during noxious stimulation resulting in a desynchronization among cortical brain areas. Fentanyl successfully inhibited the extracellular and intracellular responses induced by noxious stimulation and naloxone reversed the fentanyl' effects. Parecoxib, on the other hand, had no effect on neuronal response.

Previous studies have shown that EEG waves are desynchronized when a noxious stimulus is applied to a lightly anesthetized human or animal [13,16]. EEG desynchronization along with a significant decrease in total power and δ and θ band frequency have also been observed after noxious stimulation in isoflurane anesthetized goats [13]. Results from the present study show that all cortical areas investigated were activated by noxious stimulation (Fig. 1). This non-specificity in cortical activation may be restricted to anesthetized animals. Previous studies show that this type of total cortical activation can also be mimicked by stimulation of the ascending arousal system during anesthetic treatment



[17,18]. For example, brief electrical stimulation of the pedunculo-pontine tegmentum (PPT) or locus coeruleus under ketamine-xylazine anesthesia typically induced a prolonged period of desynchronized EEG activity that was paralleled by a persistent depolarized state of the membrane potential. These conditions are similar to that observed in the natural awake state [17,18]. Our results are consistent with those observed with the PPT stimulation in that the membrane potential was locked into a persistent Up state when nociceptive stimulation was applied to the tail. These findings indicate that the ascending arousal system can be activated directly by nociceptive stimulation. In other words, if the nociceptive stimulation had only activated the spinothalamic

tract, then the entire cerebral cortex would not have been activated. Thus, the present data, together with existing information indicate that the desynchronization of EEG or depolarized membrane potential may be an effective indicator of pain during anesthesia.

Traditional NSAIDs such as acetylsalicylic acid non-specifically inhibit both COX-1 and COX-2. Parecoxib on the other hand, is a new selective COX-2 inhibitor, able to maintain analgesic effects without altering the homeostatic functions of COX-1 [11,19]. In addition, parecoxib is injected and is therefore suitable for patients who cannot take oral medication. In this study, we systematically compared the analgesic effects of parecoxib and fentanyl through electrophysiological




Figure 5 The effects of parecoxib and fentanyl on MU (Multi-Unit) and Non-uniform index of MU recordings. A: Spike mean firing rate from MU recordings were not altered in the presence of nociceptive stimulation. B: The non-uniform index of MU was decreased when nociceptive stimulus was applied (control), indicating that the neuronal firing pattern was more uniform under nociceptive stimulation than during the resting state. Fentanyl partially inhibited this response while parecoxib had no effect. *P < 0.05, **P < 0.01, #P < 0.001.

techniques. We demonstrated that fentanyl successfully inhibited the electrophysiological profile evoked by nociceptive stimulation, and this inhibition was reversed by naloxone. Parecoxib, on the other hand, had no effect. Recent studies have demonstrated that parecoxib is effective in treating postoperative pain [20,21]. In addition, the analgesic effects of parecoxib have been demonstrated through fMRI studies showing a reduction in acute mechanical pain and mechanical hyperalgesia induced brain activity [22]. Moreover, other reports on

postoperative analgesic effects of parecoxib, have found a decrease in pain score and in opioid consumption [20]. There are a number of possible reasons for the negative results in our study. First, surgery injury leads to inflammation and COX-2 is highly expressed in injured tissue, while acute mechanical stimuli (used in this study) only activates the A_δ and C fibers without abnormally high expression of COX-2. Our results are in line with the results of recent studies reporting no improvement of postoperative pain management after





Figure 6 Group data for intracellular recordings at S1. The time ratio of Down state to Up state decreased (A, control), while transition frequency of Up state and the distance of cumulative V_m distribution increased (B and C, control) in response to nociceptive stimulation. The response of all measures to nociceptive stimulation was not inhibited by parecoxib (left panels, $n = 8$). Fentanyl (right panels, $n = 12$) inhibited all responses and naloxone reversed fentanyl's action ($n = 9$). * $P < 0.05$, ** $P < 0.01$, # $P < 0.001$.



discectomy with parecoxib [23]. These studies found that the timing of parecoxib administration is vital to its effectiveness: peri-operative administration was more effective than postoperative administration [23,24]. Another study found that parecoxib inhibited behavioral changes seen in carrageenan evoked inflammation, but had no effect in acute pain models such as acetic acid-induced writhing and the formalin test [25]. COX-2 mRNA and protein in spinal cord was increased for 3-6

h after injection of carrageenan [26], while acetic acid and formalin induced increases in COX-2 mRNA and protein lasted less than 60 min. In our study, the analgesic effect of parecoxib was tested 20 min after administration when there may not have been abnormally high levels of COX-2 expressed. Therefore, it is possible that the negative parecoxib findings in the present study may be due timing of drug administration. Our results, together with the previous findings suggest that the first

Table 1 Blood gas, Glucose and Hemotacrit values

	Control	Fentanyl
pH	7.31 ± 0.02	7.28 ± 0.02
PaCO ₂ (mmHg)	43 ± 2	48 ± 3
PaO ₂ (mmHg)	98 ± 3	99 ± 4
SO ₂	96.8 ± 0.4	96.8 ± 0.6
Base excess (mmol/L)	-5.3 ± 0.9	-4.6 ± 0.9
Glucose (g/dL)	15.0 ± 0.5	14.2 ± 0.3
Hematocrit (%)	48.2 ± 1.5	45.9 ± 1.1

dose of parecoxib should be administered before or during a procedure, and that treatment should continue post operation. Our study also suggests that parecoxib should not be used for analgesia in mechanical pain without inflammation.

The inhibitory effect of opioids on dorsal horn projection neurons has been well studied. For example, morphine can inhibit voltage-gated calcium channels of dorsal root ganglion neurons [27]. In addition, topical spinal application of 10 μ M morphine produces an inhibitory effect on projection neurons of lumbar spinal cord [28,29]. Previous electrophysiological studies have suggested that ACC pyramidal cells undergo rapid and prolonged depolarization after the induction of acute pain, such as by digit amputation [30]. It is conceivable that cortical desynchronization induced by mechanical stimulation results in the enhancement Ca^{2+} of influx through voltage-gated Ca^{2+} channels and NMDA receptors. Opioid agonists were shown to inhibit voltage-gated Ca^{2+} currents in neurons and reduce excitation by synaptic inhibition [31]. Recent study has shown that low doses of fentanyl could block the induction of LTP in rat spinal cord in vivo [32], which can be explained by the action of opioid receptors on synaptic transmission: presynaptic inhibition of Ca^{2+} channels and membrane hyperpolarization by activation K^+ channels [31]. So the cortical desynchronization might also be blocked by the above mechanisms of fentanyl. In addition, glutamatergic transmission can also be inhibited by opioid agonist, for example, morphine intra-basolateral amygdaloid nucleus injection reduces pain-induced aversion [33]. It is well known that, fentanyl has significant side effects, including respiratory depression, which may confound its analgesic effects [9]. We therefore monitored heart rate and respiratory rate to exclude the possibility of these effects as confounding variable. The present results show that heart and respiratory rates were not greatly affected by fentanyl administration (Fig. 7), and there was no effect on blood gas analysis (table 1). However, there was a slight respiratory depression with fentanyl administration, possibly leading to disruption in internal homeostasis [9]. In the future it will be necessary to exclude the influence of respiratory depression on fentanyl's analgesic effects.

Synaptic and cellular mechanisms of LTP in the ACC are well studied, which is believed to potentiate the sensory responses after injury. Field recording and whole cell patch-clamp recording techniques have been used to study the LTP of excitatory synapse responses in the ACC [34,35]. NMDA receptors play an important role in prefrontal cortex including ACC LTP [34], while NR2A and NR2B subunits contribute to most of the NMDA receptor currents [35,36]. Gene knockout and pharmacological methods have shown that calcium-activated adenylyl cyclases, especially subtype 1 and 8 (AC1, AC8), contribute to the induction of LTP in the ACC [37]. In addition, activity-dependent immediate early genes such as Egr1 and CREB (Cyclic-AMP Response Element Binding proteins) are suggested to contribute to synaptic plasticity [38]. Furthermore, the enhanced synaptic responses persist for a long period in the ACC neurons after tissue inflammation or amputation [34,39]. The animal model of neuropathic pain shows that post-synaptic AMPA receptor mediated responses are enhanced in the ACC synapses, such nerve injury triggered enhancement of postsynaptic responses could be abolished in AC1 knockout mice [40], suggesting that AC1 may serve as a potential therapeutic target for treating neuropathic pain. Recent research has examined potential changes in NR2B receptors in the ACC with inflammatory pain, which showed that NR2B receptor mediated responses are enhanced after peripheral inflammation. With selective forebrain NMDA NR2B overexpression in the case of tissue injury and inflammation, the inflammatory pain and allodynia were significantly enhanced [39,41], providing evidence that NMDA NR2B receptor is critical for chronic pain. Changes in NMDA NR2B mediated EPSCs by nerve injury could also be abolished in AC1 lacking mice. ACC is a critical structure for generating emotional response and encoding pain affects [42,43]. Short and long-term plasticities of ACC synapses have been believed to be the cellular basis for the development of chronic pain [44,45]. Lesions of the ACC in patients and inhibition of NMDA receptor/cAMP pathways in the ACC have been shown to be analgesic in chronic pain conditions [46], providing key basic evidence for future chronic pain treatment.

Conclusion

The cortical activity of the ACC and S1 is desynchronized during noxious mechanical stimulation, while the neural membrane potential is locked in a persistent Up state. Our study provides a useful model for the detailed study of the electrophysiology of the cortex under different states. The activated cortical response to nociceptive stimulation in our study provides a measure for monitoring analgesia during surgery. The results suggest that

desynchronization of EEG recordings during surgery should be avoided. Although parecoxib was not effective in inhibiting the cortical response to acute pain in our study, other studies with an alternative dosing schedule should be conducted.

Materials and methods

Animal preparation

Adult Sprague-Dawley rats weighing 240-330 g were group housed, 5 per cage and maintained under a 12-h dark-light cycle at $22 \pm 1^\circ\text{C}$. Food and water were available ad libitum. All procedures complied with the Animal Use Committee of Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences. Following initial anesthesia with urethane (1.5 g/kg, i.p.), animals were mounted on a stereotaxic apparatus. Lidocaine was applied locally to the scalp prior to surgery. A sufficient level of anesthesia was maintained throughout surgery such that animals had no spontaneous whisker or eyelid movement. A ceramic pressure sensor was used to monitor respiration and heart rate. Rectal temperature was maintained at $37\text{-}38^\circ\text{C}$ through an auto feedback electrical heating blanket. Supplementary doses (one-fifth of initial dose) of urethane were given when necessary. Small craniotomies were made for microelectrode implantation. According to the rat atlas of Paxinos and Watson [47], stereotaxic coordinates were: (1) for SI, 2.5 mm posterior to Bregma (-2.5 A) and 5.0 mm lateral to midline (L); (2) for ACC, 2.0 A, 0.8 L and 1.0-3.0 mm ventral to the brain surface; (3) for V1, -7.0 A, 3.0 L; (4) for PtA, -4.5 A, 4.0 L; and FrA, -5.0 A, 2.0 L. A 2.0 mm diameter incision was made at the corresponding areas by dental drill, and epidural mater was removed to expose the cortex.

Experimental procedure

For extracellular recordings a tungsten microelectrode (impedance of approximately $450\text{ K}\Omega$) was slowly lowered into the ACC with a microdriver (Narishige) under a dissecting microscope while a reference electrode was placed just below the scalp. The extracellular signal (0.1 to 3000 Hz) was acquired at a sampling frequency of 20 KHz and then passed through an amplifier (MODEL 2000, A-M system), digitalized by a MICRO 1401 A/D converter (Cambridge Electronic Design) and then stored with Spike 2 software (Cambridge Electronic Design).

Intracellular recordings from barrel cortical neurons were performed by sharp glass microelectrodes pulled from glass capillaries with filament (outer diameter of 1.5 mm). The electrode (filled with 2 M potassium acetate) had a tip resistance of 60-100 M Ω . After removing the epidural mater, the electrode was advanced to the surface of brain tissue with a motorized manipulator

(Narishige) under a dissecting microscope. The cortical exposure was sealed with low-melting agar to reduce fluctuation of brain tissue when the electrode was lowered into the brain. The electrode was slowly advanced with the microdriver at 3-5 μm steps. Intracellular recordings began after ACC extracellular activity was enhanced by tail pinch. The electrical signal was amplified by an AxoClamp2B amplifier (Molecular Devices), converted by a MICRO 1401 and stored for offline analysis. Data from neurons with a resting membrane potential below -60 mV and spontaneous spike amplitude > 50 mV were included in the study.

After acquiring a stable extracellular and intracellular recording, a neuronal response was elicited by pinching the tail using small forceps (constant force: $\sim 350\text{ g/mm}^2$) for about 15 s. The stimulus was applied 2 or 3 times with an inter-stimulus interval of 5 min. The effect of parecoxib (20 mg/kg, i.m.) or fentanyl (30 $\mu\text{g/kg}$, i.m.) on the mechanical stimulation induced response was tested, and repeated stimulus was applied 20 min or 5 min after each drug injection, respectively. Naloxone (30 $\mu\text{g/kg}$, i.m.) was used to reverse fentanyl's action. In addition, abdominal aorta blood was collected and measured for blood gas analysis from eight rats before or 5-10 min after fentanyl injection.

Data analysis

Raw signals were visually inspected and artifacts were deleted before analysis. MATLAB (Mathworks, OriginLab 8 (OriginLab) and Spike 2 software were used for data analysis. FFT (fast Fourier transform, Hanning window) was performed on the EEG data (filtered out at 1-100 Hz) to obtain a power spectrum from which we calculated the integrations of power within the following bands [48]: total, 1-100 Hz; δ , 1-4 Hz; θ , 4-8 Hz; α , 8-12 Hz; β , 12-25 Hz; γ , 25-100 Hz. For a more specific description, we plotted the cumulative power distribution (CPD) curve based on the power spectrum by directly acquiring MF and SEF (frequency below which 50% and 95% of total power resides, respectively). We then measured the "distance" of cumulative power distribution, defined by the area of the region between any two given cumulative curves. This measure quantifies the change of power distribution under two corresponding experimental conditions. A similar analysis was used in describing the change of the membrane potential distribution from intracellular recordings.

Multi-unit data were extracted from the extracellular signal through a band-pass filter (600~3000 Hz), and appropriate thresholds were manually selected for spike discrimination. To describe the uniformity of the spike train in the absence and the presence of the nociceptive stimulus, we defined a "Non-uniform Index" of any given time period T with a spike number of N as

$$\text{Nouniform Index} = \frac{1}{N-2} \left(\sum_{i=1}^{N-1} \left(\frac{t_{i+1}-t_i}{T/(N-1)} - 1 \right)^2 \right)^{\frac{1}{2}}$$

where t_i stands for the time of the i th spike. This index adequately caught the distinct characteristics of the “Up and Down” in contrast to the persistent “Up” firing patterns, regardless of their overall mean firing rates.

To collect data from intracellular recordings, we first distinguished the Up and Down states by comparing the average membrane potential during a given time window (~10 ms) with that of a much larger time span (~3 s). We then counted the frequency of membrane potential transitions between the two states. In addition, as described above for extracellular recording analysis, the distance between two cumulative curves was used to quantify differences between experimental groups.

Statistical analysis

Data were normally distributed and presented as mean \pm SEM. Statistical significance was tested with paired t -tests using $p < 0.05$ as statistically significant.

List of abbreviations

EEG: electroencephalogram; S1: primary somatosensory cortex; S2: secondary somatosensory cortex; ACC: anterior cingulate cortex; LTP: long-term potentiation; NSAIDs: anti-inflammatory drugs; COX-2: cyclooxygenase-2; PtA: parietal association cortex; V1: primary visual cortex; FrA: frontal association cortex; CPD: cumulative power distribution; MF: medial frequency; SEF: 95% spectral edge frequency; MU: Multi-unit; V_m : membrane potential; AMPA: α -amino-3-hydroxyl-5-methyl-4-isoxazole-propionate; NMDA: N-methyl-D-aspartic acid.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (No. 30872443), Shuguang Program (08SG18) of Shanghai Municipal Education Commission and Rising Star Program of Shanghai (No. 09QH1401800). Thanks for Dr. Min Zhuo's suggestions on revising the manuscript (Professor of Physiology, University of Toronto, Canada).

Authors' contributions

YZP participated in the design of the study, carried out the experiment, and drafted the manuscript. XXL assisted with the electrophysiological recording. YWW conceived of the study, assisted with the data analysis, and wrote the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 9 November 2009

Accepted: 21 January 2010 Published: 21 January 2010

References

1. Ohara PT, Vit JP, Jasmin L: **Cortical modulation of pain.** *Cell Mol Life Sci* 2005, **62**:44-52.
2. Talbot JD, Marrett S, Evans AC, Meyer E, Bushnell MC, Duncan GH: **Multiple representations of pain in human cerebral cortex.** *Science* 1991, **251**:1355-8.
3. Apkarian AV, Bushnell MC, Treede RD, Zubieta JK: **Human brain mechanisms of pain perception and regulation in health and disease.** *Eur J Pain* 2005, **9**:463-84.
4. Steriade M, Nunez A, Amzica F: **A novel slow (< 1 Hz) oscillation of neocortical neurons in vivo: depolarizing and hyperpolarizing components.** *J Neurosci* 1993, **13**:3252-65.
5. Steriade M, Timofeev I, Grenier F: **Natural waking and sleep states: a view from inside neocortical neurons.** *J Neurophysiol* 2001, **85**:1969-85.
6. Sanchez-Vives MV, McCormick DA: **Cellular and network mechanisms of rhythmic recurrent activity in neocortex.** *Nat Neurosci* 2000, **3**:1027-34.
7. Wei F, Vadakkan KI, Toyoda H, Wu LJ, Zhao MG, Xu H, Shum FW, Jia YH, Zhuo M: **Calcium calmodulin-stimulated adenylyl cyclases contribute to activation of extracellular signal-regulated kinase in spinal dorsal horn neurons in adult rats and mice.** *J Neurosci* 2006, **26**:851-61.
8. Ikeda H, Stark J, Fischer H, Wagner M, Drdla R, Jager T, Sandkuhler J: **Synaptic amplifier of inflammatory pain in the spinal dorsal horn.** *Science* 2006, **312**:1659-62.
9. Wheeler M, Oderda GM, Ashburn MA, Lipman AG: **Adverse events associated with postoperative opioid analgesia: a systematic review.** *J Pain* 2002, **3**:159-80.
10. Giron I, Milne B, Hong M: **Cyclooxygenase-2 inhibitors in postoperative pain management: current evidence and future directions.** *Anesthesiology* 2003, **99**:1198-208.
11. Gajraj NM: **Cyclooxygenase-2 inhibitors.** *Anesth Analg* 2003, **96**:1720-38.
12. Puolakka PA, Puura AI, Pirhonen RA, Ranta AU, Autio V, Lindgren L, Rorarius MG: **Lack of analgesic effect of parecoxib following laparoscopic cholecystectomy.** *Acta Anaesthesiol Scand* 2006, **50**:1027-32.
13. Antognini JF, Wang XW, Carstens E: **Isoflurane action in the spinal cord blunts electroencephalographic and thalamic-reticular formation responses to noxious stimulation in goats.** *Anesthesiology* 2000, **92**:559-66.
14. Nagata T, Suzuki H, Zhang R, Ozaki M, Kawakami Y: **Mechanical stimulation activates small fiber mediated nociceptive responses in the nucleus gigantocellularis.** *Exp Brain Res* 2003, **149**:505-11.
15. Kuo CC, Chiou RJ, Liang KC, Yen CT: **Differential involvement of the anterior cingulate and primary sensorimotor cortices in sensory and affective functions of pain.** *J Neurophysiol* 2009, **101**:1201-10.
16. Bischoff P, Kochs E, Haferkorn D, Schulte am Esch J: **Intraoperative EEG changes in relation to the surgical procedure during isoflurane-nitrous oxide anesthesia: hysterectomy versus mastectomy.** *J Clin Anesth* 1996, **8**:36-43.
17. Rudolph M, Pelletier JG, Pare D, Destexhe A: **Characterization of synaptic conductances and integrative properties during electrically induced EEG-activated states in neocortical neurons in vivo.** *J Neurophysiol* 2005, **94**:2805-21.
18. Steriade M, Amzica F, Contreras D: **Synchronization of fast (30-40 Hz) spontaneous cortical rhythms during brain activation.** *J Neurosci* 1996, **16**:392-417.
19. Romsing J, Moiniche S: **A systematic review of COX-2 inhibitors compared with traditional NSAIDs, or different COX-2 inhibitors for post-operative pain.** *Acta Anaesthesiol Scand* 2004, **48**:525-46.
20. Barton SF, Langeland FF, Snabes MC, LeComte D, Kuss ME, Dhadda SS, Hubbard RC: **Efficacy and safety of intravenous parecoxib sodium in relieving acute postoperative pain following gynecologic laparotomy surgery.** *Anesthesiology* 2002, **97**:306-14.
21. Desjardins PJ, Grossman EH, Kuss ME, Talwalker S, Dhadda S, Baum D, Hubbard RC: **The injectable cyclooxygenase-2-specific inhibitor parecoxib sodium has analgesic efficacy when administered preoperatively.** *Anesth Analg* 2001, **93**:721-7.
22. Maihofner C, Ringler R, Herrndobler F, Koppert W: **Brain imaging of analgesic and antihyperalgesic effects of cyclooxygenase inhibition in an experimental human pain model: a functional MRI study.** *Eur J Neurosci* 2007, **26**:1344-56.

23. Grundmann U, Wornle C, Biedler A, Kreuer S, Wrobel M, Wilhelm W: **The efficacy of the non-opioid analgesics parecoxib, paracetamol and metamizol for postoperative pain relief after lumbar microdiscectomy.** *Anesth Analg* 2006, **103**:217-22.
24. Riest G, Peters J, Weiss M, Dreyer S, Klassen PD, Stegen B, Bello A, Eikermann M: **Preventive effects of perioperative parecoxib on post-discectomy pain.** *Br J Anaesth* 2008, **100**:256-62.
25. Padi SS, Jain NK, Singh S, Kulkarni SK: **Pharmacological profile of parecoxib: a novel, potent injectable selective cyclooxygenase-2 inhibitor.** *Eur J Pharmacol* 2004, **491**:69-76.
26. Beiche F, Scheuerer S, Brune K, Geisslinger G, Goppelt-Struebe M: **Up-regulation of cyclooxygenase-2 mRNA in the rat spinal cord following peripheral inflammation.** *FEBS Lett* 1996, **390**:165-9.
27. Wu ZZ, Chen SR, Pan HL: **Differential sensitivity of N- and P/Q-type Ca²⁺ channel currents to a mu opioid in isolectin B4-positive and -negative dorsal root ganglion neurons.** *J Pharmacol Exp Ther* 2004, **311**:939-47.
28. Reeve AJ, Dickenson AH, Kerr NC: **Spinal effects of bicuculline: modulation of an allodynia-like state by an A1-receptor agonist, morphine, and an NMDA-receptor antagonist.** *J Neurophysiol* 1998, **79**:1494-507.
29. Chen YP, Chen SR, Pan HL: **Effect of morphine on deep dorsal horn projection neurons depends on spinal GABAergic and glycinergic tone: implications for reduced opioid effect in neuropathic pain.** *J Pharmacol Exp Ther* 2005, **315**:696-703.
30. Wu MF, Pang ZP, Zhuo M, Xu ZC: **Prolonged membrane potential depolarization in cingulate pyramidal cells after digit amputation in adult rats.** *Mol Pain* 2005, **1**:23.
31. Ingram SL: **Cellular and molecular mechanisms of opioid action.** *Prog Brain Res* 2000, **129**:483-92.
32. Benrath J, Brechtel C, Martin E, Sandkuhler J: **Low doses of fentanyl block central sensitization in the rat spinal cord in vivo.** *Anesthesiology* 2004, **100**:1545-51.
33. Deyama S, Yamamoto J, Machida T, Tanimoto S, Nakagawa T, Kaneko S, Satoh M, Minami M: **Inhibition of glutamatergic transmission by morphine in the basolateral amygdaloid nucleus reduces pain-induced aversion.** *Neurosci Res* 2007, **59**:199-204.
34. Wei F, Li P, Zhuo M: **Loss of synaptic depression in mammalian anterior cingulate cortex after amputation.** *J Neurosci* 1999, **19**:9346-54.
35. Zhao MG, Toyoda H, Lee YS, Wu LJ, Ko SW, Zhang XH, Jia Y, Shum F, Xu H, Li BM, Kaang BK, Zhuo M: **Roles of NMDA NR2B subtype receptor in prefrontal long-term potentiation and contextual fear memory.** *Neuron* 2005, **47**:859-72.
36. Wu LJ, Xu H, Ren M, Cao X, Zhuo M: **Pharmacological isolation of postsynaptic currents mediated by NR2A- and NR2B-containing NMDA receptors in the anterior cingulate cortex.** *Mol Pain* 2007, **3**:11.
37. Liauw J, Wu LJ, Zhuo M: **Calcium-stimulated adenylyl cyclases required for long-term potentiation in the anterior cingulate cortex.** *J Neurophysiol* 2005, **94**:878-82.
38. Ko SW, Vadakkan KI, Ao H, Gallitano-Mendel A, Wei F, Milbrandt J, Zhuo M: **Selective contribution of Egr1 (zif/268) to persistent inflammatory pain.** *J Pain* 2005, **6**:12-20.
39. Wei F, Wang GD, Kerchner GA, Kim SJ, Xu HM, Chen ZF, Zhuo M: **Genetic enhancement of inflammatory pain by forebrain NR2B overexpression.** *Nat Neurosci* 2001, **4**:164-9.
40. Xu H, Wu LJ, Wang H, Zhang X, Vadakkan KI, Kim SS, Steenland HW, Zhuo M: **Presynaptic and postsynaptic amplifications of neuropathic pain in the anterior cingulate cortex.** *J Neurosci* 2008, **28**:7445-53.
41. Wu LJ, Toyoda H, Zhao MG, Lee YS, Tang J, Ko SW, Jia YH, Shum FW, Zerbiniatti CV, Bu G, Wei F, Xu TL, Muglia LJ, Chen ZF, Auberson YP, Kaang BK, Zhuo M: **Upregulation of forebrain NMDA NR2B receptors contributes to behavioral sensitization after inflammation.** *J Neurosci* 2005, **25**:11107-16.
42. Tang J, Ko S, Ding HK, Qiu CS, Calejesan AA, Zhuo M: **Pavlovian fear memory induced by activation in the anterior cingulate cortex.** *Mol Pain* 2005, **1**:6.
43. Rainville P, Duncan GH, Price DD, Carrier B, Bushnell MC: **Pain affect encoded in human anterior cingulate but not somatosensory cortex.** *Science* 1997, **277**:968-71.
44. Shyu BC, Vogt BA: **Short-term synaptic plasticity in the nociceptive thalamic-anterior cingulate pathway.** *Mol Pain* 2009, **5**:51.
45. Zhuo M: **Cortical excitation and chronic pain.** *Trends Neurosci* 2008, **31**:199-207.
46. Zhuo M: **Neuronal mechanism for neuropathic pain.** *Mol Pain* 2007, **3**:14.
47. Paxinos G, Watson C: **The Rat Brain in Stereotaxic Coordinates.** New York: Academic Press 1998.
48. John ER: **The neurophysics of consciousness.** *Brain Res Brain Res Rev* 2002, **39**:1-28.

doi:10.1186/1744-8069-6-3

Cite this article as: Peng et al: Effects of Parecoxib and Fentanyl on nociception-induced cortical activity. *Molecular Pain* 2010 **6**:3.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

