

# Role of SERK genes in plant environmental response

Marcelo O. Santos<sup>1</sup> and Francisco J.L. Aragão<sup>2,\*</sup>

<sup>1</sup>Departamento de Biologia; ICB; Universidade Federal de Juiz de Fora; Juiz de Fora, MG Brazil; <sup>2</sup>Embrapa Recursos Genéticos e Biotecnologia; Brasília, DF Brazil

**Key words:** SERK, fungus tolerance, environmental stress, brassinosteroids, SAR

In plants, cell signaling connects the environmental input to the intracellular responses in plants. Exogenous signals play an important role in cell metabolism leading to growth and defense responses. Some of these stimuli induce anatomical and physiological modifications that are generally modulated by gene expression. *SERK* belongs to a small family of genes that code for a transmembrane protein involved in signal transduction and that have been strongly associated with somatic embryogenesis and apomixis in a number of plant species. Recent studies corroborate its role in somatic embryogenesis and suggest a broader range of functions in plant response to biotic and abiotic stimuli. This mini-review aims to present new data on *SERK* and discuss its involvement in plant development as well as in response to environmental stress.

## Role of SERK in Plant Tissue Organogenesis and Somatic Embryogenesis

During somatic embryogenesis, biochemical and morphological changes occur throughout the development of induced tissues, which is closely related to alterations in gene expression. Several genes are differentially expressed during somatic embryogenesis induction, while others are expressed during differentiation from embryo maturation up to full plant development. Among the genes involved in the induction of somatic embryogenesis, the *Somatic Embryogenesis Receptor Kinase (SERK)* gene is claimed to have an important role. *SERK* gene was first isolated from carrot embryogenic cells, hailed as a molecular marker for somatic embryogenesis<sup>1,2</sup> and associated with somatic embryogenesis (SE) in a number of species including *Dactylis glomerata*,<sup>2</sup> *Arabidopsis thaliana*,<sup>3</sup> *Medicago truncatula*,<sup>4</sup> *Helianthus annuus*,<sup>5</sup> *Ocotea catharinensi*,<sup>6</sup> *Citrus unshiu*<sup>7</sup> and *Theobroma cacao*<sup>8</sup> and lately for apomixes.<sup>9,10</sup> *SERK* is a leucine-rich repeat (LRR) transmembrane protein kinase that enhances the ability of the apical meristem in *Arabidopsis* to form somatic embryos.<sup>3</sup> LRR kinases transmit their signal by forming homodimers or heterodimers with other RLKs, in response to binding by a ligand. This ligand-induced dimerization causes phosphorylation of the intracellular kinase domains of the receptor-like kinases (RLKs), which activates the

next stages of the signal transduction pathway (Fig. 1). There is potential for different levels of complexity in the signaling through variation in the binding partners of different RLKs.<sup>11</sup>

In sunflower, *SERK* expression is correlated to induction of two different developmental pathways, somatic embryogenesis and shoot organogenesis.<sup>5</sup> In spite of its role in plant development, different genes from the family showed a different expression pattern during somatic or zygotic embryogenesis<sup>12</sup> and the expression of these genes was differentially detected in different tissues in several plant species, such as in the apical meristem, root and leaves,<sup>5,8,11,13,14</sup> suggesting additional roles for this gene. T-DNA mutational insertions showed the redundancy of two *SERK* homologs and its role in male sporogenesis in *Arabidopsis*.<sup>15,16</sup>

Somatic embryogenesis is an artificial simulation of natural embryogenesis and may be induced by different signalling stresses such as osmotic pressure, ABA induction and SA pathway, which are generally activated by auxin, mainly 2,4-D, which promotes oxidative stress in plants or similar molecules. The *SERK* pattern of expression in cultures suggests that it is part of a signaling pathway that mediates developmental changes in cells in response to culture conditions. These developmental changes involve the initiation of cell division and cellular reprogramming. Cell fate involving organogenesis or somatic embryogenesis might be conditioned by the status of *SERK* gene expression, and it is supposed to be a key factor in differentiation.<sup>5,11</sup> *SERK* gene expression was strongly related to auxin elicitation, a known somatic embryogenesis inducer.<sup>4,17,18</sup>

Brassinosteroid is an essential class of plant growth regulator involved in signal transduction during different plant growth events<sup>19</sup> that include salinity stress,<sup>20</sup> osmotic stress during seed germination<sup>21</sup> or disease resistance.<sup>22</sup> It was shown that BRASSINOSTEROID-INSENSITIVE1 (BRI1) receptor co-precipitates and interacts in vivo with SERK1.<sup>23</sup> *SERK* autophosphorylates in the threonine serine site<sup>24</sup> and is involved in transphosphorylation of BRI, which is enhanced under brassinosteroid treatment.<sup>25</sup> Although *SERK* gene belongs to a family that may show redundant roles and differential expression, some evolutionary changes in protein function were observed and differences in spatial expression may not be the only way to control its role.<sup>26</sup>

## SERK as a Broader Signal in Response to Biotic and Abiotic Stress


In addition to the study of plant development and totipotency of cells, studies have been carried out with plants exposed to

\*Correspondence to: Francisco J.L. Aragão; Email: aragao@cenargen.embrapa.br

Submitted: 08/19/09; Accepted: 08/22/09

Previously published online:

www.landesbioscience.com/journals/psb/article/9900



**Figure 1.** Somatic embryogenic receptor kinase (SERK) signaling. The SERK gene is expressed under auxin or other Plant Growth Regulator (PRG) stimuli. The SERK protein, a membrane receptor, phosphorylates the brassinosteroid receptor under appropriate external signal, to upregulate pathogen-related (PR) genes or disease response genes, increasing plant pathogen tolerance. SERK is involved in the expression of stress-response genes, and developmental pathways such as somatic embryogenesis or shoot organogenesis.

different sources of stress, involving plant growth regulators, elicitors and signal transduction (Fig. 1). These plant stresses can be separated into two main classes: biotic stresses including herbivores, fungal and bacterial attack or virus infection, and a second class that includes abiotic stresses such as drought, extreme temperature, salinity, light etc. Environmental interactions are mediated by elicitors signaling molecules through the transduction pathway.<sup>27</sup>

Recent advances in research on gene regulation have led to the emergence of post-transcriptional gene silencing (PTGS) or epigenetic variation as additional levels of gene regulation.<sup>28</sup> In recent work we have produced a stable PTGS silencing state in transgenic lettuce carrying a *SERK* gene antisense cassette.<sup>29</sup> These lettuce plants exhibited *LsSERK* gene silencing, showed reduced somatic embryogenesis ability and became more susceptible to Sclerotinia attack. Thus, the elimination of *LsSERK* via RNA silencing

corroborates the hypothesis that *SERK* is involved in somatic embryogenesis and also in plant defence, since these plants had less ability to form somatic embryos or to resist fungus attack.<sup>29</sup>

Similarly, increased somatic embryogenesis ability was induced by overexpressing *OsSERK1*, controlled by the constitutive CaMV35S promoter, in rice plants.<sup>13</sup> These transgenic rice plants showed an increase resistance to *Magnapothe grisea*.<sup>13</sup> Reinforcing the idea that *SERK* is involved in response to biotic stress, transgenic rice expressing the bacterial flagellin (N1141) gene triggered the activation of innate immune response and increased resistance to *M. grisea*.<sup>30</sup> The *OsBISERK* gene showed upregulation under benzothiadiazole (BTH) treatment, which also leads to an increase in rice resistance to *M. grisea*.<sup>31</sup> It has been shown that BTH induces systemic acquired resistance (SAR) and activates the gene involved in disease resistance.<sup>32</sup> In *Vitis vinifera* differential expression of *SERK* genes was observed concomitantly to some PR proteins upregulated by 2,4-D, a known somatic embryogenesis inducer.<sup>14</sup>

On the other hand, it was demonstrated that the auxin naphthalene acetic acid (NAA) was able to upregulate *SERK* gene in *Medicago truncatula*.<sup>4</sup> In rice some defense signaling molecules, such as SA and JA, induced *SERK* upregulation and a slight induction by ABA.<sup>13</sup> SERK3/BAK1 proteins act on BR signaling and were associated as a component of pathogen-associated molecular pattern (PAMP)-triggered immunity by flagellin elicitation (PTI).<sup>33,34</sup>

## Concluding Remarks

Current data collectively lead to the idea that *SERK* may be involved in plant signaling and multiple other functions, and not only in plant development. *SERK* is highly expressed in embryogenic tissues and some undifferentiated cells,<sup>35</sup> preceding and coinciding with early somatic embryogenesis.<sup>36</sup> Plant cell pluripotency may be associated with chromatin folding<sup>37</sup> and the pattern of DNA methylation, which determine plant cell differentiation.<sup>38,39</sup> Additional investigation of *SERK* gene methylation patterns during plant development and under biotic and abiotic stress response would increase knowledge on its role in plant environmental response.

## References

- Schmidt ED, Guzzo F, Toonen MA, de Vries SC. A leucine-rich repeat containing receptor-like kinase marks somatic plant cells competent to form embryos. *Development* 1997; 124:2049-62.
- Somleva MN, Schmidt EDL, de Vries SC. Embryogenic cells in *Dactylis glomerata* L. (Poaceae) explants identified by cell tracking and by *SERK* expression. *Plant Cell Rep* 2000; 19:718-26.
- Hecht V, Vielle-Calzada JP, Hartog MV, Schmidt ED, Boutilier K, Grossniklaus U, et al. The Arabidopsis SOMATIC EMBRYOGENESIS RECEPTOR KINASE 1 gene is expressed in developing ovules and embryos and enhances embryogenic competence in culture. *Plant Physiol* 2001; 127:803-16.
- Nolan KE, Irwanto RR, Rose RJ. Auxin upregulates *MtSERK1* expression in both *Medicago truncatula* root-forming and embryogenic cultures. *Plant Physiol* 2003; 133:218-30.
- Thomas C, Meyer D, Hember C, Steinmetz A. Spatial expression of a sunflower *SERK* gene during induction of somatic embryogenesis and shoot organogenesis. *Plant Physiol Biochem* 2004; 42:35-42.
- Santa-Catarina C, Hanai LR, Dornelas MC, Viana AM, Floh EIS. SERK gene homolog expression, polyamines and amino acids associated with somatic embryogenic competence of *Ocotea catharinensis* Mez. (Lauraceae). *Plant Cell Tiss Org Cult* 2004; 79:53-61.
- Shimada T, Hirabayashi T, Endo T, Fujii H, Kita M, Omura M. Isolation and characterization of the somatic embryogenesis receptor-like kinase gene homologue (*CitSERK1*) from *Citrus unshiu* Marc. *Sci Hort* 2005; 103:233-8.
- Santos MO, Romano E, Yotoko KSC, Tinoco MLP, Dias BBA, Aragão FJL. Characterisation of the cacao somatic embryogenesis receptor-like kinase (SERK) gene expressed during somatic embryogenesis. *Plant Sci* 2005; 168:723-9.
- Tucker MR, Araujo AC, Paech NA, Hecht V, Schmidt ED, Rossell JB, et al. Sexual and apomictic reproduction in Hieracium subgenus pilosella are closely interrelated developmental pathways. *Plant Cell* 2003; 15:1524-37.
- Albertini E, Marconi G, Reale L, Barcaccia G, Porceddu A, Ferranti F, et al. SERK and APOSTART. Candidate genes for apomixis in *Poa pratensis*. *Plant Physiol* 2005; 138:2185-99.
- Nolan KE, Kurdyukov S, Rose RJ. Expression of the SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASE1 (*SERK1*) gene is associated with developmental change in the life cycle of the model legume *Medicago truncatula*. *J Exp Bot* 2009; 60:1759-71.
- Baudino S, Hansen S, Brettschneider R, Hecht VF, Dresselhaus T, Lörz H, et al. Molecular characterisation of two novel maize LRR receptor-like kinases, which belong to the *SERK* gene family. *Planta* 2001; 213:1-10.

13. Hu H, Xiong L, Yang Y. Rice *SERK1* gene positively regulates somatic embryogenesis of cultured cell and host defense response against fungal infection. *Planta* 2005; 222:107-17.
14. Maillot P, Lebel S, Schellenbaum P, Jacques A, Walter B. Differential regulation of SERK, LEC1-like and pathogenesis-related genes during indirect secondary somatic embryogenesis in grapevine. *Plant Physiol Biochem* 2009; 47:743-52.
15. Albrecht C, Russinova E, Hecht V, Baaijens E, de Vries S. The *Arabidopsis thaliana* SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASES1 and 2 control male sporogenesis. *Plant Cell* 2005; 17:3337-49.
16. Colcombet J, Boisson-Dernier A, Ros-Palau R, Vera CE, Schroeder JI. Arabidopsis SOMATIC EMBRYOGENESIS RECEPTOR KINASES1 and 2 are essential for tapetum development and microspore maturation. *Plant Cell* 2005; 17:3350-61.
17. Sharma SK, Millam S, Hein I, Bryan GJ. Cloning and molecular characterisation of a potato *SERK* gene transcriptionally induced during initiation of somatic embryogenesis. *Planta* 2008; 228:319-30.
18. Singla B, Khurana JP, Khurana P. Characterization of three somatic embryogenesis receptor kinase genes from wheat, *Triticum aestivum*. *Plant Cell Rep* 2008; 27:833-43.
19. Li J, Chory J. Brassinosteroid action in plants. *J Exp Bot* 1999; 50:275-82.
20. Anuradha S, Rao SSR. Effect of brassinosteroids on salinity stress induced inhibition of seed germination and seedling growth of rice (*Oryza sativa* L.). *Plant Growth Regul* 2001; 33:151-3.
21. Vardhini VB, Rao SSR. Amelioration of osmotic stress by brassinosteroids on seed germination and seedling growth of three varieties of sorghum. *Plant Growth Regul* 2003; 41:25-31.
22. Nakashita H, Yasuda M, Nitta T, Asami T, Fujioka S, Arai Y, et al. Brassinosteroid functions in a broad range of disease resistance in tobacco and rice. *Plant J* 2003; 33:887-98.
23. Karlova R, Boeren S, Russinova E, Aker J, Vervoort J, de Vries S. The Arabidopsis SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASE1 Protein Complex Includes BRASSINOSTEROID-INSENSITIVE1. *The Plant Cell* 2006; 18:626-38.
24. Shah K, Schmidt ED, Vlak JM, de Vries SC. Expression of the *Daucus carota* somatic embryogenesis receptor kinase (DcSERK) protein in insect cells. *Biochimie* 2001; 83:415-21.
25. Karlova R, Boeren S, van Dongen W, Kwaaitaal M, Aker J, Vervoort J, et al. Identification of in vitro phosphorylation sites in the *Arabidopsis thaliana* somatic embryogenesis receptor-like kinases. *Proteomics* 2009; 9:368-79.
26. Albrecht C, Russinova E, Kemmerling B, Kwaaitaal M, de Vries SC. Arabidopsis SOMATIC EMBRYOGENESIS RECEPTOR KINASE proteins serve brassinosteroid-dependent and -independent signaling pathways. *Plant Physiol* 2008; 148:611-9.
27. de Wit PJ. How plants recognize pathogens and defend themselves. *Cell Mol Life Sci* 2007; 64:2726-32.
28. Hollick JB. Sensing the epigenome. *Trends Plant Sci* 2008; 13:398-404.
29. Santos MO, Romano E, Vieira LS, Baldoni AB, Aragão FJL. Suppression of SERK gene expression affects fungus tolerance and somatic embryogenesis in transgenic lettuce. *Plant Biol* 2009; 11:83-9.
30. Takakura Y, Che FS, Ishida Y, Tsutsumi F, Kurotani K, Usami S, et al. Expression of a bacterial flagellin gene triggers plant immune responses and confers disease resistance in transgenic rice plants. *Mol Plant Pathol* 2008; 9:525-9.
31. Song D, Li G, Song F, Zheng Z. Molecular characterization and expression analysis of *OsBISERK1*, a gene encoding a leucine-rich repeat receptor-like kinase, during disease resistance responses in rice. *Mol Biol Rep* 2009; 35:275-83.
32. Grolach J, Volrath S, Knauf-Beite G, Hengy G, Beckhove U, Kogel K, et al. Benzothiadiazole, a novel class of inducers of systemic acquired resistance, activates gene expression and disease resistance in wheat. *Plant Cell* 1996; 8:629-43.
33. Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nürnberger T, Jones JD, et al. A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. *Nature* 2007; 448:497-500.
34. Heese A, Hann DR, Gimenez-Ibanez S, Jones AME, He K, Li J, et al. The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. *Proc Natl Acad Sci USA* 2007; 104:12217-22.
35. Kwaaitaal MA, de Vries SC, Russinova E. *Arabidopsis thaliana* Somatic Embryogenesis Receptor Kinase 1 protein is present in sporophytic and gametophytic cells and undergoes endocytosis. *Protoplasma* 2005; 226:55-65.
36. Salaj J, von Recklinghausen IR, Hecht V, de Vries SC, Schel JH, van Lammeren AA. *AtSERK1* expression precedes and coincides with early somatic embryogenesis in *Arabidopsis thaliana*. *Plant Physiol Biochem* 2008; 46:709-14.
37. Verdeil JL, Alemanno L, Niemenak N, Tranbarger TJ. Pluripotent versus totipotent plant stem cells: dependence versus autonomy? *Trends Plant Sci* 2007; 12:245-52.
38. Gehring M, Henikoff S. DNA methylation dynamics in plant genomes. *Biochim Biophys Acta* 2007; 1769:276-86.
39. Gehring M, Reik W, Henikoff S. DNA demethylation by DNA repair. *Trends Genet* 2009; 25:82-90.