

RESEARCH ARTICLE

Open Access

# Chromosome 7p linkage and association study for diabetes related traits and type 2 diabetes in an African-American population enriched for nephropathy

Tennille S Leak<sup>1,2</sup>, Carl D Langefeld<sup>3</sup>, Keith L Keene<sup>1,4,8</sup>, Carla J Gallagher<sup>1,6,7</sup>, Lingyi Lu<sup>3</sup>, Josyf C Mychaleckyj<sup>4,5</sup>, Stephen S Rich<sup>4,5</sup>, Barry I Freedman<sup>9</sup>, Donald W Bowden<sup>1,6,9</sup>, Michèle M Sale<sup>1,4,8,10\*</sup>

## Abstract

**Background:** Previously we performed a linkage scan of 638 African American affected sibling pairs (ASP) with type 2 diabetes (T2D) enriched for end-stage renal disease (ESRD). Ordered subset linkage analysis (OSA) revealed a linkage peak on chromosome 7p in the subset of families with earlier age of T2D diagnosis.

**Methods:** We fine mapped this region by genotyping 11 additional polymorphic markers in the same ASP and investigated a total of 68 single nucleotide polymorphisms (SNPs) in functional candidate genes (*GCK1*, *IL6*, *IGFBP1* and *IGFBP3*) for association with age of T2D diagnosis, age of ESRD diagnosis, duration of T2D to onset of ESRD, body mass index (BMI) in African American cases and T2D-ESRD in an African American case-control cohort. OSA of fine mapping markers supported linkage at 28 cM on 7p (near D7S3051) in early-onset T2D families (max. LOD = 3.61,  $P = 0.002$ ). SNPs in candidate genes and 70 ancestry-informative markers (AIMs) were evaluated in 577 African American T2D-ESRD cases and 596 African American controls.

**Results:** The most significant association was observed between ESRD age of diagnosis and SNP rs730497, located in intron 1 of the *GCK1* gene (recessive T2D age-adjusted  $P = 0.0006$ ). Nominal associations were observed with *GCK1* SNPs and T2D age of diagnosis (BMI-adjusted  $P = 0.014$  to  $0.032$ ). Also, one *IGFBP1* and four *IGFBP3* SNPs showed nominal genotypic association with T2D-ESRD ( $P = 0.002$ - $0.049$ ). After correcting for multiple tests, only rs730497 remained significant.

**Conclusion:** Variant rs730947 in the *GCK1* gene appears to play a role in early ESRD onset in African Americans.

## Background

A genome wide linkage scan was performed on 638 African American affected sibling pairs (ASPs) with type 2 diabetes (T2D) from 247 families; 166 families contained at least one ASP concordant for diabetic end-stage renal disease (T2D-ESRD) [1]. Ordered subset analysis (OSA) revealed a linkage peak on chromosome 7p in the subset of T2D families with an early age of diagnosis (29% of pedigrees, max. LOD = 3.85,  $P = 0.003$  for the change in LOD score) [1]. T2D-ESRD subsets with lower body mass index (BMI) (64% of pedigrees, max. LOD = 3.93,  $P$

= 0.010) and longer duration from T2D diagnosis to ESRD onset (37% of pedigrees, max. LOD = 3.59,  $P = 0.010$ ) also showed evidence for linkage at this locus [2].

Fine mapping of this region using the same African American families was undertaken in order to localize the peak and to determine the primary phenotype of association. There are several plausible diabetes and nephropathy candidate genes in the resulting region of interest, including glucokinase isoform 1 (*GCK1*), interleukin-6 (*IL6*), insulin growth factor binding protein 1 (*IGFBP1*) and insulin growth factor binding protein 3 (*IGFBP3*).

Mutations of *GCK* have been identified in subjects with maturity-onset diabetes of the Young (MODY)

\* Correspondence: msale@virginia.edu

<sup>1</sup>Center for Human Genomics, Wake Forest University School of Medicine, Winston-Salem, NC, USA

[3,4]. However, single nucleotide polymorphisms (SNPs) in this gene have also been shown to be associated with typical T2D [5-7]. A dinucleotide (CA)<sub>n</sub> repeat element located approximately 10 kb 3' to the coding region has also shown association with T2D across several diverse ethnic populations [8-10].

The cytokine interleukin 6 (IL-6) is an essential regulator of the acute phase response associated with T2D and diabetic nephropathy [11,12]. Variants in the *IL6* promoter region have been shown to affect promoter strength [13].

The insulin like growth factors 1 and 2 circulate tightly bound to IGFBP-1 and IGFBP-3. A cross-sectional study of T2D cases and controls found that polymorphisms in *IGFBP3* was associated with levels of HbA<sub>1c</sub> [14]. Also, *in vivo* and *in vitro* studies have demonstrated that IGFBP-3 is a potent insulin antagonist [15].

We fine mapped the region of interest using additional microsatellite markers, and investigated associations with SNPs in four plausible candidate genes with age of T2D diagnosis, age of ESRD diagnosis, duration of T2D to onset of ESRD, BMI and T2D-ESRD in an African American cohort.

## Methods

### Selection criteria and recruitment of African American families

This study was conducted under Institutional Review Board approval from Wake Forest University School of Medicine and adhered to the tenets of the Declaration of Helsinki. All participants provided written informed consent. Clinical characteristics and recruitment of African American patients have been described previously [1,2]. DNA samples were collected from self-described African American families with multiple T2D affected members. Briefly, families were originally identified through a proband with impaired renal function associated with T2D. Medical records were reviewed to verify the etiology of the nephropathy. Impaired renal function was attributed to diabetes in the presence of the following criteria: serum creatinine  $\geq$  1.5 mg/dl, diabetes for >10 years or presence of proliferative diabetic retinopathy, and/or proteinuria  $\geq$  500 mg/24 h or >100 mg/dl, in the absence of other known causes of renal failure. T2D was diagnosed in patients developing diabetes after the age of 35 years and treated at the time of recruitment with oral hypoglycemic agents, insulin, or diet and exercise, where treatment was considered permanent (i.e., excluding steroid-induced diabetes and gestational diabetes). The family set for the genome-wide scan comprised 247 African American families with 638 ASPs, totaling 675 individuals.

### Case-control subjects

Identification, clinical characteristics, and recruitment of African Americans and European American patients and controls have been described previously [16]. Briefly, 577 unrelated African American patients with T2D, born in North Carolina, South Carolina, Georgia, Tennessee or Virginia, were recruited from dialysis facilities. T2D was diagnosed if patients reported an initial diagnosis of diabetes after 35 years of age, received dietary therapy or hypoglycemic agents in the absence of insulin alone for at least 1 year after diagnosis, and were currently receiving diabetes medications. Case subjects had severe diabetes accompanied by nephropathy, and T2D at least 5 years prior to initiating renal replacement therapy, background or greater diabetic retinopathy and/or > 3+ proteinuria on urinalysis in the absence of other causes of nephropathy. Nondiabetic control subjects were recruited from the same geographic region: 596 were African American and 39 European American without a prior diagnosis of T2D or renal disease, born in the same southeastern states, were recruited. DNA extraction was performed using the PureGene system (Gentra Systems, Minneapolis, MN). DNA was also obtained from 44 Yoruba Nigerians (YRI) from the National Institute of General Medical Sciences Human Variation Collection (Coriell Cell Repositories, Camden, NJ).

For admixture analyses, 39 unrelated European American controls were recruited using the same criteria as African American controls. DNA was also obtained from 44 YRI from the National Institute of General Medical Sciences Human Variation Collection (Coriell Cell Repositories, Camden, NJ).

### Genotyping for fine mapping

Eleven fine mapping microsatellites at an average spacing of 4 cM were selected from the Marshfield Center of Medical Genetics database <http://research.marshfield-clinic.org/genetics/> and the Genome Database <http://www.gdb.org/gdb/>. The polymerase chain reaction (PCR) was used to incorporate fluorescently labeled primers from Integrated DNA Technologies, Inc. (Coralville, IA) and Applied Biosystems (Foster City, CA) into PCR products, which were then electrophoresed on an ABI PRISM 3700 DNA Analyzer and results analyzed with Genescan Analysis Software (Applied Biosystems, Foster City, CA). Electropherograms were imported into GeneMapper (Applied Biosystems, Foster City, CA) for binning and allele calling. Familial relationships and Mendelian inconsistencies were checked using Prest [17] and PedCheck [18]. Inconsistent familial relationships were discarded from further analysis.

### Ordered subsets linkage analysis

A series of OSA [19] were computed to investigate the influence of a pedigree's mean age of diagnosis, BMI and duration of T2D to ESRD onset. The statistical significance of the change in the LOD score was evaluated by a permutation test under the null hypothesis that the ranking of the covariate is independent of the LOD score of the family on the target chromosome. Thus, the families were randomly permuted with respect to the covariate ranking, and an analysis proceeded as above for each permutation of these data. The resulting empirical distribution of the change in the LOD scores yielded a chromosome-wide  $P$  value ( $\Delta P$ ).

### Diabetes candidate gene selection and genotyping

Plausible functional candidate genes (*IL6*, *GCK1*, *IGFBP1* and *IGFBP3*) that lie under the intersection of the optimal linkage peaks for early age of T2D, lower BMI and longer duration to ESRD was selected (Figure 1). This regions roughly spans the distance between D7S513 and D7S1818. In the absence of a comprehensive African American SNP database of allele frequencies, we used the genotypic data of the YRI and CEPH Europeans (CEU) from the International HapMap project (HapMap #20, NCBI B35 assembly). Using the largest gene transcript plus 5 kb upstream and downstream of the gene, we selected YRI HapMap SNPs with a minor allele frequency (MAF)  $\leq 0.05$ , excluded SNPs with an Illumina designability score  $< 1.0$ , then used the aggressive tagging option of Tagger [20] implemented in Haploview 3.2 [21]. Next, we selected CEU HapMap SNPs with MAF  $\leq 0.05$ , forced the exclusion of Illumina-undesignable SNPs and inclusion of the already selected YRI tagSNPs, then used the aggressive tagging option of Tagger [20] to select any additional tags required to capture CEU variation. In addition, we added previously associated SNPs from the literature, yielding a total of 68 SNPs: glucokinase isoform 1 (*GCK1*, 24 SNPs); interleukin-6 (*IL6*, 17 SNPs); insulin growth factor binding protein 1 (*IGFBP1*, 16 SNPs) and insulin growth factor binding protein 3 (*IGFBP3*, 11 SNPs).

Genotyping was performed by Illumina Genotyping Services (Illumina Inc., San Diego, CA). Six SNPs were genotyped using iPLEX methodology on a MassARRAY system (Sequenom Inc., San Diego, CA) [22]: *IL6* rs1800797; *GCK1* rs741038, rs2971672 and *IGFBP1*: rs1496495, rs1908751, rs9658239. Primer sequences are available on request. The genotyping success rates were  $>96.5$  in cases and  $>96.1$  in controls. For quality control purposes each plate contained 2 duplicate samples and 4 inter-plate controls. Concordance between duplicate samples included in the genotyping was  $>98.9\%$ .

### Genotyping for admixture analyses

Seventy biallelic Admixture Informative Markers (AIMs), selected to maximize European and African allele frequency differences and sample all non-acrocentric arms of the autosomal genome, were genotyped by Illumina Inc.'s Custom Genotyping Service (Illumina Inc., San Diego, CA) in 577 African American cases, 596 African American controls, 44 YRI and 39 European Americans. Primer sequences are available on request. Genotyping success rates for AIMs were  $>97.4\%$  in African American cases and  $>95.6\%$  in African American controls.

### Statistical Analysis

Characteristics were compared and tested for significant differences using  $\chi^2$  tests for categorical variables, and one-way ANOVA test for continuous variables. Hardy-Weinberg equilibrium (HWE) was assessed using the  $\chi^2$  goodness-of-fit statistic at a significance level of  $\leq 0.01$ . Haplotype block structure was established using Haploview 3.2 [21], using the block definition from Gabriel *et al.* [23].


Tests of association under the three *a priori* genetic tests of association (dominant, additive and recessive) are reported. Due to a lack of validity of the large sample  $\chi^2$  test statistic, only the dominant model was considered for SNPs with ten or fewer individuals that were homozygous for the minor allele.

Quantitative trait (duration of T2D to ESRD onset, BMI, age at T2D and ESRD onset) association analysis for all SNPs was performed using a series of analysis of variance (ANOVA) tests implemented in QSNPGWA [24].

SNPs that showed nominal evidence for association with age of T2D onset were adjusted by BMI. Additionally, ESRD age, duration of T2D to ESRD onset and BMI were adjusted for age of T2D onset. All analyses were adjusted using linear regression on *a priori* genotypic models, and conducted using Stata 10 (College Station, TX).

Unadjusted odds ratios (ORs) and 95% confidence intervals (CI) in the case-control population were computed using SNPGWA to test all SNPs for genotypic association with T2D-ESRD [25]. SNPs that showed nominal evidence for association were then tested using a permutation test (1,000 permutations) as implemented in SNPGWA.

Individual ancestral proportions were calculated using an expectation-maximization (EM) algorithm (FRAPPE) [26] under a two-population model. Logistic regression tests of genetic models included adjustments for individual estimates of African ancestry, implemented in the program SNPADMIX [25].



**Figure 1 Chromosome 7p Ordered Subset Analysis.** Chromosome 7p fine mapping linkage results showing the Ordered Subset Analyses of early age at type 2 diabetes diagnosis (red), lower body mass index (green) and longer duration of type 2 diabetes mellitus to development of end stage renal disease (blue).

PS power and sample size program specifying a two-sided test at an alpha level of  $\leq 0.05$ , to detect an OR of 1.30 was utilized to approximate study power [27].

We report nominally significant associations at the  $P \leq 0.05$  level. To correct for multiple tests at the gene level the conservative Bonferonni method was used, with a  $P$ -value  $\leq 0.0007$  considered significant evidence for association when assuming independence based on LD.

## Results

### Fine mapping linkage analyses

OSA including the new fine mapping microsatellite data continued to support linkage at 28 cM on chromosome 7p, closest to D7S3051, in early-onset T2D families. Twenty-one percent of the pedigrees account for a maximum LOD = 3.61 ( $P = 0.002$ ) with a mean age at diagnosis  $29.5 \pm 3.0$  years vs. mean age at diagnosis of the unlinked families  $44.5 \pm 7.3$ . Although the max. LOD score dropped (from 3.85,  $P = 0.003$  in the genome scan) the LOD-1 interval increased from 11.5-38.2 cM to 11.5-49.5 cM. There was no significant difference ( $P = 0.56$ ) between mean BMI in the linked early-onset families ( $31.5 \text{ kg/m}^2 \pm 7.62$ ) versus the unlinked ( $31.8 \text{ kg/m}^2 \pm 7.51$ ) pedigrees. The evidence for linkage in subsets of T2D-ESRD families with lower BMI (64% of pedigrees; max. LOD = 2.20,  $P = 0.11$ ) and longer duration of diabetes before onset of ESRD (56% of pedigrees; max. LOD = 2.81,  $P = 0.20$ ) was no longer significant (Figure 1).

### Case-control population

Characteristics of the study population used for tests of association with positional candidate genes are shown in

Table 1. The mean age of examination for the non-diabetic controls is 7.5 years later than the mean age of onset for diabetes in the T2D-ESRD participants. The estimated mean proportion of African ancestry in the case subjects was  $81.7 \pm 13.3$  in T2D-ESRD cases and  $79.1 \pm 13.1$  in controls ( $P = 0.0008$ ).

### Hardy-Weinberg Equilibrium and linkage disequilibrium

All SNPs in candidate genes were examined for departures from HWE assumptions in cases and controls separately. Deviations from HWE ( $P \leq 0.01$ ) were observed for *IL6* rs1474347 in controls ( $P = 0.002$ ) and *IGFBP1* rs1496495 in cases ( $P = 0.001$ ). Although inconsistent with HWE proportions, these SNPs were retained for exploratory analyses, however there was no evidence for association of these SNPs with any of the traits investigated.

The 24 *GCK1* SNPs fell into five blocks of LD, with pairwise values of  $D'$  values ranging from 0.41-1.00, and  $r^2 = 0.003$ -0.76.  $D'$  values between the 17 *IL6* SNPs ranged from 0.34-1.00, with  $r^2$  values of 0.0002-0.82, resulting in one high LD block. The 16 *IGFBP1* SNPs fell within three blocks of high LD, with SNPs in the 3'UTR contained in block 2 ( $D' = 0.88$ -0.99;  $r^2 = 0.02$ -0.34) and the 11 genotyped *IGFBP3* SNPs fell within two blocks of LD (data not shown).

### Analyses with age at diagnosis of T2D and age at ESRD

Association analyses revealed two *GCK1* SNPs nominally associated with T2D age at diagnosis and five *GCK1* SNPs associated with age at ESRD onset (Table 2). The most significant result from the entire dataset was the association with intron 1 SNP rs730497 and age at ESRD onset ( $P = 0.0004$ ; T2D age-adjusted  $P = 0.0006$ ). The homozygote recessive genotype ("AA") for *IL6* SNP rs2069849, located in exon 5, was nominally associated with age at ESRD onset (T2D age-adjusted  $P = 0.013$ ). SNPs in *IGFBP1* and *IGFBP3* were not associated with T2D age at diagnosis or ESRD onset (Additional file 1, Table S1 and S2).

### Analyses with duration of T2D to ESRD onset

Association results for duration of T2D to ESRD onset are presented in Additional file 1, Table S3. *IL6* SNP rs1554606 was nominally associated (T2D age-adjusted  $P = 0.006$ ) with shorter duration of T2D to ESRD onset (Table 2;  $15.8 \pm 13.0$  years for "AA" genotype, compared with  $20.0 \pm 14.1$  years for "CC"/"CA"). Results are further summarized in Table 2.

### Analyses with BMI

Association results for BMI are shown in Table 2. *IGFBP1* and *IGFBP3* SNPs were not associated when adjusted for age of T2D diagnosis ( $P$  value = 0.067 and 0.120 respectively). Single SNP association results for

**Table 1 Characteristics of African American subjects**

Trait	T2D-ESRD Cases		Controls	
	n*		n*	
% Female (n)	577	61% (352)	596	51% (306)
Age at exam (years ± SD)	541	62.2 ± 10.3‡	448	49.3 ± 9.8
Age at T2D diagnosis (years ± SD)	544	41.8 ± 11.6	-	-
Age at ESRD diagnosis (years ± SD)	560	59.0 ± 10.5	-	-
BMI (kg/m <sup>2</sup> )	559	29.6 ± 7.0†	450	29.8 ± 7.1
T2D duration to ESRD onset (years ± SD)	511	20.3 ± 10.5	-	-
Mean African Ancestry	577	81.7 ± 13.3‡	596	79.1 ± 13.1

\*n = number with data available. ‡ Mean differs significantly between cases and controls P-value < 0.001. †Reflects dry weight on dialysis. All cases were diagnosed with T2D and ESRD.

BMI are presented thoroughly in Additional file 1, Table S4.

#### Analyses with T2D-ESRD in the African American case-control population

Single SNP genotypic association results ( $P \leq 0.05$ ) for T2D-ESRD are shown in Additional file 1, Table S5. Nominal evidence of association was observed with *IGFBP3*: two promoter SNPs: rs903889 (dominant  $P = 0.014$ ; admixture-adjusted  $P_a = 0.029$ ), rs924140 (additive  $P = 0.019$ ;  $P_a = 0.033$ ) and two SNPs located in intron 3; rs10255707 (dominant  $P = 0.005$ ;  $P_a = 0.01$ ) and rs3110697 (recessive  $P = 0.002$ ;  $P_a = 0.003$ ). Nominal evidence of association was also detected with *IGFBP1* rs9658233 located in the 3' UTR region. *GCK1* and *IL6* SNPs showed no evidence of association with T2D-ESRD (Additional file 1, Table S6). Genotype frequencies and counts for each SNP are shown in Additional file 1, Table S7.

Tagger [20] results indicated that 2- or 3-SNP haplotypes were not required to capture ungenotyped YRI and CEU HapMap SNPs in these genes.

#### Discussion

Evidence of linkage on chromosome 7p in families with age of T2D diagnosis, ESRD onset, lower BMI and longer duration to ESRD was detected in an earlier study of African American families [1,2]. These results generated our interest in genotyping additional polymorphic markers to refine the linkage peak and explore positional candidate genes. Fine mapping analyses continued to support evidence for linkage at 7p in a subset of T2D families with an early age of diagnosis. Although this linkage peak does not appear to have been reported in other populations, two recent reports detected modest evidence of linkage to T2D in the same region in African American families from the GENNID Cohort [28] and an African American population from South Carolina [29].

The present study demonstrates that *GCK1* rs730497 recessive "AA" genotype is strongly associated with early age ESRD onset (Table 2). *GCK1*, a strong functional candidate for T2D, appears to play a role in susceptibility to early age of ESRD in African Americans. We observed an approximately 6 year shorter duration of T2D to ESRD and early age of T2D onset with recessive "AA" homozygotes compared with "GG" homozygotes and heterozygotes (Additional file 1, Tables S1 and S3). Earlier reports suggest that nephropathy is rarely observed in persons with *GCK1* mutations [30]. MODY is characterized as non-ketotic T2D that is inherited by an autosomal dominant manner [31-33], with early onset (usually before the age of 25). MODY is often asymptomatic mild hyperglycemia, which can go undetected for prolonged periods of time which can lead to diabetic complication later in adulthood. Hence, it is plausible that in the presence of hyperglycemia and increased BMI (mean BMI 30.8 kg/m<sup>2</sup> ± 7.1; Additional file 1, Table S4) along with the rs730497 recessive genotype results in ESRD progression. In addition, several reports have shown that MODY genes segregate in late onset T2D cases and families [34-36]. In an effort to reduce the likelihood of including potential MODY2 patients, only individuals with T2D onset over the age of 35 were ascertained as probands.

Recently, Paré *et al.* [37] reported that variant rs730497 is associated with an increase of 0.03% HbA<sub>1c</sub> levels ( $P = 2.8 \times 10^{-12}$ ) in non-diabetic European Americans from the Women's Genome Health Study (WGHS) and participants from the Boston metropolitan area. The MAF of the rs730497 A "risk" allele was similar in African American cases (0.20) and previously reported European Americans from WGHS (0.17) [37]. Several studies have shown a strong correlation between increased levels of HbA<sub>1c</sub> and nephropathy [38]. The relationship between rs730497 minor "A" allele and ESRD progression is unknown, however, marker rs730497 is located 4.8 kb from exon 1 and 30.7 kb from exon 2. The slightly more common G allele

**Table 2 Nominally significant (P < 0.05) single-SNP genotypic tests of association with T2D age at diagnosis, ESRD age at onset, BMI on dialysis, and duration of T2D to ESRD onset.**

Phenotype	Gene	Marker	Major/Minor alleles	MAF Case	1/1 Mean ± SD (n)	1/2 Mean ± SD (n)	2/2 Mean ± SD (n)	Dominant P-value	Additive P-value	Recessive P-value
ESRD age	<i>IL6</i>	Rs2069849	G/A	0.15	59.0 ± 10.5 (419)	59.8 ± 10.4 (140)	53.7 ± 10.4 (16)	0.896	0.581	<b>0.040</b> (0.013) <sub>a</sub>
Duration of T2D to ESRD	<i>IL6</i>	rs1554606	C/A	0.34	20.2 ± 13.7 (255)	19.7 ± 14.7 (245)	15.8 ± 13.0 (68)	0.247	0.052	<b>0.007</b> (0.006) <sub>a</sub>
T2D age	<i>GCK1</i>	rs2908296	C/A	0.35	41.6 ± 11.0 (235)	42.8 ± 12.2 (277)	38.0 ± 10.3 (63)	0.733	0.313	<b>0.007</b> (0.014) <sub>b</sub>
T2D age	<i>GCK1</i>	rs12673242	A/G	0.37	41.3 ± 11.4 (236)	43.2 ± 11.9 (258)	39.0 ± 11.2 (81)	0.38	0.606	<b>0.023</b> (0.032) <sub>b</sub>
ESRD age	<i>GCK1</i>	rs2908296	C/A	0.35	59.2 ± 10.0 (235)	59.4 ± 11.2 (277)	56.1 ± 8.7 (63)	0.632	0.157	<b>0.027</b> (0.133) <sub>a</sub>
ESRD age	<i>GCK1</i>	rs2971676	G/A	0.24	59.4 ± 10.4 (328)	59.0 ± 10.8 (217)	54.4 ± 8.8 (30)	0.328	0.092	<b>0.02</b> (0.035) <sup>a</sup>
ESRD age	<i>GCK1</i>	rs2971675	C/A	0.23	59.5 ± 10.5 (338)	58.7 ± 10.8 (210)	54.5 ± 8.4 (27)	0.2	0.065	<b>0.033</b> (0.029) <sub>a</sub>
ESRD age	<i>GCK1</i>	rs730497	G/A	0.2	58.9 ± 10.4 (367)	60.0 ± 10.6 (180)	51.5 ± 8.6 (25)	0.989	0.206	<b>0.0004</b> (0.0006) <sup>a</sup>
ESRD age	<i>GCK1</i>	rs2908289	G/A	0.27	58.5 ± 10.5 (315)	60.6 ± 10.7 (211)	55.1 ± 8.8 (49)	0.223	0.827	<b>0.008</b> (0.129) <sub>a</sub>
BMI	<i>IGFBP1</i>	rs1908751	C/T	0.33	29.8 ± 7.3 (254)	29.2 ± 6.7 (243)	31.6 ± 7.1 (60)	0.838	0.436	<b>0.043</b> (0.067) <sub>a</sub>
BMI	<i>IGFBP3</i>	rs10255707	G/A	0.04	29.9 ± 7.1 (511)	27.3 ± 5.4 (33)	23.5 ± 7.4 (2)	<b>0.022</b> (0.120) <sup>a</sup>	-	-
Duration of T2D to ESRD	<i>IGFBP3</i>	rs3110697	G/A	0.35	21.3 ± 15.7 (234)	18.5 ± 12.7 (270)	19.8 ± 14.2 (63)	<b>0.027</b> (0.067) <sup>a</sup>	0.055	0.586

Test models refer to the minor allele. \*Only the dominant model was considered where the minor allele homozygote count was <10. Bold: P-values < 0.05. Unadjusted P values without parentheses and P values with parentheses adjusted for <sup>a</sup>T2D age- and <sup>b</sup>BMI.

appears to be ancestral since it is present in the chimpanzee sequence.

Nominal associations with MODY gene *GCK1* was also observed with T2D age at diagnosis (Table 2). Four *GCK1* SNPs were found to be associated with T2D risk (recessive model: OR ranging from 1.36-1.87) in a case-control population from Finland [39]. Bonnycastle *et al.* [39] observed an 87% increased T2D risk with marker rs882020. Marker rs2908296 is approximately 14.7 kb from rs882020; it is plausible that these variants may tag or be in LD with the causal variant(s). The reported associated SNPs (rs2284769, rs12534623, rs2268573 and rs882020) [39] and the dinucleotide repeat (D7S531) in Mauritian Creoles [8], African Americans [9], and South Indians [10], were not directly investigated in our African American population due to our HapMap-coverage based SNP selection process, however the 24 *GCK1* SNPs genotyped captured all 32 HapMap *GCK1* SNPs with MAF > 5% with r<sup>2</sup> values > 0.80.

*IL6* SNP rs2069849, located in exon 5, was nominally associated with earlier age of ESRD onset. This variant causes a synonymous coding change for phenylalanine at position 201 (F201), which may alter mRNA stability and translation. Promoter polymorphisms rs1800795 and rs1800796 have shown association with T2D and

diabetic nephropathy respectively [40-42]. We investigated these two SNPs in our African American population (Additional file 1, Table S6) and, consistent with a study conducted in a larger population of T2D cases and non-T2D controls [43], did not observe association with T2D-ESRD. The MAF for promoter SNPs rs1800797, rs1800796 and rs1800795 in African American (controls: 0.07 0.09; 0.07, respectively; cases 0.05; 0.11; 0.06, respectively) are comparable with the YRI frequencies (MAF = 0.0; 0.09; 0.0), but contrast with those of the European American controls (MAF = 0.48; 0.04; 0.53). The lack of association may be due to the reduced prevalence of the risk alleles in our African American population.

Modest associations with T2D-ESRD were also detected with one *IGFBP1* (rs9658233) and four *IGFBP3* SNPs (rs10255707, rs3110697, rs924140 and rs903889). The major "T" allele for SNP rs9658233, located in the downstream region of *IGFBP1*, appears to confer risk with T2D-ESRD, with OR between 1.25-1.35 (Additional file 1, Table S5). In contrast, having at least one copy of the minor allele for *IGFBP3* variants rs3110697 and rs924140 appear to protect against T2D-ESRD, with OR between 0.60-0.82 (Additional file 1, Table S5). The frequencies of rs3110697 and rs924140 located in intron 3

and the promoter region of *IGFBP3* and rs9658233 of *IGFBP1* more closely resemble the frequencies of the HapMap YRI samples, suggesting African influence for these markers. Several genome wide association studies performed in European populations have replicated associations with SNPs in *IGF2BP2* [44-47], implicating a role for the IGF pathway and T2D susceptibility.

We also note in the results that 2- and 3-SNP moving windows analyses were not required to capture ungenotyped YRI and CEU HapMap SNPs in *IL6*, *GCK1*, *IGFBP1* and *IGFBP3*.

One limitation of this study is that all controls did not have measures of diabetes status or renal impairment. Serum glucose values were obtained for 42.9% (256 individuals) of the African American controls (mean 93.6 mg/dl). Five of these individuals had glucose levels greater than 126 mg/dl. Four of these measures were non-fasting and the fifth had unknown dietary status, suggesting the overall misclassification rate of controls is likely to be below 2%. While this has not impacted our ability to detect an association, this may have reduced our power to detect more subtle influences of additional variants, and underestimate effect sizes.

While adjustments for admixture reduced the strength for the observed associations with T2D-ESRD (Additional file 1, Table S5), associations with *IGFBP1* rs9658233 and *IGFBP3* (rs10255707, rs3110697, rs924140 and rs903889) remained nominally significant. Multiple comparisons were corrected by conservative Bonferroni method ( $P \leq 0.00005$ ) however, when assuming independence based on LD, one would consider a  $P$ -value  $\leq 0.0007$  as evidence for association, only SNP rs730497 remained significant (T2D age-adjusted  $P = 0.0006$ ).

We estimate 65.7% power to detect an OR of 1.30 at a significance level of 0.05 in the case-control cohort [27].

We have confirmed linkage to early-onset T2D on chromosome 7p, and examined common variants across positional candidate genes *GCK1*, *IL6*, *IGFBP1* and *IGFBP3*. A variant of the *GCK1* gene, rs730497, appears to play a role in susceptibility to earlier age of ESRD onset in African Americans. This may not represent the "true" casual variant but may be in LD with a functional SNP. Denser SNP genotyping and direct sequencing along with functional studies investigating intron 1 of the *GCK1* gene are imperative to investigate the etiology of ESRD risk. It is plausible that multiple rare variants in this region of the gene may contribute to this phenotype. In addition, other candidates under the LOD-1 interval, includes the growth factor receptor-bound protein 10 (*GRB10*) [48,49] and IGF2 mRNA binding protein 3 (*IGF2BP3*), warrant investigation.

## Conclusion

Our findings support that intron 1 of the *GCK1* play a role with ESRD risk. Further studies in other populations and functional studies may be warranted.

**Additional file 1: Supplementary tables.** Supplemental tables S1-S6. Click here for file [http://www.biomedcentral.com/content/supplementary/1471-2350-11-22-S1.DOC]

## Acknowledgements

We thank the patients and individuals recruited as controls for their participation. We also thank the staff of the Southeastern Kidney Council/ESRD Network 6, Pamela Hicks, Joyce Byers, Mitzie Spainhour, Matt Stiegert, Christopher Vallis, Candace Gordon, Shelly Smith, and Mark Hansen and colleagues at Illumina Inc. This work was supported by grants R01 DK066358, and the Wake Forest University General Clinical Research Center M01 RR07122; MMS is supported by a Career Development Award from the American Diabetes Association. Programs developed at Wake Forest University Health Sciences and used in this manuscript are freely available at [http://www.phs.wfubmc.edu/web/public\\_bios/sec\\_gene/downloads.cfm](http://www.phs.wfubmc.edu/web/public_bios/sec_gene/downloads.cfm).

## Author details

<sup>1</sup>Center for Human Genomics, Wake Forest University School of Medicine, Winston-Salem, NC, USA. <sup>2</sup>Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, USA. <sup>3</sup>Department of Biostatistical Sciences, Wake Forest University School of Medicine, Winston-Salem, NC, USA. <sup>4</sup>Center for Public Health Genomics, University of Virginia, Charlottesville, VA, USA. <sup>5</sup>Department of Public Health Sciences, University of Virginia, Charlottesville, VA, USA. <sup>6</sup>Department of Biochemistry, Wake Forest University School of Medicine, Winston-Salem, NC, USA. <sup>7</sup>Milton S Hershey Medical Center, Pennsylvania State University, Hershey, PA, USA. <sup>8</sup>Department of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, VA, USA. <sup>9</sup>Department of Internal Medicine, Wake Forest University School of Medicine, Winston-Salem, NC, USA. <sup>10</sup>Department of Medicine, University of Virginia, Charlottesville, VA, USA.

## Authors' contributions

TSL carried out the genotyping of additional polymorphic markers, prepared linkage data for analysis, performed association analysis, and drafted the manuscript. CDL developed software programs SNP-GWA and QSNP-GWA, and directed LL in conducting linkage analyses. KLK performed the admixture analysis. JCM assisted with statistical analysis. CJG assisted in preparing linkage data for analysis. SSR, BIF and DWB participated in the design of the study and participant recruitment. MMS conceived of the study, participated in its design and coordination and helped draft the manuscript. All authors read and approved the final manuscript.

## Competing interests

The authors declare that they have no competing interests.

Received: 30 July 2009

Accepted: 8 February 2010 Published: 8 February 2010

## References

1. Sale MM, Freedman BI, Langefeld CD, Williams AH, Hicks PJ, Colicigno CJ, Beck SR, Brown WM, Rich SS, Bowden DW: A genome-wide scan for type 2 diabetes in African-American families reveals evidence for a locus on chromosome 6q. *Diabetes* 2004, **53**(3):830-837.
2. Bowden DW, Colicigno CJ, Langefeld CD, Sale MM, Williams A, Anderson PJ, Rich SS, Freedman BI: A genome scan for diabetic nephropathy in African Americans. *Kidney Int* 2004, **66**(4):1517-1526.
3. Froguel P, Vaxillaire M, Sun F, Velho G, Zouali H, Butel MO, Lesage S, Vionnet N, Clement K, Fougereousse F, et al: Close linkage of glucokinase locus on chromosome 7p to early-onset non-insulin-dependent diabetes mellitus. *Nature* 1992, **356**(6365):162-164.



4. Froguel P, Zouali H, Vionnet N, Velho G, Vaxillaire M, Sun F, Lesage S, Stoffel M, Takeda J, Passa P, et al: **Familial hyperglycemia due to mutations in glucokinase. Definition of a subtype of diabetes mellitus.** *N Engl J Med* 1993, **328**(10):697-702.
5. Hattersley AT, Turner RC: **Mutations of the glucokinase gene and type 2 diabetes.** *Q J Med* 1993, **86**(4):227-232.
6. Laakso M, Malkki M, Kekalainen P, Kuusisto J, Mykkanen L, Deeb SS: **Glucokinase gene variants in subjects with late-onset NIDDM and impaired glucose tolerance.** *Diabetes Care* 1995, **18**(3):398-400.
7. Takekawa K, Ikegami H, Fukuda M, Ueda H, Kawaguchi Y, Fujioka Y, Fujisawa T, Ogiwara T: **Early-onset type 2 (non-insulin-dependent) diabetes mellitus is associated with glucokinase locus, but not with adenosine deaminase locus, in the Japanese population.** *Diabetes Res Clin Pract* 1994, **23**(3):141-146.
8. Chiu KC, Province MA, Dowse GK, Zimmet PZ, Wagner G, Serjeantson S, Permutt MA: **A genetic marker at the glucokinase gene locus for type 2 (non-insulin-dependent) diabetes mellitus in Mauritian Creoles.** *Diabetologia* 1992, **35**(7):632-638.
9. Chiu KC, Province MA, Permutt MA: **Glucokinase gene is genetic marker for NIDDM in American blacks.** *Diabetes* 1992, **41**(7):843-849.
10. McCarthy ML, Hitchins M, Hitman GA, Cassell P, Hawrami K, Morton N, Mohan V, Ramachandran A, Snehalatha C, Viswanathan M: **Positive association in the absence of linkage suggests a minor role for the glucokinase gene in the pathogenesis of type 2 (non-insulin-dependent) diabetes mellitus amongst south Indians.** *Diabetologia* 1993, **36**(7):633-641.
11. Fernandez-Real JM, Broch M, Vendrell J, Gutierrez C, Casamitjana R, Pugeat M, Richart C, Ricart W: **Interleukin-6 gene polymorphism and insulin sensitivity.** *Diabetes* 2000, **49**(3):517-520.
12. Vozarova B, Fernandez-Real JM, Knowler WC, Gallart L, Hanson RL, Gruber JD, Ricart W, Vendrell J, Richart C, Tataranni PA, et al: **The interleukin-6 (-174) G/C promoter polymorphism is associated with type-2 diabetes mellitus in Native Americans and Caucasians.** *Hum Genet* 2003, **112**(4):409-413.
13. Bennermo M, Held C, Stemme S, Ericsson CG, Silveira A, Green F, Tornvall P: **Genetic predisposition of the interleukin-6 response to inflammation: implications for a variety of major diseases?** *Clin Chem* 2004, **50**(11):2136-2140.
14. Garay-Sevilla ME, Nava LE, Malacara JM, Wrobel K, Perez U: **Advanced glycosylation end products (AGEs), insulin-like growth factor-1 (IGF-1) and IGF-binding protein-3 (IGFBP-3) in patients with type 2 diabetes mellitus.** *Diabetes Metab Res Rev* 2000, **16**(2):106-113.
15. Kim HS, Ali O, Shim M, Lee KW, Vuguin P, Muzumdar R, Barzilai N, Cohen P: **Insulin-like growth factor binding protein-3 induces insulin resistance in adipocytes in vitro and in rats in vivo.** *Pediatr Res* 2007, **61**(2):159-164.
16. Sale MM, Smith SG, Mychaleckyj JC, Keene KL, Langefeld CD, Leak TS, Hicks PJ, Bowden DW, Rich SS, Freedman BI: **Variants of the transcription factor 7-like 2 (TCF7L2) gene are associated with type 2 diabetes in an African-American population enriched for nephropathy.** *Diabetes* 2007, **56**(10):2638-2642.
17. McPeck MS, Sun L: **Statistical tests for detection of misspecified relationships by use of genome-screen data.** *Am J Hum Genet* 2000, **66**(3):1076-1094.
18. O'Connell JR, Weeks DE: **PedCheck: a program for identification of genotype incompatibilities in linkage analysis.** *Am J Hum Genet* 1998, **63**(1):259-266.
19. Hauser ER, Watanabe RM, Duren WL, Bass MP, Langefeld CD, Boehnke M: **Ordered subset analysis in genetic linkage mapping of complex traits.** *Genet Epidemiol* 2004, **27**(1):53-63.
20. de Bakker PI, Yelensky R, Pe'er I, Gabriel SB, Daly MJ, Altshuler D: **Efficiency and power in genetic association studies.** *Nat Genet* 2005, **37**(11):1217-1223.
21. Barrett JC, Fry B, Maller J, Daly MJ: **Haploview: analysis and visualization of LD and haplotype maps.** *Bioinformatics* 2005, **21**(2):263-265.
22. Buetow KH, Edmonson M, MacDonald R, Clifford R, Yip P, Kelley J, Little DP, Strausberg R, Koester H, Cantor CR, et al: **High-throughput development and characterization of a genomewide collection of gene-based single nucleotide polymorphism markers by chip-based matrix-assisted laser desorption/ionization time-of-flight mass spectrometry.** *Proc Natl Acad Sci USA* 2001, **98**(2):581-584.
23. Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J, DeFelice M, Lochner A, Faggart M, et al: **The structure of haplotype blocks in the human genome.** *Science* 2002, **296**(5576):2225-2229.
24. Lewis JP, Palmer ND, Hicks PJ, Sale MM, Langefeld CD, Freedman BI, Divers J, Bowden DW: **Association analysis in African Americans of European-derived type 2 diabetes single nucleotide polymorphisms from whole-genome association studies.** *Diabetes* 2008, **57**(8):2220-2225.
25. Leak TS, Perlegas PS, Smith SG, Keene KL, Hicks PJ, Langefeld CD, Mychaleckyj JC, Rich SS, Kirk JK, Freedman BI, et al: **Variants in intron 13 of the ELMO1 gene are associated with diabetic nephropathy in African Americans.** *Ann Hum Genet* 2009, **73**(2):152-159.
26. Tang H, Peng J, Wang P, Risch NJ: **Estimation of individual admixture: analytical and study design considerations.** *Genet Epidemiol* 2005, **28**(4):289-301.
27. Dupont WD, Plummer WD Jr: **Power and sample size calculations. A review and computer program.** *Control Clin Trials* 1990, **11**(2):116-128.
28. Elbein SC, Das SK, Hallman DM, Hanis CL, Hasstedt SJ: **Genome-wide linkage and admixture mapping of type 2 diabetes in African American families from the American Diabetes Association GENNID (Genetics of NIDDM) Study Cohort.** *Diabetes* 2009, **58**(1):268-274.
29. Sale MM, Lu L, Spruill JJ, Fernandes JK, Lok KH, Divers J, Langefeld CD, Garvey WT: **Genome-wide linkage scan in Gullah-speaking African American families with type 2 diabetes: the Sea Islands Genetic African American Registry (Project SuGAR).** *Diabetes* 2009, **58**(1):260-267.
30. Olek K: **Maturity-onset diabetes of the young: an update.** *Clin Lab* 2006, **52**(11-12):593-598.
31. Froguel P, Velho G: **Molecular Genetics of Maturity-onset Diabetes of the Young.** *Trends Endocrinol Metab* 1999, **10**(4):142-146.
32. Velho G, Froguel P: **Genetic, metabolic and clinical characteristics of maturity onset diabetes of the young.** *Eur J Endocrinol* 1998, **138**(3):233-239.
33. Winter WE: **Molecular and biochemical analysis of the MODY syndromes.** *Pediatr Diabetes* 2000, **1**(2):88-117.
34. Damcott CM, Hoppman N, Ott SH, Reinhart LJ, Wang J, Pollin TI, O'Connell JR, Mitchell BD, Shuldiner AR: **Polymorphisms in both promoters of hepatocyte nuclear factor 4-alpha are associated with type 2 diabetes in the Amish.** *Diabetes* 2004, **53**(12):3337-41.
35. März W, Nauck M, Hoffmann MM, Nagel D, Boehm BO, Koenig W, Rothenbacher D, Winkelmann BR: **G(-30)A polymorphism in the pancreatic promoter of the glucokinase gene associated with angiographic coronary artery disease and type 2 diabetes mellitus.** *Circulation* 2004, **109**(23):2844-9.
36. Karim MA, Wang X, Zhang Z, Bowden DW, Freedman BI, Elbein SC: **Association of an insulin gene promoter insertion polymorphism with type 2 diabetes and diabetic nephropathy in African Americans.** *Mol Genet Metab* 2004, **83**(4):344-6.
37. Paré G, Chasman DI, Parker AN, Nathan DM, Miletich JP, Zee RY, Ridker PM: **Novel association of HK1 with glycated hemoglobin in a non-diabetic population: a genome-wide evaluation of 14,618 participants in the Women's Genome Health Study.** *PLoS Genet* 2008, **4**(12):e1000312.
38. Roohk HV, Zaidi AR: **A review of glycated albumin as an intermediate glycation index for controlling diabetes.** *J Diabetes Sci Technol* 2008, **2**(6):1114-1121.
39. Bonycastle LL, Willer CJ, Conneely KN, Jackson AU, Burrill CP, Watanabe RM, Chines PS, Narisu N, Scott LJ, Enloe ST, et al: **Common variants in maturity-onset diabetes of the young genes contribute to risk of type 2 diabetes in Finns.** *Diabetes* 2006, **55**(9):2534-2540.
40. Buraczynska M, Jozwiak L, Ksiazek P, Borowicz E, Mierzicki P: **Interleukin-6 gene polymorphism and faster progression to end-stage renal failure in chronic glomerulonephritis.** *Transl Res* 2007, **150**(2):101-105.
41. Kitamura A, Hasegawa G, Obayashi H, Kamiuchi K, Ishii M, Yano M, Tanaka T, Yamaguchi M, Shigeta H, Ogata M, et al: **Interleukin-6 polymorphism (-634C/G) in the promoter region and the progression of diabetic nephropathy in type 2 diabetes.** *Diabet Med* 2002, **19**(12):1000-1005.
42. Mittal RD, Manchanda PK: **Association of interleukin (IL)-4 intron-3 and IL-6 -174 G/C gene polymorphism with susceptibility to end-stage renal disease.** *Immunogenetics* 2007, **59**(2):159-165.
43. Qi L, van Dam RM, Meigs JB, Manson JE, Hunter D, Hu FB: **Genetic variation in IL6 gene and type 2 diabetes: tagging-SNP haplotype**



analysis in large-scale case-control study and meta-analysis. *Hum Mol Genet* 2006, **15**(11):1914-1920.

44. Sladek R, Rocheleau G, Rung J, Dina C, Shen L, Serre D, Boutin P, Vincent D, Belisle A, Hadjadj S, *et al*: **A genome-wide association study identifies novel risk loci for type 2 diabetes.** *Nature* 2007, **445**(7130):881-885.
45. **Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls.** *Nature* 2007, **447**(7145):661-678.
46. Cauchi S, Meyre D, Durand E, Proenca C, Marre M, Hadjadj S, Choquet H, De Graeve F, Gaget S, Allegaert F, *et al*: **Post genome-wide association studies of novel genes associated with type 2 diabetes show gene-gene interaction and high predictive value.** *PLoS ONE* 2008, **3**(5):e2031.
47. Zeggini E, Weedon MN, Lindgren CM, Frayling TM, Elliott KS, Lango H, Timpson NJ, Perry JR, Rayner NW, Freathy RM, *et al*: **Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes.** *Science* 2007, **316**(5829):1336-1341.
48. Di Paola R, Ciociola E, Boonyasrisawat W, Nolan D, Duffy J, Miscio G, Cisternino C, Fini G, Tassi V, Doria A, *et al*: **Association of hGrb10 genetic variations with type 2 diabetes in Caucasian subjects.** *Diabetes Care* 2006, **29**(5):1181-1183.
49. Rampersaud E, Damcott CM, Fu M, Shen H, McArdle P, Shi X, Shelton J, Yin J, Chang YP, Ott SH, *et al*: **Identification of novel candidate genes for type 2 diabetes from a genome-wide association scan in the Old Order Amish: evidence for replication from diabetes-related quantitative traits and from independent populations.** *Diabetes* 2007, **56**(12):3053-3062.

#### Pre-publication history

The pre-publication history for this paper can be accessed here:<http://www.biomedcentral.com/1471-2350/11/22/prepub>

doi:10.1186/1471-2350-11-22

**Cite this article as:** Leak *et al*: Chromosome 7p linkage and association study for diabetes related traits and type 2 diabetes in an African-American population enriched for nephropathy. *BMC Medical Genetics* 2010 **11**:22.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

