

Published in final edited form as:

Metabolism. 2009 June ; 58(6): 854–859. doi:10.1016/j.metabol.2009.02.012.

Adherence to Mediterranean Diet is favorably associated with metabolic parameters in HIV positive patients with the HAART-induced metabolic syndrome and lipodystrophy

S Tsiodras¹, KA Poulia³, M Yannakoulia³, SN Chimienti¹, S Wadhwa¹, AW Karchmer¹, and CS Mantzoros²

¹ Division of Infectious Diseases, Department of Internal Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, US

² Division of Endocrinology, Department of Internal Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, US

³ Department of Nutrition and Dietetics, Harokopio University, Athens, Greece

Abstract

Objective—To investigate whether closer adherence to a Mediterranean dietary pattern is associated with metabolic aspects of the Highly Active Antiretroviral Therapy (HAART)-induced metabolic syndrome (fat redistribution, insulin resistance, dyslipidemia) in HIV positive patients.

Design—Cross sectional study.

Methods—227 HIV- infected patients were evaluated during a single outpatient visit to the General Clinical Research Center of Beth Israel Deaconess Medical Center. Usual dietary intake and physical activity habits were evaluated; the Mediterranean Diet Score (MedDietScore) was calculated. Dual-energy x-ray absorptiometry, CT findings anthropometrics and data from the study interviews and questionnaires were used for the assessment of body composition using specific criteria. A complete metabolic profile was available for all subjects.

Results—In the entire study sample, a weak inverse association was found between insulin resistance, estimated using the homeostasis model (HOMA-IR), and MedDietScore (standardized $\beta = -0.15$, $p = 0.03$). Interaction models revealed that this was largely driven by an inverse association in patients with fat redistribution (FR) (standardized $\beta = -0.13$, $p = 0.02$). Moreover, MedDietScore was positively correlated with HDL-cholesterol (standardized $\beta = 0.15$, $p = 0.01$) and marginally negatively associated with circulating triglyceride levels (standardized $\beta = -0.16$, $p = 0.13$) in this group of patients.

Corresponding author: Christos S. Mantzoros, M.D., Division of Endocrinology, ST 816, Beth Israel Deaconess Medical Center, Harvard Medical School, 330 Brookline Ave., Boston, Massachusetts 02215, U.S.A., Fax: (617) 667-8634, cmantzor@bidmc.harvard.edu. The first three authors contributed equally to the work presented herein.

Conflicts of interest: None

Authors' contributions: Dr. Mantzoros designed this study. Drs. Tsiodras, Chimienti and Wadhwa collected the data. Ms. Poulia and Dr. Yannakoulia analyzed the dietary information, performed the statistical analysis and drafted the paper. Drs. Mantzoros and Tsiodras contributed to subsequent revisions of the paper and to the responses to the reviewers' comments. Drs. Karchmer and Mantzoros secured funding for this study. All authors critically revised the manuscript and approved the final version submitted.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conclusions—Adherence to a Mediterranean dietary pattern was favorably related to cardiovascular risk factors in HIV-positive patients with FR. Further clinical studies are needed, to confirm our data in different populations and to explore the underlying mechanisms.

Keywords

Dietary pattern; Mediterranean diet; fat redistribution; HAART-induced metabolic syndrome

Introduction

Highly Active Antiretroviral Therapy (HAART)-induced lipodystrophy and metabolic syndrome, characterized by fat redistribution (FR), glucose intolerance, insulin resistance (IR) and dyslipidemia are well recognized among HIV-infected patients on HAART. Both protease inhibitor (PIs) and non-PIs containing regimens i.e. nucleoside or non-nucleoside reverse transcriptase inhibitors (NRTI's and NNRTI's) have each been implicated in the pathogenesis of these metabolic disturbances, and the exact underlying mechanisms are still under investigation. Intakes of macronutrients and specific food groups have been studied in relation to their effect on the development of metabolic abnormalities in this syndrome [1]. Specifically, increased consumption of saturated fat has been associated with hypertriglyceridemia among HIV-infected patients with metabolic abnormalities [2]. In a small study of HIV patients with FR, intakes of dietary protein, animal protein and trans fatty acids were positively, whereas intake of soluble fiber was negatively correlated with dyslipidemia [3]. Moreover, dietary vitamin E intake has been found to be negatively associated with diastolic blood pressure (DBP), body fat and, possibly, IR [4].

Although it is widely recognized that foods and nutrients are not eaten in isolation, but in the context of whole diets, no prior study has to date examined potential associations between aspects of the HAART-induced metabolic syndrome and dietary patterns. Dietary pattern analysis, with the use of diet scores, has been used as an alternative, holistic approach to examine the relationship between diet and disease prevention or treatment [5-7]. Adherence to a Mediterranean dietary pattern, in specific, characterized by abundant intake of plant foods, such as whole grain cereals, fruits, vegetables, legumes and olive oil, moderate intakes of fish and dairy products; and low intakes of red meat, saturated fats, and sweets, has been associated with decreased all-cause mortality, better health status and improvement of cardiovascular risk factors [8-15]. The aim of the present study was to investigate whether closer adherence to a Mediterranean dietary pattern might be related to metabolic aspects of the HAART-induced metabolic syndrome in HIV-positive patients.

Methods

Study cohort

We evaluated 227 consecutively enrolled HIV-infected subjects during a single outpatient visit to the General Clinical Research Centre of Beth Israel Medical Center. The sample of this study constitutes approximately 10% of the entire population admitted at two ambulatory care clinics of an urban major academic medical center (approx 2,000 patients) and is representative of the clinics population. Inclusion criteria were age ≥ 16 years, documented HIV infection and ≥ 6 months of cumulative exposure to any antiretroviral regimen. The Institutional Review Board at Beth Israel Deaconess Medical Center approved the study and all subjects gave written informed consent before participation.

Dietary assessment

A validated self-administered food frequency questionnaire (FFQ) (Block 98 revision of Block/NCI Health Habits and History Questionnaire; Block Dietary Data System, Berkeley, CA) [16,17] was used for the assessment of the usual dietary intake. Estimates of energy, macro- and micro-nutrient, as well as food group consumption for the year preceding the study were obtained from the analysis of 217 completed questionnaires. Intake of selected food items has been used for the calculation of the Mediterranean Diet Score (MedDietScore), based on the rationale of the Mediterranean dietary pyramid [14]: the score ranges from 0 to 55; higher scores indicate closer adherence to this dietary pattern.

Exercise assessment

Current exercise was evaluated using 3 multiple-choice questions regarding the type and intensity of exercise (i.e., type of exercise: 1, walking on level ground/swimming; 2, running, aerobic classes, or use of cardiovascular machines, treadmill, or stationary bike; and 3, weight training; intensity: 1, slight; 2, moderate; and 3, heavy), exercise frequency (from 0 to 7 sessions/week), and session duration (<15, 15–29, 30–59, 60–89, or >90 min) [4,18]. Cumulative indexes for either aerobic or total (aerobic and/or resistance) exercise were calculated as number of sessions per week \times duration (min) per session \times exercise intensity. All subjects completed the exercise questionnaire.

Body composition, clinical and biochemical assessment

Body composition was evaluated in all subjects using dual-energy x-ray absorptiometry whole-body scanner (Hologic QDR-2000 version 5.73A; Hologic). Anthropometric measurements were also performed, namely body weight, height, waist and hip circumferences. Each subject received a complete physical examination with emphasis given to body FR. Single slice CT scan was used to assess the cross-sectional areas of abdominal subcutaneous and abdominal visceral fat (Sensation 4 and Sensation 16, Siemens, Forchheim, Germany). A single slice CT section was obtained at the level of the L4 interspace using 120 kVp and 100 mA. Calculation of area was done by semiautomatic measurements of pixels, with density within specific attenuation numbers. Fat was defined as having attenuation number -150 to -15 , and soft tissues as -15 to $+100$ Hounsfield units (HU). All recorded data including anthropometric measurements, BMI, the widest diameter of the “buffalo hump” (when present), standardized digital photographs of selected body region (face, arms, legs, chest, abdomen side view, gluteus regions), DEXA scan results, and the single slice CT of the lumbar spine were presented to a specific external Fat Redistribution Adjudication Committee that used specific strict criteria [19] to support the classification of study subjects as having fat redistribution or not. The adjudication committee comprised of three clinical investigators who were not involved with subject interviews, data collection or analysis or in the clinical care of the study subjects performed the evaluation [19]. After the committee members assigned a first preliminary classification of each subject (according to medical chart information and data from study questionnaires and interviews) they subsequently proceeded to a second classification according to documented physical examination findings and digital photographs. This was performed according to published criteria [19]. Final classification of each subject was verified according to the diagnostic test criteria specified in the same publication [19]. All committee members were provided with identical information on which to base their decisions. The final classification of each subject required the unanimous agreement of the committee. Subjects whose classification was not unanimously agreed on ($n=9$) were not included in the analysis. This procedure was followed to obtain an objective assessment of the “fat redistribution subjects. Duration of illness was estimated from the time of diagnosis by the serological testing of each patient that was recorded in the medical chart.

Blood was drawn from each subject on the morning of the study visit after an overnight fast. Serum samples were immediately frozen at -70°C . Hormonal analyses were performed simultaneously on a subsequent day. The core lab of the Clinical Research center at the Beth Israel Deaconess Medical Center performed the insulin measurements. Glucose levels were measured by the Beth Israel Deaconess Medical Center Clinical Laboratory (Roche/Hitachi, Indianapolis, IN). Insulin levels were measured using a commercially available RIA (DSL-1600; Diagnostic Systems Laboratories, Inc., Webster, TX) with a sensitivity of 1.3 $\mu\text{IU/ml}$ and inter- and intra-assay coefficients of variation between 4.7% and 12.2% and 4.5% and 8.3%, respectively. Insulin resistance was estimated using the homeostasis model (HOMA) with the following formula: Insulin resistance (IR) = (fasting insulin \times fasting glucose)/22.5. Fasting lipoprotein profile was measured in the Clinical Chemistry laboratory at Children's Hospital in Boston as previously described. [20]. Both CD4 counts and viral load were measured in the BIDMC Clinical Laboratory (via flow cytometry / three-color CD4 reagent from Becton Dickinson and Co., Franklin Lakes, NJ for CD4 and ultrasensitive PCR (Amplicor HIV-1 monitor test, version 1.5; Roche-Cobas, Branchburg, NJ for HIV RNA measurements.

Statistical analysis

All data were expressed as mean values \pm standard deviation (Mean \pm SD). Kruskal-Wallis tests followed by post hoc (Mann-Whitney) analyses were used for comparisons of continuous variables among the four FR subgroups [non-FR, fat accumulation (FA), fat wasting (FW), and mixed FR]. Associations between MedDietScore and several aspects of HAART-induced metabolic syndrome (expressed as continuous variables) were investigated using bivariate analysis and multivariate linear regression. Variables that were not normally distributed were logarithmically transformed. SPSS version 10.0 for Windows software (SPSS, Inc., Chicago, IL) was used for data analysis. A P value of 0.05 was used to test for statistical significance, and all statistical tests were two tailed.

Results

Of the total sample, 133 (61%) patients had FR: 42 FA (19.3%), 56 mixed-FR (25.7%) and 35 FW (16.1%). Significant body composition differences were found among groups, with the FA group exhibiting significantly higher Body Mass Index (BMI), percent body fat and waist-to-hip ratio (WHR) compared to the non-FR group (Table 1). Fasting insulin levels and the HOMA-IR index were significantly higher in the FA and mixed-FR patients, compared to the non-FR patients, whereas high-density lipoprotein (HDL) cholesterol levels were lower in the mixed FR and the FW groups compared to the non-FR and FA groups. Subjects in the FR group had higher poultry intake and tended to have higher red meat/red meat products intake compared to the non-FR group ($p = 0.03$ and $p = 0.10$, respectively), but no other differences were detected with respect to food consumption or adherence to the Mediterranean diet among FR groups (Table 2).

In the entire study sample, we found a trend for a weak association between HOMA-IR index and MedDietScore (Spearman $r = -0.12$, $p = 0.07$). This association became statistically significant when we adjusted for potential confounders, namely age, sex, energy intake, BMI, WHR, smoking and exercise habits, CD4 cell count, total PI (months of use), total NRTI (months of use), total NNRTI (months of use) and duration of illness (standardized $\beta = -0.15$, $p = 0.03$). As the addition of FR in the above model tended to weaken the significance of the association (standardized $\beta = -0.11$, $p = 0.10$), we evaluated the effect of a potential interaction between the MedDietScore and the presence of FR in predicting HOMA-IR levels. The interaction term was a statistically significant predictor of HOMA-IR ($p = 0.03$) and, therefore, all subsequent multivariate analyses were stratified by FR group. No similar statistically significant interaction was found between the MedDietScore and the presence of FW.

Multivariate regression analysis in the FR group, adjusting for age, sex, total energy intake, BMI, WHR, revealed an inverse relationship between MedDietScore and HOMA-IR index (standardized $\beta = -0.13$, $p = 0.02$). The association remained significant after further controlling for smoking, physical activity, CD4 cell count, total PI (months of use), total NRTI (months of use), total NNRTI (months of use) and duration of illness (standardized $\beta = -0.23$, $p = 0.02$). Moreover, the degree of adherence to the Mediterranean Diet was found to be positively related with HDL-cholesterol (standardized $\beta = 0.15$, $p = 0.01$) and marginally negatively associated with blood triglyceride (TG) levels (standardized $\beta = -0.16$, $p = 0.13$), after adjustment for all the above-mentioned confounders. No similar associations were detected in the non-FR group, apart from a significant negative correlation between MedDietScore and DBP (standardized $\beta = -0.32$, $p = 0.01$).

Discussion

Adoption of a dietary pattern close to the Mediterranean dietary pattern has been associated with favorable effects on lipoprotein levels, markers of endothelium function, IR and the metabolic syndrome in non HIV-infected populations [21]. Furthermore, available evidence indicates that intensive lifestyle interventions are safer and better or of comparable effectiveness to drugs in reducing the prevalence of the metabolic syndrome and and/or preventing diabetes [22]. In HIV-infected patients, the role of lifestyle changes is increasingly recognized for the management of the metabolic abnormalities. Among the potential targets for dietary modification are polyunsaturated fatty acids, dietary fiber intake, alcohol and cholesterol intake, i.e. dietary factors that could affect insulin resistance and blood lipid abnormalities [23]. Current lifestyle recommendations for the treatment of dyslipidemia in HIV-infected adults do not differ from recommendations for the rest of the population at high cardiovascular risk [24]. The potential effect of the adherence to a Mediterranean-type dietary pattern on the parameters of the metabolic syndrome has not been previously examined in HIV-positive patients with the HAART-induced metabolic syndrome. According to our study, adherence to this dietary pattern was found to be favorably associated with important cardiovascular risk factors, namely HOMA-IR and HDL – cholesterol, in patients with HAART induced metabolic changes and FR, after controlling for factors considered to be significant in the pathogenesis of the syndrome, such as age and BMI. Thus, dietary interventions enhancing adherence to a Mediterranean dietary pattern may be proven to be an effective strategy for the reduction of dyslipidemia and insulin resistance in HIV – infected patients with FR.

Interestingly, associations between diet and metabolic factors were observed mainly in the FR group. Results from the Fat Redistribution and Metabolic Change in HIV Infection (FRAM) study have suggested that less leg subcutaneous adipose tissue and more visceral adipose tissue are important risk factors for adverse metabolic profile in HIV-men and women [25,26]. In our study, we pooled FA, mixed-FR and FW under the comprehensive “FR” variable, but we also evaluated “FW” separately. No interaction between insulin resistance and the degree of adherence to the Mediterranean diet was detected when lipoatrophy or fat wasting was considered separately. Due to the relatively small number of subjects in each group, further studies in this direction should be performed with larger numbers of patients. The fact that no major effect of diet was found in HIV-patient without FR could be potentially explained by several mechanisms. These patients have a metabolic profile closer to the normal population thus the effects would not be expected to be strong. They may also differ from the rest of the patients regarding other factors affecting the metabolic syndrome (e.g. they were younger, had shorter duration of illness, had used less NRTIs). It might also be possible that patients who have a predisposition to develop the metabolic syndrome may also be predisposed to have fat redistribution changes.

The inclusion of the MedDietScore in the prediction model of HOMA-IR levels changed the % variability only by 1% compared to the inclusion of FR only (data not shown). This is similar to the variance reported in recent population based diet studies [13]. On the other hand, prudent dietary patterns may affect circulating levels of adipocyte secreted hormones which are of major importance in mediating some of the changes observed [27,28]. In addition, dietary patterns may also affect the development of lipohypertrophy and/or regional body composition changes, and adipose tissue changes may be equally or more important predictors of adipocytokines and/or metabolic abnormalities in these patients as they precede or coincide with the metabolic changes observed. Limitations of this work include its cross-sectional, observational design, which does not allow us to establish firm cause-effect relationships and/or to elucidate underlying mechanisms. Our study may only raise hypotheses that could be further investigated by future prospective cohort studies and randomized clinical trials. Therefore, one cannot rule out the possibility that the effects presented herein could be observed in the metabolic syndrome in general and they are not uniquely related to HIV infection or its treatment. Furthermore, interventions based on diet should be considered in context to other therapeutic approaches for the metabolic syndrome, such as lipid lowering agents, exercise and drug switching.

Another limitation of the study is that the components of the index (MedDietScore) are equally weighted and similarly scored from 1 to 5. This may affect its accuracy as not all foods or food groups might influence the investigated health outcomes in the same way [14]. Although a misclassification would be possible, it would have been expected to suppress effect estimates. Also, misreporting of food items consumed could have influenced the calculation of the diet score and could, thus, have biased the results towards the null, but neither of the above factors could have strengthened and/or made significant the associations reported herein. Moreover, patients with FR may have been more accurate in reporting their dietary habits, due to higher motivation given the metabolic problems they were already facing. It is possible that this could have potentially resulted in a bias towards demonstrating stronger associations among this group. Although we examined separately the effects of the three major classes of antiretroviral medications, some of the individual medications may differ regarding their metabolic effects. Much larger studies will be necessary to further elucidate the role of individual medications and their interaction with dietary intake.

Finally, in the present study, IR was assessed using HOMA and not the euglycaemic-hyperinsulinaemic clamp, i.e. the gold-standard method. HOMA, however, is considered appropriate for cross-sectional epidemiological studies since it is strongly related to clamp-measured IR in both non-diabetic and diabetic subjects [29].

In conclusion, this is the first study revealing that adherence to the Mediterranean diet is favorably related to cardiovascular risk factors in a sample of HIV-positive patients with the HAART-induced metabolic syndrome studied cross-sectionally. The patients who benefited the most by higher adherence to this dietary pattern were found to be those with FR. As the Mediterranean diet has proven to be a metabolically favorable dietary pattern in the long term, its adoption may play a role in the prevention and treatment of the HAART-induced metabolic syndrome. Further clinical trials to confirm these findings and to investigate the mechanisms underlying its protective effects, are needed.

Acknowledgments

Financial Support: National Institutes of Health (grant M01-RR 01032 to the Beth Israel Deaconess Medical Center General Clinical Research Center and DK 58785, DK 799 and DK 0811913 to Dr. Mantzoros); American Diabetes Association (clinical research grant); Merck Research laboratories; and a discretionary grant from Beth Israel Deaconess Medical Center.

References

1. Hadigan C. Dietary habits and their association with metabolic abnormalities in human immunodeficiency virus-related lipodystrophy. *Clin Infect Dis* 2003;37:S101–4. [PubMed: 12942382]
2. Joy T, Keogh HM, Hadigan C, et al. Dietary fat intake and relationship to serum lipid levels in HIV-infected patients with metabolic abnormalities in the HAART era. *Aids* 2007;21:1591–600. [PubMed: 17630554]
3. Shah M, Tierney K, Adams-Huet B, et al. The role of diet, exercise and smoking in dyslipidaemia in HIV-infected patients with lipodystrophy. *HIV Med* 2005;6:291–8. [PubMed: 16011535]
4. Gavrilu A, Tsiodras S, Doweiko J, et al. Exercise and vitamin E intake are independently associated with metabolic abnormalities in human immunodeficiency virus-positive subjects: a cross-sectional study. *Clin Infect Dis* 2003;36:1593–601. [PubMed: 12802761]
5. Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* 2002;13:3–9. [PubMed: 11790957]
6. Jacobs DR Jr, Steffen LM. Nutrients, foods, and dietary patterns as exposures in research: a framework for food synergy. *Am J Clin Nutr* 2003;78:508S–513S. [PubMed: 12936941]
7. Jacques PF, Tucker KL. Are dietary patterns useful for understanding the role of diet in chronic disease? *Am J Clin Nutr* 2001;73:1–2. [PubMed: 11124739]
8. Assmann G, de Backer G, Bagnara S, et al. International consensus statement on olive oil and the Mediterranean diet: implications for health in Europe. The Olive Oil and the Mediterranean Diet Panel. *Eur J Cancer Prev* 1997;6:418–21. [PubMed: 9466113]
9. Hu FB. The Mediterranean diet and mortality--olive oil and beyond. *N Engl J Med* 2003;348:2595–6. [PubMed: 12826632]
10. Trichopoulos D, Lagiou P. Dietary patterns and mortality. *Br J Nutr* 2001;85:133–4. [PubMed: 11242479]
11. Cereda E, Malavazos AE, Favaro C, Pagani AM. Modified Mediterranean diet and survival: evidence for diet linked longevity is substantial. *Bmj* 2005;330:1329. author reply 1329-30. [PubMed: 15933361]
12. Chrysohou C, Panagiotakos DB, Pitsavos C, Das UN, Stefanadis C. Adherence to the Mediterranean diet attenuates inflammation and coagulation process in healthy adults: The ATTICA Study. *J Am Coll Cardiol* 2004;44:152–8. [PubMed: 15234425]
13. Mantzoros CS, Williams CJ, Manson JE, Meigs JB, Hu FB. Adherence to the Mediterranean dietary pattern is positively associated with plasma adiponectin concentrations in diabetic women. *Am J Clin Nutr* 2006;84:328–35. [PubMed: 16895879]
14. Panagiotakos DB, Pitsavos C, Arvaniti F, Stefanadis C. Adherence to the Mediterranean food pattern predicts the prevalence of hypertension, hypercholesterolemia, diabetes and obesity, among healthy adults; the accuracy of the MedDietScore. *Prev Med* 2007;44:335–40. [PubMed: 17350085]
15. Trichopoulou A, Bamia C, Norat T, et al. Modified Mediterranean diet and survival after myocardial infarction: the EPIC-Elderly study. *Eur J Epidemiol*. 2007
16. Block G, Coyle LM, Hartman AM, Scoppa SM. Revision of dietary analysis software for the Health Habits and History Questionnaire. *Am J Epidemiol* 1994;139:1190–6. [PubMed: 8209877]
17. Block G, Woods M, Potosky A, Clifford C. Validation of a self-administered diet history questionnaire using multiple diet records. *J Clin Epidemiol* 1990;43:1327–35. [PubMed: 2254769]
18. Mantzoros CS, Georgiadis EI. Body mass and physical activity are important predictors of serum androgen concentrations in young healthy men. *Epidemiology* 1995;6:432–5. [PubMed: 7548356]
19. Addy CL, Gavrilu A, Tsiodras S, Brodovicz K, Karchmer AW, Mantzoros CS. Hypoadiponectinemia is associated with insulin resistance, hypertriglyceridemia, and fat redistribution in human immunodeficiency virus-infected patients treated with highly active antiretroviral therapy. *J Clin Endocrinol Metab* 2003;88:627–36. [PubMed: 12574192]
20. Ridker PM, Hennekens CH, Buring JE, Rifai N. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 2000;342:836–43. [PubMed: 10733371]
21. Serra-Majem L, Roman B, Estruch R. Scientific evidence of interventions using the Mediterranean diet: a systematic review. *Nutr Rev* 2006;64:S27–47. [PubMed: 16532897]

22. Esposito K, Ciotola M, Giugliano D. Mediterranean diet and the metabolic syndrome. *Mol Nutr Food Res* 2007;51:1268–74. [PubMed: 17879992]
23. Hadigan C, Meigs JB, Corcoran C, et al. Metabolic abnormalities and cardiovascular disease risk factors in adults with human immunodeficiency virus infection and lipodystrophy. *Clin Infect Dis* 2001;32:130–9. [PubMed: 11118392]
24. Dube MP, Stein JH, Aberg JA, et al. Guidelines for the evaluation and management of dyslipidemia in human immunodeficiency virus (HIV)-infected adults receiving antiretroviral therapy: recommendations of the HIV Medical Association of the Infectious Disease Society of America and the Adult AIDS Clinical Trials Group. *Clin Infect Dis* 2003;37:613–27. [PubMed: 12942391]
25. Wohl D, Scherzer R, Heymsfield S, et al. The associations of regional adipose tissue with lipid and lipoprotein levels in HIV-infected men. *J Acquir Immune Defic Syndr* 2008;48:44–52. [PubMed: 18360291]
26. Currier J, Scherzer R, Bacchetti P, et al. Regional adipose tissue and lipid and lipoprotein levels in HIV-infected women. *J Acquir Immune Defic Syndr* 2008;48:35–43. [PubMed: 18197118]
27. Tsiodras S, Mantzoros C. Leptin and Adiponectin in the HIV Associated Metabolic Syndrome: Physiologic and Therapeutic Implications. *Am J Infect Dis* 2006;2:141–152. [PubMed: 17183414]
28. Yannakoulia M, Yiannakouris N, Melistas L, Kontogianni MD, Malagaris I, Mantzoros CS. A dietary pattern characterized by high consumption of whole-grain cereals and low-fat dairy products and low consumption of refined cereals is positively associated with plasma adiponectin levels in healthy women. *Metabolism* 2008;57:824–30. [PubMed: 18502266]
29. Bonora E, Targher G, Alberiche M, et al. Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity: studies in subjects with various degrees of glucose tolerance and insulin sensitivity. *Diabetes Care* 2000;23:57–63. [PubMed: 10857969]

Table 1

Characteristics of the patients according to their fat redistribution status (Data are presented as mean values \pm standard deviations or as frequencies).

	Non-FR group (n = 85)	FA group (n = 42)	FR-group Mixed FR group (n = 56)	FW group (n = 35)	Overall P
<i>Demographic and lifestyle characteristics</i>					
Age (yrs)	42 \pm 8	45 \pm 7	46 \pm 9 ^a	45 \pm 7	0.03
Sex (% female)	10.6	28.6 ^{a,c}	17.9 ^c	2.9	0.01
Race (% white)	71.4	54.8 ^f	91.1 ^{d,e}	88.6 ^a	<0.001
Exercise (no of sessions/week)	4.7 \pm 2.6	4.1 \pm 2.8	3.7 \pm 2.8 ^b	5.3 \pm 2.3	0.03
Alcoholic drinks (no of drinks/week)	2.2 \pm 3.7	1.3 \pm 2.3	1.7 \pm 4.0	3.3 \pm 7.4	0.21
Current smoker (%)	44.7	31.0	32.1	40.0	0.34
<i>Anthropometric, body composition and metabolic variables</i>					
BMI (kg/m ²)	23.7 \pm 2.5	30.9 \pm 7.0 ^{d,f}	24.5 \pm 2.9 ^e	22.7 \pm 2.6 ^e	<0.001
Waist to hip ratio	0.92 \pm 0.06	0.98 \pm 0.09 ^d	0.99 \pm 0.07 ^{d,f}	0.94 \pm 0.04	<0.001
Percent body fat	19.8 \pm 6.1	29.5 \pm 8.4 ^d	19.2 \pm 6.5 ^{e,f}	12.7 \pm 4.5 ^d	<0.001
Systolic blood pressure (mmHg)	127 \pm 19	132 \pm 19	130 \pm 18	132 \pm 18	0.51
Diastolic blood pressure (mmHg)	73 \pm 10	78 \pm 13	76 \pm 9	75 \pm 11	0.06
Total Cholesterol (mg/dL)	204 \pm 52	227 \pm 64	219 \pm 75	212 \pm 55	0.21
LDL-cholesterol (mg/dL)	120 \pm 43	129 \pm 44	111 \pm 48	117 \pm 33	0.22
HDL-Cholesterol (mg/dL)	43 \pm 14	41 \pm 11 ^c	32 \pm 9 ^{d,e}	32 \pm 10 ^d	<0.001
Triglycerides (mg/dL)	193 \pm 158	286 \pm 232	436 \pm 489 ^d	352 \pm 240	<0.001
Glucose (mg/dL)	85.6 \pm 12.7	113.9 \pm 60.4 ^{c,d}	93.5 \pm 33.0 ^b	87.0 \pm 14.1	<0.001
Insulin (μ U/mL)	10.4 \pm 9.6	22.3 \pm 21.0 ^{c,d}	23.3 \pm 23.3 ^a	22.3 \pm 26.9	<0.001
Insulin resistance, HOMA	2.4 \pm 3.2	6.9 \pm 8.1 ^d	6.1 \pm 8.9 ^a	4.8 \pm 5.2 ^d	<0.001
<i>Variables related to HIV-infection and antiretroviral therapy</i>					

	Non-FR group (n = 85)			FR-group Mixed FR group (n = 56)		FW group (n = 35)		Overall P
	FA group (n = 42)							
Duration of illness (months)	101 ± 55	116 ± 56	126 ± 48 ^a	127 ± 40	0.01			
CD4 cells count (cells/mm ³)	484 ± 281	568 ± 350	506 ± 315	502 ± 263	0.55			
Viral Load (copies/mL)	6551 ± 20377	10162 ± 24710	15792 ± 48096	33881 ± 104306	0.08			
Total PI use (months)	30 ± 28	40 ± 35	44 ± 29	44 ± 33	0.04			
Total NRTI use (months)	96 ± 91	104 ± 59	141 ± 72 ^{b,d}	143 ± 71 ^a	<0.001			
Total NNRTI use (months)	10 ± 11	16 ± 19	14 ± 17	12 ± 14	0.23			

FR= Fat redistribution, FA = fat accumulation, FW = fat wasting, BMI = Body Mass Index, HOMA = homeostasis model, LDL = Low Density Lipoprotein, HDL = High Density Lipoprotein, HIV = Human Immunodeficiency Virus, PI = Protease Inhibitors, NRTI = Nucleoside Reverse Transcriptase Inhibitors, NNRTI = Non-Nucleoside Reverse Transcriptase Inhibitors

- ^a Statistically different from non-FR Group (p<0.05);
- ^b Statistically different from FA Group (p<0.05);
- ^c Statistically different from FW group (p<0.05);
- ^d Statistically different from non-FR (p≤0.01);
- ^e Statistically different from FA Group (p≤0.001);
- ^f Statistically different from FW group (p≤0.001).

Table 2

Consumption of main food groups (as servings/week unless otherwise specified) and Mediterranean Diet Score of the study participants by fat redistribution status (Data are presented as mean values \pm standard deviation).

	Non-FR group (n = 81)		FR-group		
	Total group (n = 128)	FA group (n = 39)	Mixed FR group (n = 55)	FW group (n = 34)	
Non-refined cereals	7.0 \pm 9.9	5.8 \pm 6.4	5.8 \pm 6.9	7.6 \pm 9.9	
Potatoes	4.1 \pm 4.0	5.3 \pm 5.8	4.5 \pm 4.7	4.8 \pm 3.8	
Fruits	19.1 \pm 15.8	22.3 \pm 16.5	15.2 \pm 13.0	17.0 \pm 13.8	
Vegetables	9.0 \pm 8.1	10.2 \pm 10.1	10.4 \pm 9.5	11.3 \pm 8.6	
Legumes	2.1 \pm 2.4	1.6 \pm 1.4	1.9 \pm 1.5	2.6 \pm 2.9	
Fish	4.4 \pm 5.8	7.0 \pm 11.2	5.4 \pm 10.0	4.6 \pm 4.6	
Red meat / red meat products	5.9 \pm 5.5	9.6 \pm 12.8	6.0 \pm 3.7	9.4 \pm 14.7	
Poultry	3.9 \pm 3.6	6.1 \pm 6.2	4.6 \pm 4.4	5.2 \pm 5.6	
Full-fat dairy	7.8 \pm 8.2	7.9 \pm 8.1	7.1 \pm 7.7	10.0 \pm 9.9	
Alcoholic beverages (ml ethanol/day)	12.1 \pm 19.4	11.4 \pm 24.1	10.4 \pm 19.3	11.0 \pm 27.0	13.4 \pm 24.5
MedScore (0 – 55) (range)	26.3 \pm 5.1 (15 - 37)	25.8 \pm 4.9 (13 - 39)	25.7 \pm 4.3 (18 - 35)	25.7 \pm 5.2 (13 - 36)	26.0 \pm 5.2 (19 - 39)

FR= Fat redistribution, FA = fat accumulation, FW = fat wasting, MedScore = Mediterranean Diet Score

^aComparisons vs. non-FR group, p = 0.10;

^bComparisons vs. non-FR group, p = 0.03.