

RESEARCH

Open Access

Differential genome-wide gene expression profiling of bovine largest and second-largest follicles: identification of genes associated with growth of dominant follicles

Ken-Go Hayashi, Koichi Ushizawa, Misa Hosoe, Toru Takahashi*

Abstract

Background: Bovine follicular development is regulated by numerous molecular mechanisms and biological pathways. In this study, we tried to identify differentially expressed genes between largest (F1) and second-largest follicles (F2), and classify them by global gene expression profiling using a combination of microarray and quantitative real-time PCR (QPCR) analysis. The follicular status of F1 and F2 were further evaluated in terms of healthy and atretic conditions by investigating mRNA localization of identified genes.

Methods: Global gene expression profiles of F1 (10.7 +/- 0.7 mm) and F2 (7.8 +/- 0.2 mm) were analyzed by hierarchical cluster analysis and expression profiles of 16 representative genes were confirmed by QPCR analysis. In addition, localization of six identified transcripts was investigated in healthy and atretic follicles using in situ hybridization. The healthy or atretic condition of examined follicles was classified by progesterone and estradiol concentrations in follicular fluid.

Results: Hierarchical cluster analysis of microarray data classified the follicles into two clusters. Cluster A was composed of only F2 and was characterized by high expression of 31 genes including IGFBP5, whereas cluster B contained only F1 and predominantly expressed 45 genes including CYP19 and FSHR. QPCR analysis confirmed AMH, CYP19, FSHR, GPX3, PIGF, PLA2G1B, SCD and TRB2 were greater in F1 than F2, while CCL2, GADD45A, IGFBP5, PLAUR, SELP, SPP1, TIMP1 and TSP2 were greater in F2 than in F1. In situ hybridization showed that AMH and CYP19 were detected in granulosa cells (GC) of healthy as well as atretic follicles. PIGF was localized in GC and in the theca layer (TL) of healthy follicles. IGFBP5 was detected in both GC and TL of atretic follicles. GADD45A and TSP2 were localized in both GC and TL of atretic follicles, whereas healthy follicles expressed them only in GC.

Conclusion: We demonstrated that global gene expression profiling of F1 and F2 clearly reflected a difference in their follicular status. Expression of stage-specific genes in follicles may be closely associated with their growth or atresia. Several genes identified in this study will provide intriguing candidates for the determination of follicular growth.

Background

The final stage of bovine follicular development occurs in a wave-like fashion [1,2]. During a wave, increase of follicle-stimulating hormone (FSH) induces recruitment of a cohort of follicles beyond 4 mm in diameter and usually a single follicle is selected as a dominant follicle

(DF) [3,4]. Although the DF continues to grow by transition of gonadotropin dependency from FSH to luteinizing hormone (LH) and secretes large quantities of estradiol (E₂), the remaining subordinate follicles (SFs) cease to grow, then undergo atresia [5]. It is well documented that increased expression of LH receptor (LHR) in granulosa cells (GC) and specific changes of intrafollicular factors such as the insulin-like growth factor (IGF) and inhibin-activin-follistatin systems play a critical role

* Correspondence: tatoru@affrc.go.jp

Reproductive Biology Research Unit, Division of Animal Science, National Institute of Agrobiological Sciences, Tsukuba 305-8602, Japan

in E₂ production in the DF [6,7]. Therefore, regulatory mechanisms of follicular development are closely associated with complex interactions between follicular local paracrine/autocrine factors and endocrine hormones.

Increasing evidence using global gene expression analysis such as a DNA microarray, suppression subtractive hybridization and serial analysis of gene expression have identified numerous genes in various aspects of bovine follicular development [8-18]. Some studies compared the gene expression profiles between DF and SF around the time of follicular selection. They showed that DF up-regulates genes regulating E₂ synthesis, anti-apoptosis, cell proliferation and gene transcription. Conversely, SF enhanced the expression of genes associated with pro-apoptosis and cell death compared with the DF [8,9,13,14]. Recent studies found that 93 mostly novel genes were differently expressed in the GC of newly selected DF compared with SF and/or growing cohort follicles whereas most of these genes were down-regulated in the GC of preovulatory follicles during final maturation before the LH surge [15,17]. Growth of a DF during 2-5.5 days following follicular wave emergence was associated with a decrease in genes encoding proliferation and pro-apoptotic factors and an increase in genes regulating anti-apoptotic factors [12]. An increase in follicular diameter during follicular growth was accompanied by alteration of gene expression regulating some growth factors and cytokines [16,18]. Ndiaye *et al.* identified a subset of novel genes down-regulated in preovulatory follicles after human chorionic gonadotropin (hCG) stimulation compared with DF, which may contribute to ovulation and luteinization [11].

These previous studies lead us to suggest that gene expression profiles in individual follicles reflect their developmental status, thus each follicle can be classified by differences in gene expression profiles. On the other hand, details of the genetic processes and biological pathways regulating bovine follicular development still remain to be elucidated. We consider that investigating the global gene expression of follicles after selection can help to understand the molecular mechanisms responsible for the regulation and control of follicular development and atresia. Therefore, in this study, we tried to classify the largest (F1) and second-largest (F2) follicles according to differences in gene expression profiles and to identify differentially expressed genes between the groups using a combination of microarray analysis and quantitative real-time PCR (QPCR) analysis. In addition, spatial expression profiles of several identified genes were investigated using *in situ* hybridization in healthy and atretic follicles classified based on follicular fluid (FF) concentration of steroids.

Methods

Experiment 1: classification of F1 and F2 and identification of genes by microarray analysis and QPCR analysis

Sample collection and RNA extraction

Paired ovaries were obtained from four pregnant Japanese Black cows in the institute ranch less than 10 min after slaughtering. These cows were pregnant and slaughtered for another study. Both F1 and F2 were dissected from the ovaries. Then, the surrounding stroma and theca externa were removed from the follicular walls. We collected three F1 and three F2 from four cows because two cows had both F1 and F2 collected whereas one cow had only a F1 collected and another cow had only a F2 collected. The follicles were snap-frozen and stored at -80°C until RNA extraction. Total RNA from the follicular wall (i.e., granulosa plus theca interna) was extracted from each follicle using ISOGEN (NipponGene, Tokyo, Japan) according to the manufacturer's instructions. All procedures for animal experiments were carried out in accordance with guidelines approved by the Animal Ethics Committee of the National Institute of Agrobiological Sciences for the use of animals.

Microarray analysis

A custom-made bovine oligonucleotide microarray fabricated by Agilent Technologies (Santa Clara, CA, USA) was used in this study. Sixty-mer nucleotide probes for customized microarray were synthesized on a glass slide. The annotated bovine oligonucleotide array represented 10263 sequences 4466 of which were known bovine genes, 5697 were unknown sequences and possible candidates for novel bovine genes, and 100 internal references.

We performed one-color microarray using five follicles (three F1 and two F2). Fluorescence-labeled (Cy3) cRNA probes were prepared from 150-300 ng of total RNA of each follicle using a Low RNA Input Linear Amplification Kit (Agilent Technologies). Labeled cRNA probes (750 ng each) were hybridized to the customized microarray in hybridization buffer (Gene Expression Hybridization Kit, Agilent Technologies) at 60°C for 17 h. After hybridization, the arrays were washed with 6 × SSC, 0.005% Triton X-102 at room temperature for 10 min, followed by 5-min washes in 0.1 × SSC, 0.005% Triton X-102 at 4°C. Hybridized arrays were blow dried with N₂ gas and scanned using an Agilent Microarray Scanner (Agilent Technologies), and Feature Extraction ver. 9.1 (Agilent Technologies) was used for image analysis and data extraction. Gene expression datasets were normalized using the median of the signal intensity for 100 *GAPDH* genes on a microarray platform as internal control.

After normalization, 3308 genes were left to use for further analysis. The relative abundance of individual genes between follicles was calculated by dividing the normalized value of the genes between each follicle. We used the normalized microarray data of genes that showed an expression level of more than 20-fold between at least two follicles for subsequent hierarchical cluster analysis. The data were transformed \log_2 values and hierarchical cluster analysis was performed using the TIGR MultiExperiment Viewer 4.0 (MeV 4.0) software program [19]. Two parameters (average linkage and cosine correlation) were selected for constructing the hierarchical tree. Compliance with Minimum Information About a Microarray Experiment (MIAME) [20] was assured by depositing all the data in the Gene Expression Omnibus (GEO) repository [21]. The GEO accession numbers are as follows. Platform: GPL9136; Samples: GSM453634, GSM453635, GSM453636, GSM453637 and GSM453638; Series: GSE18145.

Quantitative real-time RT-PCR analysis

To validate the results of microarray analysis, we confirmed mRNA expression of 16 representative genes using QPCR analysis. All six follicles were used in QPCR analysis. The procedures for QPCR were previously described [22]. Briefly, single-strand cDNA was reverse-transcribed from 50 ng of total RNA using MultiScribe™ reverse transcriptase with a random primer, dNTP mixture, $MgCl_2$ and RNase inhibitor (Applied Biosystems, Foster City, CA, USA). The reverse transcription cycle consisted of 10 min annealing at 25°C, 30 min cDNA synthesis at 48°C and 5 min inactivation at 95°C. The primers were designed using the Primer Express computer software program (Applied Biosystems) based on the bovine sequences. The primer sequences for each gene are given in Table 1. Each QPCR reaction (25 μ l) contained 1 μ l cDNA template, 0.5 μ l forward primer (20 μ M), 0.5 μ l reverse primer (20 μ M), 12.5 μ l Power SYBR® Green PCR Master Mix (Applied Biosystems) and 10.5 μ l nuclease-free water. The thermal cycling conditions included one cycle at 50°C for 2 min, one cycle at 95°C for 10 min, and 40 cycles at 95°C for 15 s and 60°C for 1 min. Each cDNA template was analyzed for quantitation in duplicate. QPCR and the resulting relative increase in reporter fluorescent dye emission were monitored in real time using an Mx3000P QPCR system (Stratagene, La Jolla, CA, USA). The relative difference in the initial amount of each mRNA species (or cDNA) was determined by comparing the cycle threshold values. To quantify the mRNA concentrations, standard curves for each gene were generated by serial dilution of the plasmid containing its cDNA. The melting curve for detecting the SYBR Green-based objective amplicon were confirmed because SYBR Green also detects double-stranded DNA

including primer dimers, contaminating DNA and PCR products from misannealed primers. Contaminating DNA or primer dimers appear as a peak separate from the desired amplicon peak.

Experiment 2: localization of characteristic genes identified in experiment 1 in healthy and atretic follicles using *in situ* hybridization

Sample collection and storage

Ovaries containing follicles more than 8 mm in diameter were obtained from Japanese Black cows at local slaughterhouse. We used only follicles which have a transparent follicular wall and fluid and did not show any aspect of cystic follicles. Eleven follicles were collected and 200 μ l of FF was aspirated from each follicle by a syringe fitted with a 27G needle. The FF was snap-frozen and stored at -30°C until hormone determinations. The follicles were dissected from the ovaries and fixed in 10% formalin, embedded in paraffin wax, and stored at 4°C until *in situ* hybridization.

Steroid hormone determinations

Concentrations of E_2 and P_4 in the FF samples were determined directly in duplicate using a time-resolved fluoroimmunoassay (TR-FIA). The TR-FIA for E_2 and P_4 was performed as previously described by our laboratory [23,24]. The FF samples were diluted to 100-, 2000- and 5000-fold for E_2 determination and 25-fold for P_4 determination using charcoal-treated plasma (collected from adult Japanese-Black cows). Ranges of the standard curves were 5-200 pg/ml for E_2 and 0.33-36 ng/ml for P_4 . The intra- and interassay coefficients of variation were 8.2 and 11.4% for E_2 , and 8.5 and 10.5% for P_4 , respectively.

***In situ* hybridization**

We classified follicles into two groups based on relative levels of FF concentrations of E_2 and P_4 ($E_2/P_4 \geq 1$: healthy; $E_2/P_4 < 1$: atretic). Six representative genes differently expressed between F1 and F2 in experiment 1 were selected for *in situ* hybridization: anti-Mullerian hormone (*AMH*), cytochrome P450, family XIX (*CYP19*), growth arrest and DNA-damage-inducible, alpha (*GADD45A*), IGF binding protein 5 (*IGFBP5*), placental growth factor (*PLGF*) and thrombospondin 2 (*TSP2*). In these genes, *CYP19* and *IGFBP5* were selected as markers of healthy or atretic follicles since mRNA expression of *CYP19* and *IGFBP5* were up-regulated in the bovine DF and SF, respectively [25,26].

Digoxigenin (DIG)-labeled antisense and sense cRNA probes were prepared as previously described [27,28]. For hybridization, follicles were sectioned into 7 μ m-thick sections. We performed *in situ* hybridization using an automated Ventana HX System Discovery with a RiboMapKit and a BlueMapKit (Roche Diagnostics, Basel, Switzerland) as previously described by our laboratory [27,28]. Briefly, the sections were hybridized

Table 1 Details of the primers used for quantitative real-time RT-PCR analysis

Gene name	GeneBank accession number	Primer	Sequences	Position
AMH	NM_173890	Forward	5'-ACACCGGCAAGCTCCTCAT-3'	1647-1665
		Reverse	5'-CACCATGTTTGGGACGTGG-3'	1714-1696
CCL2	NM_174006	Forward	5'-CGCTCAGCCAGATGCAATTA-3'	110-129
		Reverse	5'-GCCTCTGCATGGAGATCTTCTT-3'	186-165
CYP19	NM_174305	Forward	5'-TCCATGGGATTTTCCAGGC-3'	2050-2068
		Reverse	5'-TGGTGGCTTGCTTTTCCAAC-3'	2123-2103
FSHR	NM_174061	Forward	5'-AATCTACCTGCTGCTCATAGCCTC-3'	1300-1323
		Reverse	5'-TTTGCCAGTCGATGGCATAG-3'	1376-1357
GADD45A	NM_001034247	Forward	5'-CCGCATTCATCACAGTGGAA-3'	592-611
		Reverse	5'-CATCACCGTTCAGGGAGATTAATC-3'	704-681
GPX3	NM_174077	Forward	5'-GCTTCCCCTGCAACCAAT-3'	357-375
		Reverse	5'-TCGAACATACTTGAGGGTGGCT-3'	433-412
IGFBP5	NM_001105327	Forward	5'-ACTGTGACCCGAAAGGGTTCT-3'	682-702
		Reverse	5'-TTCATCCCCTACTTGTCCACG-3'	778-758
PIGF	NM_173950	Forward	5'-TGAATGACTACTCCCTCCATG-3'	877-898
		Reverse	5'-GGTCTGTCTTTCTCTCACGTTTC-3'	957-933
PLAUR	NM_174423	Forward	5'-CGCGGCCCTATGAATCAAT-3'	730-748
		Reverse	5'-CTGATGGTGAGCTTGGGTTCC-3'	800-779
PLA2G1B	NM_174646	Forward	5'-GGCCTTCATCTGCAACTGTGA-3'	358-378
		Reverse	5'-TGTGCTCCTTGTGTATGGCA-3'	428-408
SCD	NM_173959	Forward	5'-ATTCCCAGCTGGCTTTTTC-3'	659-678
		Reverse	5'-TCTTTGACAGCTGGGTGTTTG-3'	729-708
SELP	NM_174183	Forward	5'-GTCAAGCAGGGCCACTGACTAT-3'	1700-1720
		Reverse	5'-TCACTAAGCCTGTTGATCCAGCTG-3'	203-2182
SPP1	NM_174187	Forward	5'-AGCCCTGAGCAAACAGACGAT-3'	304-324
		Reverse	5'-GCGTCGTCGGAGTCATTAGAGT-3'	380-359
TIMP1	NM_174471	Forward	5'-CTATGCTGCTGGTTGTGAGGAAT-3'	508-530
		Reverse	5'-TGAGTGTGCTCTGCAGTTTG-3'	582-562
TRB2	NM_178317	Forward	5'-GACCTCAAGCTTCGAAATTCA-3'	525-546
		Reverse	5'-CGTCATCTCCCCGAGAATAT-3'	621-601
TSP2	NM_176872	Forward	5'-GGAAAACAAGTCATGGCGGA-3'	3845-3864
		Reverse	5'-TTGAGAGAAGACAAACAGACCCAG-3'	3928-3902
GAPDH	U85042	Forward	5'-ACCCAGAAGACTGTGGATGG-3'	444-463
		Reverse	5'-CAACAGACACGTTGGGAGTG-3'	621-602

with DIG-labeled probes in RiboHybe (Roche Diagnostics) hybridization solution at 65°C (*PIGF*) or 61°C (*AMH*, *CYP19*, *GADD45A*, *IGFBP5* and *TSP2*) for 6 hours, then washed for 3 × 6 min in RiboWash (Roche Diagnostics) at 65°C and fixed in RiboFix (Roche Diagnostics) at 37°C, 10 min. The hybridization signals were detected with a rabbit polyclonal anti-digoxin antibody HRP conjugate (Dako Cytomation, Carpinteria, CA, USA) using an AmpMapKit (Roche Diagnostics). The hybridized slides were observed with a Leica DMRE HC microscope (Leica Microsystems, Wetzlar, Germany) and a Nikon Digital Sight DS-Fi1-L2 (Nikon, Tokyo, Japan).

Statistical analysis

In experiment 1, the expression ratio of each gene to *GAPDH* mRNA was calculated to adjust for variations

in the QPCR reaction. The follicular diameter and the QPCR data in experiment 1 and concentrations of E₂ and P₄ and E₂/P₄ ratio in FF in experiment 2 were analyzed by Mann-Whitney's U test. Results were presented as the mean ± SEM. Statistical significance was considered to be at *P* < 0.05.

Results

Experiment 1: classification of F1 and F2 and identification of genes by microarray analysis and QPCR analysis

Mean diameter of F1 and F2 were 10.7 ± 0.7 and 7.8 ± 0.2 mm, respectively (*P* < 0.05).

Hierarchical cluster analysis of microarray data

The expression level of 76 genes was enhanced between at least two follicles by more than 20-fold. Using the

microarray data of these 76 genes, we performed a hierarchical cluster analysis and constructed a cluster heat map (Figure 1). As can be seen from the dendrogram of the sample axis, clustering analysis distinctly separated the five follicles into two clusters (A and B) based on their microarray expression profiles. Cluster A included two follicles that were both F2, whereas cluster B contained the remaining three follicles that were all F1. Cluster analysis also identified two major clusters in the gene axis. One cluster contained 31 genes that were relatively highly expressed in cluster A, while the other contained 45 genes that were relatively highly expressed in cluster B. The details of highly expressed genes in clusters A and B are listed in Table 2 and 3, respectively.

Quantitative PCR analysis of representative highly expressed genes in F1 and F2

Figure 2 shows the results of QPCR analysis of the eight representative genes that were highly expressed in F2 (cluster A) compared with F1 (cluster B) in microarray analysis. Messenger RNA expression for chemokine ligand 2 (*CCL2*), *GADD45A*, *IGFBP5*, plasminogen activator urokinase receptor (*PLAUR*), secreted phosphoprotein 1 (*SPP1*), selectin P (*SELP*), tissue inhibitor of matrix metalloproteinase-1 (*TIMP1*) and *TSP2* was greater in the F2 than in the F1 ($P < 0.05$). The results of QPCR analysis of the eight representative genes that were highly expressed in the F1 as compared with the F2 in microarray analysis are shown in Figure 3. The expression of *AMH*, *CYP19*, *FSHR*, glutathione peroxidase 3 (*GPX3*), *PIGF*, phospholipase A2 group 1B (*PLA2G1B*), stearoyl-CoA desaturase (*SCD*) and tribbles homolog 2 (*TRB2*) mRNA was greater in the F1 than in the F2 ($P < 0.05$).

Experiment 2: localization of characteristic genes identified in experiment 1 in healthy and atretic follicles using in situ hybridization

Follicular fluid concentrations of E_2 and P_4 in follicles

We classified follicles into healthy or atretic based on the relative concentrations of E_2 and P_4 in FF (healthy: E_2/P_4 ratio ≥ 1 , atretic: E_2/P_4 ratio < 1). From a total of 11 follicles, eight were categorized into healthy while the other three were atretic. Table 4 shows the characteristics of the follicles used in experiment 2. Healthy follicles had higher E_2 and lower P_4 concentrations in FF than atretic follicles. The E_2/P_4 ratio in FF was significantly higher in healthy follicles than in atretic follicles.

In situ hybridization of representative genes identified in experiment 1

Figure 4 shows mRNA localization for *GADD45A*, *IGFBP5* and *TSP2* in healthy and atretic follicles by *in situ* hybridization. These genes were highly expressed in




Figure 1 Hierarchical cluster analysis of 76 differentially expressed genes in largest (F1) and second-largest follicles (F2). These genes were enhanced between at least two follicles by more than 20-fold. Red scale indicates relative higher expression level and green scale indicates relative lower expression level. The expression levels were transformed to \log_2 values. Dendrograms of sample axis (above matrix) and gene axis (to the left of matrix) represent overall similarities in gene expression profiles. Five follicles were classified into two major clusters (A and B). The follicles divided into cluster A were all F2 and the follicles divided into cluster B were all F1. The cluster A was characterized by highly expression of 31 genes, whereas the cluster B was predominately expressed 45 genes.

Table 2 List of differentially expressed genes in cluster A as compared with cluster B.

Accession No.	Gene symbol	Gene name
NM_001101080	ADAMTS1	ADAM metalloproteinase with thrombospondin type 1 motif, 1
NM_206843	AGOUTI	AGOUTI protein
NM_001098982	CCDC80	Coiled-coil domain containing 80
NM_174006	CCL2	Chemokine (C-C motif) ligand 2
NM_001008670	CRABP2	Cellular retinoic acid binding protein 2
NM_174290	CRYAB	Crystallin, alpha B
NM_001034247	GADD45A	Growth arrest and DNA-damage-inducible, alpha
NM_001077112	GSTA3	Glutathione S-transferase, alpha 3
NM_001105327	IGFBP5	Insulin-like growth factor binding protein 5
NM_174357	IL1RN	Interleukin 1 receptor antagonist
NM_001033610	KRT8	Keratin 8
NM_001001138	LOC407171	Fc gamma 2 receptor
NM_174384	LOXL4	Lysyl oxidase-like 4
NM_174132	OLR1	Oxidized low density lipoprotein (lectin-like) receptor 1
NM_176855	OXT	Oxytocin
NM_001101883	PDK4	Pyruvate dehydrogenase kinase, isozyme 4
NM_174141	PENK	Proenkephalin
NM_174423	PLAUR	Plasminogen activator, urokinase receptor
NM_001034681	RTKN	Rhotekin
NM_174183	SELP	Selectin P
NM_174137	SERPINE1	Serpin peptidase inhibitor, clade E (nexin, plasminogen activator inhibitor type 1), member 1
NM_001075764	SFRP4	Secreted frizzled-related protein 4
NM_174601	SLC1A5	Solute carrier family 1 (neutral amino acid transporter), member 5
NM_174187	SPP1	Secreted phosphoprotein 1
NM_003254	TIMP1	TIMP metalloproteinase inhibitor 1
NM_176872	TSP2	Thrombospondin 2
AW430112		Transcription factor B1, mitochondrial
BE721140		Transcribed locus
BP101259		Caldesmon, smooth muscle
XM_587930		Similar to ATP-binding cassette sub-family G member 1 (ABCG1), mRNA.
XM_869699		Similar to tumor necrosis factor receptor superfamily, member 12A

F2 than in F1 in microarray and QPCR analysis of experiment 1. *IGFBP5* mRNA was localized in the GC and theca layer (TL) of atretic follicles but not in healthy follicles (Figure 4E, F, G and 4H). *GADD45A* (Figure 4A, B, C and 4D) and *TSP2* (Figure 4I, J, K and 4L) mRNA were found in both GC and TL of atretic follicles but they were expressed in only GC of healthy follicles. No significant signals were detected with any sense probes (Figure 4B, D, F, H, J and 4L).

Localization of *AMH*, *CYP19* and *PIGF* mRNA in healthy and atretic follicles are shown in Figure 5. These genes were expressed more in the F1 than in the F2 in experiment 1. *AMH* (Figure. 5A, B, C and 5D) and *CYP19* (Figure. 5E, F, G and 5H) mRNA was localized in GC of healthy as well as atretic follicles. *PIGF* mRNA was found in GC and TL of only healthy follicles but not atretic follicles (Figure. 5I, J, K and 5L). No significant signals were detected with any sense probes (Figure. 5B, D, F, H, J and 5L).

Discussion

In this study, as expected, hierarchical cluster analysis of the microarray data classified F1 and F2 according to differences in gene expression profiles. In each follicular group, characteristic genes determining their developmental status were expressed. The F1 showed greater expression of genes responsible for enhancement of follicular E_2 production than the F2. These genes were gonadotropin receptor (*FSHR*), steroidogenic enzymes (*CYP17*, *CYP19* and *HSD17B1*) and inhibin-activin-follistatin system (*INHA*, *INHBA* and *FST*). It is well demonstrated that mRNA expression for *FSHR*, *CYP17*, *CYP19*, *INHA* and *INHBA* increases with the progress of bovine follicular development and is greater in DF than SF [8,12,29]. On the other hand, the F2 had greater expression of *IGFBP5* mRNA than the F1. *IGFBP5* mRNA expression dramatically increased in bovine atretic follicles compared with the DF [26]. Intrafollicular levels of IGFBP proteolytic activity and *IGFBPs* gene expression


Table 3 List of differentially expressed genes in cluster B as compared with cluster A.

Accession No.	Gene symbol	Gene name
NM_173890	AMH	Anti-Mullerian hormone
NM_174518	CITED1	Cbp/p300-interacting transactivator, with Glu/Asp-rich carboxy-terminal domain, 1
NM_174304	CYP17	Cytochrome P450, subfamily XVII
NM_174305	CYP19	Cytochrome P450, family XIX, aromatase
NM_001034410	CXCL14	Chemokine (C-X-C motif) ligand 14
NM_182786	FOS	V-fos FBJ murine osteosarcoma viral oncogene homolog
NM_174061	FSHR	Follicle stimulating hormone receptor
NM_175801	FST	Follistatin
NM_001083674	GCLC	Glutamate-cysteine ligase, catalytic subunit
NM_174077	GPX3	Glutathione peroxidase 3
NM_174546	GUCA1A	guanylate cyclase activator 1A (retina)
NM_001102365	HSD17B1	Hydroxysteroid (17-beta) dehydrogenase 1
NM_174555	IGFBP2	Insulin-like growth factor binding protein 2, 36 kDa
NM_174094	INHHA	Inhibin, alpha
NM_174363	INHBA	Inhibin, beta A (activin A, activin AB alpha polypeptide)
NM_173950	PIGF	Placental growth factor
NM_174646	PLA2G1B	Phospholipase A2, group IB (pancreas)
NM_174753	PTH1H	Parathyroid hormone-like hormone
NM_173957	RGN	Regucalcin (senescence marker protein-30)
NM_173959	SCD	Stearoyl-CoA desaturase (delta-9-desaturase)
NM_174670	SERPINF2	Serpin peptidase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 2
NM_001098036	SLC39A14	Solute carrier family 39 (zinc transporter), member 14
NM_001025326	SRGN	Serglycin
NM_001076470	TMEM20	Transmembrane protein 20
NM_001007813	TNFAIP6	Tumor necrosis factor, alpha-induced protein 6
NM_178317	TRB2	TRB-2 protein
AW315959		13940 MARC 4BOV Bos taurus cDNA 5', mRNA sequence.
AW325368		16365 MARC 4BOV Bos taurus cDNA 5', mRNA sequence.
BE684800		186519 MARC 4BOV Bos taurus cDNA 5', mRNA sequence.
B1536463		393463 MARC 4BOV Bos taurus cDNA 5', mRNA sequence.
B1536468		393469 MARC 4BOV Bos taurus cDNA 5', mRNA sequence.
B1537443		397313 MARC 4BOV Bos taurus cDNA 5', mRNA sequence.
BP102158		Transcribed locus
BP103904		BP103904 ORCS bovine liver cDNA Bos taurus cDNA clone ORCS25139 3', mRNA sequence.
BP104736		BP104736 ORCS bovine liver cDNA Bos taurus cDNA clone ORCS26135 3', mRNA sequence.
BP105513		BP105513 ORCS bovine liver cDNA Bos taurus cDNA clone ORCS27141 3', mRNA sequence.
BP107839		BP107839 ORCS bovine utero-placenta cDNA Bos taurus cDNA clone ORCS11248 3', mRNA sequence.
BP108716		Isolate UoG-BovSAGE-UK2 unknown mRNA
BP110155		Testis derived transcript (3 LIM domains)
BP110180		Transcribed locus
BP110819		BP110819 ORCS bovine utero-placenta cDNA Bos taurus cDNA clone ORCS11012 5', mRNA sequence.
BP111150		BP111150 ORCS bovine utero-placenta cDNA Bos taurus cDNA clone ORCS11443 5', mRNA sequence.
XM_614289		Similar to glucocorticoid induced transcript 1 (GLCCI1), mRNA.
XM_864694		Similar to tolloid-like 2, transcript variant 2 (TLL2), mRNA.

are important for bioavailability of free IGF within the follicle and play a crucial role for determining follicular dominance and fate [7,30]. Therefore, we evaluated the F1 were selected DF and the F2 were unselected SF.

Our evaluation of follicular status was confirmed to investigate *CYP19* and *IGFBP5* mRNA localization in healthy and atretic follicles in experiment 2 using *in situ*

hybridization. *CYP19* mRNA was abundantly expressed in healthy follicles but it was also expressed in atretic follicles while *IGFBP5* mRNA was detected only in atretic follicles. Both *CYP19* and *IGFBP5* mRNA is hormonally regulated in bovine follicular cells [31-34], in addition, *in situ* hybridization is not quantitative and not be as sensitive as QPCR. Thus, small amounts of




CYP19 and *IGFBP5* mRNA may be detected or regulated in atretic and healthy follicles, respectively. Since we did not perform sample collection at a specific phase of follicular wave in experiment 1, detailed growth profiles of the follicles we used were unclear. However, our results demonstrate that randomly collected follicles can be divided into several groups by similarities of gene expression profiles among the follicles and suggest that gene expression profiles of examined follicles are closely associated with their development status.

Confirmation of microarray data by QPCR analysis successfully identified a set of genes differentially expressed between the F1 and F2. In addition, possible involvement of these genes in follicular development and/or atresia was further demonstrated to investigate mRNA localization in healthy and atretic follicles. The F1 showed greater expression of genes involved in follicular growth and survivability (*AMH*, *PLA2G1B*, *SCD2* and *TRB2*) than the F2. High expression of these genes may be closely associated with the establishment and maintenance of follicular dominance. Although the functional role of AMH in antral follicle development is poorly understood, recent studies showed that both intrafollicular AMH concentration and *AMH* mRNA expression were highest in small antral follicles and then

decreased with follicular growth, suggesting the involvement of AMH in bovine follicular recruitment and/or selection [16,35,36]. A recent study showed a significant decrease of *AMH* mRNA expression in late atretic follicles compared with healthy follicles [36], which is consistent with our present result. Furthermore, in agreement with previous studies [16,37], our *in situ* hybridization study showed that *AMH* mRNA was localized in only GC. High expression and clear localization of *AMH* mRNA in the GC of healthy large follicles implies that this growth factor has a plausible effect on the development of DF after follicular selection as well as recruitment.

PLA2 enzymes including PLA2G1B hydrolyze fatty acids from the sn-2 position of phospholipids with concomitant formation of lysophospholipids, which serve as precursor for lipid mediators such as lysophosphatidic acid (LPA) [38,39]. Released LPA has diverse biological activities including cell proliferation and differentiation, suppression of apoptosis and cytoskeleton modulation in reproductive tissues [39]. Because Diouf *et al.* reported that *PLA2G1B* mRNA expression in the GC of bovine preovulatory follicle decreased after hCG injection [40], PLA2G1B may mainly contribute to generation of LPA during DF growth before the LH surge.



SCD is a rate-limiting enzyme that catalyzes the synthesis of monounsaturated fatty acids, mainly palmitic and oleic acid [41]. Consistent with our result, *SCD* mRNA expression in bovine follicles was found to be highest in GC of DF than in cohort follicles before selection or SF [11,17]. Expression of *SCD2* is hormonally regulated during follicular development because both *SCD2* mRNA and protein expression in rat large follicles were stimulated by gonadotropin and IGF-I treatment [42]. Increase of monounsaturated acids synthesized by *SCD2* activation during DF growth may be required to maintain membrane fluidity [43] and a major lipid reserve of oocytes [44].

Members of the TRB family including TRB2 interact and modulate the activity of mitogen-activated protein kinase (MAPK) which regulates cell proliferation,

differentiation, apoptosis and survival [45]. These MAPK cascade protein levels were greater in DF than in SF [46]. In addition, it has been reported that *TRB2* mRNA was constantly expressed between bovine small follicles and DF [11]. These studies and our present result suggest the potential role of TRB2 in the regulation of MAPK cascades in the growing DF.


The F2 are characterized by high expression of the genes involved in immune reaction (*CCL2*, *SELP* and *SPP1*). In bovine follicles, expression of *CCL2* and *SPP1* mRNAs and SELP protein was up-regulated in association with follicular development and ovulation [16,18,47]. Our results raise the possibility that these immune-related genes may be involved in bovine follicular atresia as well as follicular development and ovulation. Both *CCL2* and *SELP* mediate induction of leukocyte emigration into extravascular inflammatory sites [48]. Although *SPP1*, also known as osteopontin, has diverse physiological functions, one of its potent actions is recruitment and retention of macrophages and T cells to inflamed sites [49]. Since number of leukocytes, lymphocytes and activated macrophages are increased in atretic follicles [50], *CCL2*, *SELP* and *SPP1* participate in the regulation of inflammatory processes during follicular atresia to attract white blood cells.

Table 4 Follicular fluid concentrations of estradiol (E₂) and progesterone (P₄) in examined follicles used in experiment 2.

Follicle	E ₂ (ng/ml)	P ₄ (ng/ml)	E ₂ /P ₄ ratio
Healthy	180.0 ± 44.9	15.9 ± 15.3	4.3 ± 0.8
Atretic	41.4 ± 5.3*	387.7 ± 121.7*	0.1 ± 0.1*

Values are mean ± SEM


* $P < 0.05$ vs. healthy follicle



The F2 are also characterized by high expression of genes regulating tissue remodeling (*TIMP1* and *PLAUR*). Both plasminogen activator (PA)-plasmin and matrix metalloproteinase (MMP) systems play a crucial role in the degradation and remodeling of extracellular matrix associated with follicular development, ovulation and atresia [51]. Urokinase PA (uPA) receptor is a specific cell surface receptor for uPA and its principal role is to localize pericellular plasmin activity to induce extracellular matrix degradation [52]. A previous study showed that there was no difference in mRNA expression levels of uPA between non-atretic and atretic bovine follicles, whereas atretic follicles had lower FF protein level and mRNA expression of a PA inhibitor and higher FF plasmin activity than non-atretic follicles [53]. Therefore, the follicular PA-plasmin system may be primarily regulated by changes in their receptors and inhibitors' expressions. *TIMP-1* is an intrinsic inhibitor of MMPs and preferentially binds to MMP-9 [54]. A previous study demonstrated that MMP-9 proenzyme (proMMP-9) protein in FF was detected only in atretic follicles but not in healthy follicles in cattle [55]. Atretic

follicles may balance MMPs and *TIMP-1* in response to an increase in proMMP-9 to control extracellular matrix degradation by MMP-9.

In the present study, we identified differential expression of two anti-apoptosis factors (*GADD45A* and *GPX3*) between the groups. Expression of *GADD45A* mRNA was greater in the F2 than in the F1. *GADD45A* controls cell cycle arrest, apoptosis induction and DNA damage repair in response to DNA damaging agents and growth arrest signals of genotoxic stress [56]. Our result implies that the atretic follicles suffered more severe DNA damage than healthy follicles. Indeed, we found in experiment 2 that the atretic follicles expressed *GADD45A* mRNA in both GC and TL whereas the healthy follicles expressed it only in GC. This result supports our microarray result and suggests an increase in the requirement of *GADD45A* activity for progression of apoptotic cell death in GC and TC during follicular atresia. On the other hand, *GPX3* mRNA was found to be more greatly expressed in the F1 than in the F2. Glutathione peroxidase protects cells against oxidative damage to catalyze the reduction of free hydrogen



peroxide and other hydroperoxides [57]. High oxidative stress can trigger apoptosis of follicular cells and induce atresia [58]. In cultured swine GC, *GPX3* mRNA expression was upregulated by FSH treatment [59]. Thus GPX3 could prevent cell apoptosis from oxidative stress during growth of the healthy follicles. It is likely that follicular oxidative stress-response enzymes are expressed in a stage-dependent manner since mRNA expression of other anti-oxidative stress enzymes in bovine GC was increased in atretic DF than in healthy DF [60].

Providing a sufficient blood supply is essential for follicular growth [61,62]. A morphological study has demonstrated that bovine healthy DF has a high density and well developed capillaries in TL whereas atretic follicles has sparse and poorly developed capillaries [63]. In the present study, two genes regulating angiogenesis, *PIGF* and *TSP2*, were differentially expressed between the groups. *PIGF* was expressed most in F1 than in F2 and localized in both GC and TL of healthy follicles but not detected in atretic follicles. *PIGF* is a member of the

vascular endothelial growth factor family and stimulates the proliferation of endothelial cells and supports angiogenesis [64,65]. Therefore, *PIGF* may contribute to follicular thecal angiogenesis via paracrine/autocrine action in healthy follicles as well as other angiogenic factors. In contrast to *PIGF*, *TSP-2*, a member of the TSP family, acts as a potent inhibitor of angiogenesis and induces endothelial cell apoptosis [66]. In experiment 1, *TSP2* mRNA expression was greater in the F2 than in the F1. *TSP2* mRNA level in the bovine follicles decreased in accordance with an increase in follicular diameter [67]. The same authors also showed that TSP protein was localized in both GC and TC of small follicles but in only in the GC of large follicles [67]. We demonstrated in experiment 2 that *TSP2* mRNA was localized in both GC and TL of atretic follicles while it was expressed in only GC of healthy follicles. Recent studies have demonstrated that mRNA and protein expression of *TSP-1*, another antiangiogenic TSP, is upregulated in primate GC during progression of follicular atresia [68] and

TSP1 mRNA abundance is decreased by IGF-I treatment in cultured porcine GC [69]. Thus, we speculate that *TSP2* mRNA expression is maintained at high levels in follicular cells of atretic follicles whereas it decreases in healthy follicles. Highly expressed *TSP2* mRNA in the follicles could negatively influence their angiogenesis. It may cause an insufficient supply of substrates essential for follicular growth, thereby affecting follicular hormone production and cell proliferation, and, as a result, inducing atresia.

Conclusion

Microarray and QPCR analysis enabled us to classify uncharacterized bovine follicles and to evaluate their representative follicular status according to differences in global gene expression profiles. Our present study demonstrates that the expression of stage-specific genes in F1 and F2 may be closely associated with follicular growth and atresia. Several genes identified in this study will provide information on the genomic actions of intriguing candidates for the determinant of bovine follicular development.

Acknowledgements

This study was supported by grant-in-aid from the Ministry of Agriculture, Forestry and Fisheries Japan (17-1770). KGH was supported by the Japan Society for the Promotion of Science Research Fellowships for Young Scientists (0702379).

Authors' contributions

KGH participated in the design of the study, collected the materials, carried out all experiments and drafted the manuscript. KU collected the materials, carried out the microarray experiments and analysis, and helped to carry out QPCR and *in situ* hybridization. MH was responsible for all animal care, collected the materials and carried out the microarray experiments. TT supervised the study, collected the materials and helped to draft the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 30 November 2009

Accepted: 5 February 2010 Published: 5 February 2010

References

1. Sirois J, Fortune JE: Ovarian follicular dynamics during the estrous cycle in heifers monitored by real-time ultrasonography. *Biol Reprod* 1988, **39**:308-317.
2. Ginther OJ, Knopf L, Kastelic JP: Temporal associations among ovarian events in cattle during oestrous cycles with two and three follicular waves. *J Reprod Fertil* 1989, **87**:223-230.
3. Sunderland SJ, Crowe MA, Boland MP, Roche JF, Ireland JJ: Selection, dominance and atresia of follicles during the oestrous cycle of heifers. *J Reprod Fertil* 1994, **101**:547-555.
4. Ginther OJ, Kot K, Kulick LJ, Wiltbank MC: Emergence and deviation of follicles during the development of follicular waves in cattle. *Theriogenology* 1997, **48**:75-87.
5. Ginther OJ, Wiltbank MC, Fricke PM, Gibbons JR, Kot K: Selection of the dominant follicle in cattle. *Biol Reprod* 1996, **55**:1187-1194.
6. Mihm M, Crowe MA, Knight PG, Austin EJ: Follicle wave growth in cattle. *Reprod Domest Anim* 2002, **37**:191-200.
7. Fortune JE, Rivera GM, Yang MY: Follicular development: the role of the follicular microenvironment in selection of the dominant follicle. *Anim Reprod Sci* 2004, **82-83**:109-126.
8. Sisco B, Hagemann LJ, Shelling AN, Pfeffer PL: Isolation of genes differentially expressed in dominant and subordinate bovine follicles. *Endocrinology* 2003, **144**:3904-3913.
9. Evans AC, Ireland JL, Winn ME, Lonergan P, Smith GW, Coussens PM, Ireland JJ: Identification of genes involved in apoptosis and dominant follicle development during follicular waves in cattle. *Biol Reprod* 2004, **70**:1475-1484.
10. Fayad T, Levesque V, Sirois J, Silversides DW, Lussier JG: Gene expression profiling of differentially expressed genes in granulosa cells of bovine dominant follicles using suppression subtractive hybridization. *Biol Reprod* 2004, **70**:523-533.
11. Ndiaye K, Fayad T, Silversides DW, Sirois J, Lussier JG: Identification of downregulated messenger RNAs in bovine granulosa cells of dominant follicles following stimulation with human chorionic gonadotropin. *Biol Reprod* 2005, **73**:324-333.
12. Mihm M, Baker PJ, Ireland JL, Smith GW, Coussens PM, Evans AC, Ireland JJ: Molecular evidence that growth of dominant follicles involves a reduction in follicle-stimulating hormone dependence and an increase in luteinizing hormone dependence in cattle. *Biol Reprod* 2006, **74**:1051-1059.
13. Zielak AE, Forde N, Park SD, Doohan F, Coussens PM, Smith GW, Ireland JJ, Lonergan P, Evans AC: Identification of novel genes associated with dominant follicle development in cattle. *Reprod Fertil Dev* 2007, **19**:967-975.
14. Zielak AE, Cauty MJ, Forde N, Coussens PM, Smith GW, Lonergan P, Ireland JJ, Evans AC: Differential expression of genes for transcription factors in theca and granulosa cells following selection of a dominant follicle in cattle. *Mol Reprod Dev* 2008, **75**:904-914.
15. Baker PJ, Fleming LM, Mossa F, Lonergan P, Evans AC, Mihm M: Decreased mRNA expression of dominance maker genes in preovulatory compared to newly selected dominant follicles in cattle [abstract]. *Biol Reprod* 2008, **531**.
16. Skinner MK, Schmidt M, Savenkova MI, Sadler-Riggleman I, Nilsson EE: Regulation of granulosa and theca cell transcriptomes during ovarian antral follicle development. *Mol Reprod Dev* 2008, **75**:1457-1472.
17. Mihm M, Baker PJ, Fleming LM, Monteiro AM, O'Shaughnessy PJ: Differentiation of the bovine dominant follicle from the cohort upregulates mRNA expression for new tissue development genes. *Reproduction* 2008, **135**:253-265.
18. Liu Z, Youngquist RS, Garverick HA, Antoniou E: Molecular mechanisms regulating bovine ovarian follicular selection. *Mol Reprod Dev* 2009, **76**:351-366.
19. TM4. <http://www.tm4.org/mev.html>.
20. MIAME. <http://www.mged.org/Workgroups/MIAME/miame.html>.
21. GEO. <http://www.ncbi.nlm.nih.gov/projects/geo/>.
22. Ushizawa K, Takahashi T, Hosoe M, Ishiwata H, Kaneyama K, Kizaki K, Hashizume K: Global gene expression analysis and regulation of the principal genes expressed in bovine placenta in relation to the transcription factor AP-2 family. *Reprod Biol Endocrinol* 2007, **5**:17.
23. Takahashi T, Hamanaka S, Ikeda S, Kobayashi J, Hashizume K: A direct time-resolved fluorescent immunoassay (TR-FIA) for measuring plasma progesterone concentration in sika doe (*Cervus nippon centralis*). *J Reprod Dev* 2001, **47**:119-123.
24. Takahashi T, Hamanaka S, Imai K, Hashizume K: A direct time-resolved fluoroimmunoassay (TR-FIA) for measuring plasma estradiol-17beta concentrations in cattle. *J Vet Med Sci* 2004, **66**:225-229.
25. Bao B, Garverick HA, Smith GW, Smith MF, Salfen BE, Youngquist RS: Changes in messenger ribonucleic acid encoding luteinizing hormone receptor, cytochrome P450-side chain cleavage, and aromatase are associated with recruitment and selection of bovine ovarian follicles. *Biol Reprod* 1997, **56**:1158-1168.
26. Santiago CA, Voge JL, Aad PY, Allen DT, Stein DR, Malayer JR, Spicer LJ: Pregnancy-associated plasma protein-A and insulin-like growth factor binding protein mRNAs in granulosa cells of dominant and subordinate follicles of preovulatory cattle. *Domest Anim Endocrinol* 2005, **28**:46-63.
27. Ushizawa K, Kaneyama K, Takahashi T, Tokunaga T, Tsunoda Y, Hashizume K: Cloning and expression of a new member of prolactin-related protein in

- bovine placenta: bovine prolactin-related protein-VII. *Biochem Biophys Res Commun* 2005, **326**:435-441.
28. Ushizawa K, Takahashi T, Kaneyama K, Hosoe M, Hashizume K: Cloning of the bovine antiapoptotic regulator, BCL2-related protein A1, and its expression in trophoblastic binucleate cells of bovine placenta. *Biol Reprod* 2006, **74**:344-351.
 29. Bao B, Garverick HA: Expression of steroidogenic enzyme and gonadotropin receptor genes in bovine follicles during ovarian follicular waves: a review. *J Anim Sci* 1998, **76**:1903-1921.
 30. Canty MJ, Boland MP, Evans AC, Crowe MA: Alterations in follicular IGFBP mRNA expression and follicular fluid IGFBP concentrations during the first follicle wave in beef heifers. *Anim Reprod Sci* 2006, **93**:199-217.
 31. Silva JM, Price CA: Effect of follicle-stimulating hormone on steroid secretion and messenger ribonucleic acids encoding cytochromes P450 aromatase and cholesterol side-chain cleavage in bovine granulosa cells in vitro. *Biol Reprod* 2000, **62**:186-191.
 32. Silva JM, Price CA: Insulin and IGF-I are necessary for FSH-induced cytochrome P450 aromatase but not cytochrome P450 side-chain cleavage gene expression in oestrogenic bovine granulosa cells in vitro. *J Endocrinol* 2002, **174**:499-507.
 33. Voge JL, Aad PY, Santiago CA, Goad DW, Malayer JR, Allen D, Spicer LJ: Effect of insulin-like growth factors (IGF), FSH, and leptin on IGF-binding-protein mRNA expression in bovine granulosa and theca cells: quantitative detection by real-time PCR. *Peptides* 2004, **25**:2195-2203.
 34. Voge JL, Santiago CA, Aad PY, Goad DW, Malayer JR, Spicer LJ: Quantification of insulin-like growth factor binding protein mRNA using real-time PCR in bovine granulosa and theca cells: effect of estradiol, insulin, and gonadotropins. *Domest Anim Endocrinol* 2004, **26**:241-258.
 35. Monniaux D, Clemente N, Touze JL, Belleville C, Rico C, Bontoux M, Picard JY, Fabre S: Intrafollicular steroids and anti-mullerian hormone during normal and cystic ovarian follicular development in the cow. *Biol Reprod* 2008, **79**:387-396.
 36. Rico C, Fabre S, Medigue C, Clemente N, Clement F, Bontoux M, Touze JL, Dupont M, Briant E, Remy B, Beckers JF, Monniaux D: Anti-mullerian hormone is an endocrine marker of ovarian gonadotropin-responsive follicles and can help to predict superovulatory responses in the cow. *Biol Reprod* 2009, **80**:50-59.
 37. Takahashi M, Hayashi M, Manganaro TF, Donahoe PK: The ontogeny of mullerian inhibiting substance in granulosa cells of the bovine ovarian follicle. *Biol Reprod* 1986, **35**:447-453.
 38. Eder AM, Sasagawa T, Mao M, Aoki J, Mills GB: Constitutive and lysophosphatidic acid (LPA)-induced LPA production: role of phospholipase D and phospholipase A2. *Clin Cancer Res* 2000, **6**:2482-2491.
 39. Budnik LT, Mukhopadhyay AK: Lysophosphatidic acid and its role in reproduction. *Biol Reprod* 2002, **66**:859-865.
 40. Diouf MN, Sayasith K, Lefebvre R, Silversides DW, Sirois J, Lussier JG: Expression of phospholipase A2 group IVA (PLA2G4A) is upregulated by human chorionic gonadotropin in bovine granulosa cells of ovulatory follicles. *Biol Reprod* 2006, **74**:1096-1103.
 41. Nakamura MT, Nara TY: Structure, function, and dietary regulation of delta6, delta5, and delta9 desaturases. *Annu Rev Nutr* 2004, **24**:345-376.
 42. Moreau C, Froment P, Tosca L, Moreau V, Dupont J: Expression and regulation of the SCD2 desaturase in the rat ovary. *Biol Reprod* 2006, **74**:75-87.
 43. Abel S, Smuts CM, de Villiers C, Gelderblom WC: Changes in essential fatty acid patterns associated with normal liver regeneration and the progression of hepatocyte nodules in rat hepatocarcinogenesis. *Carcinogenesis* 2001, **22**:795-804.
 44. McEvoy TG, Coull GD, Broadbent PJ, Hutchinson JS, Speake BK: Fatty acid composition of lipids in immature cattle, pig and sheep oocytes with intact zona pellucida. *J Reprod Fertil* 2000, **118**:163-170.
 45. Kiss-Toth E, Bagstaff SM, Sung HY, Jozsa V, Dempsey C, Caunt JC, Oxley KM, Wyllie DH, Polgar T, Harte M, O'Neill LA, Qvarnstrom EE, Dower SK: Human tribbles, a protein family controlling mitogen-activated protein kinase cascades. *J Biol Chem* 2004, **279**:42703-42708.
 46. Ryan KE, Casey SM, Canty MJ, Crowe MA, Martin F, Evans AC: Akt and Erk signal transduction pathways are early markers of differentiation in dominant and subordinate ovarian follicles in cattle. *Reproduction* 2007, **133**:617-626.
 47. Rohm F, Spanel-Borowski K, Eichler W, Aust G: Correlation between expression of selectins and migration of eosinophils into the bovine ovary during the periovulatory period. *Cell Tissue Res* 2002, **309**:313-322.
 48. Vestweber D, Blanks JE: Mechanisms that regulate the function of the selectins and their ligands. *Physiol Rev* 1999, **79**:181-213.
 49. Mazzali M, Kipari T, Ophascharoensuk V, Wesson JA, Johnson R, Hughes J: Osteopontin—a molecule for all seasons. *Qjm* 2002, **95**:3-13.
 50. Suzuki T, Sasano H, Takaya R, Fukaya T, Yajima A, Date F, Nagura H: Leukocytes in normal-cycling human ovaries: immunohistochemical distribution and characterization. *Hum Reprod* 1998, **13**:2186-2191.
 51. Ny T, Wahlberg P, Brandstrom IJ: Matrix remodeling in the ovary: regulation and functional role of the plasminogen activator and matrix metalloproteinase systems. *Mol Cell Endocrinol* 2002, **187**:29-38.
 52. Blasi F, Carmeliet P: uPAR: a versatile signalling orchestrator. *Nat Rev Mol Cell Biol* 2002, **3**:932-943.
 53. Cao M, Buratini J Jr, Lussier JG, Carriere PD, Price CA: Expression of protease nexin-1 and plasminogen activators during follicular growth and the periovulatory period in cattle. *Reproduction* 2006, **131**:125-137.
 54. Goldberg GI, Strongin A, Collier IE, Genrich LT, Marmer BL: Interaction of 92-kDa type IV collagenase with the tissue inhibitor of metalloproteinases prevents dimerization, complex formation with interstitial collagenase, and activation of the proenzyme with stromelysin. *J Biol Chem* 1992, **267**:4583-4591.
 55. Khandoker MA, Imai K, Takahashi T, Hashizume K: Role of gelatinase on follicular atresia in the bovine ovary. *Biol Reprod* 2001, **65**:726-732.
 56. Zhan Q: Gadd45a a p53- and BRCA1-regulated stress protein, in cellular response to DNA damage. *Mutat Res* 2005, **569**:133-143.
 57. Yu BP: Cellular defenses against damage from reactive oxygen species. *Physiol Rev* 1994, **74**:139-162.
 58. Tilly JL, Tilly KI: Inhibitors of oxidative stress mimic the ability of follicle-stimulating hormone to suppress apoptosis in cultured rat ovarian follicles. *Endocrinology* 1995, **136**:242-252.
 59. Bonnet A, Frappart PO, Dehais P, Tosser-Klopp G, Hatey F: Identification of differential gene expression in in vitro FSH treated pig granulosa cells using suppression subtractive hybridization. *Reprod Biol Endocrinol* 2006, **4**:35.
 60. Valdez KE, Cuneo SP, Turzillo AM: Regulation of apoptosis in the atresia of dominant bovine follicles of the first follicular wave following ovulation. *Reproduction* 2005, **130**:71-81.
 61. Zeleznik AJ, Schuler HM, Reichert LE Jr: Gonadotropin-binding sites in the rhesus monkey ovary: role of the vasculature in the selective distribution of human chorionic gonadotropin to the preovulatory follicle. *Endocrinology* 1981, **109**:356-362.
 62. Acosta TJ, Hayashi KG, Matsui M, Miyamoto A: Changes in follicular vascularity during the first follicular wave in lactating cows. *J Reprod Dev* 2005, **51**:273-280.
 63. Jiang JY, Macchiarelli G, Tsang BK, Sato E: Capillary angiogenesis and degeneration in bovine ovarian antral follicles. *Reproduction* 2003, **125**:211-223.
 64. Maglione D, Guerriero V, Viglietto G, Delli-Bovi P, Persico MG: Isolation of a human placenta cDNA coding for a protein related to the vascular permeability factor. *Proc Natl Acad Sci USA* 1991, **88**:9267-9271.
 65. Carmeliet P, Moons L, Lutun A, Vincenti V, Compernelle V, De Mol M, Wu Y, Bono F, Devy L, Beck H, Scholz D, Acker T, DiPalma T, Dewerchin M, Noel A, Stalmans I, Barra A, Blacher S, Vandendriessche T, Ponten A, Eriksson U, Plate KH, Foidart JM, Schaper W, Charnock-Jones DS, Hicklin DJ, Herbert JM, Collen D, Persico MG: Synergism between vascular endothelial growth factor and placental growth factor contributes to angiogenesis and plasma extravasation in pathological conditions. *Nat Med* 2001, **7**:575-583.
 66. Streit M, Riccardi L, Velasco P, Brown LF, Hawighorst T, Bornstein P, Detmar M: Thrombospondin-2: a potent endogenous inhibitor of tumor growth and angiogenesis. *Proc Natl Acad Sci USA* 1999, **96**:14888-14893.
 67. Greenaway J, Gentry PA, Feige JJ, LaMarre J, Petrik JJ: Thrombospondin and vascular endothelial growth factor are cyclically expressed in an inverse pattern during bovine ovarian follicle development. *Biol Reprod* 2005, **72**:1071-1078.
 68. Thomas FH, Wilson H, Silvestri A, Fraser HM: Thrombospondin-1 expression is increased during follicular atresia in the primate ovary. *Endocrinology* 2008, **149**:185-192.

69. Grado-Ahuir JA, Aad PY, Ranzenigo G, Caloni F, Cremonesi F, Spicer LJ: Microarray analysis of insulin-like growth factor-I-induced changes in messenger ribonucleic acid expression in cultured porcine granulosa cells: possible role of insulin-like growth factor-I in angiogenesis. *J Anim Sci* 2009, **87**:1921-1933.

doi:10.1186/1477-7827-8-11

Cite this article as: Hayashi *et al.*: Differential genome-wide gene expression profiling of bovine largest and second-largest follicles: identification of genes associated with growth of dominant follicles. *Reproductive Biology and Endocrinology* 2010 **8**:11.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

