

Structural Determinants Involved in the Formation and Activation of G Protein $\beta\gamma$ Dimers

William E. McIntire

Department of Pharmacology, University of Virginia Health System, Charlottesville, Va., USA

Key Words

G protein • Receptor • Activation • $\beta\gamma$ Dimer • Signal transduction

Abstract

Heterotrimeric G proteins, composed of an α , β and γ subunit, represent one of the most important and dynamic families of signaling proteins. As a testament to the significance of G protein signaling, the hundreds of seven-transmembrane-spanning receptors that interact with G proteins are estimated to occupy 1–2% of the human genome. This broad diversity of receptors is echoed in the number of potential heterotrimer combinations that can arise from the 23 α subunit, 7 β subunit and 12 γ subunit isoforms that have been identified. The potential for such vast complexity implies that the receptor G protein interface is the site of much regulation. The historical model for the activation of a G protein holds that activated receptor catalyzes the exchange of GDP for GTP on the α subunit, inducing a conformational change that substantially lowers the affinity of α for $\beta\gamma$. This decreased affinity enables dissociation of $\beta\gamma$ from α and receptor. The free form of $\beta\gamma$ is thought to activate effectors, until the hydrolysis of GTP by G α (aided by RGS proteins) allows the subunits to re-associate, effectively deactivating the G protein until another interaction with activated receptor.

Copyright © 2009 S. Karger AG, Basel

Introduction

Although there are many facets of G protein signaling via the $\beta\gamma$ dimer, this review will primarily focus on how the structure of a $\beta\gamma$ dimer participates in the transfer of signal from activated receptor to the heterotrimeric G protein. Structural heterogeneity of $\beta\gamma$ combinations will be discussed regarding how specificity of interactions of $\beta\gamma$ with G α and receptor determine which $\beta\gamma$ dimer isoforms are activated. A discussion of specific interactions of $\beta\gamma$ dimers with effector molecules is beyond the scope of this review, but is addressed elsewhere in this issue. However, some references to interactions with effectors are included, as the activation of effectors is by definition one of the primary measures of activity of a G protein $\beta\gamma$ dimer. Beyond the brief descriptions of the G protein activation cycle that have appeared in thousands of papers over the years, it is clear that $\beta\gamma$ dimer ‘activity’ may not simply be synonymous with dissociation from α . Thus, an important question is whether dissociation of α from $\beta\gamma$ is a requisite step in the activation of $\beta\gamma$. Also, is a $\beta\gamma$ active if it does not properly localize with its effector molecule? Does a $\beta\gamma$ need a receptor, or even an α subunit, to be activated? Does a $\beta\gamma$ always exist as dimer? The record in the literature may be limited for some of these later questions, but they represent intriguing ideas that will help refine the model of G protein signaling.

G Protein $\beta\gamma$ Heterogeneity

At least 16 α genes, 5 β genes and 12 γ genes have been identified in the human genome [1–6]. G α isoforms can be separated into four subfamilies, G_i , G_s , G_q and G_{12} ; when alternative splicing and posttranslational processing are taken into account, there are at least 23 α isoforms, which are reviewed elsewhere [5]. The first four β isoforms discovered, β_{1-4} , are highly homologous (80–90% identical) 36-kDa proteins; G β_5 , a 40-kDa protein, is only 50% identical to the first four β isoforms. Several truncated splice variants of β_3 have been characterized, β_{3s} [7], β_{3s2} [8] and β_{3v} [9]; a splice variant of β_5 which has an N-terminal extension, β_{5L} [10], has also been characterized. All 12 γ isoforms are between 7 and 8.5 kDa in size, but are much more divergent than the β isoforms. Since β and γ are believed to form a functional dimer *in vivo*, heterogeneity of $\beta\gamma$ defined as the product of the β and γ genes is likely more diverse than G α [11] even though not every possible $\beta\gamma$ combination can form. Posttranslational processing of β and γ further contribute to the structural diversity of these proteins [12], with γ isoforms receiving more extensive study than β isoforms to date.

Posttranslational Modifications

β Subunit

Several posttranslational modifications of G β have been characterized. The N-terminus of β_1 was reported to undergo removal of the methionine at position 1, followed by N-acetylation of serine at position 2 [13], the functional implications of which were unclear. Phosphorylation of β has also been reported, but interestingly, at a histidine residue [14] instead of the traditional serine, threonine or tyrosine; nucleoside diphosphate kinase (NDPK) was identified as responsible for phosphorylating histidine 266 of β_1 [15]. This phosphorylation event was predicted as a mechanism for G protein activation, which will be discussed further below. In contrast, the reversible mono-ADP-ribosylation of arginine-129 of activated or ‘free’ G β was demonstrated to reduce activity at effectors such as type 1 adenylyl cyclase [16], phosphoinositide 3-kinase- γ and phospholipase C- β_2 [17]. Mono-ADP ribosylation of β increased upon activation of a variety of cell surface receptors, including the G_q -linked thrombin receptor and the G_i -linked 5-HT serotonin receptor, and thus was predicted to be a regulatory mechanism to inhibit $\beta\gamma$ signaling. Inhibition of $\beta\gamma$ ac-

tivity by posttranslational modification has parallels to the GTPase activity of the G α subunit, and may represent an unappreciated regulatory mechanism in G protein signaling.

γ Subunit

C-Terminal Processing

The G γ subunit contains posttranslational modifications at the N- and C-terminus, and much of this processing has been demonstrated to be critical to G protein function. Most attention has focused on covalent modification at the C-terminus of γ with either of two distinct isoprenoid moieties [for review, 18]. For γ isoforms ending in amino acid CAAX, where X is serine, glutamine or methionine, CVIS in the case of γ_1 , and additionally γ_8 , and γ_{11} , the enzyme farnesyltransferase (FTase) covalently attaches a 15-carbon farnesyl group via a thioether bond to the cysteine in the CAAX motif [19]. If the X in the CAAX sequence is a leucine such as CAIL for γ_2 , or the C-terminal sequences in γ_3 , γ_4 , γ_5 , γ_7 , γ_9 , γ_{10} , γ_{12} and γ_{13} , the enzyme geranylgeranyltransferase type I (GG-Tase-I) attaches a larger 20-carbon geranylgeranyl group to the cysteine [20, 21] via the same thioether bond. After modification with either prenyl moiety, processing is similar for all γ isoforms. An endoprotease residing in the microsomal membranes cleaves the C-terminal-AAX residues [22], and the new prenylated C-terminal cysteine is carboxy methylated by a methyltransferase [23]. Both assembly of $\beta\gamma$ dimers, which occurs in the cytosol [24], and prenylation of γ are required to target $\beta\gamma$ to membranes [25]; there is some debate over which membranes, as $\beta\gamma$ has been observed in endoplasmic reticulum membranes from biochemical and immunofluorescence studies of $\beta\gamma$ expressed in mammalian cells [26]. Alternatively, live cell imaging demonstrated localization of $\beta\gamma$ predominantly to the plasma membrane [27]. Although posttranslational processing of G γ as generalized above is well documented and well accepted, it is also evident that processing exceptions may provide insights into how G $\beta\gamma$ functions *in vivo*.

For example, it is believed that the cleavage of the C-terminal-AAX amino acids is a necessary step in protein maturation prior to $\beta\gamma$ assembly [28]. However, this is not always the case; a geranylgeranylated γ_5 isoform that had not undergone cleavage of the C-terminal-AAX amino acids was characterized by mass spectrometry from a purified preparation of G protein from bovine brain [29]. This unprocessed form was discovered to predominate over the cleaved form, and be dependent on an aromatic phenylalanine residue in the CSFL C-terminal sequence

of γ_5 [30]. A recent study noting a physical interaction between proteins containing PDZ domains, which are important in the construction of elaborate scaffolding networks, and $G\gamma_{13}$ [31] made the story of this processing pattern more interesting. $G\gamma_5$, with its C-terminus ending in CSFL, is one of only four γ isoforms other than γ_{13} with a C-terminal target sequence (CT/SXX) for class I PDZ domain containing proteins. The fact that the C-terminal sequence of γ_5 can remain after maturation of a $\beta\gamma$ dimer suggests that γ_5 isoforms that do not undergo C-terminal proteolytic cleavage are differentially targeted compared to other $\beta\gamma$ isoforms. Since this form of processing appears to be unique for γ_5 , there may be a specific signaling role for $\beta\gamma$ dimers containing γ_5 in signaling complexes containing PDZ domains.

Unprenylated γ

The absence of prenylation in $G\gamma$ is also associated with unexpected signaling properties. After the observation that a fraction of $\beta_2\gamma_2$ could localize to the nucleus and regulate transcriptional activity [32], a study by Kino et al. [33] demonstrated that lack of prenylation of γ_2 , either by mutation of the C-terminal cysteine to serine or by pharmacological inhibition, resulted in increased nuclear localization of $\beta\gamma$ and increased ability to regulate transcription. Further, non-prenylated γ_5 expressed in bacteria was shown to regulate transcription by binding to the adipocyte enhancer-binding protein (AEBP1) transcriptional repressor [34]. Although prenylation of γ is not thought to be reversible, fully processed isoforms, missing a prenyl group, have been characterized by mass spectrometry in γ_2 (<1% of γ_2 observed) and γ_7 (1–5% of γ_7 observed) from G protein purified from bovine brain [12]. These exceptions in the prenylation pathway, although apparently low in occurrence, are surprising in that conventional wisdom assumes that, although prenylation is not required for assembly [28], it is a prerequisite for $\beta\gamma$ activity. Moreover, since regulation of transcription is not a classical signaling function for $\beta\gamma$, the prenylation status of $\beta\gamma$ may represent a significant point in modulation of activity of $\beta\gamma$ dimers, with respect to identity and localization of effector targets.

N-Terminal Processing

Other regions of γ that are sites for covalent modification include the N-terminus, which were initially observed to be refractory to Edman degradation in the case of γ_2 , γ_5 and γ_7 ; alternatively, γ_1 [35], γ_3 [36] and γ_{11} [37] were not found to be N-terminally blocked. Mass spectrometry was used to identify structural modifications at

the N-terminus of the γ_2 isoform, which were revealed to be cleavage of the N-terminal methionine followed by N-acetylation of alanine formerly at position 2 [38]. Yet another variant of γ_2 was characterized by Edman degradation, in which a novel N-terminal sequence was determined to be a substrate for the N-end rule ubiquitylation pathway [39]. Modification by ubiquitin was found likely to occur at lysine residues in the C-terminal region of γ ; although ubiquitylation is generally regarded as a signal for protein degradation, the authors suggest that this modification may also modulate membrane binding of $\beta\gamma$.

Phosphorylation

Interestingly, phosphorylation has only been observed in the γ_{12} isoform, in which protein kinase C was shown to phosphorylate a serine at position one in vitro [40]. Phosphorylation appeared to increase the affinity of $\beta_1\gamma_{12}$ for Go α [40], and increase the ability of $\beta_1\gamma_{12}$ to couple to receptor [41]; however, phosphorylation diminished the activity of $\beta_1\gamma_{12}$ at adenylyl cyclase type II, but not phospholipase C- β [41]. Phorbol 12-myristate 13-acetate (PMA) induced in vivo phosphorylation of γ_{12} in cultured Swiss 3T3 cells was augmented by activation of the lysophosphatidic acid receptor, and inhibited by pertussis toxin, suggesting a role for G_i proteins in the regulation of γ_{12} by protein kinase C [42]. The increase in γ_{12} phosphorylation after receptor activation is also in agreement with the observation that $\beta\gamma$ dimer, rather than heterotrimer, is the preferred substrate for protein kinase C [40]. It is possible that two of the potential signaling effects of this modification, increased G protein cycling, and targeting of specific effectors, combine to regulate signaling pathways within a cell. These examples demonstrate that posttranslational modifications of G protein β and γ subunit isoforms, while often overlooked, can be dynamic and thus represent points of signaling modulation that could affect intracellular targeting of a $\beta\gamma$ dimer, and regulation of interactions with G α , receptor, and effector molecules.

Structure of $\beta\gamma$

G protein β subunits belong to a family of WD40 repeat proteins, characterized by a repeating motif of 27–45 amino acids punctuated at the C-terminal end by the Trp-Asp (WD) dipeptide sequence [43]. The solutions to the crystal structures of a G protein and its constituent subunits were watershed events in the G protein field, providing for the first time a three-dimensional model of the heterotrimeric signaling molecule. This review will




Fig. 1. Clam shell model of G protein opening after activation using the crystal structure of Wall et al. [45]. Left structure: Relationship of inactive GDP-bound form of $G_{i1} \alpha\beta_1\gamma_2$ to the plasma membrane, with stylized lipids added to the structure. Right structure: $\beta\gamma$ is in same conformation, but $G_{i1} \alpha$ has been rotated approximately 90° counterclockwise; note that although lipids

maintain their proximity, the face of the β propeller and switch I and II regions of $G_{i1} \alpha$, both occluded at left, are now sterically free to interact with effectors, and are thus 'active'. (Although the same structure was used for simplicity, a truly active $G_{i1} \alpha$ would have conformational changes in the switch regions, and GTP γ S bound instead of GDP.)

begin with those structures, and also discuss biochemical data that have served to complement the crystal structures of the G protein $\beta\gamma$ dimer. The structure of the $\beta_1\gamma_1$ dimer of transducin was published by Sonddek et al. [44] in 1996. At about the same time, the crystal structures of the $\beta_1\gamma_2$ dimer associated with the $G_{i1} \alpha$ subunit, and the $\beta_1\gamma_1$ dimer associated with a chimera of the $G_t \alpha$ and $G_{i1} \alpha$ subunits, were published by Wall et al. [45] and Lambright et al. [46], respectively. According to the structures, the 340-amino-acid β subunit forms a toroidal structure defined by seven propeller blades (fig. 1), with each blade comprised of a series of β -sheets. The N-terminus of the β subunit is an α -helix which interacts with the α -helical structure of the N-terminus of the approximately 70-amino-acid γ subunit to form a coiled-coil domain (fig. 1), which had been predicted from earlier biochemical studies [47]. The remainder of the γ subunit exists in an α helical structure that makes extensive contacts with the blades of the β torus (fig. 1). Crystal structures of $\beta_1\gamma_1$ and $\beta_1\gamma_2$ were highly homologous; further, a crystal structure of the more divergent β_5 complexed with an

RGS9 protein [48] revealed that β_1 and β_5 form very similar toroidal structures. This suggests that, even considering the high degree of diversity among the β and γ subunit isoforms, the overall structure of a $\beta\gamma$ dimer is highly conserved.

Although the $\beta_1\gamma_2$ [45] and $\beta_1\gamma_1$ [46] dimers had been crystallized with G α subunits, revealing important sites of interaction, data elucidating the dynamic nature of these interactions was derived from the comparison of the crystal structures of $G_t \alpha$ in the inactive GDP-bound form [49] and the active GTP γ S-bound form [50]. The general architecture of the G α subunit consists of a GTPase domain that is homologous to the monomeric GTP-binding proteins (such as ras), and a helical domain that is only found in heterotrimeric G protein α subunits. Two regions in the GTPase domain that interact with the γ -phosphate of GTP, coined switch I and switch II (fig. 1), stand out in the $G_t \alpha$ crystal structures in that they undergo a conformational change depending on the nucleotide bound. A third switch region found in the helical domain, switch III, also undergoes a conformational

change, apparently dependent upon the conformational change of switch II [49]. These conformational changes, reviewed in more detail elsewhere [51], were critical to understanding the mechanism of activation of G proteins, as the crystal structures of two heterotrimeric G proteins revealed that although no contacts were observed between α and γ , many of the contact sites between α and β reside in the switch I and switch II regions (fig. 1) [46, 45]. Thus, the logical conclusion based on the crystal structures, and also predicted from earlier biochemical experiments that reported decreased affinity between activated α and $\beta\gamma$ [52], was that nucleotide exchange in a G protein could induce conformational changes in the $\beta\gamma$ -binding site of G α , resulting in subunit dissociation and activation of both α and $\beta\gamma$.

The model of G protein activation based on the crystal structures will continue to benefit from biochemical studies for a number of reasons. For one, purified G proteins could only be crystallized after removal of post-translational farnesyl or geranylgeranyl lipid modifications [45, 46]. The three-dimensional structure of the G protein did, however, strongly suggest that the lipid modified N- and C-termini of α and γ , respectively, were proximal to one another [46]. Even after crystallization with an intact γ including the prenylated C-terminus, conformational instability of the C-terminal region precluded assignment of a static structure to this region of γ [53]. In addition, no crystal structure exists for a G protein in the *empty* state, that is, with no nucleotide bound; this state represents the transition between receptor-dependent release of GDP and binding GTP. Furthermore, a seven-transmembrane-spanning receptor has not been crystallized with a G protein; what is known of receptor:G protein interactions has been derived from biochemical data. Putative points of contact between G protein and receptor on the heterotrimer include the N-terminus [54] and C-terminus [55] of G α , the C-terminus of γ [56–58] and the C-terminus of β [59]. Thus, the mechanism of signal transfer from activated receptor in a membrane environment to G protein continues to be a point of conjecture based on empirical evidence.

Mechanism of Activation

Based largely on the crystal structures of heterotrimeric G proteins and monomeric GTP-binding proteins, several studies from Henry Bourne's laboratory, Onrust et al. [60] in 1997 and Iiri et al. [61] in 1998, sought to explain the molecular mechanism of activation of G protein

by receptor through the 'lever' hypothesis. Such a hypothesis was necessary because the distance between the intracellular loops of a receptor and GDP were thought to be too far for direct interaction (fig. 1), thus the requirement of receptor to 'act at a distance' [62]. Integral to the hypothesis was the interaction between activated receptor and the $\beta 6$ strand/ $\alpha 5$ helix region of G α , which contains a loop to which GDP binds. Further, the switch I region and the $\beta 3/\alpha 2$ loop of G α , in addition to containing $\beta\gamma$ -binding sites, also form a lip that provides a secure binding site for GDP. Thus, two important events in receptor activation of G protein occur when: (1) receptor induces a conformational change in the $\beta 6$ strand/ $\alpha 5$ helix region of G α , which would subsequently alter the GDP-binding site on the $\beta 6/\alpha 5$ loop, and (2) the postulated insertion of the intracellular loops of the receptor into a crevice between α and $\beta\gamma$, which by employing $\beta\gamma$ as a lever to pry α and $\beta\gamma$ apart, could lead to an allosterically induced conformational change in the switch I/ $\beta 3\alpha 2$ loop ensconcing GDP, allowing release of nucleotide.

This hypothesis was tested biochemically by the creation of a mutant $G_s \alpha$ subunit that bound the $\beta\gamma$ dimer in a conformation that mimicked the G protein structure described above in the absence of receptor [63]. Transfecting cells with $\beta\gamma$ and the mutant $G_s \alpha$ increased the activity of the mutant $G_s \alpha$ at adenylyl cyclase, compared with the mutant $G_s \alpha$ alone, further supporting the hypotheses that $\beta\gamma$ acts as a lever to affect GDP release in the course of G protein activation. An iteration of the lever hypothesis is the 'gear-shift' hypothesis [64], which holds that the N-terminus of the γ subunit acts as a gear-shift by interacting with the helical domain of the α subunit, thereby stabilizing the transitory nucleotide free *empty* state, and facilitating nucleotide exchange.

Studies with rhodopsin and synthetic G protein peptides suggest that the C-terminus of γ and the C-terminus of α interact with the activated receptor in a sequential, interdependent manner, and this binding facilitates the conformational change that allows release of GDP [65]. A later study refined this model using synthetic peptides and time-resolved near infrared light scattering, indicating that the prenylated C-terminus of γ was the first point of contact with activated receptor, followed by the C-terminus of α in a brief transitory state that enabled GDP release [66]. A conformational switch in $\beta\gamma$, first proposed in 1995 [67], was envisioned as a mechanism for this sequence of events. Such a switch was predicted because in both detergent solubilized heterotrimeric transducin and $\beta_1\gamma_1$, the C-terminus of γ was resistant to car-

boxypeptidase Y, suggesting inaccessibility to other proteins; however, a 12-amino-acid C-terminal farnesylated peptide from γ_1 was able to bind to and stabilize light-activated rhodopsin [67]. The details of the switch, characterized by mutational analysis and NMR structural studies, describe an unstructured C-terminus of γ that transforms into an amphipathic helix upon interaction with activated receptor [68, 69]. Three residues conserved across γ subunit isoforms, Asn62, Pro63 and Phe64, were proposed to be critical for the receptor-dependent conformational change in γ .

More recent experiments with protein in solution have begun to yield structural data on the nucleotide free *empty* state of a G protein activated by receptor. NMR studies with transducin and light-activated rhodopsin suggest that the empty state of transducin α is conformationally dynamic, a condition not attained in the absence of activated receptor [70]. The use of site-directed spin-labeling methods with $G_i \alpha_{GDP}$, $G_i \alpha_{GDP}\beta\gamma$ and $G_i \alpha_{GTP}\gamma_S$ has allowed even more precise characterization of structural changes in solution; experiments examining G protein activation by rhodopsin indicate that structural changes in $G_i \alpha$ are propagated via the switch I region to the α F-helix, in concert with movement of the α 5 helix, which form part of the nucleotide-binding pocket [71]. While this conformational change facilitates GDP release, another interesting structural change resulting from the *empty* state is the formation of new contacts between α and $\beta\gamma$, suggested by the crystal structure. The presence of these new contacts between α and $\beta\gamma$ in the *empty* state further support a receptor-dependent conformation of the *empty* heterotrimer distinct from $G_i \alpha_{GDP}\beta\gamma$.

Subunit Dissociation

Although receptor-dependent nucleotide exchange has been well established, the issue of subunit dissociation has been somewhat more controversial. In early experiments characterizing the biochemical nature of G proteins, purified transducin, G_i and G_s proteins activated with various combinations of aluminum, magnesium, fluoride, GDP, or GTP analogues and detergent could be dissociated into their constituent α and $\beta\gamma$ dimers using a number of separation techniques [72–74]. The extrapolation that this phenomenon occurred in vivo was attractive for several reasons. For one, subunit dissociation provided a bifurcation of signal after activation of receptor; both α and $\beta\gamma$ were free to regulate downstream effectors independently. The potential for α and $\beta\gamma$ dimers to form new G protein

combinations as a result of activation of several different receptors was yet another point where a cell could regulate signaling specificity. The molecular underpinnings to support the idea that subunit dissociation represented the mechanism of $\beta\gamma$ activation, and reassociation of G protein served to inhibit $\beta\gamma$ signaling, came from mutational studies that concluded that regions of $\beta\gamma$ that activated effectors were found to overlap with G α -binding sites (fig.1) [75, 76]. These findings suggested that a $\beta\gamma$ dimer in a heterotrimeric complex was in fact inactive, and that activation of $\beta\gamma$ was synonymous with release from G α upon activation by receptor.

Interestingly, a study by Bonacci et al. [77] found that the region of the β subunit that interacts with the switch II region of G α could be targeted with peptides and small organic molecules to selectively disrupt interactions between $\beta\gamma$ dimers and effectors; this strategy has been used to inhibit inflammation in vivo by blocking $\beta\gamma$ -mediated activation of PI3-kinase γ [78], and thus represents a promising area for pharmacological intervention.

Further evidence for subunit dissociation was revealed in the crystal structure of a $G_q \alpha$ -p63RhoGEF complex [79], in which p63RhoGEF interacted with the α 2/ β 4 region, containing switch II, and the α 3/ β 5 regions of $G_q \alpha$. In addition, the crystal structure of activated $G_s \alpha$ in a complex with the catalytic domains of adenylyl cyclase also revealed effector-binding sites on $G_s \alpha$ to be the switch II region and the α 3- β 5 loop [80]. Since $\beta\gamma$ binds the switch II region, the binding of p63RhoGEF and adenylyl cyclase to $G_q \alpha$ and $G_s \alpha$, respectively, appears to be mutually exclusive to the binding of $\beta\gamma$, and supports the notion that subunit dissociation is a consequence of G protein activation. However, biochemical and kinetic arguments have been advanced to suggest that subunit dissociation is not necessary for G protein activation [81]. For example, expression of a fusion protein of G protein α and β subunits in yeast signaled as well as α and β subunits co-expressed individually [82]; since the fusion protein did not allow complete physical separation of α and β , a conformational change was proposed as a means for activation of the heterotrimer leading to signaling.

Fortunately, in the last several years sophisticated imaging techniques such as FRET (fluorescence resonance energy transfer) have emerged that allow the question of subunit dissociation to be more fully evaluated in live cells. FRET studies with receptor-dependent activation of fluorescently tagged G protein in *Dictyostelium discoideum* concluded that subunits did in fact dissociate upon activation [83], although a conformational change in the heterotrimer could not be completely ruled out. A similar

FRET study with fluorescently tagged $G_i \alpha$ and $\beta_1\gamma_2$ in HEK cells concluded that the heterotrimer underwent a molecular rearrangement upon activation by receptor, but did not dissociate [84]. A comparison of heterotrimers consisting of $\beta_1\gamma_2$ and other members of the G_i subfamily using FRET found $G_O \alpha$ appeared to dissociate from $\beta_1\gamma_2$ upon receptor stimulation, whereas $G_{i1,2,3} \alpha$ or $G_z \alpha$ did not [85]; these differences in subunit dissociation were traced to several distinct regions in $G_{i1} \alpha$. Because the FRET signal is dependent on both proximity and orientation between two fluorophores, it became clear that the distinction between conformational change and physical dissociation of G protein subunits was not to be unequivocally resolved by FRET.

Another imaging technique called FRAP (fluorescence recovery after photobleaching) was more suited to address complete physical dissociation of G protein subunits, as fluorescence recovery is dependent on the movement of fluorescently tagged proteins into and out of bleached regions of the plasma membrane. In one FRAP study which examined G protein subunit dissociation by evaluation of the ability of fluorescently tagged, immobile $G_{OA} \alpha$, $G_{i3} \alpha$ and $G_s \alpha$ subunits to constrain fluorescently tagged $\beta_1\gamma_2$ [86], results were similar to the FRET studies described above. Consistent with the FRET experiments, receptor activation resulted in dissociation of $G_{OA} \alpha$ from $\beta_1\gamma_2$; however, in these experiments, $G_{i3} \alpha$ was also shown to dissociate from $\beta_1\gamma_2$. In contrast, the $G_s \alpha \beta_1\gamma_2$ heterotrimer did not dissociate upon receptor stimulation [86]. Further evidence that subunit dissociation is not always necessary for G protein activation was provided by the discovery that the stability of complexes of effectors and heterotrimeric G proteins persisted even after activation of receptor [87]. Moreover, effector activity of $G_q \alpha$ was demonstrated to be augmented by $\beta\gamma$, independent of receptor activation [88].

If subunit dissociation does not occur, one striking paradox is how a $\beta\gamma$ dimer can signal with effector-binding sites occluded by G α . Part of the answer is likely related to the three switch regions of G α (fig. 1) which undergo a conformational change upon exchange of GDP for GTP (GTP γ S in the crystal structure) that leads to a decreased affinity of G α for $\beta\gamma$. However, an additional contact site suggested by the crystal structure between α and γ is the N-terminus of α and the C-terminus of γ , both modified by lipid. Biochemical evidence also supports this interaction, as myristoylated $G_O \alpha$ has a higher affinity for $\beta\gamma$ than $G_O \alpha$ with an unmodified N-terminus [89]; interactions between the γ C-terminal prenyl group and a N-terminal lipid of α may also be strength-

ened if both modifications are inserted into the lipid membrane (fig. 1).

Thus, one model, often referred to as the 'clam shell', predicts that upon activation, α and $\beta\gamma$ pull apart, perhaps with the lipid moieties inserted into the membrane serving as a hinge in the heterotrimer, enough for effector molecules to interact with the binding regions of $\beta\gamma$ or α (fig. 1). This activation without complete subunit dissociation may be manifested in subtle conformational changes that may be difficult to detect using cellular imaging techniques. The rearrangement, but not dissociation of $G_i \alpha:\beta\gamma$ upon activation [84] discussed earlier may be an example of the clam shell model. Lack of, or more likely incomplete subunit dissociation implies that particular $\beta\gamma$ dimers from heterotrimers such as G_s may not readily reassociate with other G α isoforms after activation by receptor. In a model where a receptor has unlimited access to all G proteins expressed in a cell [90], the question of preserving signaling specificity may be related to the extent to which a G protein dissociates upon activation, and subsequently, the potential to re-association with other subunit combinations.

Specificity of β/γ Interactions

Years of experimental data have led to the consensus that β and γ isoforms form a functional dimer that does not dissociate under physiological conditions. Thus, the activity of a $\beta\gamma$ dimer is derived by the identity of both β and γ isoforms. To examine the functional diversity of $\beta\gamma$ dimers, many experiments have aimed to discover which $\beta\gamma$ dimers can physically form. In cell types with a restricted assortment of β and γ isoforms, such as expression of β_1 and γ_1 in rods and β_3 and γ_8 in cones, this approach was sufficient to estimate probable G protein subunit interactions. A more difficult question, especially in the context of expression of several β and γ isoforms in a single cell, is which $\beta\gamma$ dimers actually form? Useful answers have come from studies dissecting receptor signaling pathways, discussed below; however, the most direct answer is revealed by the purification of G protein $\beta\gamma$ isoforms from a specific tissue or cell type. Unfortunately, these are not practical experiments to undertake for the multitude of cell types and tissues that constitute an organism.

β_1 and β_2 Interactions with γ Isoforms

Several systems have been used to study β and γ interactions, including the yeast two-hybrid [91], in vitro

translation [92–94] and transfected cell assays [95]. The conclusions generally held that for the first two β isoforms, β_1 was the least restricted in its interactions with other γ isoforms, and β_2 failed to interact with γ_1 and γ_{11} . The region of γ_1 that inhibited dimer formation with β_2 was ultimately determined to be the 3 amino acids at positions 38–40 of γ_1 [96]; conversely, replacement of these residues in γ_2 with the analogous residues of γ_1 also inhibited dimer formation with β_2 . More recent live cell-imaging techniques have confirmed much of the earlier literature, although in a noteworthy finding, β_1 appeared to display the strongest preference for γ_{12} [97].

β_3 Interactions with γ Isoforms

Interpretation of results with β_3 is more complex. Unlike dimers containing β_1 or β_2 , dimers containing β_3 are much less resistant to complete proteolysis by trypsin [7], and the β_3 subunit has displayed weak or absent capacity to interact with γ subunits, including γ_1 and γ_2 [92]. The β_3 isoform does contain an additional tryptic cleavage site at lysine 177; however, in experiments with the more specific protease Arg-C, complete digestion of $\beta_3\gamma$ dimers was also observed [93]. On the other hand, β_3 has been shown to co-precipitate with γ_5 , $\gamma_{8\text{cone}}$ or γ_{12} [7], and a $\beta_3\gamma_2$ dimer with activity at receptor and effector has been purified using a baculovirus expression system [98]. The anomalous results obtained from studies with β_3 suggest that the crystal structure of a G protein $\beta\gamma$ dimer based on the β_1 isoform may not necessarily predict subtle structural variations of $\beta\gamma$ dimers containing other even highly homologous β isoforms.

β_3 Splice Variant Interactions with γ Isoforms

Several β_3 splice variants have been characterized: β_{3s} , β_{3s2} and β_{3v} . β_{3s} , distinguished from β_3 by deletion of 41 amino acids, or one entire WD-40 domain affecting blades three and four of the torus, is similar to β_3 in γ -binding specificity [7]. However, purification of the β_{3s} isoform from mammalian or insect cell expression systems has not been reported; in contrast to other $\beta\gamma$ combinations, β_{3s} was poorly extracted with 1% (v/v) Genapol, 1% (w/v) CHAPS, or 1% (w/v) cholate when expressed in Sf9 insect cells with either the γ_2 , γ_5 or γ_7 isoforms [98]. The β_{3s2} splice variant, characterized by a similar deletion that affects blades five and six, was also not found to be substantially different in γ -binding preference from β_3 [8]. However, the β_{3v} splice variant, which lacks the last three blades of the β torus and has a unique C-terminal region, dimerizes with only the γ_3 and γ_{12} isoforms [9]. The β_{3s} and β_{3s2} splice variants have been

found to be associated with a C825T polymorphism in the β_3 gene that is associated with increased risk of hypertension [99]. Functional studies on the β_{3s} protein suggest that the loss of 41 amino acids results in enhanced receptor-dependent G protein signaling [7], which may contribute to the etiology of hypertension.

β_4 Interactions with γ Isoforms

The β_4 isoform has been shown to interact similarly with all γ isoforms in precipitation experiments from in vitro expression systems [94, 100]. However, when $\beta\gamma$ dimers were precipitated from bovine lung using specific γ antibodies, β_4 clearly showed a preference for association with γ_5 and γ_{12} over γ_2 and γ_3 [101]. This is a good example of the difference between what $\beta\gamma$ dimers can form in vitro and which dimers do actually form in vivo. Factors that influence such specificity in $\beta\gamma$ formation may include chaperone proteins, which have recently been suggested to play a role in dimer formation between specific β and γ isoforms. For example, the Cytosolic Chaperonin Complex (CCT), which has been demonstrated to bind G β during $\beta\gamma$ dimer biosynthesis, interacts most strongly with the β_1 and β_4 subunits, weakly with β_5 and β_{3s} , and intermediately with β_2 and β_3 [102]. In addition, the Dopamine Receptor-interacting Protein 78 (DRIP78) has been characterized as a chaperone for G γ , and also exhibits specificity in binding with highest affinity to γ_2 and γ_3 , compared to γ_1 , γ_7 and γ_{11} [103]. More detail on this emerging field as it relates to $\beta\gamma$ dimer formation can be found in a recent review by Willardson and Howlett [104].

β_5 Interactions with γ Isoforms

β_5 occupies a special niche in the G β family both structurally and functionally. Early in its characterization, it was noted that dimers containing β_5 and γ_2 [105], or any of several other γ isoforms [M.B. Jones, unpubl. observation] were highly unstable in certain detergents; this observation will be discussed further below. It was also observed that in addition to G γ subunits, the β_5 isoform could bind to members of the R7 subfamily of Regulators of G Protein Signaling (RGS) proteins, and the crystal structure of a complex of β_5 and RGS9 was recently solved [48]. The β_5 RGS literature is beyond the scope of this review, but see Berman and Gilman [4] and De Vries et al. [106] for other reviews. In terms of forming a conventional $\beta\gamma$ dimer, β_5 was shown to interact preferentially with γ_4 by the yeast two-hybrid system [91], and live cell-imaging techniques suggested $\beta_5\gamma_2$ and $\beta_5\gamma_7$ are favored dimer combinations [107], although in the same

system, β_5 could interact with an RGS7 protein as well. The specificity observed in formation of $\beta\gamma$ dimers with different β and γ isoforms suggests the capacity for extensive modulation of G protein signaling. Further, the fact that the β_5 isoform can bind γ subunits, or alternatively members of the R7 subfamily of RGS proteins, indicates that other signaling pathways may be highly integrated into receptor-dependent $\beta\gamma$ signaling.

Specificity of $\alpha/\beta\gamma$ Interactions

The G α isoform is necessary for the activation of a G $\beta\gamma$ dimer, since specific localization of $\beta\gamma$ to the plasma membrane has been reported to require both prenylation and heterotrimer formation [26]. Further, all three subunits are required for transfer of signal from activated receptor to G protein [108]. In addition, the identity of the G α subunit may also determine the cellular mobility of an activated $\beta\gamma$ dimer via its propensity to undergo subunit dissociation upon activation (discussed above). Considering the large number of $\alpha:\beta:\gamma$ subunit combinations that could potentially form, relatively little is known as to which heterotrimers can form, not to mention which actually do form in vivo. Limited subunit expression has provided information on probable heterotrimer combinations in different physiological systems. For example, in the visual system, G $_t \alpha\beta_1\gamma_1$ is likely to predominate in rods, whereas G $_t \alpha\beta_3\gamma_8$ is most prevalent in cones; in taste receptor cells, the most abundant isoforms are α -gustducin, β_1 , β_3 and γ_{13} [109].

Regulation of Heterotrimer Composition

From empirical observations, many studies characterizing various G α isoforms have used $\beta_1\gamma_2$ as the archetypal $\beta\gamma$ dimer. The $\beta_4\gamma_2$ dimer was also observed to form a heterotrimer with G $_{OA} \alpha$ [110]. Specificity has been reported in the $\beta\gamma$ dimers associated with different G $_O$ isoforms purified from bovine brain, with the G $_{OA}$ heterotrimers containing much more γ_7 than the G $_{OC}$ heterotrimers [111]; G $_{OC} \alpha$ is distinguished structurally from G $_{OA} \alpha$ by deamidation of Asn346 and Asn347 at the C-terminus [112]. However, $\beta\gamma$ purified from G $_{OA}$ interacted equally well with the α subunits purified from G $_{OA}$ and G $_{OC}$, as judged by γ_7 immunoreactivity [113], suggesting that in this case, differences in the $\beta\gamma$ composition observed between G $_{OA}$ and G $_{OC}$ were due to restricted expression of isoforms within cells or tissues. Transcriptional regulation is another mechanism that may regulate combinations of G α and $\beta\gamma$ dimers; one study

noted that G β_4 mRNA was regulated by expression of G $_s \alpha$, G $_{i3} \alpha$ and G $_{i11} \alpha$ [114]. Reduction of G $_{olf} \alpha$ protein was also correlated with deletion of the G γ_7 gene in mice [115]. Furthermore, genetic deletion of the γ_1 gene in mice resulted in a greater than 25-fold reduction in G $_t \alpha$ and β_1 protein levels in the retina, although interestingly, mRNA levels for G $_t \alpha$ and β_1 were similar to wild-type mice [116]. These studies suggest that heterotrimer composition and formation are highly regulated at the levels of transcription, translation and posttranslational processing.

In one study that used immunofluorescence microscopy to compare the ability of a panel of $\beta\gamma$ dimers to target mutant G $_s \alpha$ and G $_q \alpha$ to plasma membranes as an indication of heterotrimer formation, $\beta\gamma$ dimers containing β_1 or β_2 were equally effective at interacting with either G $_s$ or G $_q \alpha$ isoforms [117]; $\beta\gamma$ dimers containing β_3 did not interact well with G $_s \alpha$ or G $_q \alpha$, and $\beta\gamma$ dimers containing β_4 were able to interact with G $_s \alpha$, but not G $_q \alpha$. Interactions between G $_s \alpha$ or G $_q \alpha$ and $\beta\gamma$ dimers containing G β_5 were not observed [117]. On the other hand, a study that used live cell-imaging techniques observed that both G $_O \alpha$ and G $_q \alpha$ could target $\beta_5\gamma_2$ to the plasma membrane [107]. Moreover, there is also a report of G $_q \alpha$ from brain extract binding a $\beta_5\gamma_2$ affinity column [118]. The presence of a receptor may also facilitate interactions between specific G α and $\beta\gamma$ isoforms. For example, in contrast to co-localization studies with only G protein subunits [117], purified $\beta_4\gamma_2$ was able to couple G $_q \alpha$ to the M $_1$ muscarinic receptor [119], and $\beta_3\gamma_2$ was able to couple G $_s \alpha$ to the β_1 -adrenergic and adenosine A $_{2A}$ receptors [98]. Furthermore, the $\beta_5\gamma_2$ dimer has also been demonstrated to couple G $_q \alpha$ to the M $_1$ muscarinic receptor [120], and weakly couple G $_s \alpha$ to the β_1 -adrenergic receptor [98].

Receptor-Dependent Translocation of $\beta\gamma$

The $\beta\gamma$ dimer has been shown to translocate upon receptor stimulation. In perfused rat hearts, stimulation of the β_1 -adrenergic receptor induced the translocation of the β_3 subunit from cytosol to membranes [121]; no such effect was observed for the β_1 or β_2 subunits, which were predominantly in the membrane fraction. It should be noted that the γ isoforms identified with the β_1 , β_2 and β_3 subunits were not characterized, and thus it is possible that both β and γ isoforms contributed to the translocation of $\beta_3\gamma$.

The γ subunit is also a determinant of which $\beta\gamma$ dimers undergo receptor-dependent translocation. Live cell-imaging experiments revealed that $\beta\gamma$ dimers containing γ_1 , γ_{11} and γ_9 translocate rapidly to the Golgi membranes upon receptor stimulation, $\beta\gamma$ dimers containing γ_5 and γ_{10} translocate slowly, and $\beta\gamma$ dimers containing γ_2 , γ_3 , γ_4 , γ_7 , γ_8 and γ_{12} do not translocate [122, 123]; the translocation was observed to be reversed upon addition of a receptor antagonist. Interestingly, although γ_1 , γ_{11} and γ_9 are all farnesylated, the geranylgeranylated γ_{13} also translocated rapidly to the endoplasmic reticulum [123]. This behavior of γ_{13} reflects the study's finding that $\beta\gamma$ translocation occurs as a reversible, diffusion-mediated process that is related to the amino acid sequence of the γ isoform, and not the identity of the prenyl group [123]. Within the family of $\beta\gamma$ dimers that translocate, the α subunit also has influence on the rate of translocation, primarily related to the nucleotide exchange rates of the α subunits [124]. One important point to make with respect to receptor-dependent $\beta\gamma$ translocation is that it can occur independently of G α [122], and thus represents an example of complete G protein subunit dissociation (see discussion above) determined by the nature of the γ isoform in a heterotrimer. Subsequent studies have observed that even in the absence of receptor activation, there is a basal level of heterotrimer shuttling between plasma and intracellular membranes [125]. The discovery of receptor-dependent $\beta\gamma$ translocation is important in that it increases the complexity of the spatial dimension to $\beta\gamma$ signaling, and suggests that cellular localization of $\beta\gamma$ dimers after G protein activation is tightly controlled by the identity of the β and γ isoforms in a G protein.

Exceptions to the Rule

Receptor-Independent G Protein Activation

The review to this point has discussed the conventional wisdom regarding the molecular determinants that influence activation of a $\beta\gamma$ dimer, which is usually preceded by partial or complete dissociation from G α . As discussed above, the $\beta\gamma$ dimer is capable of dynamic translocation in the absence of G α , suggesting that activity may not be limited to the plasma membrane. Activation of a $\beta\gamma$ dimer has been proposed to occur in the absence of receptor, or nucleotide exchange, via direct interaction by an Activator of G protein Signaling protein, AGS8 [126]; this activation was also suggested to not require subunit dissociation [127]. Another mechanism

proposed for G protein activation is the absence of receptor involves phosphorylation of histidine 266 of β by NDPK; the phosphate is subsequently transferred onto the GDP bound to G α , effectively producing a GTP-bound activated α subunit without the requirement of receptor catalyzed release of GDP [15]. The discovery of G $\beta\gamma$ in the endoplasmic reticulum (ER) of Arabidopsis led one researcher to speculate that $\beta\gamma$ has signaling functions in the ER, such as regulation of PLC or IP₃ receptors [128], independent of heterotrimer formation with G α [129]. This is an intriguing hypothesis, considering that RACK1, a G β -like scaffolding protein, has been shown to bind both $\beta\gamma$ [130] and IP₃ receptors [131]. Moreover, the ability of RACK1 to regulate $\beta\gamma$ signaling, such as attenuation of PLC- β_2 activation [132], provides a potential mechanism for G $\beta\gamma$ signaling to occur without G α or activated receptor.

Stability of $\beta\gamma$ Dimers: Unconventional Roles for $\beta\dots\gamma$

Deviating further still from conventional wisdom is the biological significance of instability or low affinity between particular combinations of β and γ . The model of the tightly associated $\beta\gamma$ dimer that couples a G α subunit to an activated receptor is based largely on studies with $\beta_1\gamma_1$ or $\beta_1\gamma_2$. However, other less stable combinations such as $\beta_5\gamma_2$ and $\beta_4\gamma_{11}$ [119] are less amenable to biochemical studies, and thus, instability could easily be interpreted as incompatibility between particular combinations of β and γ isoforms. The high degree of heterogeneity inherent in the potential combinations of $\beta\gamma$ dimers, and the spectrum of biophysical properties suggest that unconventional signaling paradigms for β and γ exist. It may be useful in this speculative exercise to first consider examples of potential dissociation between β and γ that have been reported in the literature.

$\beta\gamma$ Instability in vitro

Examples of low affinity of β for γ can be found with many of the β isoforms. Lower affinity of β_3 for γ subunits has been reported in several systems examining $\beta\gamma$ formation, such as in vitro translation and yeast two-hybrid [91, 92, 94], although after purification from Sf9 cells with γ_2 , the $\beta_3\gamma_2$ dimer was completely stable in several different detergents [105]. More evidence exists for the instability of β_5 with γ subunits, and the biochemical properties of a β_5 monomer. Attempts to purify the $\beta_5\gamma_2$ dimer revealed that it was sensitive to detergent, and high

concentrations of CHAPS or cholate induced subunit dissociation [105]. Other $\beta_5\gamma$ dimers containing γ_1 , γ_7 , γ_{10} and γ_{12} could not be purified even at low (0.1% Genapol) detergent concentration [Miller B. Jones, doct. thesis]. Whereas the β_1 subunit forms unstable high-molecular-weight aggregates in the absence of γ [133], the β_5 subunit can exist as a monomer which is highly resistant tryptic cleavage [134].

Instances of $\beta\gamma$ instability related to γ isoforms have also been reported. One early study characterizing the purification of G protein from bovine brain reported monomeric γ_3 under non-denaturing conditions [36]. The γ_{11} isoform has also been noted for its propensity to dissociate from the β during purification [37], indicating a weak interaction. Parameters affecting G $\beta\gamma_{11}$ affinity were further explored with recombinant heterotrimers consisting of $G_{i1} \alpha \beta_1\gamma_{11}$ and $G_{i1} \alpha \beta_4\gamma_{11}$, both of which were stable upon purification; however, activation of the heterotrimers with GTP γ S resulted in a significant dissociation of β_4 and γ_{11} , but not β_1 and γ_{11} [119]. Immunoprecipitation experiments examining $\beta\gamma$ dimers containing the γ_{13} isoform found that β_1 , β_3 and β_4 could be immunoprecipitated with a hemagglutinin tagged γ_{13} subunit, whereas the β_2 and β_5 subunits could not [135]; the authors speculated that the $\beta\gamma$ instability was due to detergents used in the immunoprecipitation. This instability of $\beta_2\gamma_{13}$ and $\beta_5\gamma_{13}$ was not indicative of incompatibility between subunits, as these dimers were found to be effective in the activation and inhibition, respectively, of GIRK1/4 channels in transfection experiments [135]. Further, live cell-imaging techniques were used to observe that β_2 and γ_{13} could effectively translocate together following receptor activation [123].

Potential Biological Activity of β and γ Monomers

Biochemical evidence of instability in particular combinations of $\beta\gamma$ dimers in vitro logically leads to questions of whether monomeric β or γ subunits exist, and what they may be doing in a cell. Overexpression of two of the β_3 splice variants, β_{3s} and β_{3s2} , was shown to markedly stimulate MAP-kinase activation [8]; interestingly, co-expression of known γ partners for these β_3 splice variants had no further effect on MAP-kinase activity. These data suggest that the β_3 splice variants may have biological activity in the absence of γ . Moreover, the authors of the study offer the possibility that another β_3 splice variant with restricted γ -binding partners, β_{3v} , may exist as a monomer [9]. One study established that purified β_5 and γ_2 monomers can be reconstituted to form a dimer with the ability to active PLC- β_2 in vitro

[134]; the study further demonstrated that monomeric β_5 was able to functionally interact with $G_i \alpha$ and $G_o \alpha$. Another report characterized the ability γ_2 expressed in Sf9 cells to bind to a $G_o \alpha$ column, and protect the α subunit from tryptic cleavage [136]. Functional activity of a monomeric γ_5 subunit was also suggested in a study that demonstrated the ability of a bacterially expressed non-prenylated γ_5 subunit to regulate transcription by binding to the adipocyte enhancer-binding protein (AEBP1) transcriptional repressor [34]. Although β protein was also observed in co-immunoprecipitation studies with γ_5 and AEBP1 in mammalian cells, the data support the idea of transcriptional activity of a γ_5 monomer, perhaps in addition to $\beta\gamma_5$ dimers in the nucleus. Absence of prenylation, as discussed above, may not be the only signal to direct particular $\beta\gamma$ dimers to the nucleus; a study examining the localization of fluorescently tagged β_5 co-expressed in HEK-293 cells with various fluorescently tagged γ isoforms concluded that the γ_1 , γ_5 , γ_{10} and γ_{11} isoforms targeted β_5 to the nucleus, whereas γ_2 and γ_7 did not [107]. Since most of the γ isoforms that targeted β_5 to the nucleus could not be purified as a complex with β_5 even under conditions of low detergent stringency (see above), the observed differences among $\beta_5\gamma$ dimers in the context of localization suggests that dimer instability is related to $\beta\gamma$ signaling roles.

Receptor: γ Interactions

Another potential biological role for γ subunits can be inferred from studies characterizing C-terminal γ peptides with activated receptor. A C-terminal farnesylated γ_1 peptide was reported to stabilize the active form of rhodopsin [56], and a C-terminal geranylgeranylated peptide corresponding to γ_5 , but not γ_7 or γ_{12} , was able to inhibit M_2 muscarinic receptor signaling [137]. Further receptor-binding experiments revealed that the γ_5 peptide was able to stabilize a novel state of the M_2 muscarinic receptor [138]. These studies suggest an interaction between the C-terminal tail of γ and receptors. However, the notion that receptor may not interact with the C-terminal tail of γ during G protein signaling was supported by reconstitution studies that found that a $\beta_1\gamma_1$ dimer engineered with a photoreactive farnesyl analogue was cross-linked to phospholipid, but not receptor, after reconstitution of transducin with rhodopsin in membranes [139]. The discrepancy between peptide and whole protein studies was also highlighted by the fact that the M_2 muscarinic receptor more efficiently stimulated GTP hydrolysis in a $G_o \alpha\beta_1\gamma_7$ heterotrimer compared to a $G_o \alpha\beta_1\gamma_5$ heterotrimer [140]. This raises the possibility that

γ peptides may have properties distinct from the analogous $\beta\gamma$ dimer at receptor, and more importantly, begs the question regarding the biological activity of a monomeric γ subunit: If a γ peptide can functionally interact with receptor, would a γ subunit, which has also been reported to exist as a monomer in the absence of β or detergent [141], have similar binding and regulatory properties at receptor?

Receptor: β Interactions

Less complex biological systems may also inform the prospective roles of β and γ in G protein signaling. For example, in fission yeast *Schizosaccharomyces pombe*, the β subunit, which lacks an amino terminal coiled-coil domain, retains some activity when its γ partner is deleted [142]; further, when the C-terminal CAAX box from the γ subunit is fused onto the C-terminus of the β subunit, some activity is recovered, suggesting more of a targeting role for γ in that system. This result was mirrored in studies with mutant yeast γ subunits in *Saccharomyces cerevisiae*, which found that a C-terminal domain preceding the CAAX box was not required for $\beta\gamma$ coupling to receptor in vivo [143]. Recently, another study in *S. cerevisiae* also found that the RACK1 ortholog Asc1 could bind and influence the nucleotide-binding properties of $G\alpha$, and essentially perform the role of a β subunit, presumably without the presence of γ [144].

There are several instances that suggest that interactions between β and γ and receptor in higher vertebrates may be more dynamic than previously thought. Expression of β_3s , but not β_3 , was shown to be able to activate $G\alpha$ subunits in the presence of mastoparan-7 in digitonin-permeabilized COS-7 cells [7]; the authors reasoned that the β_3s protein dimerized with endogenous γ subunits. Such an experiment suggests the existence of monomeric β and γ subunits in vivo, because either there is biological activity of an expressed β_3 splice variant, or alternatively, a pool of γ isoforms ready to dimerize with an ectopically expressed β subunit. Moreover, purification of $\beta\gamma$ from rod outer segments of *Bufo marinus* yielded a small population of free β with no γ partner [145]; interestingly, the β subunit, noted for its high homology with bovine β_1 , was as effective as the purified $\beta\gamma$ at stimulating GTP γ S binding to $G\alpha$ after reconstitution with illuminated rod outer segment disc membranes. Possibly related to the results of β_1 activity in *B. marinus*, a γ_1 knockout mouse retained some ability of rhodopsin to signal through G_t in rod outer segments with only residual amount of $G_t \alpha$ and β_1 remaining [116]. Since no upregulation of other β or γ isoforms was observed, and

no other γ isoforms were detected in immunoprecipitations of β_1 , one possible explanation (not advanced by the authors) is that the residual β_1 may be able to substitute on some level for the $\beta_1\gamma_1$ dimer. Although examples of β coupling $G\alpha$ to receptor likely represent a small minority of signaling paradigms in higher vertebrates, they may shed light on the interactions of unstable $\beta\gamma$ dimer combinations with receptors and effectors, and prove useful in providing a way of dissecting the contribution of β and γ to G protein signaling.


Receptor/ $\beta\gamma$ Interactions Determine Signaling Specificity

Specificity of receptor:G protein interactions can be influenced by the identity of each of the subunit isoforms. Thus, the assembly of a $\beta\gamma$ dimer with an α subunit to form a heterotrimer of defined composition within a cell functions to impart a degree of specificity by limiting the number of signaling pathways in which the G protein can directly participate. The first evidence suggesting that the identity of the β and γ isoforms in a heterotrimer can influence receptor coupling were based on experiments using antisense oligonucleotides to attenuate expression of specific subunit isoforms. Inhibition of voltage-sensitive Ca^{2+} channels in rat GH₃ cells was reported to be mediated via coupling of the $G_{O1} \alpha\beta_3\gamma_4$ heterotrimer to the muscarinic receptor and coupling of the $G_{O2} \alpha\beta_1\gamma_3$ heterotrimer to the somatostatin receptor [146–148]. The antisense approach was further employed to suggest that in rat portal vein myocytes, the angiotensin AT_{1A} receptor couples to $G_{13} \alpha\beta_1\gamma_3$ [149], and the ET_A receptor couples to $G_{11} \alpha\beta_3\gamma_5$ [150].

Prenyl Status

The type of prenyl moiety on the γ subunit has also been shown to be critical for receptor G protein interactions. Experiments involving rhodopsin have generated the most conflicting reports; on the one hand, the farnesyl group was observed to interact more efficiently than the geranylgeranyl group with activated rhodopsin [151, 152]. Alternatively, a mutated $\beta_1\gamma_1$ dimer that incorporated the geranylgeranyl group instead of the farnesyl group was found to have a 3-fold higher affinity for rhodopsin than wild-type $\beta_1\gamma_1$ [153]. These results were mirrored in other studies that characterized $\beta_1\gamma_1$ and $\beta_1\gamma_2$ dimers that were mutated to alter the specificity of prenylation; farnesylation of $\beta_1\gamma_2$ reduced its affinity for the adenosine A₁ receptor, and conversely, geranylgeranyl-

Fig. 2. Effect of different $\beta\gamma$ isoforms on A_{2a} adenosine receptor signaling. $\beta_1\gamma_2$ couples $G_s \alpha$ poorly to the A_{2a} receptor, leading to lower activation of adenylyl cyclase; increasing levels of $\beta_4\gamma_2$, which couples $G_s \alpha$ more efficiently to the A_{2a} receptor, increases adenylyl cyclase activation and intracellular cAMP levels.



ation of $\beta_1\gamma_1$ increased its ability to couple to the same receptor [154]. Farnesylation, however, does not always result in a decreased affinity of $\beta\gamma$ for receptor. In the case of $\beta_1\gamma_{11}$, which like $\beta_1\gamma_1$ is farnesylated, high receptor coupling efficiency was observed with the α_{2A} -adrenergic receptor [155], adenosine A_1 and 5-HT $_{1A}$ receptors [156] and the M_1 muscarinic receptor [119]. These results suggest that the differences between farnesylation and geranylgeranylation of $\beta\gamma$ dimers reflect more than degrees of hydrophobicity of a membrane anchor, and the identity of the prenyl group is intimately related to activation of a G protein, and hence $\beta\gamma$, by receptor.

γ Isoform Specificity

Thus, the primary sequence of γ is also critical for determining the efficiency of receptor:G protein interactions. This point was borne out in studies with chimeras of γ_1 and γ_2 , which concluded that the C-terminal third of γ , along with the type of prenyl group, is particularly important at determining the affinity of G protein receptor interactions [58, 153]. Studies with peptides constructed with the C-terminal sequence of a geranylgeranylated γ_5 subunit reached similar conclusions after these peptides were found to inhibit M_2 muscarinic receptor signaling [137]. The γ_5 C-terminal sequence apparently confers specificity in receptor:G protein interactions, as neither γ_7 nor γ_{12} C-terminal peptides were able to inhibit muscarinic receptor signaling [137]. A specific signaling role for γ_7 has however, been assigned to the β -adrenergic receptor using a ribozyme strategy to attenuate γ_7 mRNA and protein levels [157, 158]; the β_1 isoform was also suggested as the partner for γ_7 in this signaling cas-

cade, as β_1 protein levels fell in response to loss of γ_7 protein [158]. The ribozyme approach was also used to demonstrate that the β_1 and γ_7 isoforms are involved in activation of adenylyl cyclase via coupling to the D_1 dopamine receptor, but not the D_5 dopamine receptor [159]. The identity of γ also influences how a β isoform can interact with receptors. For example, the $\beta_1\gamma_5$ dimer couples poorly to the α_{2A} -adrenergic receptor, whereas the $\beta_3\gamma_5$ dimer couples effectively; however, when the γ isoform is changed to γ_{11} , the resulting $\beta_1\gamma_{11}$ dimer couples almost as well as $\beta_3\gamma_5$ [155], suggesting that the receptor coupling efficiency of β or γ isoforms can be interrelated.

β Isoform Specificity

The identity of the β isoform has also been shown to influence interactions between receptor and G proteins. Attenuation of specific β isoforms using RNAi has indicated that the β_2 isoform, but not β_1 , is involved in C5a-mediated chemotaxis in mouse macrophages [160]. In reconstitution experiments, the M_2 muscarinic receptor was more efficient in catalyzing nucleotide exchange with an $G_O \alpha\beta_4\gamma_2$ heterotrimer compared to a $G_O \alpha\beta_1\gamma_2$ heterotrimer [161]. The β_4 subunit as a dimer with γ_2 was also noted to have a greater ability to couple $G_s \alpha$ to the adenosine A_{2A} receptor than $\beta_1\gamma_2$ [98], and shift a larger percentage of A_{2A} receptors into the high affinity binding state than $\beta_1\gamma_2$ [162]. Interestingly, the inflammatory cytokines IL-1 and TNF- α were demonstrated to upregulate G β_4 , but not G β_1 mRNA levels in human dermal microvascular endothelial cells [163]. Figure 2 illustrates how this regulation of G β_4 , and the observed differences in adenosine A_{2A} receptor coupling between $\beta_1\gamma_2$ and $\beta_4\gamma_2$

may affect second messenger levels in cells. If the $\beta_1\gamma_2$ dimer predominates over $\beta_4\gamma_2$, $G_s \alpha$ may couple poorly to the adenosine A_{2A} receptor; however, an inflammatory stimulus could produce cytokines that elevate G_{β_4} levels, thus increasing the availability of $\beta_4\gamma_2$ and enhancing the ability of $G_s \alpha$ to couple to the adenosine A_{2A} receptor, activate adenylyl cyclase and raise cAMP levels.

Conclusion

In conclusion, in the scope of this review, there are multiple factors that determine the activation of a particular $\beta\gamma$ dimer within a cell: (1) Transcription, translation and posttranslational processing of specific β and γ isoforms, together with protein:protein interactions determine the constellation and potential localization of $\beta\gamma$ dimer combinations. (2) Expression of specific $G \alpha$ subunit isoforms, likely integrated to specific β and γ iso-

form expression, determines the identity of heterotrimer combinations, thus affecting interactions with receptors, and the degree to which a G protein may dissociate upon activation, and potentially reform with different $G \alpha$ isoforms. (3) Expression and activation of specific receptors, which through preferences for binding particular heterotrimer combinations, ultimately starts the signal which leads to activation of specific $\beta\gamma$ dimers. The crystallization of a receptor G protein complex would greatly enhance our knowledge of the specificity of receptor: α : $\beta\gamma$ interactions.

Acknowledgements

The author would like to thank Dr. James Garrison for comments helpful in the preparation of the manuscript. This study was supported in part by grants RO1-DK-19952 from the National Institutes of Health and 0535350N from the American Heart Association.

References

- International Human Genome Sequencing Consortium: Initial sequencing and analysis of the human genome. *Nature* 2001;409:860–921.
- Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, Yandell M, Evans CA, Holt RA, Gocayne JD, et al: The sequence of the human genome. *Science* 2001;291:1304–1351.
- Oldham WM, Hamm HE: Heterotrimeric G protein activation by G-protein-coupled receptors. *Nat Rev Mol Cell Biol* 2008;9:60–71.
- Berman DM, Gilman AG: Mammalian RGS proteins: barbarians at the gate. *J Biol Chem* 1998;273:1269–1272.
- Downes GB, Gautam N: The G protein subunit gene families. *Genomics* 1999;62:544–552.
- Hurowitz EH, Melnyk JM, Chen YJ, Kouros-Mehr H, Simon MI, Shizuya H: Genomic characterization of the human heterotrimeric G protein alpha, beta, and gamma subunit genes. *DNA Res* 2000;7:111–120.
- Roskopf D, Koch K, Habich C, Geerdes J, Ludwig A, Wilhelms S, Jakobs KH, Siffert W: Interaction of Gbeta3s, a splice variant of the G-protein Gbeta3, with Ggamma. *Cell Signal* 2003;15:479–488.
- Roskopf D, Manthey I, Habich C, Kielbik M, Eisenhardt A, Nikula C, Urban M, Kohnen S, Graf E, Ravens U, Siffert W: Identification and characterization of G beta 3s2, a novel splice variant of the G-protein beta 3 subunit. *Biochem J* 2003;371:223–232.
- Roskopf D, Kielbik M, Manthey I, Bilmen G, Eisenhardt A, Siffert W: Characterization of the splice variant Gbeta3v of the human G-protein Gbeta3 subunit. *Biochim Biophys Acta* 2003;1626:33–42.
- Watson AJ, Aragay AM, Slepak VZ, Simon MI: A novel form of the G protein beta subunit Gbeta5 is specifically expressed in the vertebrate retina. *J Biol Chem* 1996;271:28154–28160.
- Hildebrandt JD: Role of subunit diversity in signaling by heterotrimeric G proteins. *Biochem Pharmacol* 1997;54:325–339.
- Cook LA, Schey KL, Wilcox MD, Dingus J, Ettling R, Nelson T, Knapp DR, Hildebrandt JD: Proteomic analysis of bovine brain G protein gamma subunit processing heterogeneity. *Mol Cell Proteomics* 2006;5:671–685.
- Matsuda T, Takao T, Shimonishi Y, Murata M, Asano T, Yoshizawa T, Fukada Y: Characterization of interactions between transducin alpha/beta gamma-subunits and lipid membranes. *J Biol Chem* 1994;269:30358–30363.
- Wieland T, Nurnberg B, Ulibarri I, Kaldenberg-Stasch S, Schultz G, Jakobs KH: Guanine nucleotide-specific phosphate transfer by guanine nucleotide-binding regulatory protein beta-subunits. Characterization of the phosphorylated amino acid. *J Biol Chem* 1993;268:18111–18118.
- Cuello F, Schulze RA, Heemeyer F, Meyer HE, Lutz S, Jakobs KH, Niroomand F, Wieland T: Activation of heterotrimeric G proteins by a high energy phosphate transfer via nucleoside diphosphate kinase (NDPK) B and Gbeta subunits. Complex formation of NDPK B with Gbeta gamma dimers and phosphorylation of His-266 IN Gbeta. *J Biol Chem* 2003;278:7220–7226.
- Lupi R, Corda D, Di Girolamo M: Endogenous ADP-ribosylation of the G protein beta subunit prevents the inhibition of type 1 adenylyl cyclase. *J Biol Chem* 2000;275:9418–9424.
- Lupi R, Dani N, Dietrich A, Marchegiani A, Turacchio S, Berrie CP, Moss J, Gierschik P, Corda D, Di Girolamo M: Endogenous mono-ADP-ribosylation of the free Gbetagamma prevents stimulation of phosphoinositide 3-kinase-gamma and phospholipase C-beta2 and is activated by G-protein-coupled receptors. *Biochem J* 2002;367:825–832.
- Zhang FL, Casey PJ: Protein prenylation: molecular mechanisms and functional consequences. *Annu Rev Biochem* 1996;65:241–269.
- Lai RK, Perez-Sala D, Canada FJ, Rando RR: The gamma subunit of transducin is farnesylated. *Proc Natl Acad Sci USA* 1990;87:7673–7677.

- 20 Yamane HK, Farnsworth CC, Xie HY, Howald W, Fung BK, Clarke S, Gelb MH, Glomset JA: Brain G protein gamma subunits contain an all-*trans*-geranylgeranyl cysteine methyl ester at their carboxyl termini. *Proc Natl Acad Sci USA* 1990;87:5868–5872.
- 21 Mumby SM, Casey PJ, Gilman AG, Gutowski S, Sternweis PC: G protein gamma subunits contain a 20-carbon isoprenoid. *Proc Natl Acad Sci USA* 1990;87:5873–5877.
- 22 Clarke S: Protein isoprenylation and methylation at carboxyl-terminal cysteine residues. *Annu Rev Biochem* 1992;61:355–386.
- 23 Fukada Y, Matsuda T, Kokame K, Takao T, Shimonishi Y, Akino T, Yoshizawa T: Effects of carboxyl methylation of photoreceptor G protein gamma subunit in visual transduction. *J Biol Chem* 1994;269:5163–5170.
- 24 Rehm A, Ploegh HL: Assembly and intracellular targeting of the betagamma subunits of heterotrimeric G proteins. *J Cell Biol* 1997;137:305–317.
- 25 Simonds WF, Butrynski JE, Gautam N, Unson CG, Spiegel AM: G-protein beta gamma dimers. Membrane targeting requires subunit coexpression and intact gamma C-A-A-X domain. *J Biol Chem* 1991;266:5363–5366.
- 26 Takida S, Wedegaertner PB: Heterotrimer formation, together with isoprenylation, is required for plasma membrane targeting of Gbeta gamma. *J Biol Chem* 2003;278:17284–17290.
- 27 Hynes TR, Tang L, Mervine SM, Sabo JL, Yost EA, Devreotes PN, Berlot CH: Visualization of G protein betagamma dimers using bimolecular fluorescence complementation demonstrates roles for both beta and gamma in subcellular targeting. *J Biol Chem* 2004;279:30279–30286.
- 28 Higgins JB, Casey PJ: In vitro processing of recombinant G protein gamma subunits. Requirements for assembly of an active beta gamma complex. *J Biol Chem* 1994;269:9067–9073.
- 29 Cook LA, Schey KL, Wilcox MD, Dingus J, Hildebrandt JD: Heterogeneous processing of a G protein gamma subunit at a site critical for protein and membrane interactions. *Biochemistry* 1998;37:12280–12286.
- 30 Kilpatrick EL, Hildebrandt JD: Sequence dependence and differential expression of Ggamma5 subunit isoforms of the heterotrimeric G proteins variably processed after prenylation in mammalian cells. *J Biol Chem* 2007;282:14038–14047.
- 31 Li Z, Benard O, Margolskee RF: Ggamma13 interacts with PDZ domain-containing proteins. *J Biol Chem* 2006;281:11066–11073.
- 32 Kino T, Tiulpakov A, Ichijo T, Chheng L, Kozasa T, Chrousos GP: G protein beta interacts with the glucocorticoid receptor and suppresses its transcriptional activity in the nucleus. *J Cell Biol* 2005;169:885–896.
- 33 Kino T, Kozasa T, Chrousos GP: Statin-induced blockade of prenylation alters nucleocytoplasmic shuttling of GTP-binding proteins gamma2 and beta2 and enhances their suppressive effect on glucocorticoid receptor transcriptional activity. *Eur J Clin Invest* 2005;35:508–513.
- 34 Park JG, Muise A, He GP, Kim SW, Ro HS: Transcriptional regulation by the gamma5 subunit of a heterotrimeric G protein during adipogenesis. *EMBO J* 1999;18:4004–4012.
- 35 Ovchinnikov I, Lipkin VM, Telezhinskaya IN, Shuvaeva TM, Obukhov AN: Primary structure of the gamma-subunit of the GTP-binding protein from the bovine retina (in Russian). *Bioorg Khim* 1985;11:1301–1314.
- 36 Morishita R, Kato K, Asano T: A brain-specific gamma subunit of G protein freed from the corresponding beta subunit under non-denaturing conditions. *FEBS Lett* 1994;337:23–26.
- 37 Morishita R, Ueda H, Kato K, Asano T: Identification of two forms of the gamma subunit of G protein, gamma10 and gamma11, in bovine lung and their tissue distribution in the rat. *FEBS Lett* 1998;428:85–88.
- 38 Wilcox MD, Schey KL, Busman M, Hildebrandt JD: Determination of the complete covalent structure of the gamma 2 subunit of bovine brain G proteins by mass spectrometry. *Biochem Biophys Res Commun* 1995;212:367–374.
- 39 Hamilton MH, Cook LA, McRackan TR, Schey KL, Hildebrandt JD: Gamma 2 subunit of G protein heterotrimer is an N-end rule ubiquitylation substrate. *Proc Natl Acad Sci USA* 2003;100:5081–5086.
- 40 Morishita R, Nakayama H, Isobe T, Matsuda T, Hashimoto Y, Okano T, Fukada Y, Mizuno K, Ohno S, Kozawa O, et al: Primary structure of a gamma subunit of G protein, gamma 12, and its phosphorylation by protein kinase C. *J Biol Chem* 1995;270:29469–29475.
- 41 Yasuda H, Lindorfer MA, Myung CS, Garrison JC: Phosphorylation of the G protein gamma12 subunit regulates effector specificity. *J Biol Chem* 1998;273:21958–21965.
- 42 Asano T, Morishita R, Ueda H, Asano M, Kato K: GTP-binding protein gamma12 subunit phosphorylation by protein kinase C – identification of the phosphorylation site and factors involved in cultured cells and rat tissues in vivo. *Eur J Biochem* 1998;251:314–319.
- 43 Garcia-Higuera I, Fenoglio J, Li Y, Lewis C, Panchenko MP, Reiner O, Smith TF, Neer EJ: Folding of proteins with WD-repeats: comparison of six members of the WD-repeat superfamily to the G protein beta subunit. *Biochemistry* 1996;35:13985–13994.
- 44 Sondek J, Bohm A, Lambright DG, Hamm HE, Sigler PB: Crystal structure of a G-protein beta gamma dimer at 2.1 Å resolution. *Nature* 1996;379:369–374; published erratum appears in *Nature* 1996;379:847.
- 45 Wall MA, Coleman DE, Lee E, Iniguez-Lluhi JA, Posner BA, Gilman AG, Sprang SR: The structure of the G protein heterotrimer G_i alpha 1 beta 1 gamma 2. *Cell* 1995;83:1047–1058.
- 46 Lambright DG, Sondek J, Bohm A, Skiba NP, Hamm HE, Sigler PB: The 2.0 Å crystal structure of a heterotrimeric G protein. *Nature* 1996;379:311–319.
- 47 Garritsen A, van Galen PJ, Simonds WF: The N-terminal coiled-coil domain of beta is essential for gamma association: a model for G-protein beta gamma subunit interaction. *Proc Natl Acad Sci USA* 1993;90:7706–7710.
- 48 Cheever ML, Snyder JT, Gershburg S, Siderovski DP, Harden TK, Sondek J: Crystal structure of the multifunctional Gbeta5-RGS9 complex. *Nat Struct Mol Biol* 2008;15:155–162.
- 49 Lambright DG, Noel JP, Hamm HE, Sigler PB: Structural determinants for activation of the alpha-subunit of a heterotrimeric G protein. *Nature* 1994;369:621–628.
- 50 Noel JP, Hamm HE, Sigler PB: The 2.2 Å crystal structure of transducin-alpha complexed with GTP gamma S. *Nature* 1993;366:654–663.
- 51 Sprang SR: G protein mechanisms: insights from structural analysis. *Annu Rev Biochem* 1997;66:639–678.
- 52 Katada T, Bokoch GM, Northup JK, Ui M, Gilman AG: The inhibitory guanine nucleotide-binding regulatory component of adenylate cyclase. Properties and function of the purified protein. *J Biol Chem* 1984;259:3568–3577.
- 53 Loew A, Ho YK, Blundell T, Bax B: Phosducin induces a structural change in transducin beta gamma. *Structure* 1998;6:1007–1019.
- 54 Hamm HE, Deretic D, Arendt A, Hargrave PA, Koenig B, Hofmann KP: Site of G protein binding to rhodopsin mapped with synthetic peptides from the alpha subunit. *Science* 1988;241:832–835.
- 55 Martin EL, Rens-Domiano S, Schatz PJ, Hamm HE: Potent peptide analogues of a G protein receptor-binding region obtained with a combinatorial library. *J Biol Chem* 1996;271:361–366.
- 56 Kisselev OG, Ermolaeva MV, Gautam N: A farnesylated domain in the G protein gamma subunit is a specific determinant of receptor coupling. *J Biol Chem* 1994;269:21399–21402.
- 57 Kisselev OG, Downs MA: Rhodopsin-interacting surface of the transducin gamma subunit. *Biochemistry* 2006;45:9386–9392.
- 58 Myung CS, Lim WK, Defilippo J, Yasuda H, Neubig R, Garrison JC: Regions in the G protein gamma subunit important for interaction with receptors and effectors. *Mol Pharmacol* 2005;69:877–887.

- 59 Taylor JM, Jacob-Mosier GG, Lawton RG, VanDort M, Neubig RR: Receptor and membrane interaction sites on Gbeta. A receptor-derived peptide binds to the carboxyl terminus. *J Biol Chem* 1996;271:3336–3339.
- 60 Onrust R, Herzmark P, Chi P, Garcia PD, Lichtarge O, Kingsley C, Bourne HR: Receptor and betagamma binding sites in the alpha subunit of the retinal G protein transducin. *Science* 1997;275:381–384; published erratum appears in *Science* 1997;276:341.
- 61 Iiri T, Farfel Z, Bourne HR: G-protein diseases furnish a model for the turn-on switch. *Nature* 1998;394:35–38.
- 62 Lichtarge O, Bourne HR, Cohen FE: Evolutionarily conserved Galphabeta binding surfaces support a model of the G protein-receptor complex. *Proc Natl Acad Sci USA* 1996;93:7507–7511.
- 63 Rondard P, Iiri T, Srinivasan S, Meng E, Fujita T, Bourne HR: Mutant G protein alpha subunit activated by Gbeta gamma: a model for receptor activation? *Proc Natl Acad Sci USA* 2001;98:6150–6155.
- 64 Cherfils J, Chabre M: Activation of G-protein Galpha subunits by receptors through Galpha-Gbeta and Galpha-Ggamma interactions. *Trends Biochem Sci* 2003;28:13–17.
- 65 Kisselev OG, Meyer CK, Heck M, Ernst OP, Hofmann KP: Signal transfer from rhodopsin to the G-protein: evidence for a two-site sequential fit mechanism. *Proc Natl Acad Sci USA* 1999;96:4898–4903.
- 66 Herrmann R, Heck M, Henklein P, Henklein P, Kleuss C, Hofmann KP, Ernst OP: Sequence of interactions in receptor-G protein coupling. *J Biol Chem* 2004;279:24283–24290.
- 67 Kisselev O, Pronin A, Ermolaeva M, Gautam N: Receptor-G protein coupling is established by a potential conformational switch in the beta gamma complex. *Proc Natl Acad Sci USA* 1995;92:9102–9106.
- 68 Gautam N: A conformational switch regulates receptor-G protein interaction. *Structure* 2003;11:359–360.
- 69 Kisselev OG, Downs MA: Rhodopsin controls a conformational switch on the transducin gamma subunit. *Structure (Camb)* 2003;11:367–373.
- 70 Abdulaev NG, Ngo T, Ramon E, Brabazon DM, Marino JP, Ridge KD: The receptor-bound 'empty pocket' state of the heterotrimeric G-protein alpha-subunit is conformationally dynamic. *Biochemistry* 2006;45:12986–12997.
- 71 Oldham WM, Van Eps N, Preininger AM, Hubbell WL, Hamm HE: Mapping allosteric connections from the receptor to the nucleotide-binding pocket of heterotrimeric G proteins. *Proc Natl Acad Sci USA* 2007;104:7927–7932.
- 72 Fung BK: Characterization of transducin from bovine retinal rod outer segments. I. Separation and reconstitution of the subunits. *J Biol Chem* 1983;258:10495–10502.
- 73 Codina J, Hildebrandt JD, Birnbaumer L, Sekura RD: Effects of guanine nucleotides and Mg on human erythrocyte Ni and Ns, the regulatory components of adenylyl cyclase. *J Biol Chem* 1984;259:11408–11418.
- 74 Northup JK, Smigel MD, Sternweis PC, Gilman AG: The subunits of the stimulatory regulatory component of adenylyl cyclase. Resolution of the activated 45,000-dalton (alpha) subunit. *J Biol Chem* 1983;258:11369–11376.
- 75 Ford CE, Skiba NP, Bae H, Daaka Y, Reuveny E, Shekter LR, Rosal R, Weng G, Yang CS, Iyengar R, Miller RJ, Jan LY, Lefkowitz RJ, Hamm HE: Molecular basis for interactions of G protein betagamma subunits with effectors. *Science* 1998;280:1271–1274.
- 76 Li Y, Sternweis PM, Charnecki S, Smith TF, Gilman AG, Neer EJ, Kozasa T: Sites for Galpha binding on the G protein beta subunit overlap with sites for regulation of phospholipase Cbeta and adenylyl cyclase. *J Biol Chem* 1998;273:16265–16272.
- 77 Bonacci TM, Mathews JL, Yuan C, Lehmann DM, Malik S, Wu D, Font JL, Bidlack JM, Smrcka AV: Differential targeting of Gbetagamma-subunit signaling with small molecules. *Science* 2006;312:443–446.
- 78 Lehmann DM, Seneviratne AM, Smrcka AV: Small molecule disruption of G protein beta gamma subunit signaling inhibits neutrophil chemotaxis and inflammation. *Mol Pharmacol* 2008;73:410–418.
- 79 Lutz S, Shankaranarayanan A, Coco C, Ridilla M, Nance MR, Vettel C, Baltus D, Evelyn CR, Neubig RR, Wieland T, Tesmer JJ: Structure of Galphaq-p63RhoGEF-RhoA complex reveals a pathway for the activation of RhoA by GPCRs. *Science* 2007;318:1923–1927.
- 80 Tesmer JJ, Sunahara RK, Gilman AG, Sprang SR: Crystal structure of the catalytic domains of adenylyl cyclase in a complex with Galpha.GTPgammaS. *Science* 1997;278:1907–1916.
- 81 Levitzki A, Klein S: G-protein subunit dissociation is not an integral part of G-protein action. *Chembiochem* 2002;3:815–818.
- 82 Klein S, Reuveni H, Levitzki A: Signal transduction by a nondissociable heterotrimeric yeast G protein. *Proc Natl Acad Sci USA* 2000;97:3219–3223.
- 83 Janetopoulos C, Jin T, Devreotes P: Receptor-mediated activation of heterotrimeric G-proteins in living cells. *Science* 2001;291:2408–2411.
- 84 Bunemann M, Frank M, Lohse MJ: Gi protein activation in intact cells involves subunit rearrangement rather than dissociation. *Proc Natl Acad Sci USA* 2003;100:16077–16082.
- 85 Frank M, Thumer L, Lohse MJ, Bunemann M: G Protein activation without subunit dissociation depends on a G-alpha-i-specific region. *J Biol Chem* 2005;280:24584–24590.
- 86 Digby GJ, Lober RM, Sethi PR, Lambert NA: Some G protein heterotrimers physically dissociate in living cells. *Proc Natl Acad Sci USA* 2006;103:17789–17794.
- 87 Rebois RV, Robitaille M, Gales C, Dupre DJ, Baragli A, Trieu P, Ethier N, Bouvier M, Herbert TE: Heterotrimeric G proteins form stable complexes with adenylyl cyclase and Kir3.1 channels in living cells. *J Cell Sci* 2006;119:2807–2818.
- 88 Evanko DS, Thiyagarajan MM, Takida S, Wedegaertner PB: Loss of association between activated Galpha q and Gbetagamma disrupts receptor-dependent and receptor-independent signaling. *Cell Signal* 2005;17:1218–1228.
- 89 Linder ME, Pang IH, Duronio RJ, Gordon JI, Sternweis PC, Gilman AG: Lipid modifications of G protein subunits. Myristoylation of Go alpha increases its affinity for beta gamma. *J Biol Chem* 1991;266:4654–4659.
- 90 Azpiazu I, Gautam N: A fluorescence resonance energy transfer-based sensor indicates that receptor access to a G protein is unrestricted in a living mammalian cell. *J Biol Chem* 2004;279:27709–27718.
- 91 Yan K, Kalyanaraman V, Gautam N: Differential ability to form the G protein betagamma complex among members of the beta and gamma subunit families. *J Biol Chem* 1996;271:7141–7146.
- 92 Schmidt CJ, Thomas TC, Levine MA, Neer EJ: Specificity of G protein beta and gamma subunit interactions. *J Biol Chem* 1992;267:13807–13810.
- 93 Ray K, Kunsch C, Bonner LM, Robishaw JD: Isolation of cDNA clones encoding eight different human G protein gamma subunits, including three novel forms designated the gamma 4, gamma 10, and gamma 11 subunits. *J Biol Chem* 1995;270:21765–21771.
- 94 Dings J, Wells CA, Campbell L, Cleator JH, Robinson K, Hildebrandt JD: G Protein betagamma dimer formation: Gbeta and Ggamma differentially determine efficiency of in vitro dimer formation. *Biochemistry* 2005;44:11882–11890.
- 95 Pronin AN, Gautam N: Interaction between G-protein beta and gamma subunit types is selective. *Proc Natl Acad Sci USA* 1992;89:6220–6224.
- 96 Lee C, Murakami T, Simonds WF: Identification of a discrete region of the G protein gamma subunit conferring selectivity in beta gamma complex formation. *J Biol Chem* 1995;270:8779–8784.
- 97 Mervine SM, Yost EA, Sabo JL, Hynes TR, Berlot CH: Analysis of G protein betagamma dimer formation in live cells using multicolor bimolecular fluorescence complementation demonstrates preferences of beta1 for particular gamma subunits. *Mol Pharmacol* 2006;70:194–205.

- 98 McIntire WE, MacCleery G, Garrison JC: The G protein beta subunit is a determinant in the coupling of Gs to the beta-1-adrenergic and A_{2a} adenosine receptors. *J Biol Chem* 2001;276:15801–15809.
- 99 Siffert W: G protein polymorphisms in hypertension, atherosclerosis, and diabetes. *Annu Rev Med* 2005;56:17–28.
- 100 Roszkopf D, Nikula C, Manthey I, Joisten M, Frey U, Kohnen S, Siffert W: The human G protein beta4 subunit: gene structure, expression, Ggamma and effector interaction. *FEBS Lett* 2003;544:27–32.
- 101 Asano T, Morishita R, Ueda H, Kato K: Selective association of G protein beta 4 with gamma 5 and gamma 12 subunits in bovine tissues. *J Biol Chem* 1999;274:21425–21429.
- 102 Wells CA, Dingus J, Hildebrandt JD: Role of the chaperonin CCT/TRiC complex in G protein betagamma-dimer assembly. *J Biol Chem* 2006;281:20221–20232.
- 103 Dupre DJ, Robitaille M, Richer M, Ethier N, Mamrabachi AM, Hebert TE: Dopamine receptor-interacting protein 78 acts as a molecular chaperone for Ggamma subunits before assembly with Gbeta. *J Biol Chem* 2007;282:13703–13715.
- 104 Willardson BM, Howlett AC: Function of phosphatidylinositol-3-OH kinase-like proteins in G protein signaling and chaperone-assisted protein folding. *Cell Signal* 2007;19:2417–2427.
- 105 Jones MB, Garrison JC: Instability of the G-protein beta5 subunit in detergent. *Anal Biochem* 1999;268:126–133.
- 106 De Vries L, Zheng B, Fischer T, Elenko E, Farquhar MG: The regulator of G protein signaling family. *Annu Rev Pharmacol Toxicol* 2000;40:235–271.
- 107 Yost EA, Mervine SM, Sabo JL, Hynes TR, Berlot CH: Live cell analysis of G protein beta5 complex formation, function, and targeting. *Mol Pharmacol* 2007;72:812–825.
- 108 Gilman AG: Nobel Lecture. G proteins and regulation of adenylyl cyclase. *Biosci Rep* 1995;15:65–97.
- 109 Huang L, Shanker YG, Dubauskaite J, Zheng JZ, Yan W, Rosenzweig S, Spielman AI, Max M, Margolskee RF: Ggamma13 colocalizes with gustducin in taste receptor cells and mediates IP₃ responses to bitter denatonium. *Nat Neurosci* 1999;2:1055–1062.
- 110 Ruiz-Velasco V, Ikeda SR, Puhl HL: Cloning, tissue distribution, and functional expression of the human G protein beta4 subunit. *Physiol Genomics* 2002;8:41–50.
- 111 Wilcox MD, Dingus J, Balcueva EA, McIntire WE, Mehta ND, Schey KL, Robishaw JD, Hildebrandt JD: Bovine brain G_o isoforms have distinct gamma subunit compositions. *J Biol Chem* 1995;270:4189–4192.
- 112 McIntire WE, Schey KL, Knapp DR, Hildebrandt JD: A major G protein alpha O isoform in bovine brain is deamidated at Asn346 and Asn347, residues involved in receptor coupling. *Biochemistry* 1998;37:14651–14658.
- 113 McIntire WE, Dingus J, Wilcox MD, Hildebrandt JD: The relationship of G(o)alpha subunit deamidation to the tissue distribution, nucleotide binding properties, and betagamma dimer interactions of G(o)alpha subunit isoforms. *J Neurochem* 1999;73:633–640.
- 114 Krumins AM, Gilman AG: Targeted knockdown of G protein subunits selectively prevents receptor-mediated modulation of effectors and reveals complex changes in non-targeted signaling proteins. *J Biol Chem* 2006;281:10250–10262.
- 115 Schwindinger WF, Betz KS, Giger KE, Sabol A, Bronson SK, Robishaw JD: Loss of G protein gamma 7 alters behavior and reduces striatal alpha(olf) level and cAMP production. *J Biol Chem* 2003;278:6575–6579.
- 116 Lobanova ES, Finkelstein S, Herrmann R, Chen YM, Kessler C, Michaud NA, Trieu LH, Strissel KJ, Burns ME, Arshavsky VY: Transducin gamma-subunit sets expression levels of alpha- and beta-subunits and is crucial for rod viability. *J Neurosci* 2008;28:3510–3520.
- 117 Evanko DS, Thiyagarajan MM, Siderovski DP, Wedegaertner PB: Gbeta gamma isoforms selectively rescue plasma membrane localization and palmitoylation of mutant Galphas and Galphaq. *J Biol Chem* 2001;276:23945–23953.
- 118 Fletcher JE, Lindorfer MA, DeFilippo JM, Yasuda H, Guilford M, Garrison JC: The G protein beta5 subunit interacts selectively with the Gq alpha subunit. *J Biol Chem* 1998;273:636–644.
- 119 McIntire WE, MacCleery G, Murphree LJ, Kerchner KR, Linden J, Garrison JC: Influence of differential stability of G protein betagamma dimers containing the gamma11 subunit on functional activity at the M1 muscarinic receptor, A1 adenosine receptor, and phospholipase C-beta. *Biochemistry* 2006;45:11616–11631.
- 120 Lindorfer MA, Myung CS, Savino Y, Yasuda H, Khazan R, Garrison JC: Differential activity of the G protein beta5 gamma2 subunit at receptors and effectors. *J Biol Chem* 1998;273:34429–34436.
- 121 Kageyama K, Murakami T, Iizuka K, Sato K, Ichihara K, Tokumitsu Y, Kitabatake A, Kawaguchi H: Translocation of G-protein beta3 subunit from the cytosol pool to the membrane pool by beta1-adrenergic receptor stimulation in perfused rat hearts. *Biochem Pharmacol* 1999;58:1497–1500.
- 122 Akgoz M, Kalyanaraman V, Gautam N: Receptor-mediated reversible translocation of the G protein betagamma complex from the plasma membrane to the Golgi complex. *J Biol Chem* 2004;279:51541–51544.
- 123 Saini DK, Kalyanaraman V, Chisari M, Gautam N: A family of G protein betagamma subunits translocate reversibly from the plasma membrane to endomembranes on receptor activation. *J Biol Chem* 2007;282:24099–24108.
- 124 Azpiazu I, Akgoz M, Kalyanaraman V, Gautam N: G protein betagamma11 complex translocation is induced by Gi, Gq and Gs coupling receptors and is regulated by the alpha subunit type. *Cell Signal* 2006;18:1190–1200.
- 125 Chisari M, Saini DK, Kalyanaraman V, Gautam N: Shuttling of G protein subunits between the plasma membrane and intracellular membranes. *J Biol Chem* 2007;282:24092–24098.
- 126 Sato M, Cismowski MJ, Toyota E, Smrcka AV, Lucchesi PA, Chilian WM, Lanier SM: Identification of a receptor-independent activator of G protein signaling (AGS8) in ischemic heart and its interaction with Gbetagamma. *Proc Natl Acad Sci USA* 2006;103:797–802.
- 127 Yuan C, Sato M, Lanier SM, Smrcka AV: Signaling by a non-dissociated complex of G protein betagamma and alpha subunits stimulated by a receptor-independent activator of G protein signaling, AGS8. *J Biol Chem* 2007;282:19938–19947.
- 128 Zeng W, Mak DO, Li Q, Shin DM, Foscett JK, Muallem S: A new mode of Ca²⁺ signaling by G protein-coupled receptors: gating of IP₃ receptor Ca²⁺ release channels by Gbetagamma. *Curr Biol* 2003;13:872–876.
- 129 Wang S, Narendra S, Fedoroff N: Heterotrimeric G protein signaling in the Arabidopsis unfolded protein response. *Proc Natl Acad Sci USA* 2007;104:3817–3822.
- 130 Dell EJ, Connor J, Chen S, Stebbins EG, Skiba NP, Mochly-Rosen D, Hamm HE: The betagamma subunit of heterotrimeric G proteins interacts with RACK1 and two other WD repeat proteins. *J Biol Chem* 2002;277:49888–49895.
- 131 Patterson RL, van Rossum DB, Barrow RK, Snyder SH: RACK1 binds to inositol 1,4,5-trisphosphate receptors and mediates Ca²⁺ release. *Proc Natl Acad Sci USA* 2004;101:2328–2332.
- 132 Chen S, Lin F, Hamm HE: RACK1 binds to a signal transfer region of G betagamma and inhibits phospholipase C beta2 activation. *J Biol Chem* 2005;280:33445–33452.
- 133 Schmidt CJ, Neer EJ: In vitro synthesis of G protein beta gamma dimers. *J Biol Chem* 1991;266:4538–4544.
- 134 Yoshikawa DM, Hatwar M, Smrcka AV: G protein beta 5 subunit interactions with alpha subunits and effectors. *Biochemistry* 2000;39:11340–11347.

- 135 Blake BL, Wing MR, Zhou JY, Lei Q, Hillmann JR, Behe CI, Morris RA, Harden TK, Bayliss DA, Miller RJ, Siderovski DP: Gbeta association and effector interaction selectivities of the divergent G gamma subunit G gamma 13. *J Biol Chem* 2001;276:49267-49274.
- 136 Rahmatullah M, Robishaw JD: Direct interaction of the alpha and gamma subunits of the G proteins. Purification and analysis by limited proteolysis. *J Biol Chem* 1994; 269:3574-3580.
- 137 Azpiazu I, Cruzblanca H, Li P, Linder M, Zhuo M, Gautam N: A G protein gamma subunit-specific peptide inhibits muscarinic receptor signaling. *J Biol Chem* 1999; 274:35305-35308.
- 138 Azpiazu I, Gautam N: A G protein gamma subunit peptide stabilizes a novel muscarinic receptor state. *Biochem Biophys Res Commun* 2006;341:904-910.
- 139 Hagiwara K, Wada A, Katadae M, Ito M, Ohya Y, Casey PJ, Fukada Y: Analysis of the molecular interaction of the farnesyl moiety of transducin through the use of a photoreactive farnesyl analogue. *Biochemistry* 2004;43:300-309.
- 140 Hou Y, Azpiazu I, Smrcka A, Gautam N: Selective role of G protein gamma subunits in receptor interaction. *J Biol Chem* 2000;275: 38961-38964.
- 141 Mende U, Schmidt CJ, Yi F, Spring DJ, Neer EJ: The G protein gamma subunit. Requirements for dimerization with beta subunits. *J Biol Chem* 1995;270:15892-15898.
- 142 Landry S, Hoffman CS: The git5 Gbeta and git11 Ggamma form an atypical Gbetagamma dimer acting in the fission yeast glucose/cAMP pathway. *Genetics* 2001;157: 1159-1168.
- 143 Chinault SL, Blumer KJ: The C-terminal tail preceding the CAAX box of a yeast G protein gamma subunit is dispensable for receptor-mediated G protein activation in vivo. *J Biol Chem* 2003;278:20638-20644.
- 144 Zeller CE, Parnell SC, Dohlman HG: The RACK1 ortholog Ascl functions as a G-protein beta subunit coupled to glucose responsiveness in yeast. *J Biol Chem* 2007; 282:25168-25176.
- 145 Yamazaki A, Tatsumi M, Torney DC, Bitensky MW: The GTP-binding protein of rod outer segments. I. Role of each subunit in the GTP hydrolytic cycle. *J Biol Chem* 1987; 262:9316-9323.
- 146 Kleuss C, Hescheler J, Ewel C, Rosenthal W, Schultz G, Wittig B: Assignment of G-protein subtypes to specific receptors inducing inhibition of calcium currents. *Nature* 1991;353:43-48.
- 147 Kleuss C, Scherubl H, Hescheler J, Schultz G, Wittig B: Different beta-subunits determine G-protein interaction with transmembrane receptors. *Nature* 1992;358: 424-426.
- 148 Kleuss C, Scherubl H, Hescheler J, Schultz G, Wittig B: Selectivity in signal transduction determined by gamma subunits of heterotrimeric G proteins. *Science* 1993;259: 832-834.
- 149 Macrez-Lepretre N, Kalkbrenner F, Morel JL, Schultz G, Mironneau J: G protein heterotrimer Galpha13beta1gamma3 couples the angiotensin AT1A receptor to increases in cytoplasmic Ca²⁺ in rat portal vein myocytes. *J Biol Chem* 1997;272:10095-10102.
- 150 Macrez N, Morel JL, Mironneau J: Specific galpha11beta3gamma5 protein involvement in endothelin receptor-induced phosphatidylinositol hydrolysis and Ca²⁺ release in rat portal vein myocytes. *Mol Pharmacol* 1999;55:684-692.
- 151 Kisselev O, Gautam N: Specific interaction with rhodopsin is dependent on the gamma subunit type in a G protein. *J Biol Chem* 1993;268:24519-24522.
- 152 Kisselev O, Ermolaeva M, Gautam N: Efficient interaction with a receptor requires a specific type of prenyl group on the G protein gamma subunit. *J Biol Chem* 1995;270: 25356-25358.
- 153 Jian X, Clark WA, Kowalak J, Markey SP, Simonds WF, Northup JK: Gbeta gamma affinity for bovine rhodopsin is determined by the carboxyl-terminal sequences of the gamma subunit. *J Biol Chem* 2001;276: 48518-48525.
- 154 Yasuda H, Lindorfer MA, Woodfork KA, Fletcher JE, Garrison JC: Role of the prenyl group on the G protein gamma subunit in coupling trimeric G proteins to A1 adenosine receptors. *J Biol Chem* 1996;271: 18588-18595.
- 155 Richardson M, Robishaw JD: The alpha-2A-adrenergic receptor discriminates between Gi heterotrimers of different beta-gamma subunit composition in Sf9 insect cell membranes. *J Biol Chem* 1999;274: 13525-13533.
- 156 Lim WK, Myung CS, Garrison JC, Neubig RR: Receptor-G protein gamma specificity: gamma11 shows unique potency for A₁ adenosine and 5-HT_{1A} receptors. *Biochemistry* 2001;40:10532-10541.
- 157 Wang Q, Mullah B, Hansen C, Asundi J, Robishaw JD: Ribozyme-mediated suppression of the G protein gamma7 subunit suggests a role in hormone regulation of adenylyl cyclase activity. *J Biol Chem* 1997; 272:26040-26048.
- 158 Wang Q, Mullah BK, Robishaw JD: Ribozyme approach identifies a functional association between the G protein beta1gamma7 subunits in the beta-adrenergic receptor signaling pathway. *J Biol Chem* 1999;274:17365-17371.
- 159 Wang Q, Jolly JP, Surmeier JD, Mullah BM, Lidow MS, Bergson CM, Robishaw JD: Differential dependence of the D1 and D5 dopamine receptors on the G protein gamma 7 subunit for activation of adenylylcyclase. *J Biol Chem* 2001;276:39386-39393.
- 160 Hwang JI, Fraser ID, Choi S, Qin XF, Simon MI: Analysis of C5a-mediated chemotaxis by lentiviral delivery of small interfering RNA. *Proc Natl Acad Sci USA* 2004;101: 488-493.
- 161 Hou Y, Chang V, Capper AB, Taussig R, Gautam N: G Protein beta subunit types differentially interact with a muscarinic receptor but not adenylyl cyclase type II or phospholipase C-beta 2/3. *J Biol Chem* 2001;276:19982-19988.
- 162 Murphree LJ, Marshall MA, Rieger JM, MacDonald TL, Linden J: Human A_{2A} adenosine receptors: high-affinity agonist binding to receptor-G protein complexes containing Gbeta4. *Mol Pharmacol* 2002; 61:455-462.
- 163 Nguyen DK, Montesinos MC, Williams AJ, Kelly M, Cronstein BN: Th1 cytokines regulate adenosine receptors and their downstream signaling elements in human microvascular endothelial cells. *J Immunol* 2003;171:3991-3998.