

RESEARCH

Open Access

Nitric oxide synthase modulates CFA-induced thermal hyperalgesia through cytokine regulation in mice

Yong Chen^{1*}, Michael K Boettger¹, Andreas Reif², Angelika Schmitt², Nurcan Üçeyler¹, Claudia Sommer¹

Abstract

Background: Although it has been largely demonstrated that nitric oxide synthase (NOS), a key enzyme for nitric oxide (NO) production, modulates inflammatory pain, the molecular mechanisms underlying these effects remain to be clarified. Here we asked whether cytokines, which have well-described roles in inflammatory pain, are downstream targets of NO in inflammatory pain and which of the isoforms of NOS are involved in this process.

Results: Intraperitoneal (i.p.) pretreatment with 7-nitroindazole sodium salt (7-NINA, a selective neuronal NOS inhibitor), aminoguanidine hydrochloride (AG, a selective inducible NOS inhibitor), L-N(G)-nitroarginine methyl ester (L-NAME, a non-selective NOS inhibitor), but not L-N(5)-(1-iminoethyl)-ornithine (L-NIO, a selective endothelial NOS inhibitor), significantly attenuated thermal hyperalgesia induced by intraplantar (i.pl.) injection of complete Freund's adjuvant (CFA). Real-time reverse transcription-polymerase chain reaction (RT-PCR) revealed a significant increase of nNOS, iNOS, and eNOS gene expression, as well as tumor necrosis factor-alpha (TNF), interleukin-1 beta (IL-1 β), and interleukin-10 (IL-10) gene expression in plantar skin, following CFA. Pretreatment with the NOS inhibitors prevented the CFA-induced increase of the pro-inflammatory cytokines TNF and IL-1 β . The increase of the anti-inflammatory cytokine IL-10 was augmented in mice pretreated with 7-NINA or L-NAME, but reduced in mice receiving AG or L-NIO. NNOS-, iNOS- or eNOS-knockout (KO) mice had lower gene expression of TNF, IL-1 β , and IL-10 following CFA, overall corroborating the inhibitor data.

Conclusion: These findings lead us to propose that inhibition of NOS modulates inflammatory thermal hyperalgesia by regulating cytokine expression.

Background

Several lines of evidence indicate a role for nitric oxide (NO) as a mediator of inflammation [1,2]. NO, acting as an inter- and intracellular messenger molecule in the peripheral and central nervous system, also plays a pivotal role in the development and maintenance of hyperalgesia [3-6]. NO can be synthesized by three well-characterized isoforms of NO synthase (NOS): the constitutive neuronal NOS (nNOS), endothelial NOS (eNOS), and the inducible NOS (iNOS) [7-9]. The non-selective NOS inhibitor L-N(G)-nitroarginine methyl ester (L-NAME) reduces thermal hyperalgesia in inflammatory pain models [10-12]. Further studies suggested beneficial effects of the selective NOS

inhibitors in reducing inflammatory hyperalgesia, while the baseline nociceptive responses remained unaltered [11,13-18].

Inflammatory pain hypersensitivity is the consequence of alterations in transduction sensitivity of high threshold nociceptors [19], activity-dependent changes in the excitability of spinal neurons [20], and phenotypic changes in sensory neurons innervating the inflamed tissue [21]. These changes, both at the inflamed site and throughout the nervous system, are initiated by a complex pattern of chemical signals interacting with the sensory fiber terminals. These signals originate from infective agents, damaged host cells or activated immune cells. Pro- and anti-inflammatory cytokines are small regulatory proteins that are produced by white blood cells and a variety of other cells including those in the nervous system.

* Correspondence: Chen_Y@klinik.uni-wuerzburg.de

¹Department of Neurology, University of Würzburg, Josef-Schneider-Str 11, 97080 Würzburg, Germany

Inflammatory stimuli or tissue injuries stimulate the release of cytokines, which play an essential role in inflammatory pain. Pro-inflammatory cytokines, such as tumor necrosis factor (TNF) and interleukin-1 beta (IL-1 β), reduced thermal or mechanical pain thresholds upon intraplantar application [22-24]. Pro-inflammatory cytokine antagonists were further able to reduce hyperalgesia in inflammation models, indicating that the activation of pro-inflammatory cytokines is an important step in the generation of inflammatory pain [24,25]. To limit the deleterious consequences of prolonged action of pro-inflammatory cytokines, their release is followed by the release of anti-inflammatory cytokines, such as IL-4, IL-10, and IL-13, which inhibit the production and action of the pro-inflammatory cytokines and are anti-hyperalgesic [24]. Correlations between tissue levels of cytokines and pain and hyperalgesia have been described in a number of painful states [26,27]. Although cytokines have well-described roles in inflammatory pain, it is poorly understood what regulates their production and release.

It has been largely demonstrated that inhibition of NOS attenuates inflammatory pain [11,13-18], however, the molecular mechanisms underlying these effects remain to be clarified. NO is generated in significant concentrations at sites of inflammation in which multiple hyperalgesic inflammatory mediators, such as cytokines, prostaglandin E2 (PGE2), or serotonin, are also produced [3,28]. NO may facilitate the hyperalgesia induced by those mediators using the cAMP second messenger pathway and may also have an independent cGMP-dependent hyperalgesic effect. The literature pre-dominantly documents that pro-inflammatory cytokines stimulate the production of NO, suggesting that cytokines modulate pain by regulating the release of NO [28-34]. In contrast, the effect of NO on pro-inflammatory cytokines has rarely been examined. One study reported that human immunodeficiency virus-1 (HIV-1) envelope glycoprotein gp120 stimulates pro-inflammatory cytokine-mediated pain facilitation via activation of nNOS [35]. This finding raises the intriguing possibility that reduction of inflammatory hyperalgesia with NOS inhibitors may be caused, at least in part, by reduced production of pro-inflammatory cytokines. This led us to hypothesize that cytokines, including pro- and anti-inflammatory cytokines, may be involved in pain modulation by NOS under inflammatory conditions. Here, we used a complete Freund's adjuvant (CFA)-induced inflammatory pain model in mice, to investigate whether the expression of cytokines is involved in the NOS-mediated inflammatory thermal hyperalgesia.

Results

Pretreatment with NOS inhibitors attenuates CFA-induced thermal hyperalgesia

Thermal pain thresholds were not different between groups at baseline, and not significantly changed after NS injections (Fig. 1). At 6, 16, and 24 h after i.pl. injection of CFA, significant thermal hyperalgesia was observed on the injected side (Fig. 1A).

Preemptive administration of 7-NINA, AG, or L-NAME, but not L-NIO, at a dose of 50 mg/kg, dramatically attenuated CFA-induced thermal hyperalgesia at 6, 16, and 24 h after injection (Fig 1A, $P < 0.001$). In mice receiving NS, none of the inhibitors affected pain thresholds throughout the observation period (Fig. 1B).

CFA increases both NOS and cytokine gene expression in plantar skin

The gene expression of nNOS (Fig. 2A) and eNOS (Fig. 2C) was elevated in the ipsilateral plantar skin at 6 h after CFA ($P < 0.001$), followed by a rapid decline to baseline levels at 16 and 24 h, compared to controls. iNOS gene expression was increased at 6 h and peaked at 24 h after CFA (Fig. 2B, $P < 0.01$ and $P < 0.001$).

As early as 6 h after CFA, TNF (Fig. 3A), IL-1 β (Fig. 3B), and IL-10 (Fig. 3C) gene expression in plantar skin was significantly increased compared to controls ($P < 0.001$) and remained elevated (with a decline for IL-10) until 24 h ($P < 0.01$ and $P < 0.001$). IL-1 β mRNA showed the largest increase of expression compared to control ($\times 2200\sim 3000$).

Pretreatment with the NOS inhibitors reduces the increase of TNF and IL-1 β gene expression and has a differential effect on the increase of IL-10 in plantar skin after CFA

Pretreatment with 7-NINA, AG, L-NIO, or L-NAME at a dose of 50 mg/kg did not significantly alter cytokine gene expression in plantar skin of control mice (data not shown). However, all inhibitors significantly attenuated the increase of TNF and IL-1 β in mice receiving CFA (Fig. 4A and 4B, $P < 0.05$, $P < 0.01$ and $P < 0.001$). The increase of IL-10 was augmented in mice pretreated with 7-NINA or L-NAME, but reduced in mice receiving AG or L-NIO, at 6 and 16 h after CFA (Fig. 4C, $P < 0.05$ and $P < 0.001$).

Cytokine gene expression in plantar skin is lower in NOS-KO mice after CFA compared to WT mice

Baseline gene expression of TNF was not different between nNOS-, iNOS- or eNOS-KO mice and WT mice (Fig. 5A). However, the baseline gene expression of IL-1 β was significantly higher (Fig. 5B; $P < 0.01$) and that of IL-10 lower (Fig. 5C; $P < 0.01$ and $P < 0.001$) in






Figure 1 Effects of pretreatment with the NOS inhibitors 7-NINA, AG, L-NIO and L-NAME on CFA-induced thermal hyperalgesia assessed by the paw withdrawal latency (PWL) tests. Following CFA injection, PWLs were markedly decreased within 6 h and continued until 24 h post injection in the ipsilateral hindpaw (A, $***P < 0.001$, NS + CFA vs NS + NS). Pretreatment with 7-NINA, AG and L-NAME, but not L-NIO, dramatically attenuated thermal hyperalgesia in mice receiving CFA throughout the observation period (A, $***P < 0.001$, inhibitor + CFA vs NS + CFA). None of the inhibitors altered pain thresholds in mice receiving NS (B, inhibitor + NS vs NS + NS). $n = 4$ for each group.



KO mice than in WT mice, except for IL-1 β in eNOS-KO mice, which was not different from WT (Fig. 5B).

After CFA, TNF (Fig. 5A), IL-1 β (Fig. 5B), and IL-10 (Fig. 5C) gene expression in nNOS-, iNOS- or eNOS-KO mice was lower than in WT mice at all time points ($P < 0.01$ and $P < 0.001$), except for TNF in eNOS-KO and IL-10 in iNOS-KO mice at 24 h, where they were higher (Fig. 5A; $P < 0.01$) and equal (Fig. 5C), respectively.

Discussion

The present study provided the following major findings: 1) pretreatment with 7-NINA (a selective nNOS inhibitor), AG (a selective iNOS inhibitor), or L-NAME (a non-selective NOS inhibitor), but not L-NIO (a selective eNOS inhibitor), attenuated CFA-induced thermal hyperalgesia in mice; 2) CFA caused an increase of nNOS, iNOS and eNOS, as well as of TNF, IL-1 β , and IL-10 gene expression in plantar skin; 3) preemptive systemic administration of the NOS inhibitors reduced




Figure 4 Effects of pretreatment with the NOS inhibitors 7-NINA, AG, L-NIO and L-NAME on cytokine gene expression after CFA. All treatments prevented the increase of TNF (A, $*P < 0.05$ and $***P < 0.001$; inhibitor + CFA vs NS + CFA) and IL-1 β (B, $**P < 0.01$ and $***P < 0.001$; inhibitor + CFA vs NS + CFA) gene expression induced by CFA. In contrast to the effects of AG and L-NIO, which reduced the increase of IL-10 gene expression at 6, 16, 24 h and at 6 and 16 h (C, $***P < 0.001$; AG + CFA or L-NIO + CFA vs NS + CFA), respectively, both 7-NINA and L-NAME at 6 and 16 h significantly enhanced the increase of IL-10 (C, $\#P < 0.05$ and $###P < 0.001$; 7-NINA + CFA or L-NAME + CFA vs NS + CFA), after CFA. At each time point $n = 4$.




Figure 5 Cytokine gene expression was lower in NOS-KO mice after CFA compared to WT mice. In control mice, there was no significant difference in baseline gene expression of TNF between KO and WT (A); IL-1 β in nNOS- and iNOS-KO, but not in eNOS-KO, was significantly higher than in WT (B; $^{##}P < 0.01$); IL-10 in KO was significantly lower than in WT (C; $^{**}P < 0.01$ and $^{***}P < 0.001$). After CFA, the nNOS-, iNOS- and eNOS-KO mice displayed lower TNF, IL-1 β and IL-10 gene expression (A, B and C respectively; $^{***}P < 0.001$; nNOS-, iNOS- or eNOS-KO CFA vs WT CFA), except that eNOS-KO mice had a higher TNF when compared to WT mice (A; $^{###}P < 0.001$) and there was no difference in IL-10 between iNOS-KO and WT mice (C), at 24 h. All measured samples were related to WT control. At each time point n = 4.

CFA-induced increase of TNF and IL-1 β , and produced differential effects on IL-10; 4) following CFA, nNOS-, iNOS- or eNOS-knockout (KO) mice had lower gene expression of TNF and IL-1 β , in accordance with the inhibitor data. Together, the findings imply that NOS mediates inflammatory thermal hyperalgesia by regulating cytokine expression.

Significant evidence links NO with the development and maintenance of inflammatory pain (see introduction). In addition, NO has been described either as pro- or anti-inflammatory and can produce both pro- and anti-nociceptive effects. The dual effects of NO in inflammatory pain may be related to the testing paradigms, pain models, doses of NO donors and inhibitors, etc [4]. In the present study, preemptive administration of 7-NINA, AG, or L-NAME, at a dose of 50 mg/kg, reduced thermal hyperalgesia caused by CFA, in line with recent studies [10,13,15]. Compared to the anti-hyperalgesic effects of 7-NINA and AG, we found that L-NIO, a selective eNOS inhibitor, had no significant effect on thermal hyperalgesia. This is in accordance with the finding by Tao et al. [18] who found that L-NIO did not affect carrageenan-induced thermal hyperalgesia in mice.

NO may activate sensory fibers directly, and may influence afferent fiber sensitivity indirectly by altering the availability of mediators such as calcitonin gene-related peptide (CGRP) and substance P (SP) [43]. In various pain models, increased expression of one or more of the three NOS isoforms has been shown in the spinal cord of rodents after i.pl. injection of capsaicin [44], formalin [45], carrageenan [17], or CFA [10,40]. However, to our knowledge, the expression of NOS in peripheral tissues is poorly explored. In the present study, iNOS gene expression in plantar skin was upregulated after CFA, in line with the report by De Alba et al. [13] who found that iNOS immunoreactivity in inflamed paw tissue was increased after CFA. We have extended these findings and have shown for the first time that CFA injection also increased the nNOS and eNOS gene expression. Together, these studies indicate that NOS can be induced not only in the central nervous system but also in the peripheral inflamed tissues after CFA, indicating a role in central and peripheral sensitization in inflammatory pain.

Proinflammatory cytokines like TNF and IL-1 β induce and facilitate neuropathic as well as inflammatory pain [26,46-50]. On the other hand, the anti-inflammatory cytokine IL-10, which can suppress pro-inflammatory cytokine production, exhibits antinociceptive effects in different pain models [51-54]. Both pro- and anti-inflammatory cytokines can be released by local or migrating cells, and their balance modulates pain sensitivity. We found that CFA injection resulted in a

significant elevation of TNF and IL-1 β gene expression in the inflamed plantar skin, which is in accordance with the change of TNF and IL-1 β at the protein level in the same tissue [47,49]. In addition, we detected for the first time that CFA also induced an upregulation of IL-10. Following CFA, mice pretreated with 7-NINA, AG, L-NIO or L-NAME had a reduced increase of TNF and IL-1 β . CFA-induced upregulation of IL-10 was increased in 7-NINA pretreated mice, but attenuated in mice receiving AG or L-NIO. Interestingly, pretreatment with L-NAME, a non-selective NOS inhibitor, augmented the increase of IL-10 in CFA mice. L-NAME has a higher selectivity and potency for nNOS than for eNOS and iNOS [55-58]; therefore, it might be possible that L-NAME preferentially inhibited nNOS, but not eNOS and iNOS, and thus produced the effect on IL-10, like the selective nNOS inhibitor 7-NINA. The exact mechanisms underlying the effects of the NOS inhibitors on cytokine expression are still not clear. NO activates the cyclic adenosine monophosphate (cAMP) cascade by S-nitrosylation activation of adenylate cyclase and by phosphorylation of cAMP response element-binding protein (CREB) through NO-activated cyclic guanosine monophosphate (cGMP). These second messengers can lead to the activation of transcription factors that modulate the expression of TNF, IL-1 β , and IL-10 [59-61]. In addition, the NO signaling pathway can mediate the expression of CREB target genes by controlling CREB-DNA binding, which is independent of CREB phosphorylation [62].

We then tested whether the situation of altered cytokine expression in mice receiving NOS inhibitors can be mimicked in NOS-KO mice. Following CFA, TNF and IL-1 β gene expression in nNOS-, iNOS- or eNOS-KO mice was significantly lower than in WT mice. Our data suggest that both inhibition of NOS (using the selective NOS inhibitors) and disruption of the NOS gene (using knock-out mice) can produce similar effects on TNF and IL-1 β gene expression. In a similar pattern, the reduced gene expression of IL-10 was concordant between the iNOS inhibitor AG treated- and iNOS-KO-mice, and the eNOS inhibitor L-NIO treated- and eNOS-KO-mice. Different from mice pretreated with the nNOS inhibitor 7-NINA, which had an augmented IL-10 gene expression, nNOS-KO animals showed lower expression of IL-10. The factors that may account for this discrepancy in IL-10 gene expression between the nNOS inhibitor pretreated mice and the nNOS-KO animals are still unknown. Previous studies indicated that eNOS and iNOS can compensate for the function of nNOS in nNOS-KO mice in the carrageenan [18] and CFA [63] models of inflammation, respectively. We speculate that nNOS may be compensated by iNOS and eNOS in nNOS-KO

mice here, which may lead to a lower gene expression of IL-10.

Considering that TNF and IL-1 β contribute to the generation of inflammatory pain and that IL-10 has anti-hyperalgesic effects, it is conceivable that both the attenuation of the increase of TNF and IL-1 β and an augmentation of the increase of IL-10 by 7-NINA and L-NAME may contribute to the mechanism by which pretreatment with 7-NINA and L-NAME reduces CFA-induced thermal hyperalgesia. In the same model, however, changes in hyperalgesia were absent [10] or less pronounced [63] in nNOS-KO mice. One possible explanation for these differences is that disruption of the nNOS gene not only reduces TNF and IL-1 β , but also decreases IL-10. Similarly, CFA-caused thermal hyperalgesia was not significantly altered in mice either receiving the eNOS inhibitor L-NIO (in the present study) or disrupting the eNOS gene [63] because the hyperalgesic TNF, IL-1 β and the anti-hyperalgesic IL-10 were simultaneously reduced for both treatments. Interestingly, compared to the unchanged hyperalgesia in the eNOS inhibitor L-NIO-treated mice after CFA, thermal hyperalgesia was significantly attenuated by the iNOS inhibitor AG, although AG also simultaneously reduced TNF, IL-1 β and IL-10. Different from what we observed in the iNOS inhibitor AG-treated mice, the attenuation of CFA-induced thermal hyperalgesia was less pronounced in iNOS-KO mice [63]. The mechanisms underlying these paradoxical phenomena are still unknown. Inflammation triggers a bi-directional activation of neurons and immune cells with subsequent balanced release of both pro-inflammatory and anti-inflammatory cytokines which can act locally or at a distance. Loss of this pro- and anti-inflammatory balance may underlie different pain states [27]. Differences in this balance in mice receiving different NOS inhibitors or in mice lacking different NOS genes may underlie the differential effects on behavior. Similar discrepancies in pain behavior between NOS inhibitor-treated mice and NOS-KO mice have been observed by others, and are conceivable given the probably large amount of uncontrolled compensatory changes in KO mice [10,17,18]. Last, our study does not exclude the possibility that other cytokines or chemokines and other pain mediators, such as bradykinin or prostaglandins, play a role in the modulation of inflammatory pain by NOS.

Conclusion

In summary, several lines of evidence indicate that cytokines regulation is a novel mechanism by which inhibition of NOS modulates CFA-induced inflammatory thermal hyperalgesia. Furthermore, our present data and studies from other groups [28-34] suggest that there

might be a feedback loop between NO and cytokines which modulates inflammatory pain.

Methods

Animals

Experiments were performed on adult (4-7 months, 25-30 g body weight) male mice of C57BL/6J background. These included wild type mice, mice deficient for iNOS [36], eNOS [37], and mice deficient for nNOS of 129S4BL/6J background (The Jackson Laboratory, Bar Harbor, Maine, USA) [38]. All mice were bred at the animal facilities of the University of Würzburg. The animals were housed on a light:dark cycle of 14:10 h with standard rodent chow and water available *ad libitum*. All experiments were approved by the Bavarian state authorities and performed in accordance with the European Communities Council Directive of November 24, 1986 (86/609/EEC) for the care and use of laboratory animals.

Drugs and drug administration

Intraplantar (i.pl.) injections were performed with a Hamilton syringe coupled to a 30-gauge needle under light ether anesthesia. Control mice received 0.5 ml of normal saline (NS) by intraperitoneal (i.p.) injection, and, 30 min later, 10 μ l of NS intraplantarly into the surface of one hind paw (control group: NS + NS). For induction of hindpaw inflammation, mice received i.pl. injection of 10 μ l of complete Freund's adjuvant (CFA, diluted 1:1 with PBS, 2 mg/ml; Mycobacterium tuberculosis; Difco Laboratories, Detroit, MI) 30 min after the i.p. NS injection (CFA group: NS + CFA). For the inhibitor experiments, the inhibitors including 7-nitroindazole sodium salt (7-NINA, a selective nNOS inhibitor; A.G. Scientific, Inc. Göttingen, Germany), aminoguanidine hydrochloride (AG, a selective iNOS inhibitor; Sigma, Munich, Germany), L-N(5)-(1-imi-noethyl)-ornithine (L-NIO, a selective eNOS inhibitor; Biotium Inc., Hayward, CA) and L-N(G)-nitroarginine methyl ester (L-NAME, a non-selective NOS inhibitor; Sigma) dissolved in NS except for 7-NINA in 20% DMSO, were i.p. injected at a dose of 50 mg/kg 30 min prior to i.pl. injection of CFA or NS (inhibitor pretreated CFA and control groups: inhibitor + CFA and inhibitor + NS).

Mice deficient for nNOS, iNOS and eNOS (KO mice) were i.p. injected with 0.5 ml of NS 30 min before i.pl. administration of 10 μ l of CFA or NS (NOS-KO CFA and control groups).

Behavioral testing

Sensitivity to noxious heat was assessed using the device of Hargreaves et al. [39] purchased from Ugo Basile (Comerio, Italy). A radiant heat source was focused on

the plantar surface of the hindpaw; the latency from the initiation of the radiant heat until paw withdrawal (paw withdrawal latency, PWL) was measured automatically. A maximal cutoff of 20 s was used to prevent tissue damage. The injected paw was tested two times; the mean withdrawal latency was calculated. The interval between two trials on the same paw was at least 3 min. Mice were tested 1 d before i.p.l. injection of CFA or NS to determine baseline thresholds, and then at 6, 16 and 24 h after injection.

Quantitative real-time PCR

Plantar skin of the hindpaw was removed 24 h after behavioral testing from control, CFA, and inhibitor pretreated CFA and control groups. In separate mice without behavioral testing, including CFA and inhibitor pretreated CFA groups, plantar skin was harvested at 6 and 16 h after injection. In addition, the same tissue was also dissected out from nNOS-, iNOS-, and eNOS-KO mice at 24 h after NS injection (NOS-KO control groups), and at 6, 16 and 24 h after CFA (NOS-KO CFA groups). At each time point, specimens from one treatment group were pooled ($n = 4$), immediately shock frozen in liquid nitrogen and stored at -80°C before further processing. Tissue homogenization and RNA isolation were performed as described previously [40]. The frozen tissue was incubated in TRIzol reagent[®] (Invitrogen, Karlsruhe, Germany) and homogenized with a Polytron homogenizer (Kinematica, Luzern, Switzerland). Afterwards chloroform was added and the samples were centrifuged at 13,000 g and 4°C for 15 min. Then, the upper phase was mixed with glycogen and propanol. After incubation over night at -20°C the samples were washed with 75% ethanol and the extracted RNA was dissolved in 33 μl of diethylpyrocarbonate (DEPC) treated water. The total RNA yield was photometrically quantified (Eppendorf, Hamburg, Germany).

Relative NOS and cytokine mRNA expressions were quantified with real-time PCR using the fluorescent TaqMan technology. 500 ng of total RNA were reverse transcribed (TaqMan Reverse Transcription Reagents, Applied Biosystems, Germany) using random hexamers. PCR primers and probes specific for mouse nNOS (Assay-ID: Mm00435175_m1), iNOS (Mm00440485_m1), eNOS (Mm00435204_m1), TNF (Mm00443258_m1), IL-1 β (Mm00434228_m1), IL-10 (Mm00439616_m1) and 18s rRNA were obtained from TaqMan Predeveloped Assay Reagents for gene expression (Applied Biosystems, Germany). 18s rRNA was used as an endogenous control. PCR was performed with equal amounts of cDNA in the GeneAmp 7700 sequence detection system (Applied Biosystems, Germany) using TaqMan Universal PCR Master Mix (Applied Biosystems). Reactions (total volume 50 μl)

were incubated at 50°C for 2 min, at 95°C for 10 min followed by 40 cycles of 15 s at 95°C and 1 min at 60°C . The comparative Ct method was used for relative quantification of gene expression. The amount of NOS and cytokine mRNAs, normalized to the endogenous control (18s rRNA) and relative to a calibrator (tissue from wild-type (WT) control animals), is given by $2^{-\Delta\Delta\text{Ct}}$, with Ct indicating the cycle number at which the fluorescence signal of the PCR product crosses an arbitrary threshold set within the exponential phase of the PCR, and $\Delta\Delta\text{Ct} = [(Ct_{\text{target}}(\text{unknownsample}) - Ct_{\text{end.control}}(\text{unknownsample}))] - [(Ct_{\text{target}}(\text{calibratorsample}) - Ct_{\text{end.control}}(\text{calibratorsample}))]$. The absolute value of the calibrator was set one and all measured samples were related to this sample. In order to guarantee primer specificity and to exclude genomic contamination, negative controls without cDNA template were run on each RT-PCR well plate.

To test the quality of the total RNA, the samples were photometrically quantified (Eppendorf, Hamburg, Germany) and the 260/280 ratio was measured for the integrity of the extracted RNA. In addition, agarose gel electrophoresis of the housekeeping gene glyceraldehyde-3-phosphate dehydrogenase (GAPDH) PCR product was analyzed. PCR amplification of GAPDH was performed at 94°C for 5 min and 35 cycles of 94°C for 45 sec, 58°C for 45 sec and 72°C for 45 sec, followed by 72°C for 10 min, using sequence specific primers (Sigma, Munich, Germany) 5'-TCACCACCA TGGAGAAGGCTG-3' (sense) and 5'-CCCTGTT GCTGTAGCCATATTC-3' (antisense). PCR products were loaded on 1% agarose gel containing ethidium bromide, and bands were visualised under UV rays. To assess the kinetics of RT-PCR, TNF primer was selected for RT-PCR amplification with decreasing concentrations of samples (diluted by 1:1, 1:2, 1:4, and 1:6). These data as above are provided in Additional files 1 and 2.

Data Analysis

For statistical analysis, SPSS software (Version 11.5; Chicago, IL) was used. As described previously [41,42], each sample was measured in triplicate, and data are given as means \pm SD. The data were analyzed by one-way analysis of variance (ANOVA) followed by least significant difference (LSD) *post hoc* test to determine differences between groups. $P < 0.05$ was considered to be statistically significant.

Additional file 1: Six RNA samples were randomly selected to show the quality of total RNA. The result of gel electrophoresis using GAPDH primers gave an expected band at 666 bp (A), and the ratio of OD260/OD280 was around 1.9, indicating that the samples were highly purified and largely intact (A and B).
Click here for file
[http://www.biomedcentral.com/content/supplementary/1744-8069-6-13-S1.PPT]

Additional file 2: Six RNA samples and TNF primer were randomly selected to show the kinetics of RT-PCR. TNF RT-PCR amplification plots with decreasing concentrations of samples suggested that Ct-values increase with further dilutions at 1:1, 1:2, 1:4 and 1:6. Click here for file [<http://www.biomedcentral.com/content/supplementary/1744-8069-6-13-S2.PPT>]

Acknowledgements

The study was supported by research funds from the University of Würzburg. We thank Barbara Gado and Lydia Biko for expert technical help.

Author details

¹Department of Neurology, University of Würzburg, Josef-Schneider-Str 11, 97080 Würzburg, Germany. ²Department of Psychiatry and Psychotherapy, University of Würzburg, Fuchsleinstr 15, 97080 Würzburg, Germany.

Authors' contributions

YC designed and carried out the studies, the data analyses, and drafted the manuscript. MKB and NÜ helped to design the studies and draft the manuscript. AR and AS contributed NOS knockout mice and participated in manuscript editing. CS coordinated and supervised the experiments, analyzed the data and wrote the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 23 November 2009

Accepted: 2 March 2010 Published: 2 March 2010

References

- Guzik TJ, Korbut R, Adamek-Guzik T: Nitric oxide and superoxide in inflammation and immune regulation. *J Physiol Pharmacol* 2003, **54**:469-487.
- Laroux FS, Lefer DJ, Kawachi S, Scalia R, Cockrell AS, Gray L, Van der Heyde H, Hoffman JM, Grisham MB: Role of nitric oxide in the regulation of acute and chronic inflammation. *Antioxid Redox Signal* 2000, **2**:391-396.
- Aley KO, McCarter G, Levine JD: Nitric oxide signaling in pain and nociceptor sensitization in the rat. *J Neurosci* 1998, **18**:7008-7014.
- Luo ZD, Cizkova D: The role of nitric oxide in nociception. *Curr Rev Pain* 2000, **4**:459-466.
- Malmberg AB, Yaksh TL: Spinal nitric oxide synthesis inhibition blocks NMDA-induced thermal hyperalgesia and produces antinociception in the formalin test in rats. *Pain* 1993, **54**:291-300.
- Meller ST, Gebhart GF: Nitric oxide (NO) and nociceptive processing in the spinal cord. *Pain* 1993, **52**:127-136.
- Forstermann U, Gath I, Schwarz P, Closs EI, Kleinert H: Isoforms of nitric oxide synthase: properties, cellular distribution and expressional control. *Biochem Pharmacol* 1995, **50**:1321-1332.
- Mayer B: Biochemistry and molecular pharmacology of nitric oxide synthase. *Nitric oxide in the nervous system* New York, Academic Press/Vincent SR 1995, 21-42.
- Paakkari I, Lindsberg P: Nitric oxide in the central nervous system. *Ann Med* 1995, **27**:369-377.
- Chu YC, Guan Y, Skinner J, Raja SN, Johns RA, Tao YX: Effect of genetic knockout or pharmacologic inhibition of neuronal nitric oxide synthase on complete Freund's adjuvant-induced persistent pain. *Pain* 2005, **119**:113-123.
- Osborne MG, Coderre TJ: Effects of intrathecal administration of nitric oxide synthase inhibitors on carrageenan-induced thermal hyperalgesia. *Br J Pharmacol* 1999, **126**:B1840-1846.
- Semos ML, Headley PM: The role of nitric oxide in spinal nociceptive reflexes in rats with neurogenic and non-neurogenic peripheral inflammation. *Neuropharmacology* 1994, **33**:1487-1497.
- De Alba J, Clayton NM, Collins SD, Colthup P, Chessell I, Knowles RG: GW27 a novel and highly selective inhibitor of the inducible isoform of nitric oxide synthase (iNOS), shows analgesic effects in rat models of inflammatory and neuropathic pain. *Pain* 2007, **120**:170-181.
- Handy RL, Moore PK: Effects of selective inhibitors of neuronal nitric oxide synthase on carrageenan-induced mechanical and thermal hyperalgesia. *Neuropharmacology* 1998, **37**:37-43.
- LaBuda CJ, Koblisch M, Tuthill P, Dolle RE, Little PJ: Antinociceptive activity of the selective iNOS inhibitor AR-C102222 in rodent models of inflammatory, neuropathic and post-operative pain. *Eur J Pain* 2006, **10**:505-512.
- Tang Q, Svensson CI, Fitzsimmons B, Webb M, Yaksh TL, Hua XY: Inhibition of spinal constitutive NOS-2 by 1400W attenuates tissue injury and inflammation-induced hyperalgesia and spinal p38 activation. *Eur J Neurosci* 2007, **25**:2964-2972.
- Tao F, Tao YX, Mao P, Zhao C, Li D, Liaw WJ, Raja SN, Johns RA: Intact carrageenan-induced thermal hyperalgesia in mice lacking inducible nitric oxide synthase. *Neuroscience* 2003, **120**:847-854.
- Tao F, Tao YX, Zhao C, Doré S, Liaw WJ, Raja SN, Johns RA: Differential roles of neuronal and endothelial nitric oxide synthases during carrageenan-induced inflammatory hyperalgesia. *Neuroscience* 2004, **128**:421-430.
- Kidd BL, Urban LA: Mechanisms of inflammatory pain. *Br J Anaesth* 2001, **87**:3-11.
- Ji RR, Woolf CJ: Neuronal plasticity and signal transduction in nociceptive neurons: implications for the initiation and maintenance of pathological pain. *Neurobiol Dis* 2001, **8**:1-10.
- Neumann S, Doubell TP, Leslie T, Woolf CJ: Inflammatory pain hypersensitivity mediated by phenotypic switch in myelinated primary sensory neurons. *Nature* 1996, **384**:360-364.
- Cunha F, Poole S, Lorenzetti B, Ferreira S: The pivotal role of tumor necrosis factor alpha in the development of inflammatory hyperalgesia. *Br J Pharmacol* 1992, **107**:660-664.
- Ferreira S, Lorenzetti B, Bristow A, Poole S: Interleukin-1β as a potent hyperalgesic agent antagonized by a tripeptide analogue. *Nature* 1988, **334**:698-700.
- Verri WA Jr, Cunha TM, Parada CA, Poole S, Cunha FQ, Ferreira SH: Hypernociceptive role of cytokines and chemokines: targets for analgesic drug development?. *Pharmacol Ther* 2006, **112**:116-138.
- Poole S, Cunha FQ, Ferreira SH: Hyperalgesia from subcutaneous cytokines. *Cytokines and Pain* Basel, Birkhäuser/Watkins LR, Maier SF 1999, 59-87.
- Üçeyler N, Sommer C: Cytokine-induced pain: basic science and clinical implications. *Rev Analg* 2007, **9**:87-103.
- Üçeyler N, Sommer C: Cytokine regulation in animal models of neuropathic pain and in human diseases. *Neurosci Lett* 2008, **437**:194-198.
- Ferreira SH: Inflammatory pain: the role of cytokines and its control by drugs which release nitric oxide. *Ann Ist Super Sanita* 1993, **29**:367-373.
- Hartrick CT: Increased production of nitric oxide stimulated by interferon-gamma from peripheral blood monocytes in patients with complex regional pain syndrome. *Neurosci Lett* 2002, **323**:75-77.
- Meller ST, Dykstra C, Grzybycki D, Murphy S, Gebhart GF: The possible role of glia in nociceptive processing and hyperalgesia in the spinal cord of the rat. *Neuropharmacology* 1994, **33**:1471-1478.
- Plunkett JA, Yu CG, Easton JM, Bethea JR, Yezierski RP: Effects of interleukin-10 (IL-10) on pain behavior and gene expression following excitotoxic spinal cord injury in the rat. *Exp Neurol* 2001, **168**:144-154.
- Sung CS, Wen ZH, Chang WK, Chan KH, Ho ST, Tsai SK, Chang YC, Wong CS: Inhibition of p38 mitogen-activated protein kinase attenuates interleukin-1β-induced thermal hyperalgesia and inducible nitric oxide synthase expression in the spinal cord. *J Neurochem* 2005, **94**:742-752.
- Sung CS, Wen ZH, Chang WK, Ho ST, Tsai SK, Chang YC, Wong CS: Intrathecal interleukin-1β administration induces thermal hyperalgesia by activating inducible nitric oxide synthase expression in the rat spinal cord. *Brain Res* 2004, **1015**:145-153.
- Xu XJ, Hao JX, Olsson T, Kristensson K, van der Meide PH, Wiesenfeld-Hallin Z: Intrathecal interferon-gamma facilitates the spinal nociceptive flexor reflex in the rat. *Neurosci Lett* 1994, **182**:263-266.
- Holguin A, O'Connor KA, Biedenkapp J, Campisi J, Wieseler-Frank J, Milligan ED, Hansen MK, Spataro L, Maksimova E, Bravmann C, Martin D, Fleshner M, Maier SF, Watkins LR: HIV-1 gp120 stimulates proinflammatory

- cytokine-mediated pain facilitation via activation of nitric oxide synthase-1 (nNOS). *Pain* 2004, **110**:517-530.
36. Laubach VE, Shesely EG, Smithies O, Sherman PA: Mice lacking inducible nitric oxide synthase are not resistant to lipopolysaccharide-induced death. *Proc Natl Acad Sci USA* 1995, **92**:10688-10692.
 37. Huang PL, Huang Z, Mashimo H, Bloch KD, Moskowitz MA, Bevan JA, Fishman MC: Hypertension in mice lacking the gene for endothelial nitric oxide synthase. *Nature* 1995, **377**:239-242.
 38. Huang PL, Dawson TM, Brecht DS, Snyder SH, Fishman MC: Targeted disruption of the neuronal nitric oxide synthase gene. *Cell* 1993, **75**:1273-1286.
 39. Hargreaves K, Dubner R, Brown F, Flores C, Joris J: A new and sensitive method for measuring thermal nociception in cutaneous hyperalgesia. *Pain* 1988, **32**:77-88.
 40. Chen Y, Sommer C: Activation of the nociceptin opioid system in rat sensory neurons produces antinociceptive effects in inflammatory pain: involvement of inflammatory mediators. *J Neurosci Res* 2007, **85**:1478-1488.
 41. Kleinschnitz C, Brinkhoff J, Zelenka M, Sommer C, Stoll G: The extent of cytokine induction in peripheral nerve lesions depends on the mode of injury and NMDA receptor signaling. *J Neuroimmunol* 2004, **149**:77-83.
 42. Kleinschnitz C, Hofstetter HH, Meuth SG, Bräuninger S, Sommer C, Stoll G: cell infiltration after chronic constriction injury of mouse sciatic nerve is associated with interleukin-17 expression. *Exp Neurol* 2006, **200**:480-485.
 43. Handy RL: Nitric oxide and inflammatory pain. *Pain and Neurogenic Inflammation* Basel, BirkhäuserBrain SD, Moore PK 1999, 81-102.
 44. Wu J, Fang L, Lin Q, Willis WD: Nitric oxide synthase in spinal cord central sensitization following intradermal injection of capsaicin. *Pain* 2001, **94**:47-58.
 45. Lam HH, Hanley DF, Trapp BD, Saito S, Raja S, Dawson TM, Yamaguchi H: Induction of spinal cord neuronal nitric oxide synthase (NOS) after formalin injection in the rat hind paw. *Neurosci Lett* 1996, **210**:201-204.
 46. Poole S, Cunha FQ, Ferreira SH: Bradykinin, cytokines and inflammatory hyperalgesia. *Pain and Neuroimmune Interactions* New York, Kluwer Academic/Plenum PublishersSaadé NE, Apkarian AV, Jabbur SJ 2000, 31-54.
 47. Safieh-Garabedian B, Poole S, Allchorne A, Winter J, Woolf CJ: Contribution of interleukin-1 beta to the inflammation-induced increase in nerve growth factor levels and inflammatory hyperalgesia. *Br J Pharmacol* 1995, **115**:1265-1275.
 48. Schäfers M, Svensson CI, Sommer C, Sorkin LS: Tumor necrosis factor-alpha induces mechanical allodynia after spinal nerve ligation by activation of p38 MAPK in primary sensory neurons. *J Neurosci* 2003, **23**:2517-2521.
 49. Woolf CJ, Allchorne A, Safieh-Garabedian B, Poole S: Cytokines, nerve growth factor and inflammatory hyperalgesia: the contribution of tumour necrosis factor alpha. *Br J Pharmacol* 1997, **121**:417-424.
 50. Zelenka M, Schäfers M, Sommer C: Intraneural injection of interleukin-1beta and tumor necrosis factor-alpha into rat sciatic nerve at physiological doses induces signs of neuropathic pain. *Pain* 2005, **116**:257-263.
 51. Karam MC, Hamdan HG, Abi Chedid NA, Bodman-Smith KB, Baroudy GM: Interleukin-10 reduces hyperalgesia and the level of Interleukin-1beta in BALB/c mice infected with *Leishmania major* with no major effect on the level of Interleukin-6. *J Neuroimmunol* 2007, **183**:43-49.
 52. Ledeboer A, Jekich BM, Sloane EM, Mahoney JH, Langer SJ, Milligan ED, Martin D, Maier SF, Johnson KW, Leinwand LA, Chavez RA, Watkins LR: Intrathecal interleukin-10 gene therapy attenuates paclitaxel-induced mechanical allodynia and proinflammatory cytokine expression in dorsal root ganglia in rats. *Brain Behav Immun* 2007, **21**:686-698.
 53. Milligan ED, Langer SJ, Sloane EM, He L, Wieseler-Frank J, O'Connor K, Martin D, Forsayeth JR, Maier SF, Johnson K, Chavez RA, Leinwand LA, Watkins LR: Controlling pathological pain by adenovirally driven spinal production of the anti-inflammatory cytokine, interleukin-10. *Eur J Neurosci* 2005, **21**:2136-2148.
 54. Wagner R, Janjigian M, Myers RR: Anti-inflammatory interleukin-10 therapy in CCI neuropathy decreases thermal hyperalgesia, macrophage recruitment, and endoneurial TNF- α expression. *Pain* 1998, **74**:35-42.
 55. Buckner CK, Saban R, Castleman WL, Will JA: Analysis of leukotriene receptor antagonists on isolated human intralobar airways. *Ann N Y Acad Sci* 1988, **524**:181-186.
 56. Furfine ES, Harmon MF, Paith JE, Garvey EP: Selective inhibition of constitutive nitric oxide synthase by L-NG-nitroarginine. *Biochemistry* 1993, **32**:8512-8517.
 57. Garvey EP, Tuttle JV, Covington K, Merrill BM, Wood ER, Baylis SA, Charles IG: Purification and characterization of the constitutive nitric oxide synthase from human placenta. *Arch Biochem Biophys* 1994, **311**:235-241.
 58. Southan GJ, Szabó C: Selective pharmacological inhibition of distinct nitric oxide synthase isoforms. *Biochem Pharmacol* 1996, **51**:383-394.
 59. Gee K, Angel JB, Ma W, Mishra S, Gajanayaka N, Parato K, Kumar A: Intracellular HIV-Tat expression induces IL-10 synthesis by the CREB-1 transcription factor through Ser133 phosphorylation and its regulation by the ERK1/2 MAPK in human monocytic cells. *J Biol Chem* 2006, **281**:31647-31658.
 60. Jongeneel CV: Transcriptional regulation of the tumor necrosis factor alpha gene. *Immunobiology* 1995, **193**:210-216.
 61. Watkins LR, Hansen MK, Nguyen KT, Lee JE, Maier SF: Dynamic regulation of the proinflammatory cytokine, interleukin-1beta: molecular biology for non-molecular biologists. *Life Sci* 1999, **65**:449-481.
 62. Riccio A, Alvania RS, Lonze BE, Ramanan N, Kim T, Huang Y, Dawson TM, Snyder SH, Ginty DD: A nitric oxide signaling pathway controls CREB-mediated gene expression in neurons. *Mol Cell* 2006, **21**:283-294.
 63. Boettger MK, Üceyler N, Zelenka M, Schmitt A, Reif A, Chen Y, Sommer C: Differences in inflammatory pain in nNOS-, iNOS- and eNOS-deficient mice. *Eur J Pain* 2007, **11**:810-818.

doi:10.1186/1744-8069-6-13

Cite this article as: Chen *et al.*: Nitric oxide synthase modulates CFA-induced thermal hyperalgesia through cytokine regulation in mice. *Molecular Pain* 2010 **6**:13.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

