



Published in final edited form as:

Bioorg Med Chem Lett. 2010 April 1; 20(7): 2174–2177. doi:10.1016/j.bmcl.2010.02.041.

Synthesis and SAR of *N*-(4-(4-alkylpiperazin-1-yl)phenyl)benzamides as muscarinic acetylcholine receptor subtype 1 (M₁) antagonists

Nicole R. Miller^{a,†}, R. Nathan Daniels^{c,e,†}, David Lee^e, P. Jeffrey Conn^{a,b,c,d}, and Craig W. Lindsley^{a,b,c,d,e}

^aDepartment of Pharmacology, Vanderbilt University Medical Center, Nashville, TN 37232, USA

^bVanderbilt Program in Drug Discovery, Vanderbilt University Medical Center, Nashville, TN 37232, USA

^cVanderbilt MLPCN Specialized Chemistry Center, Vanderbilt University Medical Center, Nashville, TN 37232, USA

^dVanderbilt Institute of Chemical Biology, Vanderbilt University Medical Center, Nashville, TN 37232, USA

^eDepartment of Chemistry, Vanderbilt University, Nashville, TN 37232, USA

Abstract

This Letter describes the synthesis and SAR, developed through an iterative analog library approach, of a novel series of selective M₁ mAChR antagonists, based on an *N*-(4-(4-alkylpiperazin-1-yl)phenyl)benzamide scaffold for the potential treatment of Parkinson's disease, dystonia and other movement disorders. Compounds in this series possess M₁ antagonist IC₅₀s in the 350 nM to >10 μM range with varying degrees of functional selectivity versus M₂-M₅.

There are five subtypes of muscarinic acetylcholine receptors (mAChR1-5 or M₁-M₅), members of the G Protein-Coupled Receptor (GPCR) family A, that mediate the metabotropic actions of the neurotransmitter acetylcholine.^{1,2} M₁, M₃ and M₅ activate phospholipase C and calcium mobilization through G_q whereas M₂ and M₄ block the action of adenylyl cyclase through G_{i/o}.^{1,2} The cholinergic system, mediated by mAChRs, plays a critical role in a wide variety of CNS and peripheral functions including memory and attention mechanisms, motor control, nociception, regulation of sleep wake cycles, cardiovascular function, renal and gastrointestinal function to mention only a few.¹⁻⁴ As a result, agents that can selectively modulate the activity of mAChRs have the potential for therapeutic use in multiple peripheral and central pathological states. Due to high sequence conservation within the orthosteric binding site of the five mAChR subtypes, it has been historically difficult to develop mAChR subtype-selective ligands.¹⁻⁵ Based on brain expression and cellular localization, data from mAChR knock-out mice and clinical trials with muscarinic agents, the M₁ subtype is an attractive molecular target for the treatment of CNS disorders. M₁ has been implicated in the

Correspondence to: Craig W. Lindsley.

[†]these authors contributed equally

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

pathologies of Alzheimer's disease (AD), Parkinson's disease (PD) and dystonia due to its role in cognition and motor control.⁶

The majority of reported muscarinic antagonists are unselective, such as a scopolamine, **1**.⁷ Recently, pirenzapine, **2** has emerged as a relatively selective M₁ receptor antagonist (20- to 50-fold versus M₂-M₅) and there are numerous reports of moderately selective M₃ antagonists (20- to 50-fold versus M₂) such as **3**.⁸ Interestingly, the most selective M₁ antagonist, MT7, **4**, the 65 amino acid peptide, (>1,000-fold versus M₂-M₅) was derived from venom extracts of the green mamba snake (Fig. 1).⁹ From an M1 functional screen within the MLSCN, we identified M₁ antagonists such as **5** (M₁ IC₅₀ of 441 nM and with >340-fold selectivity versus M₄, but modest selectivity versus M₂, M₃ and M₅ (7.9-fold, 7-fold, and 2.4-fold, respectively)) and **6** (M₁ IC₅₀ of 5.0 μM and with >30-fold selective versus M₂-M₅).¹⁰⁻¹² Based on the M₁ selectivity of **6**, attractive physiochemical properties (MW < 350, clogP 3.6) and the fact that it was the only benzamide-containing analog in the series, we initiated a library synthesis effort¹³ to develop SAR around **6**.

As shown in Scheme 1, the first round of library synthesis focused on benzamide analogs of **6**. Commercially available 3-chloro-(4-(4-ethylpiperazin-1yl)aniline **7** was acylated under standard conditions employing polymer-supported reagents and scavengers¹³ to afford a 24-member library of analogs **8**, along with resynthesized **6**. All analogs were then purified by mass-guided HPLC to analytical purity.¹⁴ To effectively screen small libraries of potential mAChR ligands, we have adopted a strategy to triage compounds in single-point screens (at 10 μM) at M₁, M₃ and M₅ – the G_q-coupled mAChRs – to identify active and selective compounds prior to running full concentration-response curves (CRCs).¹⁵ Figure 2 shows the 10 μM single-point screens for the first 25-member library of benzamide analogs **8**.

As Shown in Table 1, re-synthesized **6** displayed comparable potency and mAChR selectivity to the original sample (M₁ IC₅₀ = 3.2 μM, IC₅₀ >> 10 μM for M₂-M₅). Functionalized benzamide analogs **8** possessed a wide range of M₁ potency and mAChR selectivity, and we initially evaluated analogs **8** against M₁, M₃ and M₅. Substitution in the 2-position, **8a** (2-Cl) and **8b** (2-OMe) possessed submicromolar M₁ IC₅₀s (960 nM and 820 nM, respectively), but also showed low micromolar activity at M₃ and M₅. A pentafluorophenyl congener **8e** (Fig. 3A) proved to be a submicromolar antagonist of both M₁ and M₅ (IC₅₀s of 350 nM and 830 nM, respectively). Substitution at the 4-position, as with the 4-OMe derivative **8f**, was comparable to the original **6**. Interestingly, a 2,5-bisCF₃ analog **8h** had an M₁ IC₅₀ of 490 nM, with ~ 9-fold functional selectivity versus M₃ and M₅ (Fig. 3B). Intrigued by this potent and selective M₁ antagonist, we screened against M₂ and M₄ as well, but found that **8h** possessed only 3- to 4-fold selectivity versus the G_{i/o}-coupled mAChRs (Table 1). **8i**, a 3,5-bisCF₃ analog possessed a unique profile as a dual M₁/M₄ antagonist (IC₅₀s of 2.6 μM and 3.7 μM, respectively), with little effect on an ACh EC₈₀ at 10 μM on M₂, M₃ or M₅. Finally, a 3,4-difluoro **8j** derivative was also comparable to the original **6**. While this library afforded interesting results, further optimization was required.

Having surveyed the amide moiety while maintaining the *N*-ethyl piperazine, we next generated two-dimensional libraries wherein the nature of the alkyl group was varied (**9-12**) while also surveying diverse benzamides to generate analogs **9a-f**, **10 a-f**, **11a-f** and **12a-f** (Scheme 2).

Application of the same strategy to triage compounds in single-point screens (at 10 μM) at M₁, M₃ and M₅ to identify active and selective compounds prior to running full (CRCs) was employed, but >75% of these new analogs possessed no M₁ antagonist activity. The SAR for this series was incredibly shallow, with only an *N*-propyl congener with the 3,5-

dichlorobenzamide moiety **11i** displaying reasonable activity (M_1 $IC_{50} = 3.7 \mu M$, $IC_{50} > 10 \mu M$ for M_3 and M_5), and all other analogs possessing M_1 IC_{50} s in the 6-9 μM range.

In summary, a two-dimensional parallel synthesis library campaign was performed around **6**, an M_1 antagonist identified in a functional HTS screen. SAR for this series was shallow, but we were able to improve the M_1 antagonist activity of **6** into the 350 to 500 nM range with analogs **8**, while maintaining good mAChR selectivity. Interestingly, **8i** is the first reported dual M_1/M_4 -preferring antagonist, which compliments the prototypical M_1/M_4 -preferring agonist xanomeline. Other chemical series from our M_1 functional screen are currently under chemical optimization, and further refinements will be reported in due course.

Acknowledgments

The authors thank the NIH, NIMH and the MLSCN for funding of the Vanderbilt Screening Center for GPCRs, Ion Channels & Transporters (3U54MH074427), the Chemistry Supplement (3U54MH074427-02S1) and the XO1 (1XO1MH077606-01) M_1 antagonist screening application which enabled this work. The authors also wish to thank the second generation of the MLSCN, coined the MLPCN, and the Vanderbilt Specialized Chemistry Center (U54MH084659-01).

References and Notes

- (a) Bonner TI, Buckley NJ, Young AC, Brann MR. *Science* 1987;237:527. [PubMed: 3037705] (b) Bonner TI, Young AC, Buckley NJ, Brann MR. *Neuron* 1988;1:403. [PubMed: 3272174]
- Felder CC, Bymaster FP, Ward J, DeLapp N. *J Med Chem* 2000;43:4333. [PubMed: 11087557]
- Bymaster FP, McKinzie DL, Felder CC, Wess J. *Neurochem Res* 2003;28:437. [PubMed: 12675128]
- Birdsall NJM, Farries T, Gharagozloo P, Kobayashi S, Lazareno S, Sugimoto M. *Mol Pharmacol* 1999;55:778. [PubMed: 10101037]
- Eglen RM, Choppin A, Dillon MP, Hedge S. *Curr Opin Chem Biol* 1999;3:426. [PubMed: 10419852]
- Birdsall NJM, Nathanson NM, Schwarz RD. *TRENDS Pharm Sci* 2001;22:215.
- (a) Burke RE. *Mov Disord* 1986;1:135. [PubMed: 2904117] (b) Buckley NJ, Bonner TI, Buckley CM, Brann MR. *Mol Pharmacol* 1989;35:469. [PubMed: 2704370] (c) Walebroeck M, Tastenoy M, Camus J, Christophe J. *Mol Pharmacol* 1990;36:267. (d) Dei S, Bellucci C, Buccioni M, Ferraroni M, Guandalini L, Manetti L, Martini E, Marucci G, Matucci R, Nesi M, Romanelli MN, Scapecchi S, Teodori E. *J Med Chem* 2007;50:1409. [PubMed: 17305327]
- Hammer R, Berrie CP, Birdsall NJM, Burgen ASV, Hulme EC. *Nature* 1980;283:90. [PubMed: 7350532]
- Bradley KN. *Pharmacol & Therap* 2000;85:87. [PubMed: 10722122]
- For information on the Molecular Library Production Center Network see: <http://mli.nih.gov/mli/mlpcn/>
- Lewis LM, Sheffler D, Williams R, Bridges TA, Kennedy JP, Brogan JT, Mulder MJ, Williams L, Nalywajko NT, Niswender C, Weaver CD, Conn PJ, Lindsley CW. *Bioorg Med Chem Lett* 2008;18:885. [PubMed: 18178088]
- Sheffler DJ, Williams R, Bridges TM, Lewis LM, Xiang Z, Zheng F, Kane AS, Byum NE, Jadhav S, Mock MM, Zheng F, Lewis LM, Jones CK, Niswender CM, Weaver CD, Conn PJ, Lindsley CW, Conn PJ. *Mol Pharmacol* 2009;76:356. [PubMed: 19407080]
- Kennedy JP, Williams L, Bridges TM, Daniels RN, Weaver D, Lindsley CW. *J Comb Chem* 2008;10:345. [PubMed: 18220367]
- Leister WH, Strauss KA, Wisnoski DD, Zhao Z, Lindsley CW. *J Comb Chem* 2003;5:322. [PubMed: 12739949]
- Details of the calcium mobilization assays: Chinese Hamster Ovary (CHO-K1) cells stably expressing human (h) M_1 , hM_3 , and hM_5 were used for calcium mobilization assays. hM_2 and hM_4 were adapted to this assay and signaling pathway after stably transfecting G_{q15} chimeric G protein. To measure agonist-induced calcium mobilization and determine effect of novel compounds, stable muscarinic cell lines plated overnight in Costar 96-well cell culture plates (Corning) were incubated with 50 μ L

of 2 μ M Fluo-4 AM diluted in assay buffer [HBSS (Invitrogen) supplemented with 20mM HEPES and 2.5mM probenecid, pH 7.4] for 45min at 37°C. Dye was then removed and replaced with assay buffer. Cells were pre-incubated with 10 μ M or a concentration-response curve of novel compound, followed by a sub-maximal concentration of Acetylcholine or Carbachol. The signal amplitude was first normalized to baseline and then expressed as a percentage of the maximal response to acetylcholine.




Figure 1.
Structures of representative mAChR antagonists.




Figure 2. Single-point EC₈₀ plus 10 μ M compound triage screen at M₁, M₃ and M₅ to select compounds for full CRCs.






Figure 3. CRCs for M₁, M₃ and M₅ for (A) compound **8e** (M₁ IC₅₀ = 350 nM) and (B) compound **8h** (M₁ IC₅₀ = 490 nM), showing ~9-fold functional selectivity versus M₃ and M₅.


**Scheme 1.**

Library synthesis of first generation analogs **8**. All library compounds were purified by mass-guided HPLC to >98% purity.¹⁴


**Scheme 2.**

Library synthesis of second generation analogs **9a-f**, **10a-f**, **11a-f** and **12a-f**. All library compounds were purified by mass-guided HPLC to >98% purity.¹⁴

Table 1

and mAChR activities of analogues **8**.


Ar	M1 IC ₅₀ (μM) ^a	M2 IC ₅₀ (μM) ^a	M3 IC ₅₀ (μM) ^a	M4 IC ₅₀ (μM) ^a	M5 IC ₅₀ (μM) ^a
	3.2	>10	>10	>10	>10
	0.96	ND	8.2	ND	2.3
	0.82	ND	5.6	ND	1.3




Ar

M1 IC₅₀ (μM)^a M2 IC₅₀ (μM)^a M3 IC₅₀ (μM)^a M4 IC₅₀ (μM)^a M5 IC₅₀ (μM)^a


2.9 6.9 >10 3.7 >10



Ar





M1 IC ₅₀ (μM) ^a	M2 IC ₅₀ (μM) ^a	M3 IC ₅₀ (μM) ^a	M4 IC ₅₀ (μM) ^a	M5 IC ₅₀ (μM) ^a
2.1	ND	>10	ND	3.5



Ar


M1 IC ₅₀ (μM) ^a	M2 IC ₅₀ (μM) ^a	M3 IC ₅₀ (μM) ^a	M4 IC ₅₀ (μM) ^a	M5 IC ₅₀ (μM) ^a
0.35	ND	3.7	ND	0.83






Ar

M1 IC ₅₀ (μM) ^a	M2 IC ₅₀ (μM) ^a	M3 IC ₅₀ (μM) ^a	M4 IC ₅₀ (μM) ^a	M5 IC ₅₀ (μM) ^a
3.2	ND	>10	ND	>10



Ar

M1 IC ₅₀ (μM) ^a	M2 IC ₅₀ (μM) ^a	M3 IC ₅₀ (μM) ^a	M4 IC ₅₀ (μM) ^a	M5 IC ₅₀ (μM) ^a
0.49	2.7	4.2	1.5	4.1



Ar

M1 IC₅₀ (μM)^a M2 IC₅₀ (μM)^a M3 IC₅₀ (μM)^a M4 IC₅₀ (μM)^a M5 IC₅₀ (μM)^a

4.7 >10 >10 >10 >10