

Published in final edited form as:

Curr Opin Microbiol. 2010 April ; 13(2): 198–203. doi:10.1016/j.mib.2010.01.014.

## Global Regulation by the Seven-component $P_i$ Signaling System

Yi-Ju Hsieh and Barry L. Wanner\*

Department of Biological Sciences, Purdue University, West Lafayette, Indiana 47907

### Summary

This review concerns how *Escherichia coli* detects environmental inorganic orthophosphate ( $P_i$ ) to regulate genes of the phosphate (Pho) regulon by the PhoR/PhoB two-component system (TCS).  $P_i$  control by the PhoR/PhoB TCS is a paradigm of a bacterial signal transduction pathway in which occupancy of a cell surface receptor(s) controls gene expression in the cytoplasm. The  $P_i$  signaling pathway requires seven proteins, all of which probably interact in a membrane-associated signaling complex. Our latest studies show that  $P_i$  signaling involves three distinct processes, which appear to correspond to different states of the sensory histidine kinase PhoR: an inhibition state, an activation state, and a deactivation state. We describe a revised model for  $P_i$  signal transduction of the *E. coli* Pho regulon.

### Introduction

How cells respond to environmental (extracellular) signals is of fundamental importance in biology. The control of the *Escherichia coli* phosphate (Pho) regulon by extracellular inorganic orthophosphate ( $P_i$ ) is of special interest for it serves as a paradigm for a two-component system (TCS) in which signaling is mediated by an ABC (ATP-binding cassette) transporter, the Pst (phosphate-specific transport) system, in the absence of transport.

The *E. coli* Pho regulon is comprised of a large number of genes that are co-regulated by environmental  $P_i$ , the preferred P source, and that are required for assimilation of a variety of phosphorus (P) sources for growth. Signal transduction by environmental  $P_i$  requires seven proteins, which are thought to interact in a membrane-associated signaling complex. These  $P_i$  signaling proteins include: (i) two that are members of the large family of TCSs, namely the sensory histidine kinase (HK) PhoR (an integral membrane protein) and its partner DNA-binding response regulator (RR) PhoB (a transcription factor); (ii) four components of the ABC transporter Pst; and (iii) the chaperone-like PhoR/PhoB inhibitory protein called PhoU.

The PhoR HK is required for activation (phosphorylation) of the PhoB RR under conditions of  $P_i$  limitation. Other (non-partner) HKs, e. g., the CreC HK of the CreC/CreB TCS, can also activate (phosphorylate) PhoB, both *in vivo* and *in vitro*. The finding of such interactions has led to the suggestion that “cross regulation” can occur between different TCSs, which may play a role in the integration of multiple signals. For example, cross regulation of the PhoR/PhoB TCS may be important for connecting different steps of  $P_i$  metabolism [1]. Similar interactions have been seen among non-partner proteins of other TCSs (e. g., the NarX/NarL and NarQ/NarP TCSs [2]). DNA microarray studies have provided further evidence for cross

\*Corresponding author: Ph: 1-765-494-8034, Fax: 1-765-496-9028, yhsieh@purdue.edu; blwanner@purdue.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

regulation among the BaeS/BaeR, PhoR/PhoB, and CreC/CreB TCSs [3]. Other data suggest that cross regulation of the PhoR/PhoB TCS is likely to be even more extensive [4]. Thus, further studies of the Pho regulon can serve as a model for cross regulation among different TCSs.

This review covers the period from when inorganic orthophosphate ( $P_i$ ) control of the *Escherichia coli* phosphate (Pho) regulon was last reviewed in 1996 [1] through 2009. It includes new information on genes controlled by the PhoR/PhoB TCS, cross regulation and stochasticity in the control of  $P_i$ -regulated genes, and our current understanding of how environmental  $P_i$  regulates the *E. coli* Pho regulon.

## The PhoR/PhoB TCS controls genes for phosphorus assimilation

Estimates for the number of  $P_i$ -regulated genes vary widely. Proteome profiles of cells grown under  $P_i$  excess and limited conditions revealed nearly 400 proteins (almost 10% of the *E. coli* proteome) whose amounts varied in response to the environmental P source [5]. Results from DNA microarray experiments have also shown the number of PhoR/PhoB-regulated genes to be large (Y. Jiang, Y. H., and B. L. W., unpublished data). These data are consistent with computational predictions of a large number of PhoB-binding sites on the genome [6]. However, in the absence of direct evidence, it is difficult to provide a complete catalog of Pho regulon genes. To date, only 31 genes (9 transcriptional units: *eda*, *phnCDEFGHIJKLMNOP*, *phoA*, *phoBR*, *phoE*, *phoH*, *psiE*, *pstSCAB-phoU*, and *ugpBAECQ*) have been shown to be directly controlled by the PhoR/PhoB TCS (Table 1). Although strong evidence exists for several others (such as *amn*, *psiF*, *yidD*, and *yibD*), direct evidence for their control by PhoB is lacking. In this regard, expression of the acid-inducible *asr*, which had been previously reported to be transcriptionally controlled by the PhoR/PhoB TCS [17], is now known to be instead regulated by the stationary phase sigma factor RpoS [18]. Earlier interpretations from the same investigators were based on indirect effects of the PhoR/PhoB TCS under conditions of  $P_i$  limitation.

## The Pst system is the predominant system for $P_i$ uptake

Nearly all genes directly controlled by the PhoR/PhoB TCS have a role in assimilation of  $P_i$  or an alternative P source for growth (Table 1). The most strongly activated promoter *pstSp* (for the *pstSCAB-phoU* operon) governs expression of the ABC transporter Pst and PhoU [1]. It had until recently been thought that the Pst system has a role in  $P_i$  uptake only under conditions of  $P_i$  limitation. A variety of data now show that the Pst system, not the low affinity “phosphate inorganic transporter” PitA, serves as the primary  $P_i$  transporter when  $P_i$  is in excess. PitA is unlikely to act primarily as a  $P_i$  transporter, but rather as a transporter of divalent metal cations ( $Zn^{2+}$ ) that are transported in complex with  $P_i$  [19]. A primary role for PitA as a  $Zn^{2+}$ , and not a  $P_i$ , transporter is supported by the finding that *pitA* expression is activated by  $Zn^{2+}$ , and not by  $P_i$  limitation [20;21]. Likewise, *pitB* [22;23] probably has no role in  $P_i$  uptake in normal cells, as it is not expressed under normal growth conditions.

## The PhoB RR acts as a transcription factor for Pho regulon promoters

PhoB belongs to the OmpR/PhoB subfamily, the largest group of RRs. PhoB is comprised of an N-terminal receiver domain and a C-terminal DNA-binding domain. Its activity as transcription factor depends upon its state of phosphorylation (D53) of the PhoB receiver domain. Several structures of PhoB have been determined of both its receiver and DNA-binding domain (without and with  $Mg^{++}$  and DNA; [www.prfect.org/EcoliProteins](http://www.prfect.org/EcoliProteins)), including those of two “constitutively active” mutants [24-27]. NMR studies have also examined the activation mechanism for receiver domain ([28]; see also [29] in this volume) and the mechanism of DNA binding [30].

## The PhoR HK lacks a P<sub>i</sub> sensory domain

PhoR acts as the P<sub>i</sub> sensory HK and is essential for three distinct processes that control PhoB activity as a transcription factor: inhibition (prevention of PhoB phosphorylation), activation (phosphorylation of PhoB), and deactivation (dephosphorylation of phospho-PhoB). As shown in Fig. 1, PhoR is comprised of five domains (or regions). Its N-terminal transmembrane (TM) domain is required solely for association of PhoR to the membrane. Presumably, membrane localization of PhoR is necessary for interaction with the Pst transporter. PhoR acts as a sensory protein via an interaction between a cytosolic domain of PhoR (possibly its PAS domain; Y.H. and B.L.W., manuscript in preparation) and the Pst transporter (possibly the ABC component PstB; Fig. 2) and/or PhoU.

## Cross regulation of Pho regulon by non-partner HKs

PhoB can also be activated in the absence of PhoR. Activation of PhoB in the absence of PhoR is due to cross regulation (PhoB phosphorylation; [1]) by non-partner HKs such as CreC [31] or small molecule phosphoryl donor(s) such as acetyl phosphate [32]. When PhoR is absent, the non-partner HKs ArcB, CreC, KdpD, and QseC can lead to moderate activation of PhoB in response to different growth conditions, while the non-partner HKs BaeS and EnvZ can lead to low level activation [4;33]. It should be noted that these studies were carried out by examining gene expression in cultures, in which gene expression levels reflect only population averages and not the dynamics of gene expression in single cells.

## Stochastic expression of the Pho regulon

Single-cell profiling by using flow cytometry to monitor gene expression in single cells has revealed an unforeseen stochastic, “all-or-none,” character for activation of PhoB by non-partner HKs [4]. Modeling has shown that stochastic behavior can result not only from TCSs that have a positive feedback loop (i. e., phospho-PhoB leads to autoamplification of PhoB synthesis) but also from systems in which the rate of HK translation initiation is limited (as appears to be the case for PhoR [34]). Accordingly, the low amounts of PhoR resulting from low rates of PhoR translation are expected to lead to the formation of occasional cells in a population having no PhoR protein. Activation of PhoB by non-partner HKs in these cells would lead to stochastic activation of PhoB and to the emergence of multiple stable phenotypes within a population of genetically identical cells. Such behavior at the cellular level is likely to be of fundamental importance not only in the recovery of cells from periods of stress but also in persistence, host-phage interactions and pathogenesis [35-38]. While other TCSs have not been similarly tested for stochasticity, it is reasonable to propose that several are likely to exhibit similar bimodal expression patterns. Two characteristics that appear to be important for stochastic behavior are: (i) the presence of an autoregulatory loop controlling expression of the TCS; and (ii) low translation rates for the HK mRNA [34].

## The Pst transporter is required for P<sub>i</sub> signal transduction

Early studies showed that the Pst transporter is essential for detecting environmental P<sub>i</sub>. Also, recent data show that PhoR detects P<sub>i</sub> only indirectly (Y.H. and B.L.W., manuscript in preparation). Further, the Pst system but not P<sub>i</sub> uptake *per se* is essential for P<sub>i</sub> signaling by the Pst system [1]. By analogy to the ABC (MalEFGK) transporter for maltose [39], we propose that the Pst transporter exists in two distinct states: in one state, the Pst transporter is both transport and signaling active; and in the other, the Pst transporter is both transport and signaling inactive. These states would correspond to closed (transport active) conformation when P<sub>i</sub> is bound and an open (resting state) conformation in the absence of bound P<sub>i</sub>. Thus, mutations of the Pst system that abolish P<sub>i</sub> uptake without affecting P<sub>i</sub> signaling block uptake but yet allow formation of the closed and open conformations [1].

## A model for $P_i$ signaling

Mechanistically,  $P_i$  signaling is a negative process. Excess  $P_i$  is required for turning the system off. Activation is the default state and results under conditions of  $P_i$  limitation. The Pst transporter is essential for inhibition, as well as deactivation [1]. Deactivation resets the PhoR/PhoB system to its inhibition state (Fig. 2). That activation (phosphorylation) of PhoB leads to a conformational change in PhoB has been shown by examination of the structural changes brought about by phosphoryl group analog  $\text{BeF}_3^-$  [28] and the structure of constitutively active PhoB proteins [27].

Like the Pst transporter, PhoU also has an obligatory role in both inhibition and deactivation of PhoB. The finding that PhoU-like proteins from *Aquifex aeolicus* and *Thermotoga maritima* share structural similarity with proteins belonging to the eukaryotic chaperone Hsp70 family [13;14] support a chaperone-like role for PhoU. The action of PhoU as an accessory protein is fully compatible with PhoU being a chaperone. Accordingly, PhoU probably acts together with PhoR to promote autodephosphorylation of PhoB-P [40].

A caveat of  $P_i$  signaling by the proposed PhoR/PhoB/PstSCAB/PhoU complex is that individual complexes can exist in different states within a cell. Accordingly, when  $P_i$  is in excess, all complexes probably exist in the transport and signaling active state, in which PhoR would be in the inhibition state. Under conditions of  $P_i$  limitation, these complexes probably exist in different states within the same cell. That is, under these conditions, some complexes would be in the transport and signaling inactive (PhoR activation) state. Other complexes would be in the transport and signaling active (PhoR inhibition) state. The existence of complexes in both states within the same cell would be necessary to permit simultaneous activation of PhoR/PhoB-regulated genes and growth on limiting amounts of  $P_i$ .

## Conclusions

Much new information has been learned about the molecular control of the Pho regulon over the past decade, especially with respect to signaling by environmental  $P_i$ . Three areas are likely to contribute substantial new information about the Pho regulon and its control in the future (Box 1).

### Key problems for future studies of the PhoR/PhoB TCS

- The advent of genome-wide mRNA analysis by deep sequencing (RNA-seq) coupled with chromatin immunoprecipitation (ChIP-seq) can provide unprecedented sensitivity and specificity for protein-DNA interactions on a genome-wide scale [41]. Application of such technology to  $P_i$  signal transduction should provide comprehensive identification of genes controlled by the PhoR/PhoB TCS.
- Studying single-cell gene expression by the PhoR/PhoB TCS under diverse environmental conditions is likely to provide definitive results regarding the role of cross regulation among different TCSs.
- Studying the different states of the proposed seven-component  $P_i$  signaling complex is likely to require development of new technologies that enable examination of single protein complexes inside living cells that are similar to ones now being used to study activities of other machines at the single molecule level [42].

## Acknowledgments

Research from this laboratory has been supported by NIH GM62662 and GM83296.


## References and recommended reading

1. Wanner, BL. Phosphorus assimilation and control of the phosphate regulon. In: Neidhardt, FC.; Curtiss, R., III; Ingraham, JL.; Lin, ECC.; Low, KB., Jr; Magasanik, B.; Reznikoff, WS.; Riley, M.; Schaechter, M.; Umberger, HE., editors. *Escherichia coli* and *Salmonella typhimurium* cellular and molecular biology. 2. Washington, D.C.: ASM Press; 1996. p. 1357-1381.
- 2. Noriega CE, Lin HY, Chen LL, Williams SB, Stewart V. Asymmetric cross-regulation between the nitrate-responsive NarX-NarL and NarQ-NarP two-component regulatory systems from *Escherichia coli* K-12. *Mol Microbiol* 2010;75:394–412. [PubMed: 19968795] This manuscript provides evidence for an asymmetry for cross regulation in the NarX-NarL and NarQ-NarP TCSs network. While NarQ interacted similarly with both NarL and NarP RRs, NarX preferentially interacts with NarL.
3. Nishino K, Honda T, Yamaguchi A. Genome-wide analyses of *Escherichia coli* gene expression responsive to the BaeSR two-component regulatory system. *J Bacteriol* 2005;187:1763–1772. [PubMed: 15716448]
4. Zhou, L.; Gregori, G.; Blackman, JM.; Robinson, JP.; Wanner, BL. Stochastic activation of the response regulator PhoB by noncognate histidine kinases. In: Hofestädt, R., editor. *IMBio, Informationsmanagement in der Biotechnologie e V. Magdeburg*: 2005. p. 11-24.
5. VanBogelen RA, Olson ER, Wanner BL, Neidhardt FC. Global analysis of proteins synthesized during phosphorus restriction in *Escherichia coli*. *J Bacteriol* 1996;178:4344–4366. [PubMed: 8755861]
6. Yuan ZC, Zaheer R, Morton R, Finan TM. Genome prediction of PhoB regulated promoters in *Sinorhizobium meliloti* and twelve proteobacteria. *Nucleic Acids Res* 2006;34:2686–2697. [PubMed: 16717279]
7. Baek JH, Lee SY. Novel gene members in the Pho regulon of *Escherichia coli*. *FEMS Microbiol Lett* 2006;264:104–109. [PubMed: 17020555]
8. Murray EL, Conway T. Multiple regulators control expression of the Entner-Doudoroff aldolase (Eda) of *Escherichia coli*. *J Bacteriol* 2005;187:991–1000. [PubMed: 15659677]
9. Gorelik M, Lunin VV, Skarina T, Savchenko A. Structural characterization of GntR/HutC family signaling domain. *Protein Sci* 2006;15:1506–1511. [PubMed: 16672238]
10. Adams MA, Luo Y, Hove-Jensen B, He SM, van Staalduinen LM, Zechel DL, Jia Z. Crystal structure of PhnH: an essential component of carbon-phosphorus lyase in *Escherichia coli*. *J Bacteriol* 2008;190:1072–1083. [PubMed: 17993513]
11. Hove-Jensen B, Rosenkrantz TJ, Haldimann A, Wanner BL. *Escherichia coli* *phnN*, encoding ribose 1,5-bisphosphokinase activity (phosphoribosyl diphosphate forming): dual role in phosphonate degradation and NAD biosynthesis pathways. *J Bacteriol* 2003;185:2793–2801. [PubMed: 12700258]
12. Podzelinska K, He SM, Wathier M, Yakunin A, Proudfoot M, Hove-Jensen B, Zechel DL, Jia Z. Structure of PhnP, a phosphodiesterase of the carbon-phosphorus lyase pathway for phosphonate degradation. *J Biol Chem* 2009;284:17216–17226. [PubMed: 19366688]
13. Oganessian V, Oganessian N, Adams PD, Jancarik J, Yokota HA, Kim R, Kim SH. Crystal structure of the “PhoU-like” phosphate uptake regulator from *Aquifex aeolicus*. *J Bacteriol* 2005;187:4238–4244. [PubMed: 15937186]
14. Liu J, Lou Y, Yokota H, Adams PD, Kim R, Kim SH. Crystal structure of a PhoU protein homologue: a new class of metalloprotein containing multinuclear iron clusters. *J Biol Chem* 2005;280:15960–15966. [PubMed: 15716271]
15. Riley M, Abe T, Chaudhuri RR, Horiuchi T, Mori H, Perna NT, Plunkett G III, Rudd KE, Serres MH, Wanner BL, et al. *Escherichia coli* K-12: a cooperatively developed annotation snapshot--2005. *Nucleic Acids Res* 2006;34:1–9. [PubMed: 16397293]
16. Yamazaki Y, Niki H, Kato J. Profiling of *Escherichia coli* Chromosome database. *Methods Mol Biol* 2008;416:385–389. [PubMed: 18392982]



17. Suziedeliene E, Suziedelis K, Garbenciute V, Normark S. The acid-inducible *asr* gene in *Escherichia coli*: transcriptional control by the *phoBR* operon. *J Bacteriol* 1999;181:2084–2093. [PubMed: 10094685]
18. Seputiene V, Suziedelis K, Normark S, Melefors O, Suziedeliene E. Transcriptional analysis of the acid-inducible *asr* gene in enterobacteria. *Res Microbiol* 2004;155:535–542. [PubMed: 15313253]
19. Beard SJ, Hashim R, Wu G, Binet MR, Hughes MN, Poole RK. Evidence for the transport of zinc (II) ions via the *pit* inorganic phosphate transport system in *Escherichia coli*. *FEMS Microbiol Lett* 2000;184:231–235. [PubMed: 10713426]
20. Jackson RJ, Binet MR, Lee LJ, Ma R, Graham AI, McLeod CW, Poole RK. Expression of the PitA phosphate/metal transporter of *Escherichia coli* is responsive to zinc and inorganic phosphate levels. *FEMS Microbiol Lett* 2008;289:219–224. [PubMed: 19054109]
21. Graham AI, Hunt S, Stokes SL, Bramall N, Bunch J, Cox AG, McLeod CW, Poole RK. Severe zinc depletion of *Escherichia coli*: roles for high affinity zinc binding by ZinT, zinc transport and zinc-independent proteins. *J Biol Chem* 2009;284:18377–18389. [PubMed: 19377097]
22. Harris RM, Webb DC, Howitt SM, Cox GB. Characterization of PitA and PitB from *Escherichia coli*. *J Bacteriol* 2001;183:5008–5014. [PubMed: 11489853]
23. Hoffer SM, Schoondermark P, Van Veen HW, Tommassen J. Activation by Gene Amplification of *pitB*, Encoding a Third Phosphate Transporter of *Escherichia coli* K-12. *J Bacteriol* 2001;183:4659–4663. [PubMed: 11443103]
24. Sola M, Gomis-Ruth FX, Serrano L, Gonzalez A, Coll M. Three-dimensional crystal structure of the transcription factor PhoB receiver domain. *J Mol Biol* 1999;285:675–687. [PubMed: 9878437]
25. Blanco AG, Sola M, Gomis-Ruth FX, Coll M. Tandem DNA Recognition by PhoB, a Two-Component Signal Transduction Transcriptional Activator. *Structure (Camb)* 2002;10:701–713. [PubMed: 12015152]
26. Sola M, Drew DL, Blanco AG, Gomis-Ruth FX, Coll M. The cofactor-induced pre-active conformation in PhoB. *Acta Crystallogr D Biol Crystallogr* 2006;62:1046–1057. [PubMed: 16929106]
27. Arribas-Bosacoma R, Kim SK, Ferrer-Orta C, Blanco AG, Pereira PJ, Gomis-Ruth FX, Wanner BL, Coll M, Sola M. The X-ray crystal structures of two constitutively active mutants of the *Escherichia coli* PhoB receiver domain give insights into activation. *J Mol Biol* 2007;366:626–641. [PubMed: 17182055]
28. Bachhawat P, Swapna GV, Montelione GT, Stock AM. Mechanism of activation for transcription factor PhoB suggested by different modes of dimerization in the inactive and active states. *Structure* 2005;13:1353–1363. [PubMed: 16154092]
29. Gao R, Stock AM. Molecular strategies for phosphorylation mediated regulation of response regulator activity. *Curr Opin Microbiol* 2010;13
30. Yamane T, Okamura H, Ikeguchi M, Nishimura Y, Kidera A. Water-mediated interactions between DNA and PhoB DNA-binding/transactivation domain: NMR-restrained molecular dynamics in explicit water environment. *Proteins* 2008;71:1970–1983. [PubMed: 18186481]
31. Yamamoto K, Hirao K, Oshima T, Aiba H, Utsumi R, Ishihama A. Functional characterization *in vitro* of all two-component signal transduction systems from *Escherichia coli*. *J Biol Chem* 2005;280:1448–1456. [PubMed: 15522865]
32. Zundel CJ, Capener DC, McCleary WR. Analysis of the conserved acidic residues in the regulatory domain of PhoB. *FEBS Lett* 1998;441:242–246. [PubMed: 9883892]
33. Kim SK, Wilmes-Riesenberg MR, Wanner BL. Involvement of the sensor kinase EnvZ in the *in vivo* activation of the response-regulator PhoB by acetyl phosphate. *Mol Microbiol* 1996;22:135–147. [PubMed: 8899716]
- 34. Kierzek AM, Zhou L, Wanner BL. Stochastic kinetic model of two component system signalling reveals all-or-none, graded and mixed mode stochastic switching responses. *Mol Biosyst*. 2010;10:1039/b906951hThe authors describe kinetic parameters based on mathematic modeling how TCSs can exhibit all-or-none, graded or mixed mode responses. Modeling is based on data in reference 6 above.
35. Pearl S, Gabay C, Kishony R, Oppenheim A, Balaban NQ. Nongenetic individuality in the host-phage interaction. *PLoS Biol* 2008;6:e120. [PubMed: 18494559]


36. Gefen O, Balaban NQ. The importance of being persistent: heterogeneity of bacterial populations under antibiotic stress. *FEMS Microbiol Rev* 2009;33:704–717. [PubMed: 19207742]
37. Glover WA, Yang Y, Zhang Y. Insights into the molecular basis of L-form formation and survival in *Escherichia coli*. *PLoS ONE* 2009;4:e7316. [PubMed: 19806199]
38. Avery SV. Microbial cell individuality and the underlying sources of heterogeneity. *Nat Rev Microbiol* 2006;4:577–587. [PubMed: 16845428]
39. Oldham ML, Khare D, Quijcho FA, Davidson AL, Chen J. Crystal structure of a catalytic intermediate of the maltose transporter. *Nature* 2007;450:515–521. [PubMed: 18033289]
- 40. Lamarche MG, Wanner BL, Crepin S, Harel J. The phosphate regulon and bacterial virulence: a regulatory network connecting phosphate homeostasis and pathogenesis. *FEMS Microbiol Rev* 2008;32:461–473. [PubMed: 18248418] Authors review evidence for a role of the Pho regulon in bacterial virulence.
41. Hu M, Yu J, Taylor JM, Chinnaiyan AM, Qin ZS. On the detection and refinement of transcription factor binding sites using ChIP-Seq data. *Nucleic Acids Res.* 2010
42. Allemand JF, Maier B. Bacterial translocation motors investigated by single molecule techniques. *FEMS Microbiol Rev* 2009;33:593–610. [PubMed: 19243443]



**Fig. 1.**

Domain organization of PhoR. TM, transmembrane-anchoring domain; CR, positively charged linker region; PAS, Per-Arnt-Sim domain; DHp, dimerization and histidine phosphoacceptor domain; CA, a catalytic domain.



**Fig. 2.**

Model for transmembrane signal transduction by environmental  $P_i$ . The signaling processes of inhibition, activation, and deactivation are proposed to correspond to different states of PhoR: an inhibition state (PhoR<sup>I</sup>), an activation state (PhoR<sup>A</sup>), and a deactivation state (PhoR<sup>D</sup>). The  $P_i$  binding protein PstS is fully saturated when  $P_i$  is in excess. Under these conditions, a signal is propagated to PhoR leading to formation of PhoR<sup>I</sup>, which interferes with phosphorylation of PhoB. No such signal exists under conditions of  $P_i$  limitation (or absence of a Pst component), leading to formation of the default state PhoR<sup>A</sup> which acts as a phospho-donor for autophosphorylation of PhoB. Following a period of  $P_i$  limitation, PhoR<sup>D</sup> promotes dephosphorylation of phospho-PhoB. Formation of PhoR<sup>D</sup> requires an increased amount of PhoU or PstB in addition to excess  $P_i$ .

Table 1

Genes of the *E. coli* K-12 phosphate regulon

Gene	ECK number <sup>a</sup>	Product description <sup>b</sup>	References
<i>amn</i>	ECK1977	AMP nucleosidase	[7]
<i>eda</i>	ECK1851	aldolase	[8]
<i>phnC</i>	ECK4099	phosphonate transporter subunit, predicted ATP-binding component	[1]
<i>phnD</i>	ECK4098	phosphonate transporter subunit, periplasmic-binding component	[1]
<i>phnE</i>	ECK4096	phosphonate transporter subunit, membrane component	[1]
<i>phnF</i>	ECK4095	predicted transcription regulator, GntR/HutC family	[1;9]
<i>phnG</i>	ECK4094	carbon-phosphorus lyase complex subunit	[1]
<i>phnH</i>	ECK4093	carbon-phosphorus lyase complex subunit	[1;10]
<i>phnI</i>	ECK4092	carbon-phosphorus lyase complex subunit	[1]
<i>phnJ</i>	ECK4091	carbon-phosphorus lyase complex subunit	[1]
<i>phnK</i>	ECK4090	carbon-phosphorus lyase complex subunit, predicted ATP-binding component	[1]
<i>phnL</i>	ECK4089	carbon-phosphorus lyase complex subunit, predicted ATP-binding component	[1]
<i>phnM</i>	ECK4088	carbon-phosphorus lyase complex subunit, membrane component	[1]
<i>phnN</i>	ECK4087	carbon-phosphorus lyase complex subunit, predicted ATP-binding component, ribose 1,5-bisphosphokinase activity protein	[1;11]
<i>phnO</i>	ECK4086	carbon-phosphorus lyase complex subunit, predicted acyltransferase with acyl-CoA N-acyltransferase domain	[1]
<i>phnP</i>	ECK4085	carbon-phosphorus lyase complex accessory protein, phosphodiesterase activity protein	[1;12]
<i>phoA</i>	ECK0378	bacterial alkaline phosphatase	[1]
<i>phoB</i>	ECK0393	DNA-binding response regulator	[1]
<i>phoE</i>	ECK0242	outer membrane phosphoporin protein E	[1]
<i>phoH</i>	ECK1010	conserved protein with nucleoside triphosphate hydrolase domain	[1]
<i>phoR</i>	ECK0394	sensory histidine kinase	[1]
<i>phoU</i>	ECK3717	chaperone-like PhoR/PhoB inhibitory protein	[1;13;14]
<i>psiE</i>	ECK4022	predicted phosphate starvation-inducible protein E	[1]
<i>psiF</i>	ECK0379	predicted phosphate starvation-inducible protein F	[1]
<i>pstA</i>	ECK3719	phosphate transporter subunit, membrane component	[1]
<i>pstB</i>	ECK3718	phosphate transporter subunit, ATP-binding component	[1]
<i>pstC</i>	ECK3720	phosphate transporter subunit, membrane component	[1]
<i>pstS</i>	ECK3721	phosphate transporter subunit, periplasmic-binding component	[1]
<i>ugpA</i>	ECK3436	glycerol-3-phosphate transporter subunit	[1]
<i>ugpB</i>	ECK3437	glycerol-3-phosphate transporter subunit, periplasmic-binding component	[1]
<i>ugpC</i>	ECK3434	glycerol-3-phosphate transporter subunit, ATP-binding component	[1]
<i>ugpE</i>	ECK3435	glycerol-3-phosphate transporter subunit, membrane component	[1]
<i>ugpQ</i>	ECK3433	glycerol-3-phosphate transporter subunit, membrane component	[1]
<i>yibD</i>	ECK3605	predicted glycosyl transferase	[7]
<i>ytfK</i>	ECK4213	conserved protein	[7]

<sup>a</sup>ECK numbers are in accordance with Riley et al. [15]

<sup>b</sup>Product descriptions are in accordance with Riley et al. [15], the latest GenBank record for *E. coli* K-12 MG1655 (U00096 dated July 2009), and the EcoGene ([www.ecogene.org](http://www.ecogene.org)) and PEC (Profiling of *E. coli* Chromosome [16]; <http://www.shigen.nig.ac.jp/ecoli/pec/>) databases (December 2009 version).