

# Induction of *Salmonella* pathogenicity island 1 under different growth conditions can affect *Salmonella*–host cell interactions *in vitro*

J. Antonio Ibarra,<sup>1</sup> Leigh A. Knodler,<sup>1</sup> Daniel E. Sturdevant,<sup>2</sup> Kimmo Virtaneva,<sup>2</sup> Aaron B. Carmody,<sup>3</sup> Elizabeth R. Fischer,<sup>4</sup> Stephen F. Porcella<sup>2</sup> and Olivia Steele-Mortimer<sup>1</sup>

## Correspondence

Olivia Steele-Mortimer  
omortimer@niaid.nih.gov

<sup>1</sup>Laboratory of Intracellular Parasites, Rocky Mountain Laboratories, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Hamilton, MT 59840, USA

<sup>2</sup>Genomics Unit, Rocky Mountain Laboratories, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Hamilton, MT 59840, USA

<sup>3</sup>Flow Cytometry Unit, Rocky Mountain Laboratories, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Hamilton, MT 59840, USA

<sup>4</sup>Microscopy Unit, Rocky Mountain Laboratories, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Hamilton, MT 59840, USA

*Salmonella* invade non-phagocytic cells by inducing massive actin rearrangements, resulting in membrane ruffle formation and phagocytosis of the bacteria. This process is mediated by a cohort of effector proteins translocated into the host cell by type III secretion system 1, which is encoded by genes in the *Salmonella* pathogenicity island (SPI) 1 regulon. This network is precisely regulated and must be induced outside of host cells. *In vitro* invasive *Salmonella* are prepared by growth in synthetic media although the details vary. Here, we show that culture conditions affect the frequency, and therefore invasion efficiency, of SPI1-induced bacteria and also can affect the ability of *Salmonella* to adapt to its intracellular niche following invasion. Aerobically grown late-exponential-phase bacteria were more invasive and this was associated with a greater frequency of SPI1-induced, motile bacteria, as revealed by single-cell analysis of gene expression. Culture conditions also affected the ability of *Salmonella* to adapt to the intracellular environment, since they caused marked differences in intracellular replication. These findings show that induction of SPI1 under different pre-invasion growth conditions can affect the ability of *Salmonella* to interact with eukaryotic host cells.

Received 18 July 2009

Revised 15 December 2009

Accepted 17 December 2009

## INTRODUCTION

Type III secretion systems (T3SS) are important virulence determinants for many Gram-negative pathogens. These systems consist of a specialized secretion apparatus that translocates effector proteins from the bacterial cytoplasm into the host cell. Translocated T3SS effector proteins effectively allow the bacteria to ‘hijack’ many essential

**Abbreviations:** aer-LL, *Salmonella* grown with aeration to late exponential phase; FDR, false discovery rate; GSEA, gene set enrichment analysis; p.i., post-infection; PFA, paraformaldehyde; qPCR, quantitative PCR; SCV, *Salmonella*-containing vacuole; SPI, *Salmonella* pathogenicity island;  $\mu$ aer-ST, *Salmonella* grown without aeration to stationary phase; T3SS, type III secretion system; WT, wild-type.

The microarray data discussed in this paper are available from the NCBI GEO database under accession number GSE13343.

Four supplementary figures and a supplementary table are available with the online version of this paper.

intracellular processes. *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) is a facultative intracellular pathogen that causes gastroenteritis in humans and a systemic typhoid-like disease in susceptible mice. *S. Typhimurium* virulence is dependent on two T3SS (T3SS1 and T3SS2) that are used for invasion of non-phagocytic host cells and modification of the intracellular environment. T3SS1 translocates effector proteins required for invasion of non-phagocytic host cells and some post-invasion events [for a review, see the paper by McGhie *et al.* (2009)]. In contrast, T3SS2 is induced intracellularly and its effectors are translocated into the host cell across the membrane of the *Salmonella*-containing vacuole (SCV), a modified phagosome within which the bacteria survive and replicate (Steele-Mortimer, 2008b). Most of the genes encoding T3SS1 and T3SS2 structural components as well as many regulatory factors and effectors are located on *Salmonella* pathogenicity islands (SPIs) 1 and 2, respectively, although

some effector proteins are encoded on other pathogenicity islands or islets (Hansen-Wester & Hensel, 2001; Marcus *et al.*, 2000).

Various environmental signals such as oxygen concentration, osmolarity and bacterial growth state have been shown to influence the expression of SPII genes and the secretion of T3SS1 effectors (Ellermeier & Slauch, 2007). A number of *in vitro* culture conditions for obtaining SPII-induced invasive *S. Typhimurium* have been described. In aerated culture, the invasive phenotype is induced during a brief period at the end of exponential growth, or late exponential phase (Song *et al.*, 2004; Steele-Mortimer *et al.*, 1999), although other studies have shown that oxygen-limiting, or microaerophilic, conditions are required for optimal induction and that, under these conditions, invasiveness is induced during stationary phase (Bajaj *et al.*, 1996; Behlau & Miller, 1993; Jones & Falkow, 1994; Lee & Falkow, 1990; Schiemann & Shope, 1991; Temme *et al.*, 2008).

In this study, we have grown *S. Typhimurium* under two distinct 'invasion-inducing' conditions and then analysed gene expression and the ability of the bacteria to invade and replicate within cultured epithelial cells. Bacteria were grown in Luria–Bertani (LB)–Miller broth, either to late-exponential phase with aeration (aer-LL) (Steele-Mortimer *et al.*, 1999) or to stationary phase in the absence of aeration under microaerophilic conditions ( $\mu$ aer-ST). Aer-LL bacteria were more invasive than  $\mu$ aer-ST bacteria and, although transcriptome analysis did not reveal significant variation in SPII expression, single-cell analysis revealed differences in frequency of SPII-induced bacteria, which were predominant in the aer-LL culture. Moreover, co-expression of flagella and SPII genes was associated with higher invasion levels. Finally, we found that intracellular replication could be affected by pre-invasion growth conditions and this was associated with differences in intracellular expression of SPII genes. Altogether, this study suggests that pre-invasion growth conditions used to induce invasive bacteria *in vitro* can be an influential factor in the outcome of *Salmonella*–host cell interactions.

## METHODS

**Bacterial cultures and growth conditions.** *S. Typhimurium* SL1344 (Hoiseth & Stocker, 1981) was used in all experiments unless otherwise indicated.  $\Delta$ SPII::kan has been described previously (Drecktrah *et al.*, 2006). The flagellar mutants *fliC*::Tn10, *fliB*::Mud-Cm and *flgB*::Tn10 and the *fimAICDHF*::kan mutant were obtained by P22 transduction of *S. Typhimurium* SL1344 with phages kindly provided by Ed Miao (Miao *et al.*, 2006) and Andreas Bäumler (Weening *et al.*, 2005). Bacteria were initially streaked on LB–Miller agar (10 g tryptone l<sup>-1</sup>, 5 g yeast extract l<sup>-1</sup>, 10 g NaCl l<sup>-1</sup>, pH 7.0) supplemented with streptomycin (100  $\mu$ g ml<sup>-1</sup>), kanamycin (50  $\mu$ g ml<sup>-1</sup>), chloramphenicol (30  $\mu$ g ml<sup>-1</sup>), carbenicillin (50  $\mu$ g ml<sup>-1</sup>) and/or tetracycline (10  $\mu$ g ml<sup>-1</sup>). Fresh plates were streaked every week and stored at 4 °C. The  $\mu$ aer-ST bacteria were prepared by inoculating one colony into 3 ml LB–Miller broth in a 17 × 100 mm, 14 ml polypropylene round-bottom test tube

(Becton Dickinson) with a loose cap, and then incubating at 37 °C without aeration (no shaking) for 18 h. Aer-LL bacteria were prepared by inoculating one colony into 2 ml LB–Miller broth, in a 17 × 100 mm, 14 ml polypropylene round-bottom test tube (Becton Dickinson) with a loose cap, and incubating at 37 °C with aeration (shaking at 225 r.p.m.) for 16–18 h. Thereafter, a 300  $\mu$ l aliquot of this culture was immediately subcultured into 10 ml LB–Miller broth, in a 150 ml glass Erlenmeyer flask with a loose cap, and incubated at 37 °C for 3.5 h with aeration (shaking at 225 r.p.m.).

**Measurement of bacterial growth.** For measurement of bacterial growth, *Salmonella* was cultured as described, then samples were taken every hour for OD<sub>600</sub> readings in a SmartSpec 3000 spectrophotometer (Bio-Rad) and diluted in sterile 1 × PBS, pH 7.2, and plated on LB–Miller agar for enumeration of viable bacteria.

**Plasmid stability.** Plasmid stabilities in culture, aerated and non-aerated as described, or in infected cells were confirmed by plating on LB Miller agar and by comparing the c.f.u. obtained in the presence (50  $\mu$ g carbenicillin ml<sup>-1</sup>) or absence (no antibiotic) of selection.

***Salmonella* Affymetrix array.** The *S. Typhimurium* SL1344 (NCTC 13347) genome was first analysed by Integrated Genomics to remove genetic redundancy and provide annotation for intergenic and opposite-strand coding regions using the sequence contigs from the Sanger Institute ([www.sanger.ac.uk](http://www.sanger.ac.uk)). The microarray is an 11 micron, 49-5241 format array containing 108 166 probe pairs or about 9833 eleven-probe-pair probe sets specific to 4712 ORFs and 5121 intergenic and opposite-strand coding targets (Affymetrix).

**Microarray experiments.** For each condition ( $\mu$ aer-ST or aer-LL *Salmonella*) six biological replicates were prepared. Total RNA from each sample was stabilized with RNAProtect (Qiagen). The concentrate was fragmented with a QIAshredder (Qiagen) and total RNA was purified in 96-well format using the RNeasy 96 kit centrifugation system (Qiagen) as described previously (Virtaneva *et al.*, 2005). RNA samples were treated with DNA-free DNase I (Ambion) and sample quality was monitored on Agilent RNA Pico II LabChips (Agilent Technologies). Additionally, possible DNA contamination was tested by quantitative PCR (qPCR; see below). Total RNA (10  $\mu$ g) from each sample was reverse-transcribed to cDNA using 4.5  $\mu$ g random hexamer primers (Invitrogen) and 30 U SuperScript III reverse transcriptase (Invitrogen). RNA was removed by treatment with 0.25 M NaOH for 30 min at 65 °C and then neutralized with 1 M HCl. cDNA was purified using a QIAquick 96 PCR purification kit (Qiagen), quantified and fragmented with 0.6 U DNase I (Roche). An Enzo BioArray Terminal Labelling kit with Biotin-ddUTP (EnzoLite Sciences) was used to label the 3' termini of the fragmented cDNA products. Labelled and fragmented cDNA (2  $\mu$ g) was hybridized to *Salmonella* GeneChips following a previously described procedure (Virtaneva *et al.*, 2005).

**Transcriptome analysis.** Affymetrix GeneChip Operating Software (GCOS v1.4, <http://www.affymetrix.com>) was used to perform the preliminary analysis of the custom chips at the probe-set level. A pivot table with all samples was created including calls, call *P*-value and signal intensities for each gene. The pivot table was then imported into GeneSpring GX 7.3 (Agilent), where hierarchical clustering (condition tree) using a Pearson correlation similarity measure with average linkage was used to produce the dendrogram indicating that biological replicates were grouping together (data not shown). The CEL files were imported into Partek Genomics Suite and quantile normalized to produce a principal component analysis (PCA) plot as a secondary statistical check on biological replicates grouping together. A one-way ANOVA was run from these datasets to produce a false discovery rate (FDR) report of false positive reduced *P*-values for each of the treatments. Once the data had passed all preliminary statistical

tests, the biological replicates were combined. A custom Microsoft Excel worksheet was then used to combine correlated replicates of all test conditions and controls. Quality filters based upon combined calls and signal intensities were used in the worksheet in order to further evaluate individual gene comparisons. Present and marginal were treated as the same while absent calls were negatively weighted for the filters and dropped completely from quality summations. All individual genes passing the above filters and combined from all usable replicates have the ratios of test:control reported with associated probability of Student's *t*-test values. Significance analysis of microarrays (SAM) (Tusher *et al.*, 2001) was also run within the Excel sheet and a column was added with the results. An additional column was added for Partek's ANOVA *P*-values that passed the FDR.

**Gene set enrichment analysis (GSEA).** GSEA was performed using gene sets obtained from the Kyoto Encyclopaedia of Genes and Genomes (KEGG; <http://www.genome.jp/kegg/pathway.html>) and ERGO (Integrated Genomics) (Kanehisa *et al.*, 2008). These databases were curated to include the SPI1 and SPI2 regulon gene sets from previous microarray analysis (Eriksson *et al.*, 2003; Hautefort *et al.*, 2008). The enriched sets were analysed using GSEA (Mootha *et al.*, 2003; Subramanian *et al.*, 2005, 2007) and SAM (Tusher *et al.*, 2001) with 1000 permutations and an FDR of 0.25 for the significance threshold.

**qPCR analysis.** Briefly, RNA samples were obtained from separate experiments and verified on Agilent RNA Pico II LabChips. For qPCR, the CellsDirect One-Step qRT-PCR kit (Invitrogen) was used according to manufacturer's instructions. Primers and probes for different genes were designed using Primer Express v2.0 software for multiplex qPCR (Applied Biosystems) and 100% amplification efficiency and compatibility were verified with *S. Typhimurium* genomic DNA. Relative quantification of the gene expression was determined with the comparative threshold ( $\Delta\Delta C_T$ ) method (Applied Biosystems). *ftsZ* was chosen as an internal control because no significant variation in expression for this gene was observed under the conditions used [this study and Drecktrah *et al.* (2006)]. In order to detect the correlation between microarray and qPCR fold changes, the Pearson product-moment correlation coefficient (*r*) was calculated (using Microsoft Excel).

**Transcriptional fusions.** A destabilized variant of green fluorescent protein, GFP[LVA], with a half-life of approximately 40 min in *S. Typhimurium* (data not shown) was used as a reporter of gene expression (Andersen *et al.*, 1998). Promoters from the following six genes were selected to drive *gfp*[LVA] expression: *invF*, *prgH*, *ssaG*, *sopB*, *pipB* and *fliC*. *P<sub>invF</sub>* was amplified with the oligonucleotides *invF*-Xho (5'-ccg ctc gag aag aat gag gcg cca tgt-3') and *invF*-Kpn (5'-cgg ggt acc aca acg gcc tgc tgc caa-3') (engineered restriction sites are underlined). *P<sub>prgH</sub>* was amplified with *prgH*-Xho (5'-ccg ctc gag ttg taa atg ctc ttt att-3') and *prgH*-Kpn (5'-cgg ggt acc aca aag agt gtt cgg cct-3'). *P<sub>ssaG</sub>* was amplified with *ssaH*-Xho (5'-ccg ctc gag tgg tag ttt ggg act-3') and *ssaH*-Kpn (5'-cgg ggt acc ata acc gtt agc gct-3'). These amplicons were digested with *XhoI* and *KpnI* and ligated into *Sall*/*KpnI*-digested pGFP[LVA] (Clontech Laboratories). *P<sub>sopB</sub>* was amplified with the oligonucleotides *PsigD*-gfp-XbaF (5'-tgc tct aga tgt tca agc atg gaa-3') and *PsigD*-gfp-Kpn2R (5'-cgt tgt atg gta cct ttt gta gg-3'). *P<sub>pipB</sub>* was amplified with *PpipB*-gfp-XbaF (5'-tgc tct aga cgc att tat tct ggt ata-3') and *PpipB*-gfp-KpnR (5'-cgg ggt acc ggc cgc atg caa ata tct-3'). *P<sub>fliC</sub>* was amplified with *PfliC*-gfp-XbaF (5'-tgc tct aga cag tag tta agc gcg-3') and *PfliC*-gfp-KpnR (5'-cgg ggt acc cag agc gga ctg gga-3'). These three amplicons were digested with *XbaI* and *KpnI* and ligated into the corresponding sites of pGFP[LVA].

The promoter-*gfp*[LVA] fusion cassettes were next inserted into the plasmid pMPM-A3ΔPlac, made by removing the *lac* promoter from

pMPM-A3 (Mayer, 1995). *P<sub>lac</sub>* was removed by inverse PCR with the oligonucleotides delta Plac-for (5'-cgg ggt acc agt cgg gaa acc tgt cgt-3') and delta Plac-rev (5'-gga aca aaa gct ggg tac cgg g-3'). The resulting amplicon was digested with *KpnI* and self-ligated to give pMPM-A3ΔPlac. Promoter-*gfp*[LVA] fusion cassettes were excised from pGFP[LVA] by *HindIII* digestion and ligated into *HindIII*-digested pMPM-A3ΔPlac. All fusions are in the opposite direction to the *α-lacZ* gene. The reporter genes are designated: pMPMA3ΔPlac *P<sub>invF</sub>*-*gfp*[LVA]/R, pMPMA3ΔPlac *P<sub>prgH</sub>*-*gfp*[LVA]/R, pMPMA3ΔPlac *P<sub>sopB</sub>*-*gfp*[LVA]/R, pMPMA3ΔPlac *P<sub>ssaG</sub>*-*gfp*[LVA]/R, pMPMA3ΔPlac *P<sub>pipB</sub>*-*gfp*[LVA]/R and pMPMA3ΔPlac *P<sub>fliC</sub>*-*gfp*[LVA]/R. To construct a negative control reporter fusion, a fragment containing *gfp*[LVA] without any promoter was removed from pGFP[LVA] by *HindIII* digestion and ligated into pMPM-A3ΔPlac to give pMPMA3ΔPlac null-*gfp*[LVA]/R. All fusion sequences were verified by DNA sequencing (National Institute for Allergy and Infectious Diseases genomics facility, Rocky Mountain Laboratories).

**Flow cytometry.** Aliquots (100 μl) of bacterial cultures in a round-bottom 96-well plate (Falcon) were centrifuged at 2250 g for 6 min, washed with PBS and resuspended in freshly made 1% paraformaldehyde (PFA; w/v) in PBS. Bacteria were then washed twice in PBS and immunostained with an anti-*Salmonella* LPS mAb (Meridian Life Science) diluted (1:2000) in PBS for 10 min. Bacteria were again washed twice with PBS, followed by incubation with a Cy5-conjugated donkey anti-mouse antibody (1:1000 in PBS; Jackson ImmunoResearch Laboratories) for 10 min and two more PBS washes. For all measurements, a BD LSR II flow cytometer (BD Bioscience) was used and results were analysed using FlowJo software (Tree Star).

**Electron microscopy.** Samples (1 ml) of bacterial culture, containing approximately  $6 \times 10^9$  or  $1.5 \times 10^9$  c.f.u. of aer-LL or  $\mu$ aer-ST respectively, were centrifuged at 1000 g for 10 min, washed gently with 1 ml PBS and resuspended in 500 μl PBS. Droplets (10 μl) were then adsorbed for 5 min on carbon-coated Formvar grids (Ted Pella). Excess fluids were removed from the grid surface by blotting with filter paper, and the sample was stained for 2 min with 2% ammonium molybdate in double-distilled water. Excess stain was removed by blotting and the samples were dried at room temperature prior to being viewed at 80 kV in a Hitachi 7500 transmission electron microscope (Hitachi High-Technologies). Images were acquired with a digital camera system (Advanced Microscopy Techniques) and processed with Adobe Photoshop v. 7 (Adobe Systems).

#### Immunofluorescence detection of flagella and type 1 fimbriae.

Bacterial cultures were prepared using either wild-type (WT) *S. Typhimurium* or a *flgB::Tn10* mutant, both harbouring the reporter plasmid pMPMA3ΔPlac *P<sub>prgH</sub>*-*gfp*[LVA]/R, or a *fimAICDHF::kan* mutant. Bacteria were collected by centrifugation at 1000 g for 10 min and resuspended carefully in PBS. PFA was then added (1% w/v final concentration) and the sample was incubated for 10 min. Fixed bacteria were washed twice with PBS and stained with mouse anti-FliC mAb (1:100; BioLegend) and rabbit anti-*Salmonella* LPS antibodies (1:300; Difco) or rabbit polyclonal anti-FimA (1:100; provided by Andreas Bäuml) and mouse anti-*Salmonella* LPS mAb (1:2000). Secondary antibodies were AlexaFluor 568-conjugated goat anti-mouse antibodies (1:1000; Invitrogen) and Cy5-conjugated donkey anti-rabbit antibodies (1:1000; Jackson ImmunoResearch Laboratories). All antibodies were diluted in 0.1% (w/v) saponin, 10% (w/v) normal goat serum in PBS (SS-PBS). To label DNA, bacteria were incubated with 1 μg Hoechst 33342 solution ml<sup>-1</sup> (Invitrogen). After gentle washing with PBS, bacteria were resuspended in PBS and mounted on glass slides using ProLong Gold anti-fade reagent (Invitrogen). FliC-positive and FimA-positive *Salmonella*

were enumerated visually using an upright fluorescence microscope (Nikon Eclipse E800).

**Motility assays.** Motility assays on semi-solid agar were performed as described previously (Kim & Surette, 2003) with some modifications. Briefly, 0.5  $\mu$ l of overnight cultures were spotted in the middle of a swimming plate (LB–Miller, 0.25% agar) in duplicate and allowed to dry for 1 h at room temperature. Plates were incubated at 37 °C for 6 h and growth haloes were measured in mm.

Microscopy-based bacterial motility measurements were performed using *Salmonella* harbouring plasmid pFPV25.1 for the constitutive expression of *gfpmut3* under the control of the *rpsM* promoter (Valdivia & Falkow, 1996). *Salmonella* motility in swimming plates was not affected by either the presence of this plasmid or the expression of GFP (Supplementary Fig. S3).  $\mu$ aer-ST and aer-LL cultures were diluted 1:12 and 1:30, respectively, in PBS. Drops (10  $\mu$ l) of bacterial suspension on a glass slide were covered with a coverslip, which was sealed with wax, and the samples were imaged within 5 min. Images were collected using a Nikon Eclipse Ti Perfect Focus inverted microscope fitted with a Nipkow spinning disk, and a  $\times$  60, 1.4 numerical aperture oil immersion objective (Nikon). Image acquisition by a Photometrics CascadeII camera was controlled by Metamorph software (Molecular Devices). A time-lapse series of 26 sequential images (100 ms exposure) was taken over 5 s ( $n=10$  fields). To identify non-motile *Salmonella*, i.e. those that remained in the same location in images 1, 13 and 26, these three images were overlaid in ImageJ (<http://rsb.info.nih.gov/ij/>). To enumerate the total number of bacteria in the sample, *Salmonella* were fixed (1% PFA) and mounted before imaging ( $n=10$  fields). The percentage of motile bacteria was determined by subtracting from 100 the percentage of total non-motile bacteria.

**Invasion and replication assays.** These assays were done as described previously (Drecktrah *et al.*, 2006; Steele-Mortimer *et al.*, 1999; Steele-Mortimer, 2008a). Briefly, HeLa cells (human cervical adenocarcinoma epithelial, ATCC CCL-2) were grown in growth medium (GM), consisting of Eagle's medium (Mediatech) supplemented with 10% (v/v) heat-inactivated fetal bovine serum (Invitrogen), in a humidified 37 °C, 5% CO<sub>2</sub> incubator. HeLa cells were seeded 24 h prior to infection, either in 24-well plates at  $5 \times 10^4$  cells per well or in six-well plates at  $2.4 \times 10^5$  cells per well. *Salmonella* were collected by centrifugation at 5900 g for 2 min, resuspended in Hank's buffered saline solution (HBSS) and used immediately to infect epithelial cells for 10 min at 37 °C. Inoculum counts were checked by plating on LB agar plates. In some experiments, monolayers were centrifuged at 1000 g for 5 min immediately after addition of the bacteria and then incubated for a further 10 min at 37 °C. Extracellular bacteria were removed by aspiration and monolayers were washed twice with HBSS. Monolayers were then incubated at 37 °C for 20 min in GM after which time 50  $\mu$ g gentamicin ml<sup>-1</sup> was added for 30 min to kill extracellular bacteria. Subsequently GM with 10  $\mu$ g gentamicin ml<sup>-1</sup> was added for the remainder of the experiment. Monolayers were lysed in 1 ml 1% Triton X-100 (v/v), 0.1% SDS (w/v) in PBS and viable intracellular bacteria were enumerated by plating on LB agar.

**Immunofluorescence microscopy.** HeLa cells were plated 24 h prior to infection on glass coverslips in 24-well plates ( $6 \times 10^4$  cells per well). To measure lysosomal-associated membrane protein 1 (LAMP1) acquisition by the SCV, infected cells were fixed in 2.5% PFA (w/v) for 10 min at 37 °C, then permeabilized and blocked with SS-PBS for 20 min. All antibodies were diluted in SS-PBS. Primary antibodies were mouse anti-human LAMP1 (H4A3; 1:1500; Developmental Studies Hybridoma Bank) and rabbit anti-*Salmonella* LPS (1:2000; Difco). Secondary antibodies were AlexaFluor 488-conjugated goat anti-mouse IgG antibodies and AlexaFluor 568-

conjugated goat anti-rabbit IgG (1:800; Invitrogen). After extensive washing in PBS, coverslips were mounted on glass slides using Mowiol (Sigma-Aldrich). LAMP1-positive SCVs were scored visually using a Nikon Eclipse E800 upright fluorescent microscope.

For quantification of intracellular GFP-positive bacteria, HeLa cells on glass coverslips were infected with *Salmonella* containing the reporter plasmid constructs. Cells were then fixed in 2.5% PFA and processed for indirect immunofluorescence microscopy with mouse mAb against *Salmonella* LPS (1:5000; Meridian Life Science), followed by incubation with a Cy5-conjugated donkey anti-mouse antibody (1:800; Jackson ImmunoResearch Laboratories) to detect extracellular bacteria. Cells were then permeabilized and blocked with SS-PBS and total bacteria were detected with rabbit anti-*Salmonella* LPS antibodies (1:2000; Difco) followed by AlexaFluor 568-conjugated goat anti-rabbit antibodies (1:800; Invitrogen). Samples were mounted as described above. The percentage of intracellular GFP-positive *Salmonella* was enumerated visually using a Nikon Eclipse E800 upright microscope. All values represent the means  $\pm$  SD from at least three independent experiments.

**QuantiGene detection of intracellular gene expression.** Direct measurement of RNA was performed using the QuantiGene reagent system v. 2.0 (Panomics) according to the manufacturer's directions. Probe sets were custom designed by Panomics and detection efficiency was previously standardized with bacterial cultures from both growth conditions (data not shown). Briefly, HeLa cells in six-well plates ( $2.2 \times 10^5$  cells per well) were infected at an m.o.i. similar to that used for invasion assays. Infected monolayers were lysed in 0.8 ml QuantiGene lysis buffer supplemented with 150 ng Proteinase K ml<sup>-1</sup> (Panomics), and solubilized by incubation at 65 °C for 30 min. One-tenth of each sample (80  $\mu$ l) was combined with blocking buffer and probe sets for each gene in a capture plate, followed by incubation at 55 °C for 16–20 h. The wells were then washed three times with wash buffer (0.1  $\times$  SSC, 0.03% lithium lauryl sulfate). Signal for the bound mRNA was developed by sequential hybridization with branched DNA and alkaline-phosphatase-conjugated label probe (Panomics) at 55 °C for 1 h. After three washes with wash buffer, the diioxetane substrate was added to the wells and incubated at 50 °C for 1 h. Luminescence was measured over 1000 ms using a Safire<sup>2</sup> microplate reader (Tecan). Uninfected HeLa cells were used to determine the background signal and this value was subtracted from each sample. To normalize the expression of each gene, *nusG* was chosen as an internal control because no significant variation of expression for this gene was observed (Eriksson *et al.*, 2003).

## RESULTS

### Induction of invasive *S. Typhimurium*


In order to induce SPI1 gene expression, we grew *Salmonella* with aeration overnight and then diluted it in fresh medium and grew with aeration to late exponential phase. Originally we used LB broth containing 5 g NaCl l<sup>-1</sup> (Steele-Mortimer *et al.*, 1999) but have since found that increasing the NaCl concentration to 10 g l<sup>-1</sup> (LB–Miller) results in better induction of SPI1 [data not shown and Song *et al.* (2004)]. Other groups have shown that SPI1 can also be induced in stationary phase even in the absence of aeration. We hypothesized that SPI1 induction under different physiological conditions could affect the outcome of *Salmonella*–host cell interactions. In order to address this hypothesis, we grew *S. Typhimurium*

SL1344 in LB–Miller broth, either with aeration to late exponential phase (aer-LL) or without aeration to stationary phase ( $\mu$ aer-ST). As previously shown, growth curves (Fig. 1) show that the bacteria grown with aeration reach stationary phase slightly earlier (approximately 4 h) than those grown without aeration and they grow to higher cell densities (Song *et al.*, 2004).

To compare these two cultures, we first did a transcriptome analysis in order to confirm that SPI1 genes are similarly expressed and to detect differences in other gene groups. Using a custom-made microarray, 1009 genes were identified with statistically significant changes in the level of expression. Of these, 517 genes were expressed at a higher level and 492 genes were expressed at a lower level in aer-LL bacteria compared with  $\mu$ aer-ST bacteria (see Supplementary Table S1 available with the online version of this paper). These data have been submitted to the NCBI GEO database (<http://www.ncbi.nlm.nih.gov/geo/>) under

accession number GSE13343. To identify which functional groups of genes were differentially expressed in aer-LL and  $\mu$ aer-ST bacteria, we used GSEA, a computational method that determines whether defined sets of genes show statistically significant, concordant differences between two biological states (Subramanian *et al.*, 2005; Tintle *et al.*, 2008). As predicted, we did not detect significant differences in the SPI1 regulon gene set, whereas flagellar-based motility and chemotaxis gene sets were highly activated in aer-LL bacteria compared with  $\mu$ aer-ST bacteria (GSEA rank 1 and 2, data not shown). Comparison of individual genes in these functional groups (Fig. 2) confirmed that the majority of genes in the T3SS1 regulon were similarly expressed in aer-LL and  $\mu$ aer-ST bacteria. In contrast, genes involved in flagellar biosynthesis (including *flh*, *flg*, *fli* and *flj* operons) and chemotaxis (*che*, *mot* and *mcp*) had higher expression levels in aer-LL bacteria, and genes involved in type 1 fimbriae production and the stress response were more highly expressed in  $\mu$ aer-ST bacteria.


We next compared the invasiveness of aer-LL and  $\mu$ aer-ST bacteria using a gentamicin protection assay (Fig. 3). HeLa cells were infected with equivalent c.f.u. from each culture and viable intracellular bacteria were estimated at 1 h post-infection (p.i.). Cells infected with aer-LL bacteria contained four- to fivefold more *Salmonella* (approximately  $2.8 \times 10^4$  per well) than those infected with  $\mu$ aer-ST bacteria (approximately  $0.6 \times 10^4$  per well), which is within the same range as that reported previously under similar conditions (Song *et al.*, 2004). Expression of these data as percentages of the initial inoculum, to account for any differences in inoculum size, showed invasion efficiency of  $0.52\% \pm 0.09$  for aer-LL bacteria and  $0.11\% \pm 0.04$  for  $\mu$ aer-ST bacteria. The apparently low invasion efficiency is a reflection of the invasion conditions, since HeLa cells were incubated with a relatively high m.o.i. (approximately 50–60) for a short infection time (10 min). These conditions were selected in order to minimize the effect of environmental changes on gene expression while still internalizing sufficient numbers of bacteria. Given that there are approximately  $8 \times 10^4$  HeLa cells per well (data not shown), less than half of the cells are infected, even with the more invasive aer-LL bacteria, demonstrating that these infection conditions are well below saturation levels. Analysis by fluorescence microscopy showed that the distribution of bacteria within cells was not affected by growth condition and that the majority (>70%) of infected cells contained fewer than six bacteria (see Supplementary Fig. S1, available with the online version of this paper). Altogether these data show that both growth conditions produce invasive bacteria but aer-LL bacteria are significantly more invasive than  $\mu$ aer-ST bacteria.



**Fig. 1.** *S. Typhimurium* growth curves. Bacteria were grown in LB–Miller broth at 37 °C, with (●) or without (○) aeration. Bacterial growth was determined by OD<sub>600</sub> (a) or culturable bacterial numbers (b). Arrowheads and dashed vertical lines indicate the time points at which aer-LL and  $\mu$ aer-ST bacteria were harvested from aerated and non-aerated cultures, respectively. Larger symbols indicate  $n=3$ , smaller symbols  $n=2$ .

### Single-cell analysis reveals differences in SPI1 gene expression


Previous studies have shown that, under SPI1-inducing conditions, only a fraction of the population are induced



**Fig. 2.** Relative expression levels of selected genes in *S. Typhimurium*. A value of 1 indicates no detectable difference in expression between aer-LL and  $\mu$ aer-ST bacteria, values  $>1$  indicate higher expression in aer-LL bacteria and values  $<1$  indicate higher expression in  $\mu$ aer-ST bacteria. Dotted lines indicate the twofold threshold. (a) SPI1 regulon genes. (b) Motility-associated genes. (c) Stress-related genes. Black bars denote data that passed the  $P$ -value threshold. Asterisks indicate data confirmed by qPCR.

and that single-cell experiments are required to characterize the population heterogeneity (Bumann, 2002; Hautefort *et al.*, 2003; Temme *et al.*, 2008). Here, we used transcriptional fusions to a destabilized variant of GFP, GFP[LVA], to identify induced bacteria (Drecktrah *et al.*, 2006). Promoters of three SPI1-regulon genes (*invF*, *prgH* and *sopB*) were compared with two SPI2-associated genes (*ssaG* and *pipB*) and a flagellin gene (*fliC*). Plasmid-borne fusions were introduced into WT *S. Typhimurium* and the

numbers of GFP-positive bacteria were assessed by flow cytometry following growth under aerated and non-aerated conditions (Fig. 4a). Control experiments showed that under these conditions, loss of the plasmid was negligible (see Supplementary Fig. S2, available with the online version of this paper). Initial experiments, in which  $P_{prgH}$  activity was measured, confirmed that SPI1 induction peaks at approximately 3–4 h p.i. in aerated culture with approximately 80% GFP-positive bacteria. In contrast,



**Fig. 3.** Invasion of epithelial cells by SPI1-induced bacteria. To directly compare the invasiveness of aer-LL and  $\mu$ aer-ST *S. Typhimurium*, HeLa cells were infected for 10 min at 37 °C (m.o.i. 50–60). At 1 h p.i., cells were lysed and intracellular bacteria were enumerated by plating. The means  $\pm$  SD from three independent experiments are shown. (a) Invasion expressed as c.f.u. per well. (b) Invasion expressed as % of inoculum. In both cases  $P < 0.001$ , Student's *t*-test.

in non-aerated culture, GFP-positive bacteria never accounted for more than 40% of the total population.  $P_{fliC}$  activity was very similar to  $P_{prgH}$  in aerated culture but in non-aerated culture revealed very different activation levels at early time points.  $P_{fliC}$  activity was initially high (71  $\pm$  7% GFP-positive bacteria), whereas very little  $P_{prgH}$  activity could be detected (4  $\pm$  1% GFP-positive bacteria). Subsequently,  $P_{prgH}$  activity increased and  $P_{fliC}$  activity decreased too, so that by 8 h, there were similar numbers of GFP-positive bacteria, 26  $\pm$  7 and 28  $\pm$  1% respectively. Thereafter, there were no significant changes up to 24 h. When plasmids containing an SPI2 gene transcriptional fusion (*P<sub>ssaG</sub>-gfp*[LVA]) or GFP without a promoter (null-*gfp*[LVA]) were used, no significant numbers of GFP-positive bacteria were detected at any time point (Fig. 4 and data not shown). When aer-LL and  $\mu$ aer-ST *S. Typhimurium*, harvested from aerated and non-aerated cultures at 3.5 and 18 h respectively, were directly compared for promoter activities for the three SPI1-regulon genes *invF*, *prgH* and *sopB*, the aer-LL culture contained 82, 81 and 49% GFP-positive bacteria respectively compared with 40, 39 and 34% GFP-positive bacteria in the  $\mu$ aer-ST culture (Fig. 4b). Similarly, the *fliC* promoter was induced in 76  $\pm$  12 and 36  $\pm$  4% of bacteria in aer-LL and  $\mu$ aer-ST cultures, respectively. Virtually no activity could be detected in either culture for the SPI2 promoters  $P_{pipB}$  and  $P_{ssaG}$  or for the null construct. Altogether, single-cell analysis revealed a higher frequency of SPI1-induced bacteria in aer-LL compared with  $\mu$ aer-ST bacteria.

To account for the differences between the transcriptome and single-cell data, levels of induction must be higher in SPI1-induced  $\mu$ aer-ST bacteria compared with SPI1-induced aer-LL bacteria. To investigate this possibility, we compared the GFP intensity profiles in the flow


cytometry data from the above experiments. This revealed that a small, but significant, population of  $\mu$ aer-ST *Salmonella* (2–4%) had a higher GFP intensity level (RFU  $> 10^4$ ) than detected in any aer-LL bacteria (Fig. 4c). Thus, although fewer *Salmonella* are SPI1-induced in the  $\mu$ aer-ST compared with the aer-LL culture, a small percentage of 'hyper-induced' SPI1 bacteria in  $\mu$ aer-ST cultures results in similar average expression levels of SPI1 genes, as indicated by the transcriptome analysis.

### Flagella are required for efficient *Salmonella* invasion

*Salmonella* achieve motility from 5–10 peritrichous, or randomly placed, filamentous flagella that extend from the cell surface. *S. Typhimurium* has two loci encoding different flagellins, *fliC* and *fliB*, the expression of which is governed by a switch mechanism so that only one flagellin is expressed at a time [reviewed by McQuiston *et al.* (2008)]. Our transcriptome data suggested that *fliB* was induced in aer-LL bacteria whereas no difference was detected in *fliC* expression (Fig. 2b). However, a *fliC* mutant was more affected in motility and invasion of HeLa cells compared with a *fliB* mutant (data not shown and Supplementary Fig. S3, available with the online version of this paper). Since comparative analysis does not exclude the possibility that *fliC* is induced under both conditions, similar to the SPI1-regulon genes, we used the GFP-based transcriptional fusion assay to see if the SPI1-induced bacteria are also flagellated. *Salmonella* bearing the *P<sub>prgH</sub>-gfp*[LVA] construct were scored for GFP fluorescence and/or flagella using an anti-FliC antibody. Using this method, 48  $\pm$  5% of aer-LL and 23  $\pm$  3% of  $\mu$ aer-ST bacteria were GFP-positive. Of these SPI1-induced bacteria, 75  $\pm$  8% of the aer-LL and 94  $\pm$  5% of the  $\mu$ aer-ST culture were also FliC-positive, whereas only one-third of the GFP-negative bacteria were FliC-positive. The physical presence of flagella on the surface of both  $\mu$ aer-ST and aer-LL bacteria was confirmed by electron microscopy (Supplementary Fig. S4, available with the online version of this paper), which also revealed that fimbriae were only present on the surface of  $\mu$ aer-ST bacteria.

We hypothesized that the physical presence of flagella may not accurately reflect the functional competence of these structures i.e. motility. To compare the motility of aer-LL and  $\mu$ aer-ST *Salmonella*, we developed a rapid microscopy-based assay which allowed us to assess motility within minutes of harvesting the bacteria, using *S. Typhimurium* constitutively expressing GFP (Valdivia & Falkow, 1996). This assay revealed that motile bacteria predominate in the aer-LL (76  $\pm$  12%) compared with the  $\mu$ aer-ST (33  $\pm$  16%) culture (Fig. 5a) in agreement with the higher expression levels of motility and chemotaxis related genes.

The above experiments revealed a correlation between SPI1 expression and motility, with SPI1-induced motile bacteria predominating in the aer-LL culture. We next considered the contribution of flagellar-based motility in invasion of



**Fig. 4.** Single-cell analysis of promoter activity for representative SPI1 and flagellin genes. *S. Typhimurium* harbouring *gfp*[LVA] transcriptional fusions to selected promoters were analysed by flow cytometry. (a) Activity of promoters for cultures grown with (i) or without (ii) aeration. Arrowheads indicate the time points at which aer-LL and  $\mu$ aer-ST bacteria were harvested. (b) Promoter activity in aer-LL (filled bars) and  $\mu$ aer-ST (open bars) bacteria. Comparison of SPI1 regulon genes (*prgH*, *invF* and *sopB*), SPI2 regulon genes (*pipB* and *ssaG*), and the flagellin gene *fliC*. Promoterless *gfp*[LVA] (null) was included as a negative control. (c) Comparison of the frequencies of high-intensity GFP ( $>10^4$  relative fluorescent units) expressing bacteria in aer-LL (filled bars) and  $\mu$ aer-ST (open bars) bacteria. Shown are means  $\pm$  SD from at least three independent experiments. \*\*\* $P < 0.001$ , Student's *t*-test.


either  $\mu$ aer-ST or aer-LL *S. Typhimurium* into HeLa cells. To do this, we compared the invasion efficiency of a non-motile *S. Typhimurium* flagellar mutant, *fliC::Tn10 fljB::Mud-Cm*, which is deficient for both flagellin proteins, to WT *S. Typhimurium*, as well as a non-invasive  $\Delta$ SPI1 mutant (Drecktrah *et al.*, 2006). The absence of flagella and motility in the flagellar mutant was confirmed by electron microscopy and soft agar motility assay (Supplementary Fig. S3 and data not shown). To directly compare the requirement for motility in both cultures, the inoculum size of  $\mu$ aer-ST bacteria was increased (fivefold higher m.o.i. compared with aer-LL) so that similar numbers of WT bacteria were internalized (Fig. 5b). Compared with WT, both the SPI1-deletion mutant, which lacks T3SS1 and is invasion defective, and the flagellar mutant had massive defects in the numbers of internalized bacteria (approximately 3% and  $<1\%$  of WT respectively) (Khoramian-Falsafi *et al.*, 1990). We then confirmed that the invasion deficiency of the flagellar mutant could be overcome by centrifugation to increase contact with host

cells (Fig. 5b), since this can compensate for the lack of motility but not the lack of SPI1 (van Asten *et al.*, 2004). Altogether, these data show that flagella-based motility is an important component in the initiation of contact with epithelial cells for both  $\mu$ aer-ST and aer-LL *Salmonella*.

### Pre-invasion bacterial growth conditions can affect intracellular replication and intracellular virulence gene expression


The ability of *Salmonella* to survive and replicate within host cells is a multifactorial process that can be affected by a variety of bacterial genes [for review see the paper by Ibarra & Steele-Mortimer (2009)]. Since aer-LL and  $\mu$ aer-ST *Salmonella* are grown under different conditions and are induced for different gene sets, we next compared the ability of these bacteria to replicate inside HeLa cells. For these experiments, we used the gentamicin protection assay, adjusting the inoculum size to ensure that there was  $<1\%$  difference in the numbers of bacteria internalized





**Fig. 5.** Flagella-based motility is required for optimal invasion of epithelial cells. (a) Bacterial motility was assessed by fluorescence microscopy using *S. Typhimurium* constitutively expressing GFP. Shown are means  $\pm$  SD from three independent experiments.  $P < 0.05$ , Student's *t*-test. (b) Comparison of the role of flagellar-based motility in invasion. HeLa cells were infected with aer-LL or  $\mu$ aer-ST *S. Typhimurium* either with (shaded bars) or without (filled bars) centrifugation. Intracellular bacteria (1 h p.i.) were enumerated by plating. Strains were WT *S. Typhimurium* (WT), *fliC::Tn10 fliB::Mud-Cm* mutant (*fliC/fliB*) and a  $\Delta$ SPI1::*kan* mutant (SPI1). Means  $\pm$  SD from at least three independent experiments are shown. \* $P < 0.05$ , Student's *t*-test (compared with WT without centrifugation).

and no significant difference in the intracellular distribution of bacteria at 1 h p.i. (Supplementary Fig. S1). This is important since even small differences in the numbers of bacteria internalized, as opposed to the invasion efficiency, could be amplified into significant differences once replication is initiated. As shown in Fig. 6(a), intracellular *S. Typhimurium* initiated intracellular replication by 6 h p.i. irrespective of the pre-invasion growth conditions, but by 8 h, aer-LL bacteria showed a marked increase in replication compared with  $\mu$ aer-ST bacteria (51-fold increase versus 10-fold increase at 1 h).



**Fig. 6.** Intracellular replication is affected by pre-invasion growth conditions. Infected HeLa cells were lysed for enumeration of intracellular *S. Typhimurium* (a) or fixed and processed for immunofluorescence using anti-LPS and anti-LAMP1 antibodies (b). To obtain comparable invasion, cells were infected with an m.o.i. of approximately 50–60 for aer-LL and approximately 150–180 for  $\mu$ aer-ST. The numbers of intracellular aer-LL (filled symbols/bars) or  $\mu$ aer-ST (open symbols/bars) *S. Typhimurium* with coincident LAMP1 staining cell were scored visually by fluorescence microscopy. Means  $\pm$  SD from three independent experiments are shown. \* $P < 0.001$ , Bonferroni's post-hoc test.

The marked difference in intracellular replication of aer-LL and  $\mu$ aer-ST *Salmonella* suggested that pre-invasion growth conditions can affect the ability of *Salmonella* to adapt to the intracellular environment. To address this, we first asked whether there was any apparent defect in SCV biogenesis, by following the recruitment of the lysosomal membrane protein LAMP1, which occurs with well-established kinetics (Steele-Mortimer *et al.*, 1999). However, no difference in LAMP1 acquisition could be detected (Fig. 6b) suggesting that the differences in intracellular replication cannot be attributed to overt changes in vacuole biogenesis.

We next considered that intracellular virulence gene expression might be affected by the pre-invasion growth conditions. Previous experiments have shown that SPI1 genes are rapidly downregulated following internalization into host cells (Boddicker & Jones, 2004; Drecktrah *et al.*, 2006; Knodler *et al.*, 2009), so we used the GFP[LVA] transcriptional fusions to follow this process in individual bacteria following invasion of HeLa cells. As expected, the numbers of SPI1-induced GFP-positive bacteria (*prgH* and *sopB*) rapidly decreased after internalization and by 4 h p.i., virtually no GFP-positive bacteria could be detected. However, when aer-LL and  $\mu$ aer-ST bacteria harbouring the *PprgH-gfp*[LVA] fusion were compared (Fig. 7a), we observed an apparent delay in downregulation in aer-LL bacteria such that at 0.5 h p.i., more aer-LL bacteria were GFP-positive (80  $\pm$  5 % compared to 59  $\pm$  11 %). A similar, though more dramatic, trend was observed with the *PfliC-gfp*[LVA] fusion; 82  $\pm$  7 % of aer-LL bacteria compared with 38  $\pm$  7 % of  $\mu$ aer-ST bacteria were GFP-positive at 0.5 h. In contrast, we did not detect any difference in the downregulation of *sopB*, presumably because this gene is expressed at lower levels. When intracellular expression of the SPI2 regulon genes *ssaG* and *pipB* was compared, we observed induction in both cases, although there were far fewer GFP-positive bacteria when the *PpipB-gfp*[LVA] fusion was used, probably due to low expression levels of *pipB*. Comparison of numbers of GFP-positive bacteria

again revealed a slight dependence on the pre-invasion growth method, suggesting that post-invasion induction of SPI2 genes is more efficient in aer-LL bacteria. However, this was less dramatic than that seen for the *prgH* and *fliC* promoters.

Limitations of the plasmid-based GFP transcriptional fusion assay used here are that the presence of plasmid and/or the expression of GFP could affect invasion or the ability of the bacteria to respond to the intracellular environment (J. A. Ibarra, L. A. Knodler and O. Steele-Mortimer, unpublished observations; Clark *et al.*, 2009; Knodler *et al.*, 2005). Therefore, we independently confirmed these results by assessing the relative levels of RNA obtained from intracellular WT *S. Typhimurium* (Canales *et al.*, 2006; Flagella *et al.*, 2006). This method uses oligonucleotide probe sets that bind a contiguous region of the target RNA. Unfortunately, technical limitations of probe design made it impossible to make *ssaG* probes. Instead, we analysed another SPI2 gene, *ssaJ*, which is in the same operon as *ssaG* (Lober *et al.*, 2006). Probe sets were first tested on aer-LL and  $\mu$ aer-ST bacterial cultures, and gave similar results to quantitative PCR for the SPI1-regulon genes, *prgH* and *sopB*, and the flagellin gene *fliC* (data not shown). As expected, no SPI2-regulon gene expression was detected in these cultures. In intracellular bacteria, this method confirmed the results obtained with the transcriptional fusion assay for the *prgH*, *sopB* and *fliC* promoters but was unable to recapitulate any significant differences in the SPI2 gene induction (Fig. 7b). Altogether, these gene expression analyses show that the increased multiplication efficiency of aer-LL *S. Typhimurium* inside epithelial cells is associated with higher expression of SPI1 genes at early times post invasion.

## DISCUSSION


*S. enterica* serovars invade non-phagocytic host cells by inducing membrane ruffling on the host cell surface, a well-characterized process that is mediated by effector proteins translocated via T3SS1 (Patel & Galan, 2005). *In vitro*, the SPI1 regulon, which encodes T3SS1, can be induced under several different conditions including growth with aeration to late-exponential (aer-LL) and growth without aeration to stationary phase ( $\mu$ aer-ST). While the expression of SPI1 genes under similar conditions has been examined (Croinin & Dorman, 2007; Mangan *et al.*, 2006; Song *et al.*, 2004; Temme *et al.*, 2008), the effect on pathogen–host interactions of SPI1 induction under different physiological backgrounds has not been addressed. In this study, we examined aer-LL and  $\mu$ aer-ST *S. Typhimurium* cultures, and their ability to interact with cultured epithelial cells, in a comparative analysis using both population-based and single-cell methods. Our main findings are: (i) that pre-invasion conditions affect invasion, intracellular replication and post-invasion expression of SPI1 genes; (ii) that single-cell analysis of SPI1 induction reveals differences in aer-LL and  $\mu$ aer-ST cultures not revealed by transcriptome

analysis; and (iii) that co-expression of SPI1 and flagella is associated with higher invasion levels.

Here, we found that pre-invasion growth conditions can significantly affect the invasiveness of *S. Typhimurium* as well as the ability of the bacteria to replicate intracellularly. In contrast, a recent report showed that, unlike for *S. enterica* serovar Typhi, growth conditions have no effect on the invasiveness of *S. Typhimurium* (Bishop *et al.*, 2008). The differences in results are doubtless due in part to the methods or strains used, but also, Bishop *et al.* (2008) did not include late-exponential-phase bacteria in their studies. Previously, we have shown a very strong correlation between late-exponential growth and invasiveness using a different *S. Typhimurium* strain (14028s) (Steele-Mortimer *et al.*, 1999), and here we have used SL1344; however, it is possible that other strains behave differently.

Our single-cell analysis of gene expression revealed significant differences between aer-LL and  $\mu$ aer-ST *S. Typhimurium* in both the frequency of SPI1-induced bacteria and the maximum induction levels, in agreement with previous findings (Passerat *et al.*, 2009; Song *et al.*, 2004), as well as in the expression of flagellar motility. Regulation of the SPI1 regulon is complex and subjected to the actions of a number of factors, which are still not well understood but themselves could be differentially regulated under the conditions described here. Transcriptional regulators necessary for optimal induction in aerobic conditions include HilA, HilC, EnvZ/OmpR and SirA/BarA (Song *et al.*, 2004). Some may be important under both conditions, for example the positive regulator Fis, which under aerobic growth is expressed only in early exponential phase, but is expressed throughout stationary phase in non-aerobic growth conditions (Croinin & Dorman, 2007). This sustained expression of Fis in non-aerobic conditions would explain why SPI1 is induced in  $\mu$ aer-ST bacteria. Another global regulator involved in SPI1 regulation is the integration host factor (IHF), the expression of which peaks in late-exponential and stationary phases and plays a positive role in the expression of SPI1 and flagella in the early late-exponential phase of growth in aerobic conditions (Mangan *et al.*, 2006). This is consistent with our finding that in aer-LL bacteria, SPI1 and flagella genes are highly expressed and a model that proposes co-regulation of invasion and motility genes (Baxter & Jones, 2005; Kage *et al.*, 2008; Saini *et al.*, 2008). Nevertheless, other groups have found repression of motility under conditions of SPI1-induction and vice versa, and the regulatory connection between motility and virulence remains unclear (Teplitski *et al.*, 2003; Thijs *et al.*, 2007). Regardless, our data clearly define both SPI1 expression and flagellar motility as being required for optimal invasion by *Salmonella*.

We, and others, have previously shown that the mode of bacterial entry and SPI1-induction can affect the intracellular responses of *Salmonella* in phagocytic cells (Drecktrah



**Fig. 7.** Intracellular virulence gene expression can be influenced by pre-invasion growth conditions. (a) *gfp*[LVA] transcriptional fusions were used to assess promoter activity for *prgH*, *sopB*, *ssaG*, *pipB* and *fliC* following invasion of HeLa cells. GFP-positive ‘induced’ bacteria were scored visually by fluorescence microscopy. Extracellular *S. Typhimurium* were stained with anti-LPS mAbs before permeabilization and were excluded from analysis. Means  $\pm$  SD from at least three separate experiments are shown. \* $P < 0.05$ , two-way ANOVA and Bonferroni’s post-hoc analysis. (b) QuantiGene detection of *prgH*, *sopB*, *ssaJ*, *pipB* and *fliC* gene expression. The expression of each gene was normalized to *nusG* as an internal control and then compared to the value for aer-LL bacteria at 30 min. Each symbol represents the mean from one experiment in duplicate. The statistical means from three (*prgH*, *sopB* and *fliC*) or four (*ssaJ* and *pipB*) experiments are indicated. Closed symbols, aer-LL; open symbols,  $\mu$ aer-ST.

*et al.*, 2006; van der Velden *et al.*, 2000). Here, however, we report the novel observation that SPII-induction via different pre-invasion growth conditions can affect gene expression in non-phagocytic cells. Unlike phagocytic cells, non-phagocytic epithelial cells only internalize invasive, T3SS1-induced *Salmonella*. Accordingly, invasion should select specifically for SPII-induced bacteria irrespective of the pre-invasion growth conditions and thus, since only SPII-induced bacteria are internalized, these bacteria should react similarly to the intracellular environment. One caveat is that non-SPII-induced 'bystander' bacteria can be internalized into the ruffles stimulated by invasive SPII-induced bacteria (Steele-Mortimer *et al.*, 2002). In order to minimize this possibility, we did not centrifuge the bacteria onto the monolayer and after a brief 10 min invasion, we immediately removed non-internalized bacteria by washing. Under these conditions, we found that SPII genes are downregulated following invasion in agreement with qPCR showing rapidly decreasing *sopB* transcript levels in HeLa cells (Knodler *et al.*, 2009) and studies showing that most SopB is translocated early (Patel *et al.*, 2009). A recent study found that rather than being downregulated as reported here, SPII is instead induced following invasion of epithelial cells (Hautefort *et al.*, 2008). While it is difficult to reconcile these contradictions, we believe that they are most probably due to differences in methodology, including a centrifugation step to optimize internalization of bacteria, especially since this later step could increase the numbers of 'bystander' non-SPII-induced cells that are internalized. When we compared intracellular expression of SPII genes following internalization of either aer-LL or  $\mu$ aer-ST bacteria, the aer-LL bacteria exhibited sustained intracellular SPII expression that was accompanied by higher intracellular replication. While we have not shown a direct link between intracellular replication and sustained SPII expression in this study, effectors translocated by the SPII-encoded T3SS1 can have significant post-invasion roles, for example, SopB mediates sustained Akt activation in infected epithelial cells (Steele-Mortimer *et al.*, 2000) and also, though to a less obvious extent, SCV biogenesis (Hernandez *et al.*, 2004). Also, despite rapid downregulation of gene expression, the persistence of some T3SS1 effectors, such as SopB, inside host cells has been demonstrated (Drecktrah *et al.*, 2005; Hernandez *et al.*, 2004; Knodler *et al.*, 2009). Overall, our data suggest that aer-LL bacteria are better able to adapt to the intracellular environment than  $\mu$ aer-ST bacteria. This could be due to something as simple as the difference in adapting from an oxygen-rich environment (aer-LL) compared with an oxygen-poor environment ( $\mu$ aer-ST) or with multiple combined factors.

Here, we have defined pre-invasion bacterial growth status as a factor that can influence the outcome of *Salmonella*-host cell interactions *in vitro*. We conclude that, in the context of host-pathogen interactions, the physiological state of the bacteria should be considered, particularly

when different growth conditions can be used for induction of an invasive phenotype.

## ACKNOWLEDGEMENTS

We thank Seth Winfree, Patricia Fuentes, Ron Messer, Alissa Curda, Craig Martens, Greg Farneth, Damon Ellison and Luke Wicke for expert experimental and technical assistance. We are grateful to Ed Miao for kindly providing phages, to Andreas Bäumlner for providing phage stock and the anti-FimA antibody, to the Rocky Mountain Laboratories Genomics Unit for DNA sequencing analysis, to members of our lab for their helpful discussions and criticism and to the reviewers for constructive suggestions. This research was supported by the Intramural Research Program (DIR) of the National Institutes of Health (NIH), National Institute for Allergy and Infectious Diseases (NIAID).

## REFERENCES

- Andersen, J. B., Sternberg, C., Poulsen, L. K., Bjorn, S. P., Givskov, M. & Molin, S. (1998). New unstable variants of green fluorescent protein for studies of transient gene expression in bacteria. *Appl Environ Microbiol* **64**, 2240–2246.
- Bajaj, V., Lucas, R. L., Hwang, C. & Lee, C. A. (1996). Co-ordinate regulation of *Salmonella typhimurium* invasion genes by environmental and regulatory factors is mediated by control of *hila* expression. *Mol Microbiol* **22**, 703–714.
- Baxter, M. A. & Jones, B. D. (2005). The *fimYZ* genes regulate *Salmonella enterica* Serovar Typhimurium invasion in addition to type 1 fimbrial expression and bacterial motility. *Infect Immun* **73**, 1377–1385.
- Behlau, I. & Miller, S. I. (1993). A PhoP-repressed gene promotes *Salmonella typhimurium* invasion of epithelial cells. *J Bacteriol* **175**, 4475–4484.
- Bishop, A., House, D., Perkins, T., Baker, S., Kingsley, R. A. & Dougan, G. (2008). Interaction of *Salmonella enterica* serovar Typhi with cultured epithelial cells: roles of surface structures in adhesion and invasion. *Microbiology* **154**, 1914–1926.
- Boddicker, J. D. & Jones, B. D. (2004). Lon protease activity causes down-regulation of *Salmonella* pathogenicity island 1 invasion gene expression after infection of epithelial cells. *Infect Immun* **72**, 2002–2013.
- Bumann, D. (2002). Examination of *Salmonella* gene expression in an infected mammalian host using the green fluorescent protein and two-colour flow cytometry. *Mol Microbiol* **43**, 1269–1283.
- Canales, R. D., Luo, Y., Willey, J. C., Austermler, B., Barbacioru, C. C., Boysen, C., Hunkapiller, K., Jensen, R. V., Knight, C. R. & other authors (2006). Evaluation of DNA microarray results with quantitative gene expression platforms. *Nat Biotechnol* **24**, 1115–1122.
- Clark, L., Martinez-Argudo, I., Humphrey, T. J. & Jepson, M. A. (2009). GFP plasmid-induced defects in *Salmonella* invasion depend on plasmid architecture, not protein expression. *Microbiology* **155**, 461–467.
- Croinin, T. O. & Dorman, C. J. (2007). Expression of the Fis protein is sustained in late-exponential- and stationary-phase cultures of *Salmonella enterica* serovar Typhimurium grown in the absence of aeration. *Mol Microbiol* **66**, 237–251.
- Drecktrah, D., Knodler, L. A., Galbraith, K. & Steele-Mortimer, O. (2005). The *Salmonella* SPII effector SopB stimulates nitric oxide production long after invasion. *Cell Microbiol* **7**, 105–113.

- Drecktrah, D., Knodler, L. A., Ireland, R. & Steele-Mortimer, O. (2006). The mechanism of *Salmonella* entry determines the vacuolar environment and intracellular gene expression. *Traffic* **7**, 39–51.
- Ellermeier, J. R. & Schlauch, J. M. (2007). Adaptation to the host environment: regulation of the SPI1 type III secretion system in *Salmonella enterica* serovar Typhimurium. *Curr Opin Microbiol* **10**, 24–29.
- Eriksson, S., Lucchini, S., Thompson, A., Rhen, M. & Hinton, J. C. (2003). Unravelling the biology of macrophage infection by gene expression profiling of intracellular *Salmonella enterica*. *Mol Microbiol* **47**, 103–118.
- Flagella, M., Bui, S., Zheng, Z., Nguyen, C. T., Zhang, A., Pastor, L., Ma, Y., Yang, W., Crawford, K. L. & other authors (2006). A multiplex branched DNA assay for parallel quantitative gene expression profiling. *Anal Biochem* **352**, 50–60.
- Hansen-Wester, I. & Hensel, M. (2001). *Salmonella* pathogenicity islands encoding type III secretion systems. *Microbes Infect* **3**, 549–559.
- Hautefort, I., Proenca, M. J. & Hinton, J. C. (2003). Single-copy green fluorescent protein gene fusions allow accurate measurement of *Salmonella* gene expression *in vitro* and during infection of mammalian cells. *Appl Environ Microbiol* **69**, 7480–7491.
- Hautefort, I., Thompson, A., Eriksson-Ygberg, S., Parker, M. L., Lucchini, S., Danino, V., Bongaerts, R. J., Ahmad, N., Rhen, M. & Hinton, J. C. (2008). During infection of epithelial cells *Salmonella enterica* serovar Typhimurium undergoes a time-dependent transcriptional adaptation that results in simultaneous expression of three type 3 secretion systems. *Cell Microbiol* **10**, 958–984.
- Hernandez, L. D., Hueffer, K., Wenk, M. R. & Galan, J. E. (2004). *Salmonella* modulates vesicular traffic by altering phosphoinositide metabolism. *Science* **304**, 1805–1807.
- Hoise, S. K. & Stocker, B. A. (1981). Aromatic dependent *Salmonella typhimurium* are non-virulent and effective as live vaccines. *Nature* **291**, 238–239.
- Ibarra, J. A. & Steele-Mortimer, O. (2009). *Salmonella* – the ultimate insider. *Salmonella* virulence factors that modulate intracellular survival. *Cell Microbiol* **11**, 1579–1586.
- Jones, B. D. & Falkow, S. (1994). Identification and characterization of a *Salmonella typhimurium* oxygen-regulated gene required for bacterial internalization. *Infect Immun* **62**, 3745–3752.
- Kage, H., Takaya, A., Ohya, M. & Yamamoto, T. (2008). Coordinated regulation of expression of *Salmonella* pathogenicity island 1 and flagellar type III secretion systems by ATP-dependent ClpXP protease. *J Bacteriol* **190**, 2470–2478.
- Kanehisa, M., Araki, M., Goto, S., Hattori, M., Hirakawa, M., Itoh, M., Katayama, T., Kawashima, S., Okuda, S. & other authors (2008). KEGG for linking genomes to life and the environment. *Nucleic Acids Res* **36**, D480–D484.
- Khoramian-Falsafi, T., Harayama, S., Kutsukake, K. & Pechere, J. C. (1990). Effect of motility and chemotaxis on the invasion of *Salmonella typhimurium* into HeLa cells. *Microb Pathog* **9**, 47–53.
- Kim, W. & Surette, M. G. (2003). Swarming populations of *Salmonella* represent a unique physiological state coupled to multiple mechanisms of antibiotic resistance. *Biol Proced Online* **5**, 189–196.
- Knodler, L. A., Bestor, A., Ma, C., Hansen-Wester, I., Hensel, M., Vallance, B. A. & Steele-Mortimer, O. (2005). Cloning vectors and fluorescent proteins can significantly inhibit *Salmonella enterica* virulence in both epithelial cells and macrophages: implications for bacterial pathogenesis studies. *Infect Immun* **73**, 7027–7031.
- Knodler, L. A., Winfree, S., Drecktrah, D., Ireland, R. & Steele-Mortimer, O. (2009). Ubiquitination of the bacterial inositol phosphatase, SopB, regulates its biological activity at the plasma membrane. *Cell Microbiol* **11**, 1652–1670.
- Lee, C. A. & Falkow, S. (1990). The ability of *Salmonella* to enter mammalian cells is affected by bacterial growth state. *Proc Natl Acad Sci U S A* **87**, 4304–4308.
- Lober, S., Jackel, D., Kaiser, N. & Hensel, M. (2006). Regulation of *Salmonella* pathogenicity island 2 genes by independent environmental signals. *Int J Med Microbiol* **296**, 435–447.
- Mangan, M. W., Lucchini, S., Danino, V., Croinin, T. O., Hinton, J. C. & Dorman, C. J. (2006). The integration host factor (IHF) integrates stationary-phase and virulence gene expression in *Salmonella enterica* serovar Typhimurium. *Mol Microbiol* **59**, 1831–1847.
- Marcus, S. L., Brumell, J. H., Pfeifer, C. G. & Finlay, B. B. (2000). *Salmonella* pathogenicity islands: big virulence in small packages. *Microbes Infect* **2**, 145–156.
- Mayer, M. P. (1995). A new set of useful cloning and expression vectors derived from pBlueScript. *Gene* **163**, 41–46.
- McGhie, E. J., Brawn, L. C., Hume, P. J., Humphreys, D. & Koronakis, V. (2009). *Salmonella* takes control: effector-driven manipulation of the host. *Curr Opin Microbiol* **12**, 117–124.
- McQuiston, J. R., Fields, P. I., Tauxe, R. V. & Logsdon, J. M., Jr (2008). Do *Salmonella* carry spare tyres? *Trends Microbiol* **16**, 142–148.
- Miao, E. A., Alpujch-Aranda, C. M., Dors, M., Clark, A. E., Bader, M. W., Miller, S. I. & Aderem, A. (2006). Cytoplasmic flagellin activates caspase-1 and secretion of interleukin 1 $\beta$  via Ipaf. *Nat Immunol* **7**, 569–575.
- Mootha, V. K., Lindgren, C. M., Eriksson, K. F., Subramanian, A., Sihag, S., Lehar, J., Puigserver, P., Carlsson, E., Ridderstrale, M. & other authors (2003). PGC-1 $\alpha$ -responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* **34**, 267–273.
- Passerat, J., Got, P., Dukan, S. & Monfort, P. (2009). Respective roles of culturable and viable-but-nonculturable cells in the heterogeneity of *Salmonella enterica* serovar typhimurium invasiveness. *Appl Environ Microbiol* **75**, 5179–5185.
- Patel, J. C. & Galan, J. E. (2005). Manipulation of the host actin cytoskeleton by *Salmonella* – all in the name of entry. *Curr Opin Microbiol* **8**, 10–15.
- Patel, J. C., Hueffer, K., Lam, T. T. & Galan, J. E. (2009). Diversification of a *Salmonella* virulence protein function by ubiquitin-dependent differential localization. *Cell* **137**, 283–294.
- Saini, S., Brown, J. D., Aldridge, P. D. & Rao, C. V. (2008). FlhZ is a posttranslational activator of FlhD4C2-dependent flagellar gene expression. *J Bacteriol* **190**, 4979–4988.
- Schiemann, D. A. & Shope, S. R. (1991). Anaerobic growth of *Salmonella typhimurium* results in increased uptake by Henle 407 epithelial and mouse peritoneal cells *in vitro* and repression of a major outer membrane protein. *Infect Immun* **59**, 437–440.
- Song, M., Kim, H. J., Kim, E. Y., Shin, M., Lee, H. C., Hong, Y., Rhee, J. H., Yoon, H., Ryu, S. & other authors (2004). ppGpp-dependent stationary phase induction of genes on *Salmonella* pathogenicity island 1. *J Biol Chem* **279**, 34183–34190.
- Steele-Mortimer, O. (2008a). Infection of epithelial cells with *Salmonella enterica*. *Methods Mol Biol* **431**, 201–212.
- Steele-Mortimer, O. (2008b). The *Salmonella*-containing vacuole – moving with the times. *Curr Opin Microbiol* **11**, 38–45.
- Steele-Mortimer, O., Meresse, S., Gorvel, J. P., Toh, B. H. & Finlay, B. B. (1999). Biogenesis of *Salmonella typhimurium*-containing vacuoles in epithelial cells involves interactions with the early endocytic pathway. *Cell Microbiol* **1**, 33–49.

- Steele-Mortimer, O., Knodler, L. A., Marcus, S. L., Scheid, M. P., Goh, B., Pfeifer, C. G., Duronio, V. & Finlay, B. B. (2000). Activation of Akt/protein kinase B in epithelial cells by the *Salmonella typhimurium* effector sigD. *J Biol Chem* **275**, 37718–37724.
- Steele-Mortimer, O., Brumell, J. H., Knodler, L. A., Meresse, S., Lopez, A. & Finlay, B. B. (2002). The invasion-associated type III secretion system of *Salmonella enterica* serovar Typhimurium is necessary for intracellular proliferation and vacuole biogenesis in epithelial cells. *Cell Microbiol* **4**, 43–54.
- Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., Paulovich, A., Pomeroy, S. L., Golub, T. R. & other authors (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545–15550.
- Subramanian, A., Kuehn, H., Gould, J., Tamayo, P. & Mesirov, J. P. (2007). GSEA-P: a desktop application for Gene Set Enrichment Analysis. *Bioinformatics* **23**, 3251–3253.
- Temme, K., Salis, H., Tullman-Ercek, D., Levskaya, A., Hong, S. H. & Voigt, C. A. (2008). Induction and relaxation dynamics of the regulatory network controlling the type III secretion system encoded within *Salmonella* pathogenicity island 1. *J Mol Biol* **377**, 47–61.
- Teplitski, M., Goodier, R. I. & Ahmer, B. M. (2003). Pathways leading from BarA/SirA to motility and virulence gene expression in *Salmonella*. *J Bacteriol* **185**, 7257–7265.
- Thijs, I. M., De Keersmaecker, S. C., Fadda, A., Engelen, K., Zhao, H., McClelland, M., Marchal, K. & Vanderleyden, J. (2007). Delineation of the *Salmonella enterica* serovar Typhimurium HilA regulon through genome-wide location and transcript analysis. *J Bacteriol* **189**, 4587–4596.
- Tintle, N. L., Best, A. A., DeJongh, M., Van Bruggen, D., Heffron, F., Porwollik, S. & Taylor, R. C. (2008). Gene set analyses for interpreting microarray experiments on prokaryotic organisms. *BMC Bioinformatics* **9**, 469.
- Tusher, V. G., Tibshirani, R. & Chu, G. (2001). Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci U S A* **98**, 5116–5121.
- Valdivia, R. H. & Falkow, S. (1996). Bacterial genetics by flow cytometry: rapid isolation of *Salmonella typhimurium* acid-inducible promoters by differential fluorescence induction. *Mol Microbiol* **22**, 367–378.
- van Asten, F. J., Hendriks, H. G., Koninkx, J. F. & van Dijk, J. E. (2004). Flagella-mediated bacterial motility accelerates but is not required for *Salmonella* serotype Enteritidis invasion of differentiated Caco-2 cells. *Int J Med Microbiol* **294**, 395–399.
- van der Velden, A. W., Lindgren, S. W., Worley, M. J. & Heffron, F. (2000). *Salmonella* pathogenicity island 1-independent induction of apoptosis in infected macrophages by *Salmonella enterica* serotype typhimurium. *Infect Immun* **68**, 5702–5709.
- Virtaneva, K., Porcella, S. F., Graham, M. R., Ireland, R. M., Johnson, C. A., Ricklefs, S. M., Babar, I., Parkins, L. D., Romero, R. A. & other authors (2005). Longitudinal analysis of the group A *Streptococcus* transcriptome in experimental pharyngitis in cynomolgus macaques. *Proc Natl Acad Sci U S A* **102**, 9014–9019.
- Weening, E. H., Barker, J. D., Laarakker, M. C., Humphries, A. D., Tsolis, R. M. & Baumler, A. J. (2005). The *Salmonella enterica* serotype Typhimurium *lpf*, *bcf*, *stb*, *stc*, *std*, and *sth* fimbrial operons are required for intestinal persistence in mice. *Infect Immun* **73**, 3358–3366.

---

Edited by: D. L. Gally