

# Non-redundant roles for LXR $\alpha$ and LXR $\beta$ in atherosclerosis susceptibility in low density lipoprotein receptor knockout mice<sup>§</sup>

Eric D. Bischoff,<sup>1,\*</sup> Chris L. Daige,<sup>1,\*</sup> Mary Petrowski,<sup>\*</sup> Harry Dedman,<sup>\*</sup> Jennifer Pattison,<sup>†</sup> Joseph Juliano,<sup>†</sup> Andrew C. Li,<sup>†</sup> and Ira G. Schulman,<sup>2,§</sup>

Exelixis, Inc.,<sup>\*</sup> San Diego, CA; Department of Medicine,<sup>†</sup> University of California at San Diego, San Diego, CA; and Center for Molecular Design,<sup>§</sup> Department of Pharmacology, University of Virginia Health System, Charlottesville, VA

**Abstract** The liver X receptors LXR $\alpha$  and LXR $\beta$  play critical roles in maintaining lipid homeostasis by functioning as transcription factors that regulate genetic networks controlling the transport, catabolism, and excretion of cholesterol. The studies described in this report examine the individual anti-atherogenic activity of LXR $\alpha$  and LXR $\beta$  and determine the ability of each subtype to mediate the biological response to LXR agonists. Utilizing individual knockouts of LXR $\alpha$  and LXR $\beta$  in the *Ldlr*<sup>-/-</sup> background, we demonstrate that LXR $\alpha$  has a dominant role in limiting atherosclerosis in vivo. Functional studies in macrophages indicate that LXR $\alpha$  is required for a robust response to LXR ligands, whereas LXR $\beta$  functions more strongly as a repressor. Furthermore, selective knockout of LXR $\alpha$  in hematopoietic cells and rescue experiments indicate that the anti-atherogenic activity of this LXR subtype is not restricted to macrophages. These studies indicate that LXR $\alpha$  plays a selective role in limiting atherosclerosis in response to hyperlipidemia.—Bischoff, E. D., C. L. Daige, M. Petrowski, H. Dedman, J. Pattison, J. Juliano, A. C. Li, and I. G. Schulman. **Non-redundant roles for LXR $\alpha$  and LXR $\beta$  in atherosclerosis susceptibility in low density lipoprotein receptor knockout mice.** *J. Lipid Res.* 2010. 51: 900–906.

**Supplementary key words** cholesterol • hyperlipidemia • inflammation • macrophage

The contribution of elevated cholesterol levels to the development of cardiovascular disease and atherosclerosis is well documented. Nevertheless the molecular signaling pathways that regulate cholesterol homeostasis at the blood vessel wall, particularly in response to elevated cholesterol levels, remain to be fully deciphered (1). A critical

early event in the development of atherosclerosis is the recruitment of macrophages to the subendothelial space of vessel walls and the uncontrolled uptake of oxidized or aggregated low density lipoprotein particles. Continued accumulation of oxidized or aggregated LDL by macrophages and an associated inflammatory response leads to foam cell formation and the initiation of atherosclerosis (2).

The liver X receptors LXR $\alpha$  (NR1H3) and LXR $\beta$  (NR1H2), members of the nuclear hormone receptor superfamily of transcription factors, have been identified as important regulators of cholesterol homeostasis in multiple cell types, including macrophages (3). Treatment of cells with oxysterols, natural LXR ligands derived from cholesterol, or synthetic LXR agonists promotes the efflux of cholesterol by increasing expression of the ATP binding cassette transporters ABCA1 and ABCG1 and the apolipoprotein E (apoE) (4, 5). ABCA1, ABCG1, and apoE all participate in the transfer of intracellular and plasma membrane cholesterol to HDL, a process termed reverse cholesterol transport (6). Importantly, LXR agonists reduce atherosclerosis in animal models of cardiovascular disease, and upregulation of ABC transporters can be detected in the atherosclerotic lesions of treated animals (7, 8). The ability of LXRs to inhibit pro-inflammatory pathways may also contribute to the anti-atherogenic activities of these receptors (9). Previous work from our laboratory has demonstrated that LXR function in hema-

Abbreviations: ABCA1, ATP binding cassette transporter A1; ABCG1, ATP binding cassette transporter G1; apoE, apolipoprotein E; FPLC, fast-protein liquid chromatography; iNOS, inducible nitric oxide synthase; LXR, liver X receptor; MCP-1, monocyte chemoattractant protein 1; NF $\kappa$ B, nuclear factor kappa B; SREBP, sterol-regulatory element binding protein; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ .

<sup>1</sup>Present address of E. D. Bischoff and C. L. Daige: Kalypsys, Inc., San Diego, CA.

<sup>2</sup>To whom correspondence should be addressed.

e-mail: igs4c@virginia.edu

<sup>§</sup>The online version of this article (available at <http://www.jlr.org>) contains supplementary data in the form two figures.

This work was supported by funds from the University of Virginia Health System (I.G.S.) and by the National Institutes of Health Grant HL-087391 (J.P., J.J., and A.C.L.). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the National Institutes of Health.

Manuscript received 3 March 2009 and in revised form 23 March 2009.

Published, JLR Papers in Press, March 30, 2009

DOI 10.1194/jlr.M900096-JLR200

topoietic cells is necessary for the anti-atherogenic activity of LXR agonists (8), implicating macrophages as a critical site of action for the LXRs.

Each LXR subtype demonstrates a unique but overlapping expression pattern. LXR $\alpha$  is highly expressed in cell types that modulate cholesterol and fatty acid synthesis, including the liver and intestine, whereas LXR $\beta$  is ubiquitously expressed (10–12). In the liver, LXR $\alpha$  is the predominant subtype expressed, and studies indicate that LXR $\alpha$  controls fatty acid synthesis via regulation of the gene encoding the sterol-regulatory element binding protein 1c (SREBP1c) (13, 14). Additionally LXR $\alpha$  modulates cholesterol homeostasis by controlling expression of cholesterol 7 $\alpha$ -hydroxylase (11), the rate-limiting enzyme in the conversion of cholesterol to bile acids, and the cholesterol transporters ABCG5 and ABCG8 (15). In other tissues, including the intestine and macrophages, both subtypes are coexpressed and appear competent to regulate LXR target genes (4, 16, 17).

The ability to induce reverse cholesterol transport spurred great interest in the LXRs as drug targets for cardiovascular disease (13, 14, 18). Increases in plasma triglycerides and cholesterol observed in animals treated with synthetic LXR agonists, however, has raised concerns regarding the therapeutic potential of LXR-based drugs (18). As described above, the effects on lipid levels in animals treated with LXR agonists appear to be selectively mediated by LXR $\alpha$  in the liver (10, 15). In contrast, treatment with an LXR agonist reduced atherosclerosis in *apoE*<sup>-/-</sup>/*Lxr $\alpha$* <sup>-/-</sup> mice (19), suggesting that LXR $\beta$  alone is sufficient to mediate the anti-atherogenic functions of LXR activation. Not surprisingly, several investigators have suggested that LXR $\beta$ -selective activators may have an optimized therapeutic potential (10, 12). To our knowledge, the anti-atherogenic activity of such selective modulators has not yet been described.

Although the case for LXR $\beta$ -selective agonists has been put forward, the individual anti-atherogenic potential of LXR $\alpha$  and LXR $\beta$  has not yet been addressed. To examine the anti-atherogenic activity of each LXR subtype in vivo, genetic knockouts of LXR $\alpha$  or LXR $\beta$  were crossed into the LDL receptor knockout mouse (*Ldlr*<sup>-/-</sup>) background. Analysis of atherosclerosis in these strains indicates that LXR $\alpha$  plays a dominant role in limiting disease in response to elevated lipid levels and that this subtype is required for the full anti-atherogenic activity of LXR agonists. Experiments that selectively rescue LXR $\alpha$  function in bone marrow-derived or nonbone marrow-derived cells suggest that LXR activity may be required in cell types other than macrophages to limit diet-induced cardiovascular disease.

## METHODS

### Animals and diets

All animal experiments were approved by the Exelixis Institutional Animal Care and Research Advisory Committee. *Ldlr*<sup>-/-</sup>/*Lxr $\alpha$* <sup>-/-</sup> and *Ldlr*<sup>-/-</sup>/*Lxr $\beta$* <sup>-/-</sup> mice were created by mating *Lxr $\alpha$* <sup>-/-</sup> and *Lxr $\beta$* <sup>-/-</sup> mice (C57BL/6 genetic background)

with *LDLR*<sup>-/-</sup> mice (male B6.129S7-*Ldlr*<sup>tm1Her</sup>/J) purchased from Jackson Laboratories (Bar Harbor, ME). Homozygous *Lxr $\alpha$* <sup>-/-</sup>, *Lxr $\beta$* <sup>-/-</sup>, and *Lxr $\alpha$* <sup>-/-</sup>/*Lxr $\beta$* <sup>-/-</sup> knockout mice on the C57BL/6 genetic background were from a colony established and maintained at Exelixis, Inc. (San Diego, CA). Mice were fed standard chow (Lab Diet 5001) ad libitum until put on the study.

To examine atherosclerosis in *Ldlr*<sup>-/-</sup>, *Ldlr*<sup>-/-</sup>/*Lxr $\alpha$* <sup>-/-</sup>, and *Ldlr*<sup>-/-</sup>/*Lxr $\beta$* <sup>-/-</sup> mice, 8–9 week old male animals were placed on a Western diet (21% fat wt/wt, 0.15% cholesterol wt/wt; Purina Test Diets #21551) as described in the figure legends. When described in the text, the LXR agonist T0901317 was administered by oral gavage.

### Bone marrow transplantation

Bone marrow transplantations were carried as previously described (8, 20). After transplant, animals were allowed to recover on a standard diet (Lab Diet 5001) for 4 weeks before being fed the Western diet for an additional 8 weeks.

### Quantitation of atherosclerosis

Atherosclerosis in root sections and en face preparations were quantitated as previously described (8, 20).

### Lipid and lipoprotein analyses

Plasma total cholesterol and triglyceride levels were determined by colorimetric enzymatic assays that were adapted to 96-well plate as previously described (8). Terminal plasma samples from mice fed the Western diet for 20 weeks were pooled and fractionated by fast-protein liquid chromatography (FPLC) gel filtration on Superose 6 columns. Triglyceride and cholesterol concentrations of each fraction were determined as above.

### Measurement of TNF $\alpha$ plasma levels

To measure tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) protein levels, mice of the appropriate genotypes were fed the Western diets for 2 weeks and then given a single injection of LPS (25 $\mu$ g). Plasma TNF $\alpha$  levels were determined by ELISA (R and D Systems) 1.5 h after the LPS treatment.

### Macrophage isolation

Bone marrow-derived macrophages were isolated from 8–9 week old mice as previously described (21).

### Quantitative real-time PCR

Quantitative analysis of gene expression was carried out as previously described (8).


### Statistical analyses

Results were analyzed by one-way ANOVA (ANOVA) or Student's unpaired *t*-test using GraphPad Prism (GraphPad Software, Inc.).

## RESULTS

### Increased atherosclerosis in *Lxr $\alpha$* <sup>-/-</sup> mice

To define the anti-atherogenic activity of the individual LXR subtypes, single LXR $\alpha$  and LXR $\beta$  knockouts were crossed into the *Ldlr*<sup>-/-</sup> background to create *Ldlr*<sup>-/-</sup>/*Lxr $\alpha$* <sup>-/-</sup> and *Ldlr*<sup>-/-</sup>/*Lxr $\beta$* <sup>-/-</sup> double knockout strains. Animals from the three strains were placed on a Western diet (21% fat, 0.15% cholesterol), and plasma lipid levels were measured biweekly for 20 weeks (Fig. 1). Genetic deletion of LXR $\alpha$  in the *Ldlr*<sup>-/-</sup> background resulted in a dramatic decrease in



**Fig. 1.** Plasma lipid levels in double knockout mice. *Ldlr*<sup>-/-</sup> (n = 12), *Ldlr*<sup>-/-</sup>/*Lxrα*<sup>-/-</sup> (n = 9), and *Ldlr*<sup>-/-</sup>/*Lxrβ*<sup>-/-</sup> (n = 12) mice were fed a Western diet for 20 weeks, and plasma triglyceride (A) and total cholesterol (B) levels were determined.


plasma triglyceride levels at all time points, confirming previous data identifying this receptor subtype as the principal regulator of fatty acid synthesis (10, 12) (Fig. 1A). *Ldlr*<sup>-/-</sup>/*Lxrα*<sup>-/-</sup> mice also exhibited a significant decrease in total cholesterol levels beginning at week 4 (Fig. 1B). Plasma triglyceride and cholesterol levels were unchanged in *Ldlr*<sup>-/-</sup>/*Lxrβ*<sup>-/-</sup> animals for the first 12 weeks of the experiment (Fig. 1A, B). Beginning at week 12, however, there was a significant drop in lipid levels that coincided with an overall failure of these animals to thrive, including weight loss and a general decrease in activity (data not shown). FPLC analysis of pooled lipoprotein samples taken at completion of the study confirmed significant decreases in VLDL levels in both double knockout strains (see supplementary Fig. 1).

Quantitation of atherosclerosis by serial section analysis of the aortic root (Fig. 2A) demonstrated that deletion of LXRα significantly increases atherosclerosis by 25%. There was also a trend toward an increase in *Ldlr*<sup>-/-</sup>/*Lxrβ*<sup>-/-</sup> animals that did not reach statistical significance (*P* = 0.09). Immunohistochemical staining did not detect significant differences in macrophage or collagen content

when normalized to the lesion area (data not shown). To extend the analysis to additional areas beyond the aortic root, en face quantitation was carried out (Fig. 2B). Analysis of the arch and thoracic regions indicates there is an 80% increase in *Ldlr*<sup>-/-</sup>/*Lxrα*<sup>-/-</sup> samples (Fig. 2B, C) with the most dramatic differences distal to the aortic arch (Fig. 2C). Deleting LXRβ, however, had no effect. A significant increase in atherosclerosis is also selectively detected in *Ldlr*<sup>-/-</sup>/*Lxrα*<sup>-/-</sup> animals if the en face analysis is limited to aortic arch region (see supplementary Fig. II). Thus LXRα played a dominant role in limiting diet-induced atherosclerosis.

#### LXRα is required for the anti-atherogenic activity of LXR agonists

Synthetic LXR agonists have been shown to decrease atherosclerosis and promote the regression of lesions in *Ldlr*<sup>-/-</sup> mice (7, 8). To explore the contribution of each LXR subtype to the anti-atherogenic activity of LXR agonists, mice of the appropriate genotypes were exposed to the Western diet for 12 weeks and concurrently treated



**Fig. 2.** Atherosclerosis in double knockout mice. Mice were fed a Western diet for 20 weeks, and atherosclerosis was quantitated by root section (A) and en face analysis of the aorta (B) as described in the "Methods." (C) Representative Sudan IV stained aortas. \*Significantly different from *Ldlr*<sup>-/-</sup> controls.


with the LXR agonist T0901317. As expected from previous studies, plasma triglycerides were significantly lower in *Ldlr*<sup>-/-</sup>/*Lxrα*<sup>-/-</sup> mice, regardless of agonist treatment, while T0901317 elevated triglyceride levels in strains expressing LXRα (Fig. 3A). Importantly, *Ldlr*<sup>-/-</sup>/*Lxrβ*<sup>-/-</sup> mice treated with 10 mg/kg T0901317 rapidly developed massive hypertriglyceridemia (data not shown) that could not be accurately measured, and 3 out of 10 mice died within 4 weeks of initiating treatment. At week 4, the dose for this group was dropped to 3 mg/kg, and the rest of the animals survived until study completion. Nevertheless, given the small number of animals that remained in this group, the conclusions regarding the effects of T0901317 on *Ldlr*<sup>-/-</sup>/*Lxrβ*<sup>-/-</sup> mice should be interpreted with caution. The LXR agonist-dependent hypertriglyceridemia observed in these animals suggests, at least in the *Ldlr*<sup>-/-</sup> background, that LXRβ partially represses the ability of LXRα to regulate triglyceride levels. Little or no effect of T0901317 on total plasma cholesterol levels was detected (Fig. 3B).

Consistent with previous studies (7, 8, 22), treatment of *Ldlr*<sup>-/-</sup> mice with T0901317 significantly reduced atherosclerosis when quantitated by root section (Fig. 3C) or en face analysis (Fig. 3D). T0901317 also significantly reduced atherosclerosis at the aortic root in the absence of either of LXRα or LXRβ (Fig. 3C), indicating that each LXR subtype


alone can mediate an anti-atherogenic response to an LXR agonist. Once again, however, en face analysis uncovered a large increase in atherosclerosis in *Ldlr*<sup>-/-</sup>/*Lxrα*<sup>-/-</sup> animals that was unaffected by treatment with T0901317 (Fig. 3D). Thus when atherosclerosis is measured throughout the aorta, LXRα is necessary for the anti-atherogenic activity of LXR agonists. In contrast, even at a lower dose, the LXR agonist remained effective in animals lacking LXRβ (Fig. 3D).

### LXRα selective regulation of gene expression in macrophages

We previously demonstrated that macrophages are an important site of action for the anti-atherogenic activity of LXR agonists (8, 20). Therefore to explore a mechanistic basis for the selective effect of LXRα deletion on atherosclerosis, the expression of LXR target genes was examined in bone marrow-derived macrophages isolated from C57BL/6, *Lxrα*<sup>-/-</sup>, *Lxrβ*<sup>-/-</sup>, and *Lxrα*<sup>-/-</sup>/*Lxrβ*<sup>-/-</sup> mice. As shown in Fig. 4A and 4B, the ability of T0901317 to induce expression of ABCA1 and ABCG1 was selectively impaired in *Lxrα*<sup>-/-</sup> cells (compare solid black lines to dotted lines). These results indicate that LXRα is required to maximally induce genes involved in reverse cholesterol transport. The subtype selective regulation of SREBP1c in response to agonist treatment is somewhat different



**Fig. 3.** LXR agonist activity in double knockout mice. Mice were exposed to a Western diet for 12 weeks and dosed daily with vehicle or 10 mg/kg T0901317. The *Ldlr*<sup>-/-</sup>/*Lxrβ*<sup>-/-</sup> + T0901317 group was switched from 10 mg/kg to 3 mg/kg at week 4. After completion of the study plasma triglycerides (A) and total plasma cholesterol (B) were measured. \*Significantly different from strain-matched vehicle controls. \*\*Significantly different from vehicle-treated *Ldlr*<sup>-/-</sup> mice. Atherosclerosis was measured by quantitation of root sections (C) and by en face analysis of the entire aorta (D). \*Significantly different from strain-matched vehicle controls. \*\*Significantly different from vehicle-treated *Ldlr*<sup>-/-</sup> mice.



**Fig. 4.** LXR target gene expression in bone marrow derived macrophages. Bone marrow derived macrophages isolated from C57BL/6, *LXRα*<sup>-/-</sup>, *LXRβ*<sup>-/-</sup>, and *LXRα*<sup>-/-</sup>/*LXRβ*<sup>-/-</sup> mice were treated with vehicle or increasing concentrations of T0901317 for 48 h. After agonist treatment total RNA was isolated, and the mRNA levels of ABCA1 (A and D), ABCG1 (B and E), and SREBP1c (C and F) were determined by quantitative PCR. \*Significantly different from C57BL/6 controls ( $P \leq 0.05$ ). ABCA1, ATP binding cassette transporter A1; ABCG1, ATP binding cassette transporter G1; SREBP, sterol-regulatory element binding protein.


(Fig. 4C). Deletion of LXR $\alpha$  has only a small effect on the agonist-dependent induction of this target gene, whereas deletion of LXR $\beta$  actually allows a super-induction, suggesting that LXR $\beta$  functions to repress the ability of LXR $\alpha$  to induce SREBP1c.

Nuclear receptors are also known to inhibit gene expression. In the absence of activating ligands LXR can repress transcription by recruiting corepressors to promoters that contain LXR binding sites (21). Interestingly, when the mRNA levels of ABCA1, ABCG1, and SREBP1c are examined in the absence of T0901317 (Fig. 4D–F), deletion of LXR $\alpha$  has no effect on the basal/ligand-independent levels of the three target genes. On the other hand, all three mRNAs are increased in *LXRβ*<sup>-/-</sup> and *LXRα*<sup>-/-</sup>/*LXRβ*<sup>-/-</sup> cells, indicating that LXR $\beta$  plays an important role in repressing target gene expression in the absence of activating signals. LXRs can also inhibit pro-inflammatory gene expression in the presence of agonists by inhibiting the activity of nuclear factor kappa  $\beta$  (NF $\kappa$ B) (23). We did not, however, observe LXR subtype-specific differences in the ability of T0901317 to inhibit expression of monocyte chemoattractant protein 1 (MCP-1), TNF $\alpha$ , or inducible nitric oxide synthase (iNOS) in lipopolysaccharide stimulated bone marrow derived macrophages (data not shown).

#### LXR $\alpha$ functions in bone marrow- and nonbone marrow-derived cells

Macrophages as well as other immune cells play a critical role in the development of atherosclerosis (2), and the

ability of LXRs to regulate reverse cholesterol transport and inflammation in these cells is thought to underlie their anti-atherogenic activities (9). To determine if it is simply the absence of LXR $\alpha$  in the hematopoietic system that lead to the increased atherosclerosis observed in the *Ldlr*<sup>-/-</sup> background, bone marrow transplantation experiments were used to either selectively eliminate or selectively express LXR $\alpha$  in hematopoietic cells (Fig. 5). As expected, transplantation of *Ldlr*<sup>-/-</sup> mice with bone marrow from *Ldlr*<sup>-/-</sup>/*LXRα*<sup>-/-</sup> animals (hematopoietic system knockout) increased atherosclerosis relative to controls (in Fig. 5, compare groups 1 and 2;  $P = 0.03$ ). Nevertheless, atherosclerosis in *Ldlr*<sup>-/-</sup> mice transplanted with *Ldlr*<sup>-/-</sup>/*LXRα*<sup>-/-</sup> marrow was not as great as that observed in the complete LXR $\alpha$  knockout (i.e., irradiated *Ldlr*<sup>-/-</sup>/*LXRα*<sup>-/-</sup> mice that received *Ldlr*<sup>-/-</sup>/*LXRα*<sup>-/-</sup> bone marrow; in Fig. 5, compare groups 2 and 4;  $P = 0.0001$ ). Thus, eliminating LXR $\alpha$  activity in hematopoietic cells did not entirely recapitulate the phenotype of the complete knockout. Similarly, the introduction of LXR $\alpha$  positive bone marrow into *Ldlr*<sup>-/-</sup>/*LXRα*<sup>-/-</sup> mice (hematopoietic system selective expression) decreased atherosclerosis (in Fig. 5, compare groups 4 and 5;  $P = 0.0001$ ) but not to the levels measured in *Ldlr*<sup>-/-</sup> controls (in Fig. 5, compare groups 1 and 5;  $P = 0.01$ ). Taken together, the results of the bone marrow transplantation experiments indicate that LXR $\alpha$  function in cell types other than macrophages may contribute to the anti-atherogenic activity of this receptor.



**Fig. 5.** LXR $\alpha$  activity is required in bone marrow derived and nonbone marrow cells. Recipient mice (as defined in the figure) were irradiated and reconstituted with bone marrow from mice of the appropriate genotype. Four weeks after recovery from the transplant, mice were exposed to the Western diet, and atherosclerosis was quantitated by en face analysis after an additional 8 weeks. \*Statistically significant difference from  $Ldlr^{-/-}$  to  $Ldlr^{-/-}$  controls (see text for *P* values). \*\*Statistically significant difference from  $Ldlr^{-/-}/Lxra^{-/-}$  to  $Ldlr^{-/-}/Lxra^{-/-}$  controls (see text for *P* values).

## DISCUSSION

The ability of LXRs to regulate reverse cholesterol transport in macrophages has stimulated great interest in understanding the anti-atherogenic activity of the two LXR subtypes. Analysis of single genetic deletions of each receptor in the  $Ldlr^{-/-}$  background reveals a dominant effect for LXR $\alpha$  in the control of cardiovascular disease. Quantitation of atherosclerosis in animals exposed to a Western diet uncovered an 80% increase in  $Lxra^{-/-}$  animals, with dramatic increases in the lesion area in regions distal to the aortic arch. Little or no effect of deleting LXR $\beta$  was observed. Our results are consistent with reports by Bradley et al. (19) and Teupser et al. (24) that detected increased atherosclerosis in  $apoE^{-/-}/Lxra^{-/-}$  mice and decreased atherosclerosis in  $Ldlr^{-/-}$  mice over expressing LXR $\alpha$ , respectively. Neither of these studies, however, directly compared the two LXR subtypes.

Functional experiments using bone marrow-derived macrophages demonstrate that LXR $\alpha$  is quantitatively a stronger transcriptional activator of the ABCA1 and ABCG1 promoters in response to agonists than is LXR $\beta$ . The subtype selective response to agonist treatment measured in macrophages suggests an impaired ability of  $Lxra^{-/-}$  macrophages to induce gene expression and reverse cholesterol transport in response to elevated cholesterol levels. In support of the gene expression data, the anti-atherogenic activity of the agonist T0901317 is muted in the absence of LXR $\alpha$ . LXR $\beta$ , on the other hand, appears to be more responsible for setting the basal level of macrophage ABCA1 and ABCG1 expression by limiting transcription in the absence of activating signals.

The effect of the LXR agonist T0901317 on atherosclerosis in  $Ldlr^{-/-}/Lxra^{-/-}$  mice differed depending on the method of analysis used. Serial section of the aortic root indicated that T0901317 reduced atherosclerosis in

$Ldlr^{-/-}/Lxra^{-/-}$  mice (i.e., LXR $\beta$  alone can mediate an anti-atherogenic response). On the other hand, en face analysis of the aortic arch and thoracic region indicated that the agonist had no effect in the absence of LXR $\alpha$ . Additionally, the en face analysis uncovered a large increase in atherosclerosis distal to the aortic arch region in these animals. It is not unusual to see differences in atherosclerosis measured by these two methods. Serial sections provide cross-sectional information from a single area whereas en face analysis measures the surface area of the entire segment covered by lesions. In mouse models of atherosclerosis, lesions first develop in the aortic root and arch areas and then progress distally down the segment as animals age (25). The large increase in atherosclerosis observed in  $Lxra^{-/-}$  animals suggests that LXR $\alpha$  may selectively influence the kinetics of early lesion initiation and/or disease progression. After exposure to the Western diet for 20 weeks, such an acceleration of lesion progression may be easily visible at areas distal to the arch while less apparent at the older, more advanced areas measured at the aortic root. Zhu et al. (26) have also recently observed that LXR $\alpha$  is expressed at higher levels in the thoracic aorta compared with the arch region, raising the possibility that this subtype plays a unique or quantitatively dominant role in the thoracic region. LXR $\beta$  expression in these regions was not examined.

The function of LXR $\beta$  is more difficult to discern from these studies. After 12 weeks on the Western diet, there was a general failure of the  $Ldlr^{-/-}/Lxr\beta^{-/-}$  mice to thrive. Although there was no obvious pathology at necropsy 8 weeks later, lymphoid hyperplasia has been observed in older  $Lxr\beta^{-/-}$  mice (27), which, when coupled with the Western diet, may have contributed to the poor health of these animals. Interestingly, in  $Lxr\beta^{-/-}$  macrophages, the basal levels of ABCA1 and ABCG1 were elevated, suggesting a role for this receptor in limiting reverse cholesterol transport in the absence of activating signals. Additionally, treatment of  $Ldlr^{-/-}/Lxr\beta^{-/-}$  mice with T0901317 produced a massive increase in plasma triglycerides, indicating that LXR $\beta$  functions to limit the ability of LXR $\alpha$  to induce fatty acid synthesis (at least under hyperlipidemic conditions). A similar inhibitory effect of LXR $\beta$  on the agonist-dependent induction of SREBP1c was detected in macrophages. These results suggest opposing functions for the two subtypes, with LXR $\alpha$  functioning to drive transcription in the presence of agonists and LXR $\beta$  functioning to repress transcription when ligands are active.

## CONCLUSION

The ability to regulate reverse cholesterol transport and inflammation at the site of atherosclerotic lesions has made macrophages the focus of LXR studies (8, 20). To address the sites of LXR $\alpha$ -dependent anti-atherogenic activity, we used bone marrow transplantations to selectively eliminate or selectively express LXR $\alpha$  in the hematopoietic system. The introduction of LXR $\alpha$  positive bone marrow into  $Ldlr^{-/-}/Lxra^{-/-}$  mice did partially rescue the athero-

genic phenotype. The level of atherosclerosis in these “rescued” mice, however, was still significantly greater than the level of observed in mice expressing LXR $\alpha$  throughout the body. The mice used in the transplantation studies were irradiated and reconstituted with bone marrow at 8 weeks of age. Thus we cannot rule out the possibility that the absence of LXR $\alpha$  in the hematopoietic system prior to irradiation contributed to increased atherosclerosis observed in the rescued mice relative to the controls expressing LXR $\alpha$  everywhere. Nevertheless, the results of this study also suggest the possibility of an important site of action for LXR $\alpha$  that may be distinct from macrophages. Obvious candidates for such nonbone marrow–derived cells are vascular endothelial cells and smooth muscle cells, which are also components of atherosclerotic lesions. Additionally, the liver may be a critical site for the anti-atherogenic activity of LXR $\alpha$ . Analysis of plasma lipids in *Ldlr*<sup>-/-</sup>/*Lxr $\alpha$* <sup>-/-</sup> mice revealed relatively high levels of cholesterol and relatively low levels of triglycerides, raising the possibility that an aberrant pro-atherogenic lipoprotein particle may circulate in these animals. LXR $\alpha$  is also a critical regulator of apoE expression (5), and studies have suggested that apoE-deficient lipoproteins enhance macrophage foam cell formation (28). Finally, the elimination of hepatic cholesterol via excretion into the bile and catabolism to bile acids is controlled selectively by LXR $\alpha$  (11, 15). Ultimately additional cell-specific knockouts will be required to further define the critical sites of LXR $\alpha$  activity. **■**

The authors thank Lynn Hedrick for comments about the manuscript.

## REFERENCES

- Brown, M. S., and J. L. Goldstein. 2009. Cholesterol feedback: from Schoenheimer's bottle to Scap's MELADL. *J Lipid Res.* **50**(Suppl): S15–S27.
- Glass, C. K., and J. L. Witztum. 2001. Atherosclerosis. The road ahead. *Cell.* **104**: 503–516.
- Tontonoz, P., and D. J. Mangelsdorf. 2003. Liver X receptor signaling pathways in cardiovascular disease. *Mol. Endocrinol.* **17**: 985–993.
- Costet, P., Y. Luo, N. Wang, and A. R. Tall. 2000. Sterol-dependent transactivation of the ABC1 promoter by the liver X receptor/retinoid X receptor. *J. Biol. Chem.* **275**: 28240–28245.
- Laffitte, B. A., J. J. Repa, S. B. Joseph, D. C. Wilpitz, H. R. Kast, D. J. Mangelsdorf, and P. Tontonoz. 2001. LXRs control lipid-inducible expression of the apolipoprotein E gene in macrophages and adipocytes. *Proc. Natl. Acad. Sci. USA.* **98**: 507–512.
- Wang, X., and D. J. Rader. 2007. Molecular regulation of macrophage reverse cholesterol transport. *Curr. Opin. Cardiol.* **22**: 368–372.
- Joseph, S. B., E. McKilligin, L. Pei, M. A. Watson, A. R. Collins, B. A. Laffitte, M. Chen, G. Noh, J. Goodman, G. N. Hagger, et al. 2002. Synthetic LXR ligand inhibits the development of atherosclerosis in mice. *Proc. Natl. Acad. Sci. USA.* **99**: 7604–7609.
- Levin, N., E. D. Bischoff, C. L. Daige, D. Thomas, C. T. Vu, R. A. Heyman, R. K. Tangirala, and I. G. Schulman. 2005. Macrophage liver X receptor is required for antiatherogenic activity of LXR agonists. *Arterioscler. Thromb. Vasc. Biol.* **25**: 135–142.
- Zelcer, N., and P. Tontonoz. 2006. Liver X receptors as integrators of metabolic and inflammatory signaling. *J. Clin. Invest.* **116**: 607–614.
- Lund, E. G., L. B. Peterson, A. D. Adams, M. H. Lam, C. A. Burton, J. Chin, Q. Guo, S. Huang, M. Latham, J. C. Lopez, et al. 2006. Different roles of liver X receptor alpha and beta in lipid metabolism: effects of an alpha-selective and a dual agonist in mice deficient in each subtype. *Biochem. Pharmacol.* **71**: 453–463.
- Peet, D. J., S. D. Turley, W. Ma, B. A. Janowski, J.-M. A. Lobaccaro, R. E. Hammer, and D. J. Mangelsdorf. 1998. Cholesterol and bile acid metabolism are impaired in mice lacking the nuclear oxysterol receptor LXR alpha. *Cell.* **93**: 693–704.
- Quinet, E. M., D. A. Savio, A. R. Halpern, L. Chen, G. U. Schuster, J. A. Gustafsson, M. D. Basso, and P. Nambi. 2006. Liver X receptor (LXR)-beta regulation in LXRalpha-deficient mice: implications for therapeutic targeting. *Mol. Pharmacol.* **70**: 1340–1349.
- Repa, J. J., G. Liang, J. Ou, Y. Bashmakov, J. M. A. Lobaccaro, I. Shimomura, B. Shan, M. S. Brown, J. L. Goldstein, and D. J. Mangelsdorf. 2000. Regulation of mouse sterol regulatory element-binding protein-1c gene (SREBP-1c) by oxysterol receptors, LXRalpha and LXRbeta. *Genes Dev.* **14**: 2819–2830.
- Schultz, J. R., H. Tu, A. Luk, J. J. Repa, J. C. Medina, L. Li, S. Schwendner, S. Wang, M. Thoolen, D. J. Mangelsdorf, et al. 2000. Role of LXRs in control of lipogenesis. *Genes Dev.* **14**: 2831–2838.
- Yu, L., J. York, K. von Bergmann, D. Lutjohann, J. C. Cohen, and H. H. Hobbs. 2003. Stimulation of cholesterol excretion by the liver X receptor agonist requires ATP-binding cassette transporters G5 and G8. *J. Biol. Chem.* **278**: 15565–15570.
- Repa, J. J., S. D. Turley, J. A. Lobaccaro, J. Medina, L. Li, K. Lustig, B. Shan, R. A. Heyman, J. M. Dietschy, and D. J. Mangelsdorf. 2000. Regulation of absorption and ABC1-mediated efflux of cholesterol by RXR heterodimers. *Science.* **289**: 1524–1529.
- Schuster, G. U., P. Parini, L. Wang, S. Alberti, K. R. Steffensen, G. K. Hansson, B. Angelin, and J. A. Gustafsson. 2002. Accumulation of foam cells in liver X receptor-deficient mice. *Circulation.* **106**: 1147–1153.
- Groot, P. H., N. J. Pearce, J. W. Yates, C. Stocker, C. Sauermeier, C. P. Doe, R. N. Willette, A. Olzinski, T. Peters, D. d'Epagnier, et al. 2005. Synthetic LXR agonists increase LDL in CETP species. *J. Lipid Res.* **46**: 2182–2191.
- Bradley, M. N., C. Hong, M. Chen, S. B. Joseph, D. C. Wilpitz, X. Wang, A. J. Lusis, A. Collins, W. A. Hseuh, J. L. Collins, et al. 2007. Ligand activation of LXR beta reverses atherosclerosis and cellular cholesterol overload in mice lacking LXR alpha and apoE. *J. Clin. Invest.* **117**: 2337–2346.
- Tangirala, R. K., E. D. Bischoff, S. B. Joseph, B. L. Wagner, R. Walczak, B. A. Laffitte, C. L. Daige, D. Thomas, R. A. Heyman, D. J. Mangelsdorf, et al. 2002. Identification of macrophage liver X receptors as inhibitors of atherosclerosis. *Proc. Natl. Acad. Sci. USA.* **99**: 11896–11901.
- Wagner, B. L., A. F. Valledor, G. Shao, C. L. Daige, E. D. Bischoff, M. Petrowski, K. Jepsen, S. H. Baek, R. A. Heyman, M. G. Rosenfeld, et al. 2003. Promoter-specific roles for liver X receptor/corepressor complexes in the regulation of ABCA1 and SREBP1 gene expression. *Mol. Cell. Biol.* **23**: 5780–5789.
- Terasaka, N., A. Hiroshima, T. Koieyama, N. Ubukata, Y. Morikawa, D. Nakai, and T. Inaba. 2003. T-0901317, a synthetic liver X receptor ligand, inhibits development of atherosclerosis in LDL receptor-deficient mice. *FEBS Lett.* **536**: 6–11.
- Bensinger, S. J., and P. Tontonoz. 2008. Integration of metabolism and inflammation by lipid-activated nuclear receptors. *Nature.* **454**: 470–477.
- Teupser, D., D. Kretzschmar, C. Tennert, R. Burkhardt, W. Wilfert, D. Fengler, R. Naumann, A. E. Sippel, and J. Thiery. 2008. Effect of macrophage overexpression of murine liver X receptor-alpha (LXR-alpha) on atherosclerosis in LDL-receptor deficient mice. *Arterioscler. Thromb. Vasc. Biol.* **28**: 2009–2015.
- Tangirala, R. K., E. M. Rubin, and W. Palinski. 1995. Quantitation of atherosclerosis in murine models: correlation between lesions in the aortic origin and in the entire aorta, and differences in the extent of lesions between sexes in LDL receptor-deficient and apolipoprotein E-deficient mice. *J. Lipid Res.* **36**: 2320–2328.
- Zhu, M., Y. Fu, Y. Hou, N. Wang, Y. Guan, C. Tang, J. Y. Shyy, and Y. Zhu. 2008. Laminar shear stress regulates liver X receptor in vascular endothelial cells. *Arterioscler. Thromb. Vasc. Biol.* **28**: 527–533.
- Bensinger, S. J., M. N. Bradley, S. B. Joseph, N. Zelcer, E. M. Janssen, M. A. Hausner, R. Shih, J. S. Parks, P. A. Edwards, B. D. Jamieson, et al. 2008. LXR signaling couples sterol metabolism to proliferation in the acquired immune response. *Cell.* **134**: 97–111.
- Wu, D., C. Sharan, H. Yang, J. S. Goodwin, L. Zhou, G. A. Grabowski, H. Du, and Z. Guo. 2007. Apolipoprotein E-deficient lipoproteins induce foam cell formation by downregulation of lysosomal hydrolases in macrophages. *J. Lipid Res.* **48**: 2571–2578.