

Published in final edited form as:

Oral Dis. 2008 October ; 14(7): 613–619. doi:10.1111/j.1601-0825.2007.01429.x.

EXPRESSION OF PLUNC FAMILY MEMBERS IN BENIGN AND MALIGNANT SALIVARY GLAND TUMOURS

P.A. Vargas^{1,2}, P.M. Speight¹, CD Bingle³, A.W. Barrett⁴, and L Bingle^{1,*}

¹Department of Oral Pathology, School of Clinical Dentistry, University of Sheffield, Sheffield, UK.

²Department of Oral Diagnosis, Oral Pathology Section, School of Dentistry of Piracicaba, UNICAMP, Piracicaba-Sao Paulo State-Brazil.

³Academic Unit of Respiratory Medicine, University of Sheffield Medical School, Royal Hallamshire Hospital, Sheffield, UK.

⁴Department of Histopathology, Queen Victoria Hospital, East Grinstead, West Sussex, UK.

Abstract

Objectives: The aim of this study was to determine the expression of PLUNC proteins in benign and malignant salivary gland tumours and thus their potential use as diagnostic and/or prognostic tools.

Materials and Methods: A tissue microarray was assembled from 64 salivary gland tumours including adenoid cystic carcinoma, carcinoma ex-pleomorphic adenoma, mucoepidermoid carcinoma, polymorphous low-grade adenocarcinoma, pleomorphic adenoma, acinic cell carcinoma, myoepithelial carcinoma and papillary cystadenocarcinoma. Clinicopathological data were collected retrospectively and immunohistochemical analysis of 3 PLUNC proteins (SPLUNC1, SPLUNC2, and LPLUNC 1) was performed. Immunoreactivity was assessed as positive or negative.

Results: PLUNC expression was only found in mucoepidermoid carcinomas and papillary cystadenocarcinoma, all other tumours studied were negative. Mucin plugs, mucous and intermediate cells of mucoepidermoid carcinomas were positive for LPLUNC1 and SPLUNC2 but areas composed of epidermoid and clear cells were negative for all PLUNCs. Papillary cystadenocarcinoma was positive for all PLUNCs. No correlation was found with tumour grade or outcome.

Conclusions: Intense expression of two PLUNC proteins in mucous cells and mucin plugs of mucoepidermoid carcinoma and papillary cystadenocarcinoma indicate they could be used as additional diagnostic tools in some equivocal cases but further studies are needed to understand the biological processes involved in PLUNC expression.

Keywords

PLUNC; Salivary; Tumour; Mucoepidermoid; Papillary; Cystadenocarcinoma

INTRODUCTION

Salivary gland tumours are uncommon and histologically heterogeneous (Speight and Barrett, 2002). They predominantly affect major salivary glands but minor salivary glands,

*Corresponding author address: Lynne Bingle PhD Department of Oral Pathology, School of Clinical Dentistry, University of Sheffield, Claremont Crescent, Sheffield, S10 2TA l.bingle@sheffield.ac.uk Tel: 0114 271 7953 Fax: 0114 271 7894.

mainly located in the palate, can also be affected (Vargas et al, 2002). The most common salivary gland tumour is pleomorphic adenoma while mucoepidermoid carcinoma and adenoid cystic carcinoma are the most frequent malignant tumours in the major and minor salivary glands (Jones and Franklin, 2006). Salivary gland tumours show a varied and complex pattern of histopathological features and diagnosis may be difficult even for the most experienced diagnostic pathologist. The latest WHO classification (Barnes et al, 2005) contains over 40 named neoplasms, many of which show significant morphological diversity resulting in overlapping features which make differentiation between tumour types difficult. There is therefore a constant need to evaluate potential new biomarkers that may be useful in diagnosis and prognosis.

Palate Lung Nasal Clone (*PLUNC*) was first described in the nasal epithelium of the mouse embryo and the trachea and bronchi of adult mouse lung (Weston et al, 1999). We have subsequently shown that human *PLUNC* belongs to a family of 10 genes located on chromosome 20q11 (Bingle and Bingle, 2000; Bingle and Craven, 2002; Bingle et al, 2004) that are expressed in the upper airways, oro- and nasopharynx. Due to structural similarities with a family of lipid binding/transfer proteins that includes bactericidal/permeability-increasing protein (BPI) and lipopolysaccharide binding protein (LBP), we have suggested that these proteins may play a role in host defence (Bingle and Craven, 2002). *PLUNC* proteins subdivide into short (*SPLUNC*) and long (*LPLUNC*) proteins, which contain domains structurally similar to one or both domains of BPI respectively (Bingle and Craven, 2002). Although many *PLUNC*s remain poorly characterised, our analysis suggests that they are expressed in the major salivary glands, the minor mucosal glands of the oral cavity and the respiratory epithelium of the nasal, tracheal and bronchial passages (Bingle and Bingle, 2000; Bingle and Craven, 2002). Limited studies have been performed at the protein level but some *PLUNC*s have been identified in saliva, nasal fluid and pulmonary secretions. For example *SPLUNC1*, *SPLUNC2*, and *LPLUNC1* have been identified in saliva (Vitorino et al, 2004; Ramachandran et al, 2006; Guo et al, 2006) suggesting that these proteins are produced by the salivary glands. Although human *PLUNC*s have only recently been identified, a number of rodent orthologues including parotid secretory protein (*psp*) have been studied extensively (Ball et al, 2003). *psp* was identified as an abundant component of rat and mouse saliva (Owerbach and Hjorth, 1980; Mirels and Ball, 1992) and is highly expressed in salivary gland tissue. A second *psp* related protein, submandibular gland protein B (*smgb*) was also identified as being highly expressed in the salivary glands of rats (Mirels and Ball, 1992). Our recent analysis has shown that *SPLUNC2* is the orthologue of rodent *psp* (Bingle et al, 2004) and consistent with this observation *SPLUNC2* is predominantly expressed in the major salivary glands (Bingle and Craven, 2002; Geetha et al, 2005).

At the same time as we identified the human *SPLUNC1* gene, it was also identified (and named *LUNX*) as a potential marker for micrometastasis for non-small cell lung cancer (NSCLC) (Iwao et al, 2001). *LUNX* expression was also detected in colorectal cancer, esophageal cancer, hepatocellular carcinoma and breast cancer (Iwao et al, 2001). The value of *SPLUNC1/LUNX* as a diagnostic marker in NSCLC has recently been confirmed in two additional studies (Mitas et al, 2003a; Mitas et al, 2003b) and we have also shown that *SPLUNC1* protein is highly expressed in lung cancers with a glandular phenotype (Bingle et al, 2005). Differential expression of *PLUNC* family members has also been demonstrated in head and neck squamous cell carcinoma (Lemaire et al, 2003), nasopharyngeal carcinoma (Zhang et al, 2003) and breast tumours (Egland et al, 2003). Taken together these studies suggest that differential expression of *PLUNC* genes may be a characteristic of cancers and the study of these genes may prove to be of diagnostic and prognostic value. *PLUNC* proteins have not been investigated in salivary gland tumours, but their expression in other

glandular neoplasms and in normal salivary glands suggests that they should be expressed and may be useful as diagnostic markers.

The aim of this study therefore was to determine the expression of PLUNC family members in benign and malignant salivary gland tumours in an attempt to detect their potential usefulness in the diagnosis and prognosis of these lesions.

PATIENTS AND METHODS

Tissue samples

Archived formalin-fixed, paraffin-embedded tissue blocks of salivary gland tumours were obtained from the Department of Oral Pathology, School of Clinical Dentistry, University of Sheffield. All the samples were examined on haematoxylin and eosin (H&E) stained sections and classified according to the latest World Health Organization (WHO) guidelines (Barnes et al, 2005). The study was approved by the South Sheffield Research Ethics Committee.

Sixty-four cases of eight different types of salivary gland tumours were assembled in a tissue microarray format. There were thirteen adenoid cystic carcinomas (ACC), 10 carcinoma ex pleomorphic adenomas (CEPA), 10 pleomorphic adenomas (PA), 10 mucoepidermoid carcinomas (MEC), 10 polymorphous low-grade adenocarcinomas (PLGA), 9 acinic cell carcinomas (AcCC), 1 myoepithelial carcinoma (MC), 1 papillary cystadenocarcinoma (PC). We assessed recurrent tumours from 3 ACC patients and 1 MC patient. Clinical data including gender, age, tumour size, location of lesion and patient outcome (i.e. recurrence or metastasis) were collected by reviewing the pathology records (Table 1). We have previously reported DNA ploidy findings in part of this group of tumours (Vargas et al, 2007).

Tissue Microarray (TMA)

Representative tumour areas were selected and marked on H&E stained sections by using an objective marker (1.8mm, Nikon Corporation, Japan). TMA were constructed using a manual tissue arrayer (Mitogen, Beecher instruments, Inc.). Three representative cylindrical cores of 1.0mm diameter and 3mm depth were taken from each tissue block. The donor cores were then arrayed sequentially into a recipient paraffin block. Three recipient blocks were constructed for immunohistochemical analysis. An oral squamous cell carcinoma (OSCC) was inserted into the left upper corner of each block for orientation.

Generation of SPLUNC2 and LPLUNC1 antibodies

Custom antibodies were generated against SPLUNC2 and LPLUNC1 by Eurogentec (Seraing, Belgium) using established methods. Specifically we generated two antibodies by injection of two synthetic peptides into a pair of rabbits. For SPLUNC2 the peptide sequence corresponded to amino acids, 236-249-COOH: (VDNPQHKTQLQTLI) which represents the extreme C-terminal region of the protein. For LPLUNC1 the peptide sequences corresponded to amino acids, 139-154: (TIRMDTSASGPTRLV) which is located within the first BPI domain of LPLUNC1. Pooled final serum was used for affinity purification against each of the individual peptides. Antibodies were validated by western blotting using recombinant SPLUNC2 and LPLUNC1 as previously described (Bingle et al, 2005).

Immunohistochemical analysis

Immunostaining was performed on 4µm paraffin sections from the TMA blocks, mounted onto adhesive-coated glass slides (Surgipath, Richmond, USA). Slides were dried in the oven for one and a half hours to ensure the sections were firmly attached. All reactions

followed standard protocols. Sections were deparaffinized and rehydrated. Endogenous peroxidase activity was blocked by quenching the sections in 3% H₂O₂ for 20 minutes. The sections were then rinsed in PBS and where necessary (antibodies SPLUNC2, LPLUNC1) antigen retrieval was performed; tri-sodium citrate buffer for 8 minutes in a microwave oven followed by rinsing in PBS. Sections were incubated in 100% normal serum (corresponding to species of secondary antibody) for 30 minutes at room temperature in a humidified chamber. The serum was drained and replaced with primary antibody diluted in 100% normal serum (as used for blocking). The antibody dilutions used were 1:300 SPLUNC1, 1:250 SPLUNC2 and 1:100 LPLUNC1 and left on the sections for an overnight incubation at 4°C in a humidified chamber. A Vectastain kit (Vector Labs) was used for secondary antibodies and enhancement according to manufacturers instructions and a VectorRed kit (Vector Labs) was used for colour development. Expression of PLUNC family members was classified as positive or negative.

RESULTS

As previous studies have identified SPLUNC1, SPLUNC2 and LPLUNC1 in saliva, we first validated our antibodies by immunohistochemical analysis using sections of disease-free submandibular gland. This confirmed that the three proteins exhibited a different distribution in this tissue. LPLUNC1 was expressed within the ducts, with no staining found in either the mucous or the serous acini (Figure 1). SPLUNC1 staining was also seen within the ducts as well as in the cells of the mucous acini but the serous cells were negative (Figure 1). In contrast to the localisation of the other two proteins, SPLUNC2 was highly expressed in the serous cells of the gland but was negative in both the ducts and the mucous cells of the acini (Figure 1).

Having validated the antibodies on normal salivary gland tissue we applied them to the salivary gland TMAs. An oral squamous cell carcinoma (OSCC) was used as an orientation marker for TMA construction and was negative for all three PLUNC proteins.

The vast majority of the salivary gland tumours were negative for all three PLUNC proteins assessed. No immunoreactivity was seen in any cases of ACC, CEPA, PLGA, PA, AcCC or MC.

In mucoepidermoid carcinomas, LPLUNC1 (Figure 2A) displayed strong cytoplasmic positivity in the mucous cells and the mucin plugs (Figure 2B). Intermediate cells were moderately positive for LPLUNC1 (Figure 2C) and SPLUNC2 (Figure 2D). Epidermoid (Figure 2E) and clear cells (Figure 2F) within these samples were completely negative in all cases. No positive staining for SPLUNC1 was found in any case of MEC. All 10 cases exhibited a similar expression pattern. There were no correlations between PLUNC expression and tumour grade, size, age, gender or site.

The single case of PC studied showed cytoplasmic positivity for all three antibodies but staining intensity varied. LPLUNC1 showed strong cytoplasmic immunoreactivity in the mucous, cuboidal, columnar and clear cells and was also expressed in the luminal mucin (Figure 3A). SPLUNC1 showed strong immunoreactivity in the mucous, cuboidal and columnar cells (Figure 3B) while SPLUNC2 exhibited moderate positivity in these cells (Figure 3C).

DISCUSSION

This study examined the expression of three members of the PLUNC family of proteins in a range of salivary gland tumours in order to determine their potential role as diagnostic and/or prognostic tools. We focused on these particular PLUNC proteins as it has previously

been reported that they are secreted into saliva (Vitorino et al, 2004; Ramachandran et al, 2006; Guo et al, 2006) and their corresponding mRNAs have been identified in salivary glands (Bingle and Bingle, 2000; Bingle and Craven, 2002; Geetha et al, 2005). We showed that in normal submandibular gland the three proteins were readily detectable. Interestingly they also exhibited a different expression pattern where at least one family member was found in each of the three cellular compartments, namely ducts (LPLUNC1, SPLUNC1) or mucous (SPLUNC1) and serous acini (SPLUNC2).

Surprisingly only 2 of the 8 types of salivary gland tumours on our TMA demonstrated immunopositivity for any of the PLUNC family members studied. The mucin or mucous cells detected in the MECs and our single PC case were positive showing a strong relation between PLUNC expression and mucus-secreting salivary gland tumours. We have made a similar observation for SPLUNC1 in lung cancer (Bingle et al, 2005). The lack of immunoreactivity for PLUNC proteins in ACC, AcCC, CEPA, PA, PLGA and MC might be related to their histogenesis as these tumours do not produce or secrete mucin.

Papillary cystadenocarcinoma is a rare malignant neoplasm of the salivary gland mainly located in the parotid and intra-oral minor salivary glands (Harimaya et al, 2006). They are characterized by cystic structures lined by cuboidal, columnar and mucus-secreting cells and arranged in a papillary architecture (Nakagawa et al, 2002). The differential diagnosis includes papillary cystic variant of AcCC, low-grade MEC, salivary duct carcinoma, and PLGA (Foss et al, 1996). PC can usually be distinguished from these tumours using traditional stains such as H&E and PAS. However, some equivocal PC cases remain as diagnostic challenges even for senior pathologists. Slootweg (1993) reported the importance of correctly differentiating papillary PLGAs from PC in the oral cavity. Klijanienko and Vielh (1998) retrospectively compared fine needle aspiration samples from 12 PLGAs and 5 PC, both tumours showed variable proportions of malignant cells, occasionally arranged in pseudopapillary formations, whilst some PLGAs presented hyaline globules. In our study PC was immunoreactive for some PLUNC family members whereas the PLGA and AcCC cases were uniformly negative. Thus PLUNCs could aid in distinguishing PLGA and the papillary variant of AcCC from PC.

Metastases from prostate, gastrointestinal and thyroid tumors can affect the head and neck region and may also show papillary architecture. Established immunomarkers, such as PSA and thyroglobulin, could be useful in distinguishing PC from metastatic lesions but as expression of PLUNCs is limited to the lung, airways and oral cavity (Bingle and Bingle, 2000; Bingle and Craven, 2002; Bingle et al, 2004) they have the potential to give improved distinction.

MEC is the most common malignant salivary gland tumour (Speight and Barrett, 2002). Histopathologically, they are graded as low, intermediate or high-grade malignancy. H&E and other staining methods, notably, PAS and mucicarmine, are used to make a full diagnosis and grade lesions appropriately. High-grade lesions show a small number of mucous cells, which are difficult to recognize using these traditional staining methods. Mucous cells of the MEC studied were strongly positive for LPLUNC1 indicating a potential diagnostic use for this antibody in equivocal cases.

Another interesting finding was the immunoreactivity of some intermediate cells for LPLUNC1 and SPLUNC2. Based on these findings we can hypothesize that intermediate cells that are positive for PLUNC may have the potential to become mucous cells but further studies are needed to confirm this.

In conclusion, the majority of salivary gland tumours did not show immunoreactivity for PLUNC proteins but mucous cells present in MEC and PC did show intense positivity.

These immunomarkers might be used as an additional diagnostic tool to identify mucous cells in some equivocal MEC or PC cases.

Acknowledgments

This work was supported by the British Lung Foundation and the Wellcome Trust.

REFERENCES

- Ball WD, Mirels L, Hand AR. Psp and Smgb: a model for developmental and functional regulation in the rat major salivary glands. *Biochem. Soc. Trans.* 2003; 31:777–780. [PubMed: 12887304]
- Barnes, L.; Eveson, JW.; Reichart, P.; Sidransky, D. Pathology and Genetics of Head and Neck Tumours. IARC Press; Lyon: 2005. World Health Organization Classification of Tumours; p. 210
- Bingle CD, Bingle L. Characterisation of the human plunc gene, a gene product with an upper airways and nasopharyngeal restricted expression pattern. *Biochem. Biophys. Acta.* 2000; 1493:363–367. [PubMed: 11018263]
- Bingle CD, Craven CJ. PLUNC: a novel family of candidate host defence proteins expressed in the upper airways and nasopharynx. *Hum. Mol. Genet.* 2002; 11:937–943. [PubMed: 11971875]
- Bingle CD, LeClair EE, Havard S, Bingle L, Gillingham P, Craven CJ. Phylogenetic and evolutionary analysis of the PLUNC gene family. *Protein Science.* 2004; 13:422–430. [PubMed: 14739326]
- Bingle L, Cross SS, High AS, Wallace WA, Devine DA, Havard S, Campos MA, Bingle CD. SPLUNC1 (PLUNC) is expressed in glandular tissues of the respiratory tract and in cancers with a glandular phenotype. *J. Pathol.* 2005; 205:491–497. [PubMed: 15685591]
- Egland KA, Vincent JJ, Strausberg R, Lee B, Pastan I. Discovery of the breast cancer gene BASE using a molecular approach to enrich for genes encoding membrane and secreted proteins. *PNAS.* 2003; 100:1099–1104. [PubMed: 12538848]
- Foss RD, Ellis GL, Auclair PL. Salivary gland cystadenocarcinomas: a clinicopathologic study of 57 cases. *Am. J. Surg. Pathol.* 1996; 20:1440–1447. [PubMed: 8944036]
- Geetha C, Venkatesh SG, Bingle L, Bingle CD, Gorr SU. Design and validation of anti-inflammatory peptides from human Parotid Secretory Protein. *J. Dent. Res.* 2005; 84:149–153. [PubMed: 15668332]
- Guo T, Rudnick PA, Wang W, Lee CS, Devoe DL, Balgley BM. Characterization of the human salivary proteome by capillary isoelectric focusing/nanoreversed-phase liquid chromatography coupled with ESI-tandem MS. *J. Proteome Res.* 2006; 5:1469–1478. [PubMed: 16739998]
- Harimaya A, Somekawa Y, Sasaki M, Ohuchi T. Cystadenocarcinoma (papillary cystadenocarcinoma) of the submandibular gland. *J. Laryngol. Otol.* 2006; 120:1077–1080. [PubMed: 16995970]
- Iwao K, Watanabe T, Fujiwara Y, Takami K, Kodama K, Higashiyama M, Yokouchi H, Ozaki K, Monden M, Tanigami A. Isolation of a novel human lung-specific gene, LUNX, a potential molecular marker for detection of micrometastasis in non-small-cell lung cancer. *Int. J. Cancer.* 2001; 91:433–437. [PubMed: 11251963]
- Jones AV, Franklin CD. An analysis of oral and maxillofacial pathology found in adults over a 30-year period. *J. Oral. Pathol. Med.* 2006; 35:392–401. [PubMed: 16827841]
- Klijanienko J, Vielh P. Salivary carcinomas with papillae: cytology and histology analysis of polymorphous low-grade adenocarcinoma and papillary cystadenocarcinoma. *Diagn. Cytopathol.* 1998; 19:244–249. [PubMed: 9784985]
- Lemaire F, Millon R, Young J, Cromer A, Wasylyk C, Schultz I, Muller D, Marchal P, Zhao C, Melle D, Bracco L, Abecassis J, Wasylyk B. Differential expression profiling of head and neck squamous cell carcinoma (HNSCC). *Br. J. Cancer.* 2003; 89:1940–1949. [PubMed: 14612907]
- Mirels L, Ball WD. Neonatal rat submandibular gland protein SMG-A and parotid secretory protein are alternatively regulated members of a salivary protein multigene family. *J. Biol. Chem.* 1992; 267:2679–2687. [PubMed: 1370829]
- Mitas M, Cole DJ, Hoover L, Fraig MM, Mikhitarian K, Block MI, Hoffman BJ, Hawes RH, Gillanders WE, Wallace MB. Real-time reverse transcription-PCR detects KS1/4 mRNA in

- mediastinal lymph nodes from patients with non-small cell lung cancer. *Clin. Chem.* 2003a; 49:312–315. [PubMed: 12560358]
- Mitas M, Hoover L, Silvestri G, Reed C, Green M, Turrisi AT, Sherman C, Mikhitarian K, Cole DJ, Block MI, Gillanders WE. Lunx is a superior molecular marker for detection of non-small lung cell cancer in peripheral blood. *J. Mol. Diagn.* 2003b; 5:237–242. [PubMed: 14573783]
- Nakagawa T, Hattori K, Iwata N, Tsujimura T. Papillary cystadenocarcinoma arising from minor salivary glands in the anterior portion of the tongue: a case report. *Auris. Nasus. Larynx.* 2002; 29:87–90. [PubMed: 11772498]
- Owerbach D, Hjorth JP. Inheritance of a parotid secretory protein in mice and its use in determining salivary amylase quantitative variants. *Genetics.* 1980; 95:129–141. [PubMed: 6159249]
- Ramachandran P, Boonthung P, Xie Y, Sondej M, Wong DT, Loo JA. Identification of N-linked glycoproteins in human saliva by glycoprotein capture and mass spectrometry. *J. Proteome Res.* 2006; 5:1493–1503. [PubMed: 16740002]
- Slootweg PJ. Low-grade adenocarcinoma of the oral cavity: polymorphous or papillary? *J. Oral Pathol. Med.* 1993; 22:327–330. [PubMed: 8229870]
- Speight PM, Barrett AW. Salivary gland tumours. *Oral Dis.* 2002; 8:229–240. [PubMed: 12363107]
- Vargas PA, Gerhard R, Araújo Filho VJ, de Castro IV. Salivary gland tumors in a Brazilian population: a retrospective study of 124 cases. *Rev. Hosp. Clín. Fac. Med. S. Paulo.* 2002; 57:271–276.
- Vargas PA, Torres-Rendon A, Speight PM. DNA ploidy analysis in salivary gland tumours by image cytometry. *J. Oral. Pathol. Med.* 2007; 36(6):371–376. In press. [PubMed: 17559500]
- Vitorino R, Lobo MJ, Ferrer-Correia AJ, Dubin JR, Tomer KB, Domingues PM, Amado FM. Identification of human whole saliva protein components using proteomics. *Proteomics.* 2004; 4:1109–1115. [PubMed: 15048992]
- Weston WM, LeClair EE, Trzyna W, McHugh KM, Nugent P, Lafferty CM, Ma L, Tuan RS, Greene RM. Differential display identification of plunc, a novel gene expressed in embryonic palate, nasal epithelium, and adult lung. *J. Biol. Chem.* 1999; 274:13698–13703. [PubMed: 10224143]
- Zhang B, Nie X, Xiao B, Xiang J, Shen S, Gong J, Zhou M, Zhu S, Zhou J, Qian J, Lu H, He X, Li X, Hu G, Li G. Identification of tissue-specific genes in nasopharyngeal epithelial tissue and differentially expressed genes in nasopharyngeal carcinoma by suppression subtractive hybridization and cDNA microarray. *Genes Chromosomes Cancer.* 2003; 38:80–90. [PubMed: 12874788]




Figure 1. The expression of PLUNC proteins in normal, submandibular gland tissue. LPLUNC1 was localised to the cells of the ducts (LP1), SPLUNC1 was localised to the mucous acini and ducts (SP1) and SPLUNC2 was highly expressed in the serous cells of the acini (SP2). A negative section is included for comparison (NEG). The original magnification for all sections was 400.




Figure 2. Photomicrographs of mucoepidermoid carcinomas displaying cytoplasmic immunoreactivity for PLUNC family members. MEC exhibiting strong cytoplasmic positivity in the mucous cells for LPLUNC1 (A) (original magnification was x200). (B) Mucin plugs within cystic neoplastic areas positive for LPLUNC1 (original magnification x200). Intermediate cells showing moderate cytoplasmic reaction for LPLUNC1 (C) and SPLUNC2 (D) (original magnification for both LPLUNC1 and SPLUNC2 was x400). Epidermoid (E) and clear cells (F) were negative in all cases for LPLUNC1 (original magnification x400).




Figure 3. Photomicrographs showing immunoreactivity for PLUNC family members in one case of papillary cystadenocarcinoma. (A) Intraluminal papillary tumour cells showing a strong cytoplasmic positivity for LPLUNC1 in the luminal mucin, mucous, cuboidal, columnar and clear cells (original magnification x400). (B) Intense cytoplasmic immunoreactivity was seen in the mucous, cuboidal and columnar cells with SPLUNC1 antibody (original magnification x400) whilst only moderate cytoplasmic positivity was seen in these cells with the SPLUNC2 antibody (C, original magnification x400).

Table 1

Clinical data and histopathological types of the eight salivary gland tumours (n=64).

Tumours	number of cases	Gender M/F	Age (years) mean(range)	Site	Size (cm) mean(range)
ACC	13	4M/9F	57.1 (37-84)	2SMG, 11 IOSG	1.8 (0.8-3.0)
MEC	10	2M/8F	43.4 (16-73)	2PG, 1SMG, 7 IOSG	2.4 (0.9-6.5)
PA	10	4M/6F	44.5 (22-68)	7 PG, 3 IOSG	3.1 (1.5-5.5)
CEPA	10	3M/7F	54.2 (42-67)	6 PG, 1 SMG, 3 IOSG	3.7 (1.3-9.0)
PLGA	10	4M/6F	71.6 (39-91)	10 IOSG	2.6 (1.4-3.8)
AcCC	9	4M/5F	36.8 (17-77)	8 PG, 1 SMG	3.1 (1.4-8.5)
MC	1	M	83	1 IOSG	4.0
PC	1	M	74	1 IOSG	na
Total	64	23M/41F		23PG; 5SMG; 36IOSG	

na: not available; M: male; F: female; PG: parotid gland; SMG: submandibular gland; IOSG: intraoral salivary gland; for abbreviation of tumour types see text; ACC: 9 cases were cribriform/tubular, 4 solid variant, and 3 cases recurred (cribriform variant). MEC: 7 low-grade, 2 intermediate-grade and one high-grade cases. AcCC: 1 case metastasised to cervical lymph node. MC: recurrent case.