



Published in final edited form as:

Spine (Phila Pa 1976). 2010 April 1; 35(7): 803–811. doi:10.1097/BRS.0b013e3181bc9454.

Variation in eligibility criteria from studies of radiculopathy due to a herniated disc and of neurogenic claudication due to lumbar spinal stenosis: A structured literature review

S. Genevay, MD^{1,3}, S.J. Atlas, MD MPH², and J.N. Katz, MD MSc³

¹ Division of Rheumatology, University Hospitals of Geneva, Switzerland ² General Medicine Division, Massachusetts General Hospital, Boston MA, USA ³ Department of Orthopaedic Surgery and Division of Rheumatology, Immunology and Allergy, Brigham and Women's Hospital, Harvard Medical, Boston MA, USA

Abstract

Study Design—A structured literature review.

Summary of the Background Data—Widely recognized classification criteria for rheumatologic disorders have resulted in well-defined patient populations for clinical investigation.

Objectives—We sought to determine whether similar criteria were needed for back pain disorders by examining variability in eligibility criteria in published studies

Methods—Studies involving radiculopathy due to lumbar herniated disc (HD) and for neurogenic claudication due to lumbar spinal stenosis (LSS) were identified. Randomized controlled trials published between January 1, 2006 and October 1, 2008 in select peer reviewed journals were retrieved, their eligibility criteria were identified and categorized.

Results—Twelve eligible HD studies were identified. Thirteen unique categories of eligibility criteria were identified with a mean of 3.9 (+/-2.0) and a range from 0 to 8 categories per study. More categories were present for studies that included nonsurgical (5.6 +/- 2.5) treatment for studies with only surgical treatment (2.6 +/- 1.7) p= 0.04). Seven LSS studies met eligibility criteria, and 9 unique categories were identified. A mean of 5.0 (+/-2.2) categories with a range from 2 to 7 was used per study.

Conclusion—Wide variation in the number and type of eligibility criteria from randomized clinical trials of well defined back pain syndromes was identified. These results support the need for developing and disseminating international classification criteria for these clinical conditions.

Keywords

Radiculopathy; Herniated disc; Neurogenic claudication; Lumbar spine stenosis; Eligibility criteria

INTRODUCTION

The last two decades have witnessed an explosion of controlled trials involving a wide range of low back pain conditions. Prior to this, there were few high quality randomized controlled trials of common back pain treatments. Well conducted meta-analyses of low back pain treatments routinely concluded that reliable data were too sparse to draw firm conclusions¹⁻⁴.

Immense improvements in the quality and quantity of research have taken place in recent years. Major randomized controlled trials have been conducted and published in highly rated peer reviewed journals⁵⁻⁷. It is tempting to imagine that high quality meta-analyses with well grounded, clear conclusions for many low back conditions and treatments are just a few years off.

In contrast with many rheumatological conditions, such as rheumatoid arthritis⁸ or complex regional pain syndrome⁹, the literature on low back pain disorders has developed without widely recognized diagnostic and classification criteria. The development of internationally recognized classification criteria for rheumatological disorders has contributed to their pathophysiologic understanding and the development of new therapeutic targets¹⁰. Classification criteria are also critical for conducting clinical trials, epidemiologic studies and other clinical investigations. The use of established classification criteria ensures recruitment of a more clinically homogeneous sample of subjects¹⁰. Imprecise definitions of specific spinal conditions may have clinical consequences as physicians attempt to apply the results from trials with potentially different patient populations to their own patients. To document and address the potential impact of different case definitions, we examined variation in eligibility criteria in recently published studies for two seemingly well defined back pain syndromes, radiculopathy arising from a lumbar herniated disc (HD) and neurogenic claudication due to lumbar spinal stenosis (LSS). Our goals were to document variability in classification of the same entities across studies and to identify possible classification criteria for future evaluation.

METHODS

We performed a structured literature review using PubMed to identify potential clinical studies of radiculopathy due to a lumbar herniated disc and neurogenic claudication due to lumbar spinal stenosis. Two independent literature searches were conducted by one author (SG). Search terms on radiculopathy due to HD included: sciatica, lumbar radiculopathy, and lumbar disc herniation. Search term on neurogenic claudication due to LSS included: degenerative lumbar stenosis, lumbar spinal stenosis, neurogenic claudication and intermittent claudication. We focused exclusively on randomized controlled trials comparing two treatments because strict eligibility criteria are particularly important for such studies. Studies that included mixed pathological conditions were excluded. For both conditions, additional search criteria included: 1) age 19 or older and 2) English literature. As methodology in back pain literature has been said to have recently improved, the search was also restricted to manuscript published between January 1, 2006 and October 1 2008. We augmented the electronic search by including studies from the reference lists of retrieved papers. To specifically identify high quality clinical trials, the literature search was limited to publications from journals with an impact factor of at least 1. To avoid duplication we excluded secondary analyses of previously published studies if the parent study met eligibility criteria for this analysis.

Data abstraction from eligible studies involved reviewing the Methods sections to identify inclusion and exclusion criteria. Eligibility criteria were then extracted and classified into one of the following four groups: a) demographic characteristics; b) symptoms; c) physical examination signs; d) and investigative procedures. Categories were identified and classified

within each of these four broad groups. Eligibility criteria that were not directly related to the diagnosis of the underlying clinical condition but rather related to the design of the study (e.g. symptom duration, failure of previous treatments, history of previous surgery, etc.), or to the treatment provided (e.g. allergies to studied medication, specific contraindications to the procedure, etc.) were excluded. The entire selection process was reviewed with a second author (JK). Discrepancies and ambiguities were discussed and consensus was achieved.

Analyses

Eligible studies were classified into 3 treatment categories: surgical treatment only, surgical versus nonsurgical treatment, and nonsurgical treatment only (Table 1 and 2). For each clinical condition, the number of studies (mean, median, and range) reporting each identified criteria were reported. Reported criteria across studies were also compared. Unpaired t-tests for unequal variance were used to assess differences between groups.

RESULTS

Radiculopathy due to Herniated Disc

The literature search on radiculopathy due to HD retrieved 56 references. One additional study was identified by checking reference lists of the retrieved studies. After reviewing the abstracts of these 57 studies, 43 were excluded because they did not meet inclusion criteria. Two more were discarded after reviewing the entire article. Discrepancies between SG and JK arose on 3 occasions and were resolved by reanalyzing the Methods section of the specific study. The Methods section of each of the remaining 12 studies underwent review to identify eligibility criteria.

Thirteen unique categories were identified that accounted for all identified diagnostic related criteria (Table 1). No category was assigned to the demographic characteristic group. Five related to clinical symptoms: 1) presence of accompanying back pain; 2) leg pain greater than back pain; 3) characteristic of leg pain (radicular/sciatic/dermatome distribution); 4) pattern of pain radiation (anterior thigh, below the knee) and 5) pain on Valsalva manoeuvre. Three categories referred to clinical examination: 1) nerve root irritation sign (straight leg raise test [L5 or S1] or femoral stretch test [L3 or L4], contralateral straight leg raise test); 2) presence of neurological deficit; and 3) neurological deficit corresponding to pain pattern. Four categories related to investigations: 1) herniated disc on imaging; 2) herniated disc on imaging corresponding (level + side) with clinical observation; 3) specification of the kind of HD included/excluded (protrusion, extrusion, sequestration, foraminal, etc.) and 4) electromyographic findings. The final category, specification of nerve roots included, could not be specifically attributed to a specific group.

The mean number of diagnostic categories identified per study was 3.9 (standard deviation 2.0, median 4, range 0 to 8). Trials comparing surgery and nonsurgical treatments used more eligibility categories (5.6 \pm 2.5) than trials only comparing surgical techniques (2.8 \pm 1.7). Trials only comparing nonsurgical treatments had a mean of 4.3 (\pm 1.9) categories cited per article. Studies comparing only surgical techniques used fewer criteria than studies involving at least one nonsurgical treatment (2.8 \pm 1.7 vs. 4.8 \pm 1.9, $p=0.04$).

No single diagnostic eligibility category was used in all the studies (Table 3). The most commonly used categories were “radicular pattern of radiation” and “HD on imaging” (each 13/19, 68%). Congruence between pain pattern of radiation and HD on imaging were less commonly used (each 6/19, 32%), while congruence between pain pattern and neurological deficit were each used only twice (11%). There was additional variation in the definition used within a category. For example when the category “specification of nerve roots included” was used the number of defined nerve roots ranged from 1 11 (S1) to 6 12 (from L1 to S1). The

definition of the straight leg raise test varied in 2 ways: a) the kind of triggered pain (i.e. any leg pain versus radicular pain) and b) the degree of minimal/maximal angle for the test to be positive.

Neurogenic claudication due to Spinal Stenosis

The search for studies of neurogenic claudication due to LSS retrieved 20 citations. Eleven were excluded after reviewing the abstract, and an additional 2 were excluded after reviewing the entire paper. The remaining 7 eligible studies underwent review to identify eligibility criteria.

Nine unique categories were identified that accounted for all identified diagnostic related criteria (Table 2). One category was demographic: age limitation (usually older than 50). Three categories related to clinical symptoms: 1) presence of accompanying back/buttock/lumbopelvic pain; 2) characteristics of radiating pain (neurogenic claudication / radicular pain); 3) definition of neurogenic claudication (symptoms restricted to pain or include fatigue and sensory loss: factors that made it worse; factors that ease the symptoms). No category was assigned to the clinical examination group. Four categories related to investigations: 1) radiological confirmation of stenosis; 2) explicit description of radiological stenosis (e.g. minimal sagittal diameter); 3) correlation between symptoms and level of radiological stenosis; 4) specific inclusion or exclusion of some other radiological findings (e.g. degenerative spondylolisthesis, instability). The final category could not be specifically attributed to a specific group: exclusion of other spinal (congenital stenosis) or non spinal problems (e.g. vascular, articular or neurogenic).

The mean number of criteria per study was 5.0 (+/-2.2), (median 6, range 2 to 7). No differences were noticed between surgical, nonsurgical and mixed studies (means between 5 and 5.5). On average more categories were used in trials of neurogenic claudication (mean of 5 criteria used per paper of the 9 possible criteria; 57%) compared to trials on radiculopathy (mean of 3.9 criteria of a possible 13; 28%), $p < 0.02$.

The presence of imaging findings consistent with lumbar stenosis was used in all studies, while the presence of neurogenic claudication was used in 5/7 (71%) studies. The definition of neurogenic claudication varied considerably across studies (e.g. "fatigue or loss of sensation in the lower limbs aggravated by walking" 13 and "sitting as a better position for symptom severity than standing or walking" 14.) Some studies included pain located in the buttock area as a possible radiation pain likely to have claudicating pattern 13-14. This allows the inclusion of patients who would be ineligible in studies adhering to the classical definition of lumbar pain (i.e. pain located between the 12th costal margin and the buttock folds 15). Detailed radiological definition of LSS was provided in 4/7 (57%) studies. In addition, the definition of radiological LSS varied among studies. Some mentioned dural sac compression¹⁶, while others referred to precise measurement of the lumbar canal 17. Congruence between radiating pain pattern and the radiological level of stenosis was mentioned once. Of note, some radiological findings such as degenerative spondylolisthesis are mentioned in the list of exclusion criteria 18 in some studies while they are specifically mentioned as being included in others 13.

DISCUSSION

This study documented a high level of variation in diagnostic eligibility criteria for randomized controlled trials involving treatments for radiculopathy due to HD and for lumbar stenosis due to LSS. There was wide variation in the number of criteria used within each study and few criteria were used consistently across studies. These findings support our hypothesis that even for spine conditions that are thought to be well characterized, there is no single, standardized case definition used in the research literature. In addition, differences were found within

defined categories, such as the number of nerve roots included or the definition used for the straight leg raise test in the case of radiculopathy or the definition of radiological stenosis for neurogenic claudication.

Some of the categories that were created from our analysis such as “pattern of radiation” (e.g. anterior thigh, below the knee) and “specification of the nerve root included” may appear similar; the precision added by the second proposition was however considered to be crucial, justifying the separation, especially in reference to the L4 nerve root. Pain from the L4 nerve root may radiate below the knee, but the clinical assessment of L4 nerve root requires quadriceps testing for deficit and the femoral stretch test, rather than the straight leg raise test. Some discrepancies were noted in studies where eligibility criteria included pain below the knee and clinical examination related to L5 and S1 nerve root without mentioning the appropriate neurological testing for L4 nerve root ¹⁹.

In the absence of widely accepted diagnostic eligibility criteria, each group of authors devised their own construct. This makes generalizing findings across studies and to routine clinical practice a challenge. At a time when other musculoskeletal diseases are considering revision of well established sets of criteria ^{20,21}, the absence of diagnostic and/or classification criteria in the field of low back pain should be considered a major focus for international organizations and clinical investigators. We suggest this process should differentiate among several needs. For epidemiological studies, broad criteria are needed and we suggest they should exclusively rely on symptoms that can be elicited from patients as part of survey research. For clinical studies however, the focus should be put on achieving more homogenous populations. For studies in the primary care setting involving patients with acute symptoms, the addition of clinical findings would be appropriate. Finally, for most clinical trials involving treatments, confirmatory investigative findings from imaging studies or electrodiagnostic tests are needed.

A number of important limitations may limit the interpretation of our results. A broader literature search criteria would have likely retrieved more diagnostic eligibility criteria. However it would also have included a large proportion of studies without any eligibility criteria ²². We believe that finding wide variation in diagnostic eligibility criteria in recent randomized controlled trials published in high quality journals (defined as journals with higher impact factor) strengthens our conclusions. We also could have examined more spine conditions to generalize our findings across a broader range of disorders. However, focusing on two common and presumably well defined conditions, radiculopathy due to a herniated disc and neurogenic claudication due to lumbar spinal stenosis, supports the need for similar efforts for non-specific low back pain disorders as well. Some discrepancies were noted between the two reviewers in term of article selection (3 times) and in term of number of eligibility criteria (5 times). The vast majority were related to imprecision within the abstract. These discrepancies were generally resolved after considering the full text of the article. However, in some cases, imprecise or inconsistent reporting in the Methods section required subjective decisions by the reviewers. Excluding these few criteria would not alter our general findings.

In conclusion, we identified wide variation in diagnostic eligibility criteria for studies of radiculopathy due to HD and for neurogenic claudication due to LSS. These findings support efforts to convene a multidisciplinary, international effort to propose validated classification criteria for these conditions.

Key points (3-5)

- As distinct from many other musculoskeletal diseases, no widely accepted classification criteria have been developed for low back pain conditions.

- A high degree of variability in eligibility criteria was found in studies of radiculopathy due to disc herniation and of neurogenic claudication due to spinal stenosis.
- This lack of uniformity will impair the quality of knowledge of these conditions by compromising the capacity to compare studies.
- For both conditions, there is an urgent need for validated classification criteria.
- Criteria retrieved through this structured literature search could be used as a basis for the development of sets of criteria.

Acknowledgments

Pr J.N. Katz is support by NIH/NIAMS P60 AR 47782 and NIH/NIAMS K24 02123. Dr. Atlas is supported in part by a grant from (P60 AR048094) to the Multidisciplinary Clinical Research Center in Musculoskeletal Diseases from the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS).

References

1. Brosseau L, Milne S, Robinson V, et al. Efficacy of the transcutaneous electrical nerve stimulation for the treatment of chronic low back pain: a meta-analysis. *Spine* 2002;27:596–603. [PubMed: 11884907]
2. Koes BW, Bouter LM, van der Heijden GJ. Methodological quality of randomized clinical trials on treatment efficacy in low back pain. *Spine* 1995;20:228–35. [PubMed: 7716630]
3. Niemisto L, Kalso E, Malmivaara A, et al. Radiofrequency denervation for neck and back pain: a systematic review within the framework of the cochrane collaboration back review group. *Spine* 2003;28:1877–88. [PubMed: 12923479]
4. van Tulder MW, Cherkin DC, Berman B, et al. The effectiveness of acupuncture in the management of acute and chronic low back pain. A systematic review within the framework of the Cochrane Collaboration Back Review Group. *Spine* 1999;24:1113–23. [PubMed: 10361661]
5. Fairbank J, Frost H, Wilson-MacDonald J, et al. Randomised controlled trial to compare surgical stabilisation of the lumbar spine with an intensive rehabilitation programme for patients with chronic low back pain: the MRC spine stabilisation trial. *BMJ* 2005;330:1233. [PubMed: 15911537]
6. Hancock MJ, Maher CG, Latimer J, et al. Assessment of diclofenac or spinal manipulative therapy, or both, in addition to recommended first-line treatment for acute low back pain: a randomised controlled trial. *Lancet* 2007;370:1638–43. [PubMed: 17993364]
7. Peul WC, van Houwelingen HC, van den Hout WB, et al. Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med* 2007;356:2245–56. [PubMed: 17538084]
8. Arnett FC, Edworthy SM, Bloch DA, et al. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988;31:315–24. [PubMed: 3358796]
9. Stanton-Hicks M, Janig W, Hassenbusch S, et al. Reflex sympathetic dystrophy: changing concepts and taxonomy. *Pain* 1995;63:127–33. [PubMed: 8577483]
10. Singh JA, Solomon DH, Dougados M, et al. Development of classification and response criteria for rheumatic diseases. *Arthritis Rheum* 2006;55:348–52. [PubMed: 16739201]
11. Ackerman WE 3rd, Ahmad M. The efficacy of lumbar epidural steroid injections in patients with lumbar disc herniations. *Anesth Analg* 2007;104:1217–22. tables of contents. [PubMed: 17456677]
12. Ruetten S, Komp M, Merk H, et al. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study. *Spine* 2008;33:931–9. [PubMed: 18427312]
13. Malmivaara A, Slati P, Heliovaara M, et al. Surgical or nonoperative treatment for lumbar spinal stenosis? A randomized controlled trial. *Spine* 2007;32:1–8. [PubMed: 17202885]
14. Whitman JM, Flynn TW, Childs JD, et al. A comparison between two physical therapy treatment programs for patients with lumbar spinal stenosis: a randomized clinical trial. *Spine* 2006;31:2541–9. [PubMed: 17047542]

15. Dionne CE, Dunn KM, Croft PR, et al. A consensus approach toward the standardization of back pain definitions for use in prevalence studies. *Spine* 2008;33:95–103. [PubMed: 18165754]
16. Cavusoglu H, Kaya RA, Turkmenoglu ON, et al. Midterm outcome after unilateral approach for bilateral decompression of lumbar spinal stenosis: 5-year prospective study. *Eur Spine J* 2007;16:2133–42. [PubMed: 17712577]
17. Tafazal SI, Ng L, Sell P. Randomised placebo-controlled trial on the effectiveness of nasal salmon calcitonin in the treatment of lumbar spinal stenosis. *Eur Spine J* 2007;16:207–12. [PubMed: 16865379]
18. Weinstein JN, Tosteson TD, Lurie JD, et al. Surgical versus nonsurgical therapy for lumbar spinal stenosis. *N Engl J Med* 2008;358:794–810. [PubMed: 18287602]
19. Khoromi S, Cui L, Nackers L, et al. Morphine, nortriptyline and their combination vs. placebo in patients with chronic lumbar root pain. *Pain* 2007;130:66–75. [PubMed: 17182183]
20. Liao KP, Batra KL, Chibnik L, et al. Anti-cyclic citrullinated peptide revised criteria for the classification of rheumatoid arthritis. *Ann Rheum Dis* 2008;67:1557–61. [PubMed: 18234714]
21. Taylor W, Gladman D, Helliwell P, et al. Classification criteria for psoriatic arthritis: development of new criteria from a large international study. *Arthritis Rheum* 2006;54:2665–73. [PubMed: 16871531]
22. Alfonso, M.; Diaz De Rada, P.; Villas, C. The meaning of lumbar spinal stenosis. A critical review of literature. *Spineweek Geneva; Switzerland*: 2008.
23. Ryang YM, Oertel MF, Mayfrank L, et al. Standard open microdiscectomy versus minimal access trocar microdiscectomy: results of a prospective randomized study. *Neurosurgery* 2008;62:174–81. discussion 81-2. [PubMed: 18300905]
24. Brock M, Kunkel P, Papavero L. Lumbar microdiscectomy: subperiosteal versus transmuscular approach and influence on the early postoperative analgesic consumption. *Eur Spine J* 2008;17:518–22. [PubMed: 18224352]
25. Righesso O, Falavigna A, Avanzi O. Comparison of open discectomy with microendoscopic discectomy in lumbar disc herniations: results of a randomized controlled trial. *Neurosurgery* 2007;61:545–9. discussion 9. [PubMed: 17881967]
26. Ozer AF, Oktenoglu T, Sasani M, et al. Preserving the ligamentum flavum in lumbar discectomy: a new technique that prevents scar tissue formation in the first 6 months postsurgery. *Neurosurgery* 2006;59:ONS126–33. discussion ONS-33. [PubMed: 16888542]
27. Arts MP, Peul WC, Brand R, et al. Cost-effectiveness of microendoscopic discectomy versus conventional open discectomy in the treatment of lumbar disc herniation: a prospective randomised controlled trial [ISRCTN51857546]. *BMC Musculoskelet Disord* 2006;7:42. [PubMed: 16696861]
28. Aminmansour B, Khalili HA, Ahmadi J, et al. Effect of high-dose intravenous dexamethasone on postlumbar discectomy pain. *Spine* 2006;31:2415–7. [PubMed: 17023848]
29. Hoogland T, Schubert M, Miklitz B, et al. Transforaminal posterolateral endoscopic discectomy with or without the combination of a low-dose chymopapain: a prospective randomized study in 280 consecutive cases. *Spine* 2006;31:E890–7. [PubMed: 17108817]
30. Weinstein JN, Tosteson TD, Lurie JD, et al. Surgical vs nonoperative treatment for lumbar disk herniation: the Spine Patient Outcomes Research Trial (SPORT): a randomized trial. *JAMA* 2006;296:2441–50. [PubMed: 17119140]
31. Osterman H, Seitsalo S, Karppinen J, et al. Effectiveness of microdiscectomy for lumbar disc herniation: a randomized controlled trial with 2 years of follow-up. *Spine* 2006;31:2409–14. [PubMed: 17023847]
32. Friedman BW, Esses D, Solorzano C, et al. A randomized placebo-controlled trial of single-dose im corticosteroid for radicular low back pain. *Spine* 2008;33:E624–9. [PubMed: 18665021]
33. Luijsterburg PA, Verhagen AP, Ostelo RW, et al. Physical therapy plus general practitioners' care versus general practitioners' care alone for sciatica: a randomised clinical trial with a 12-month follow-up. *Eur Spine J* 2008;17:509–17. [PubMed: 18172697]
34. Gallucci M, Limbucci N, Zugaro L, et al. Sciatica: treatment with intradiscal and intraforaminal injections of steroid and oxygen-ozone versus steroid only. *Radiology* 2007;242:907–13. [PubMed: 17209164]

35. Korhonen T, Karppinen J, Paimela L, et al. The treatment of disc-herniation-induced sciatica with infliximab: one-year follow-up results of FIRST II, a randomized controlled trial. *Spine* 2006;31:2759–66. [PubMed: 17108825]
36. Finckh A, Zufferey P, Schurch MA, et al. Short-term efficacy of intravenous pulse glucocorticoids in acute discogenic sciatica. A randomized controlled trial. *Spine* 2006;31:377–81. [PubMed: 16481946]
37. Hallett A, Huntley JS, Gibson JN. Foraminal stenosis and single-level degenerative disc disease: a randomized controlled trial comparing decompression with decompression and instrumented fusion. *Spine* 2007;32:1375–80. [PubMed: 17545903]
38. Yaksi A, Ozgonenel L, Ozgonenel B. The efficiency of gabapentin therapy in patients with lumbar spinal stenosis. *Spine* 2007;32:939–42. [PubMed: 17450066]

Table 2

Eligible Randomized Control Studies of Neurogenic Claudication due to Lumbar Spinal Stenosis.

Reference	Treatments	Quoted Definition from the "Materials and Methods"	Eligibility Criteria *									Total					
			1	2	3	4	5	6	7	8	9						
Studies of different surgical techniques																	
Cavuşoğlu H Eur. Spine J. 2007 16	unilateral laminotomy versus unilateral laminectomy	"(1) symptoms of neurogenic claudication referable to the lumbar spine (Claudicant or radicular symptoms brought on either by walking or by prolonged standing, relieved by sitting or the flexed position, in the absence of vascular or neuropathic pathology), (2) radiological/neuroimaging evidence of degenerative lumbar stenosis (neurologic compression by hypertrophied (infolded) ligamentum flavum, osteophytic facet joints, and annular bulging), ... (4) the absence of associated pathology such as instability, inflammation or malignancy.... Patients presenting with mild degenerative spondylolisthesis were not excluded."			+	+	+	+	+			+	+			6	
Hallett A Spine 2007 37	Foraminotomy v. foraminotomy + PLIF v. foraminotomy + TLIF	"... had both 1) single-level degenerative disc disease and 2) evidence of associated foraminal stenosis. All the patients had suffered from some backache over the preceding 5 years, yet this was not the main presenting feature... All patients complained of unilateral or bilateral leg pain with or without positive nerve root tension signs, associated muscle weakness, and/or sensory loss... Plain radiographs and MR images were obtained in all subjects to diagnose intraforaminal or extraforaminal nerve root compromise, in association with single-level degenerative disc disease. Patients were excluded if they had 1) degenerative spondylolisthesis of Grade II or greater at the level of the degenerative disc or at an adjacent level, 2) vertebral translocation in excess of 1 cm, 3) disc space narrowing of greater than 50% proximal or distal to the level of proposed fusion...."			+										+	3	
Surgery vs. nonsurgical treatment																	
Weinstein JN NEJM 2008 18	Surgery versus nonsurgical treatment	"... history of neurogenic claudication or radicular leg symptoms ... and confirmatory cross-sectional imaging showing lumbar spinal stenosis at one or more levels; Patients with degenerative spondylolisthesis ... [and] patients with lumbar instability (which was defined as translation of more than 4 mm or 10 degrees of angular motion between flexion and extension on upright lateral radiographs) were excluded."			+										+	3	
Malmivaara A Spine 2007 13	Nonsurgical versus surgery	"... back pain radiation to lower limbs or buttocks; fatigue or loss of sensation in the lower limbs aggravated by walking... [and] spinal canal narrowing, the sagittal diameter of the dural sac being less than 10 mm2, or the planimetrically assessed cross-sectional dural area being less than 75 mm2. [and] signs and symptoms corresponding to segmental radiographic level of stenosis. The following conditions did not prevent inclusion: radiographic instability of the lumbar spine, degenerative spondylolisthesis. Patients... were ineligible in the case of spinal stenosis not caused by degeneration; spondylolysis and spondylolytic spondylolisthesis; lumbar herniated disc ...; an other spinal disorder...; intermittent claudication due to atherosclerosis; severe osteoarthritis or arthritis causing dysfunction of the lower limbs; neurologic disease causing impaired function of the lower limbs...."			+										+	+	7
Studies of different nonsurgical treatments																	
Yaksi A Spine 2007 38	Gabapentin versus placebo	"... symptoms of [neurogenic intermittent claudication] ... and diagnosed with [lumbar spinal stenosis] based on radiologic studies."			+										+	2	

Reference	Treatments	Quoted Definition from the "Materials and Methods"	Eligibility Criteria*									Total		
			1	2	3	4	5	6	7	8	9			
Tafazal SI Eur. Spine J. 2007 17	Calcitonin versus placebo	"...symptoms of neurogenic claudication and MRI proven lumbar spinal stenosis were enrolled into the study ... unilateral or bilateral leg pain made worse by walking and prolonged standing and eased by resting or leaning forwards. Inclusion criteria ... Pain including back/leg [and] Standing leg discomfort [and] Weakness, Paraesthesia and absent reflex [and] Age > 50 [and] Radiographic evidence of relative/absolute stenosis- mid sagittal diameter 13 mm or less. Exclusion criteria: ... Vascular claudication... Presence of other neurologic disease including peripheral neuropathy."	+	+	+	+	+	+				+	7	
Whitman JM Spine 2006 14	2 different physical therapy	"... pain in the lumbopelvic region and lower extremities, ≥ 50 years of age, MRI findings consistent with LSS (evidence of compression of lumbar spinal nerve root(s) by degenerative lesions of the facet joint, disc, and/or ligamentum flavum), and patient rating of sitting as a better position for symptom severity than standing or walking. Exclusion: history of ... lumbar vertebral fractures other than spondylolysis or spondylolisthesis; ... signs/symptoms suggestive of potential non benign or pathologic condition as the origin of symptoms."	+	+		+	+	+				+	+	7

* Eligibility criteria are: 1-Age limitation (usually over 50). 2-Presence of accompanying back/buttock/lumbopelvic pain. 3-Characteristics of radiation (neurogenic claudication / radicular pain). 4- Explicit description of neurogenic claudication (symptoms restricted to pain or include fatigue and sensory loss; factors that made it worse; factors that ease the symptoms). 5-Radiological confirmation of stenosis. 6- Definition of radiological stenosis (e.g. minimal sagittal diameter). 7-Correlation between symptoms and level of radiological stenosis. 8-Specific inclusion or exclusion of some other radiological findings (e.g. degenerative spondylolisthesis, instability). 9-Exclusion of other spinal (congenital stenosis) or non spinal problems (e.g. vascular, articular or neurogenic).

Table 3

Classification of diagnostic categories by frequency order of citation from selected studies

Syndrome	Number of studies citing the diagnostic criterion (in decreasing order of frequency)		
	15-19 studies	5-14 studies	1-4 studies
Radiculopathy due to HD		<ul style="list-style-type: none"> • Characteristic of leg pain • Herniated disc on imaging • Nerve root irritation sign • Neurological deficit • Herniated disc on imaging corresponding (level + side) with clinical observation • Specification of nerve roots included • Precision on the kind of HD included/excluded • Pattern of radiation 	<ul style="list-style-type: none"> • Presence of accompanying back pain. • Electromyographic findings • leg pain > back pain • Positive Valsalva manoeuvre • Neurological deficit corresponding to the pain pattern
	6-7 studies	3-5 studies	1-3 studies
Neurogenic Claudication due to LSS	<ul style="list-style-type: none"> • Radiological confirmation of stenosis 	<ul style="list-style-type: none"> • Specific inclusion or exclusion of some other radiological findings • Presence of accompanying back/buttock/lumbopelvic pain • Characteristics of radiating pain • Explicit definition of neurogenic claudication • Definition of radiological stenosis • Exclusion of other spinal or non spinal problems. 	<ul style="list-style-type: none"> • Correlation between symptoms and level of radiological stenosis • Age limitation