



Published in final edited form as:

Cancer Chemother Pharmacol. 2010 May ; 66(1): 141–150. doi:10.1007/s00280-009-1144-0.

Peroxisome Proliferator-Activated Receptor γ -Dependent Activity of Indole Ring-Substituted 1,1-Bis(3'-indolyl)-1-(*p*-biphenyl) methanes in Cancer Cells

Jingjing Guo¹, Sudhakar Chintharlapalli¹, Syng-ook Lee¹, Sung Dae Cho², Ping Lei¹, Sabitha Papineni³, and Stephen Safe^{1,3}

¹Institute of Biosciences and Technology, Texas A&M University Health Science Center, Houston, TX

²Department of Oral Pathology, School of Dentistry, Institute of Oral Sciences, Chonbuk National University, Jeonbuk, South Korea

³Department of Veterinary Physiology and Pharmacology, Texas A&M University, College Station, TX

Abstract

1,1-Bis(3-indolyl)-1-(*p*-substituted phenyl)methanes (C-DIMs) substituted in the phenyl ring with a *para*-, *t*-butyl, trifluoro or phenyl (DIM-C-*p*PhC₆H₅) group activates peroxisome proliferator-activated receptor γ (PPAR γ) in several cancer cell lines. In this study, we have examined the effects of 5,5'-dihydroxy, 5,5'-dimethyl, 5,5'-dibromo, 5,5'-dinitro and 5,5'-dimethoxyindole ring substituted analogs of DIM-C-*p*PhC₆H₅ on their activity as PPAR γ agonists. Introduction of the 5,5'-dihydroxy and 5,5'-dimethyl substituents enhanced activation of PPAR γ in colon and pancreatic cancer cells. However, activation of p21 in Panc28 pancreatic cancer cells and induction of caveolin-1 and KLF4 in colon cancer cells by the cells by these C-DIMs was structure- and cell context-dependent. The results demonstrate that DIM-C-*p*PhC₆H₅ and indole ring-substituted analogs are selective PPAR γ modulators.

Keywords

C-DIMs; PPAR γ agonists; indole ring substituents

INTRODUCTION

1,1-Bis(3'-indolyl)methane (DIM) is a metabolite of the phytochemical indole-3-carbinol and DIM has been used as scaffold to synthesize a series of 1,1-bis(3'-indolyl)-1-(*p*-substituted phenyl)methanes (C-DIMs) [1–14]. These compounds are triarylmethane derivatives which differ from DIM and ring-substituted DIMs which are diarylmethanes. Initial studies showed that some C-DIMs inhibited carcinogen-induced rat mammary tumor growth and growth of various cancer cell lines [1–5;8]. The activation of several orphan nuclear receptors by a series of C-DIMs containing various *p*-substituents has also been determined and the results showed that some analogs activated PPAR γ in breast cancer cells [8]. Subsequent studies showed that one or more of the three most active compounds, namely the *p*-trifluoromethyl (DIM-C-

Correspondence should be sent to: Stephen Safe, Distinguished Professor of Toxicology, Department of Veterinary Physiology and Pharmacology, Texas A&M University, 4466 TAMU, Vet. Res. Bldg. 410, College Station, TX 77843-4466, Tel: 979-845-5988 / Fax: 979-862-4929, ssafe@cvm.tamu.edu.

pPhCF₃), *p*-*t*-butyl (DIM-C-pPh_tBu), and *p*-phenyl (DIM-C-pPhC₆H₅) analogs also activated PPAR γ in colon, pancreatic, prostate, bladder, breast, endometrial and kidney cancer cell lines [1–6;8–12]. The PPAR γ -active C-DIMs exhibit highly tissue-specific receptor-dependent activation of responses and genes. For example, these compounds induced PPAR γ -dependent p21 gene expression in Panc28 pancreatic cancer cells and caveolin-1 in colon and bladder cancer cells [1–3]. However, most other responses such as C-DIM-induced proapoptotic and growth inhibitory effects were PPAR γ -independent.

PPAR γ -active C-DIMs also induce other proapoptotic responses that are both receptor- and ER stress-independent. For example, in SW480 colon cancer cells, DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅ do not induce typical markers of ER stress and, in both SW480 and HCT116 colon cancer cells, C-DIM compounds induce expression of NAG-1 and activating transcription factor 3 (ATF3) which are proapoptotic genes and proteins [4;15]. DIM also induces NAG-1 and ATF3 in HCT116 cells; however, the mechanism of this response has not been determined. Induction of NAG-1 in HCT116 cells by PPAR γ -active C-DIMs is dependent on PI3-K-dependent activation of early growth response-1 (Egr-1) gene which in turn activates NAG-1 through interactions with a proximal Egr-1 element in the NAG-1 promoter [4]. In contrast, induction of NAG-1 by DIM-C-pPhCF₃ in LNCaP cells is MAPK-dependent [6] and suggests that induction of some proapoptotic genes such as NAG-1 are dependent on activation of kinase pathways by C-DIMs. It is clear from studies on C-DIM compounds that DIM is an excellent scaffold from which new chemotherapeutic agents can be derived. In this study, we have synthesized a series of symmetrical 5'-indole ring substituted analogs of DIM-C-pPhC₆H₅ and have investigated their cytotoxicity and PPAR γ -dependent activity in colon and pancreatic cancer cells.

METHODS AND MATERIALS

Cell lines and reagents

SW480, HT-29 and HCT-15 human colon cancer cells, Panc-1 and Panc-28 human pancreatic cancer cells were obtained from American Type Culture Collection (Manassas, VA). SW480, HT-29, Panc-1 and Panc-28 cells were maintained in Dulbecco's modified/Ham's F-12 (Sigma Aldrich, St Louis, MO) with phenol red supplemented with 0.22% sodium bicarbonate, 0.011% sodium pyruvate, 5% fetal bovine serum and 10 ml/l 100 \times antibiotic anti-mycotic solution (Sigma). HCT-15 cells were maintained in RPMI 1640 (Sigma) supplemented with 0.22% sodium bicarbonate, 0.011% sodium pyruvate, 0.45% glucose, 0.24% *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid, 10% fetal bovine serum and 10 ml/l of 100 \times antibiotic anti-mycotic solution (Sigma). Cells were maintained at 37 $^{\circ}$ C in the presence of 5% CO₂. Reporter lysis buffer and luciferase reagent for luciferase studies were supplied by Promega (Madison, WI). β -Galactosidase (β -Gal) reagent was obtained from Tropix (Bedford, MA), and LipofectAMINE reagent was purchased from Invitrogen (Carlsbad, CA). The PPAR γ antagonist *N*-(4'-aminopyridyl)-2-chloro-5-nitrobenzamide (T007) was synthesized in our laboratory, and its identity and purity (>98%) was confirmed by gas chromatography mass spectrometry. The C-DIMs compounds were all prepared by condensation of indole or ring-substituted indoles with substituted benzaldehydes essentially as described (Qin et al., 2004).

Cell proliferation assay

This assay was performed in 12-well tissue culture plates using an initial concentration of 2 \times 10⁴ cells per well, and Dulbecco's modified Eagle's medium (DMEM)/Ham's F-12 media containing 2.5% charcoal-stripped fetal bovine serum (FBS). Cells were counted on the initial day using a Z1 cell counter (Beckman Coulter, Fullerton, CA) and then treated either with vehicle [dimethyl sulfoxide (DMSO)] or the indicated indole ring-substituted C-DIM compounds. Every 48 hr, fresh medium was added along with the indicated compounds. Cell

counts were taken after 24, 48, 72 and 96 hr. The results are expressed as means \pm standard errors for at least 3 samples for each treatment group.

Transfection Assays

The Gal4 reporter containing five Gal4 response elements (pGal4) was kindly provided by Dr. Marty Mayo (University of North Carolina, Chapel Hill, NC). Gal4DBD-PPAR γ construct was a gift of Dr. Jennifer L. Oberfield (GlaxoSmithKline Research and Development). Cells were seeded in 12-well plates, and 0.4 μ g of GAL4-Luc, 0.04 μ g of β -GAL, 0.04 μ g of GAL4DBD-PPAR γ were transfected using LipofectAMINE reagent (Invitrogen) following the manufacturer's protocol. Cells were treated either with vehicle or respective compounds suspended in complete medium after 6 hr of transfection. Cell lysates were extracted after treatment for 20 – 22 hr by adding 100 μ l of 1 \times reporter lysis buffer per well, and 30 μ l of this extract was used to quantitate the luciferase activity using Lumicount (Perkin-Elmer Life and Analytical Sciences). Each experiment was conducted in triplicate and the results were normalized to the β -GAL activity.

Western blot analysis

SW480, HT-29 and HCT-15 colon cancer cells, Panc-28 (3×10^5) pancreatic cancer cells were seeded in 6-well tissue culture plates in DMEM/Ham's F-12 medium containing 2.5% charcoal-stripped FBS. Protein was extracted from the cells treated either with vehicle or the indicated compounds for 24 hr for p21 protein, or 72 hr for caveolin-1 protein. Samples were extracted in high salt buffer [50 mmol/l *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid, 500 mmol/l NaCl, 1.5 mmol/l MgCl₂, 1 mmol/l ethyleneglycol-bis(aminoethylether)-tetraacetic acid, 10% glycerol and 1% Triton X-100 (pH 7.5) and 5 μ l/ml protease inhibitor cocktail (Sigma-Aldrich)]. Extracts were incubated at 100°C for 2 min, separated on either 10 or 12% sodium dodecylsulfate-polyacrylamide gel electrophoresis gels and then transferred to polyvinylidene difluoride membrane (Bio-Rad, Hercules, CA). The polyvinylidene difluoride membrane was blocked in 5% TBST-Blotto (10 mM Tris-HCl, 150 mM NaCl, pH 8.0, 0.05% Triton X-100 and 5% non-fat dry milk) for 30 min and then incubated in fresh 5% TBST-Blotto with 1:1000 for caveolin-1 (Santa Cruz Biotechnology, Santa Cruz, CA), 1:1000 for p21 (BD Pharmingen, Franklin Lakes, NJ) and 1:2000 for β -tubulin (Santa Cruz Biotechnology, Santa Cruz, CA) primary antibody overnight with gentle shaking at 4°C. After washing with Tris-buffered saline containing Tween-20 (TBST) for 10 min, the membrane was incubated with respective secondary antibody (1:5000) (Santa Cruz Biotechnology) in 5% TBST-Blotto for 3 hr. The membrane is then washed with TBST for 10 min, incubated with chemiluminescence reagent from Perkin-Elmer for 1 min and then exposed to Kodak X-OMAT AR autoradiography film (Eastman Kodak, Rochester, NY).

Semi-quantitative real-time polymerase chain reaction

SW480 and HT29 colon cancer cells were treated either with vehicle (DMSO) or indicated indole ring-substituted C-DIMs compounds and, after 24 hr total, RNA was extracted using RNeasy kit (Qiagen, Valencia, CA). RNA concentration was measured by UV 260:280 nm absorption ratio, and 2 μ g RNA was used to synthesize cDNA using Reverse Transcription System (Promega). Polymerase chain reaction (PCR) conditions were as follows: initial denaturation at 94°C (1 min) followed by 28 cycles of denaturation for 30 sec at 94°C, annealing for 60 sec at 55°C, and extension at 72°C for 60 sec, and a final extension step at 72°C for 5 min. mRNA levels were normalized to GAPDH as an internal housekeeping gene. Primers were obtained from IDT (Coralville, IA) and used for amplification as follows: KLF4 (sense 5'-CTA TGG CAG GGA GTC CGC TCC-3'; anti-sense 5'-ATG ACC GAC GGG CTG CCG TAC-3') and GAPDH (sense 5'-ACG GAT TTG GTC GTA TTG GGC G-3'; anti-sense 5'-

CTC CTG GAA GAT GGT GAT GG-3'). PCR products were electrophoresed on 1% agarose gels containing ethidium bromide and visualized under UV transillumination.

RESULTS

Growth Inhibition by Indole Ring-Substituted C-DIMs

It has previously been shown that DIM-C-pPhC₆H₅ (Fig. 1) inhibited growth of several different cancer cell lines and also activated PPAR γ [1;3]. In this study, we have examined the effects of various symmetrical 5,5'-indole ring-substituted analogs of DIM-C-pPhC₆H₅ and determined the effects of substituent structure on the cytotoxicity and PPAR γ agonist activity of these compounds in Panc28 pancreatic cancer cells and SW480 colon cancer cell lines.

Figure 2 summarizes the cytotoxicity of DIM-C-pPhC₆H₅ and the 5,5'-substituted nitro (NO₂), bromo (Br), hydroxyl (OH), methyl (Me) and methoxy (OMe) analogs in Panc28 cells. All compounds caused a concentration-dependent decrease in cell counts after treatment for 24, 48, 72 or 96 hr, and similar results were observed in colon cancer cells (data not shown). Table 1 summarizes the growth inhibitory IC₅₀ values at all time points. There was less than a 2-fold difference in IC₅₀ values among the six C-DIM compounds in both cell lines, suggesting that introduction of the 5,5'-substituents on the indole ring do not substantially enhance or decrease the cytotoxicity of DIM-C-pPhC₆H₅.

Activation of PPAR γ by Indole Ring-Substituted C-DIMs

The effects of the indole ring substituents on the activation of PPAR γ by DIM-C-pPhC₆H₅ and related compounds were investigated in SW480 colon cancer cells transfected with GAL4-PPAR γ /GAL4-luc constructs. Treatment of SW480 colon cancer cells with 5.0 μ M DIM-C-pPhC₆H₅ (X=H), showed that the 5,5'-dihydroxy and 5,5'-dimethyl (X=OH and Me, respectively) analogs significantly induced luciferase activity and, at a lower concentration of 2.5 μ M, transactivation was induced only with the indole ring-substituted analogs and not DIM-C-pPhC₆H₅ (Fig. 3A). The results indicate that the 5,5'-dihydroxy and 5,5'-dimethyl compounds were more active than the unsubstituted compound DIM-C-pPhC₆H₅, and the 5,5'-dinitro, -dibromo and -dimethoxy analogs were inactive at this concentration and higher doses (> 5.0 μ M) were cytotoxic. The effects of the 5,5'-dimethyl and dihydroxy substituents on activation of PPAR γ -dependent activity by DIM-C-pPhCF₃ and DIM-C-pPhtBu were compared to the results obtained for the corresponding X-DIM-C-pPhC₆H₅ compounds in SW480 cells transfected with PPAR γ -GAL4/pGAL4 (Fig. 3B). The results show that introduction of 5,5'-dimethyl or -dihydroxy groups enhanced PPAR γ -dependent activity for all three PPAR γ -active C-DIMs. In SW480 cells transfected with PPAR γ -GAL4/pGAL4, induction of luciferase activity by DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and -dimethyl analogs was inhibited after cotreatment with 10 μ M T007, a PPAR γ antagonist (Fig. 3C).

Figure 4 summarizes the structure-dependent activation of luciferase activity in Panc1 cells treated with DIM-C-pPhC₆H₅ and the 5,5'-ring substituted analogs of this C-DIM. In this cell line, 2.5 and 5 μ M concentration of the 5,5'-dihydroxy, -dimethyl and -dimethoxy analogs were active, whereas 5 μ M DIM-C-pPhC₆H₅ did not induce luciferase activity. Higher concentration of this compound only weakly induced luciferase activity (data not shown). Thus, the 5,5'-dihydroxy and 5,5'-dimethyl substituted compounds were PPAR γ agonists in both cell lines, whereas the 5,5'-dimethoxy analog activated PPAR in Panc1 but not SW480 cells (Fig. 3A).

A comparison of the activation of PPAR γ -active DIM-C-pPhCF₃, DIM-C-pPhtBu and DIM-C-pPhC₆H₅ and their 5,5'-dimethyl and 5,5'-dihydroxy derivatives in Panc1 cells is shown in Figure 4B. Treatment with 2.5 μ M concentration of the C-DIMs induced luciferase activity in Panc1 cells with 7 out of 9 compounds, only DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy

derivative of DIM-C-pPhtBu were inactive. These results were similar to those observed in SW480 cells (Fig. 3B), except that the 5,5'-dihydroxy derivative of DIM-C-pPhtBu induced transactivation in colon but not in pancreatic cancer cells.

The PPAR γ agonist activity of DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and -dimethyl derivatives in Panc1 cells transfected PPAR γ -GAL4/pGAL4 was inhibited in cells cotreated with the PPAR γ antagonist T007 (10 μ M) (Fig. 4C), and these results were similar to that observed in SW480 cells (Fig. 3).

Activation of KLF4 and Caveolin-1 by Indole Ring-Substituted C-DIMs in Colon Cancer Cells

Previous studies show that KLF4 is activated by the PPAR γ agonist methyl-2-cyano-3,11-dioxo-18 β -olean-1,12-dien-30-oate (CDODA-Me) in colon cancer cells and these induction responses are inhibited by the PPAR γ antagonist T007 [16]. Results in Figure 5A show that DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and -dimethyl analogs also induce KLF4 mRNA levels in SW480 colon cancer cells. However, the induction response is only observed after treatment of SW480 or HT-29 cells with a relatively high concentration (12.5 μ M) of these compounds. Semi-quantitative RT-PCR analysis of KLF4 mRNA induction by the C-DIM compounds alone or in combination with the PPAR γ antagonist T007 is illustrated in Figure 5B. The results confirm that the C-DIMs induce KLF4 expression in colon cancer cells; however, the lack of inhibition by T007 suggests that the induction response was PPAR γ -independent. PPAR γ -independent induction of KLF4 by the C-DIMs was similar to results of a previous report showing that induction of KLF4 mRNA in colon cancer cells by the PPAR γ agonist 15-deoxy- $\Delta^{12,14}$ -prostaglandin J2 (PGJ2) was receptor-independent [17], whereas induction of KLF4 mRNA in colon cancer cells by CDODA-Me was receptor-dependent [16].

Previous studies showed that PPAR γ -active C-DIMs also enhanced caveolin-1 protein expression in colon cancer cells (Chintharlapalli et al., 2004), and Figure 6 illustrates the effects of 5 μ M DIM-C-pPhC₆H₅ and related indole ring-substituted compounds on expression of caveolin-1 in HT-19, HCT-15 and SW480 cells. Induction of caveolin-1 was not observed in HCT-15 cells (data not shown); however all three C-DIMs alone induced caveolin-1 in HT-29 cells, whereas only DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy but not the 5,5'-dimethyl analogs induced caveolin-1 in SW480 cells (fig. 6A). The effects of T007 on induction of caveolin-1 by C-DIMs in HT-29 cells were difficult to decipher due to induction of caveolin-1 by T007 alone. However, T007 inhibited caveolin-1 induction by DIM-C-pPhC₆H₅ and the dihydroxy analog. These results show that induction of caveolin-1 by C-DIMs was dependent on cell context and also on the structure of the C-DIM (Fig. 6B), suggesting that these indole ring substituted analogs of DIM-C-pPhC₆H₅ are selective PPAR γ modulators.

Activation of p21 by Indole Ring-Substituted C-DIMs in Pancreatic Cancer Cells

Figure 7A shows that both DIM-C-pPhC₆H₅ (#9) and DIM-C-pPhCF₃ (#1) induce p21 protein expression in Panc28 cells and the induction response by the latter compound was previously reported [1]. The 5,5'-dimethyl analog of DIM-C-pPhC₆H₅ also induced p21 protein but the corresponding 5,5'-dihydroxy indole ring-substituted compound was inactive at a concentration of 12.5 μ M. However, the effects of DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs on p21 expression were unaffected after cotreatment with the PPAR γ antagonist T007. These results suggest that in Panc28 cells, DIM-C-pPhC₆H₅ and related compounds enhance PPAR γ -independent expression of p21, whereas previous studies reported induction of p21 by DIM-C-pPhCF₃ in Panc28 cells was PPAR γ -dependent [1].

DISCUSSION

PPAR γ is a member of the nuclear receptor family of ligand-activated receptors, and the transcriptionally-active complex is a heterodimer containing PPAR γ and the retinoic acid \times receptor (RXR). The thiazolidinedione class of PPAR γ agonists has been used extensively for treatment of type II diabetes, and there is evidence that these compounds may have applications for treating other diseases including atherogenesis and cancer [18;19]. PPAR γ is overexpressed in multiple tumor types, and several studies shows the effectiveness of different structural classes of PPAR γ agonists for inhibiting cancer cell growth and inducing apoptosis. Typically, these compounds induce differentiation in cancer cells and inhibit G₀/G₁ to S phase progression which is associated with decreased expression of cyclin D1 and induction of the cyclin-dependent kinase inhibitors p21 and/or p27. Mechanistic studies using PPAR γ antagonists (T00s7 and GW9662) and PPAR γ knockdown by RNA interference demonstrate that many of the growth inhibitory and proapoptotic responses induced by PPAR γ agonists are receptor-independent [4–6;14;15].

Nuclear receptor agonists typically bind structurally-diverse compounds and this has also been observed for PPAR γ . For example, PPAR γ binds and is activated by endogenous biochemicals such as fatty acids and 15-deoxy- $\Delta^{12,14}$ -prostaglandin J2 (PGJ2), flavonoids, and various synthetic compounds including substituted indoles and chromane carboxylic acids, phosphono-phosphates, PPAR γ -active C-DIMs, CDODA-Me and triterpenoids such as 2-cyano-3,12-dioxo-18 β -oleana-1,9-dien-28-oic acid (CDDO) and related derivatives [20–26]. Not surprisingly, there is evidence that among this structurally-diverse group of compounds, PPAR γ agonists exhibit tissue-/cell- and response-specific differences suggesting that these compounds are selective PPAR γ modulators. For example, induction of non-steroidal anti-inflammatory drug-activated gene-1 (NAG-1) by PGJ2 in colon cancer cells was receptor-dependent, whereas induction of this gene by other PPAR γ agonists such as thiazolidinediones and C-DIMs was receptor-independent [4;27;28]. Moreover, research in our laboratory on α - and β -CDODA-Me, which are triterpenoids that exhibit different stereochemistries at C18, and a structurally-related 2-cyanobetulinic acid derivative (CN-BA) also exhibited response- and cell context-dependent differences [16;29]. Previous studies with PPAR γ -active C-DIMs showed that DIM-C-pPhCF₃ induced p21 in Panc28 cells and all three PPAR γ -active C-DIMs induced caveolin-1 in colon cancer cells and these responses were inhibited after cotreatment with PPAR γ antagonists T007 or GW9662 [1;3]. Moreover, we have also shown that PPAR γ -active triterpenoids induce KLF4 mRNA in a receptor-dependent manner in some colon cancer cell lines [16;29]. Therefore, in this study we first examined the cytotoxicity of DIM-C-pPhC₆H₅ and symmetrical 5,5'-indole ring substituents and their activation of PPAR γ (Fig. 1). The second objective was to determine whether these compounds were selective PPAR γ modulators with respect to activation of caveolin-1, KLF-4 and p21.

Results illustrated in Figure 2 and Table 1 show that the growth-inhibitory IC₅₀ values for DIM-C-pPhC₆H₅ and several indole ring-substituted analogs in SW480 colon and Panc28 pancreatic cancer cells were similar and varied between 1–10 μ M at all time points in both cell lines. Thus, introduction of a 5-methyl, 5-nitro, 5-bromo, 5-hydroxy or 5-methoxy into both indole rings did not substantially enhance or inhibit, their cytotoxic effects compared to the unsubstituted DIM-C-pPhC₆H₅.

The comparative activation of PPAR γ by DIM-C-pPhC₆H₅ and X-DIM-C-pPhC₆H₅ analogs was determined in Panc1 and SW480 cells transfected with PPAR γ -GAL4/pGAL4 constructs (Fig. 3 – Fig 4). Panc1 cells were used instead of Panc28 cells because of the higher transfection efficiency in the former cell line. In SW480 cells, the 5,5'-dihydroxy and 5,5'-dimethyl analogs induced PPAR γ -dependent transactivation and were clearly more potent than the DIM-C-pPhC₆H₅ (Fig. 3). Moreover, 5,5'-dihydroxy and 5,5'-dimethyl analogs of DIM-C-pPhC₆H₅

and DIM-C-pPhTbu also induced transactivation in SW480 cells transfected with PPAR γ -GAL4/pGAL4 demonstrating that introduction of these substituents enhanced the PPAR γ agonist activities of other PPAR γ -active C-DIMs in SW480 (Fig. 2) and Panc1 cells (Fig. 4). With one exception, similar structure-activity relationships for activating PPAR γ were observed in SW480 and Panc1 cells (Fig. 4) where the 5,5'-dihydroxy and 5,5'-dimethyl analogs were active and the 5,5'-dinitro and 5,5'-dibromo analogs were inactive. However, it was also observed that the 5,5'-dimethoxy-derivative also induced transactivation in Panc1 (but not SW480) cells at the 2.5 and 5.0 μ M concentrations used in this study. Higher doses were not used due to cytotoxicity but it is possible that the cell-context-dependent differences observed for the 5,5'-dimethoxy derivative are due to potency differences for this compound as a PPAR γ agonist in the two cell lines. The inhibitory effects of T007, a PPAR γ antagonist confirms that DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs are PPAR γ agonists (Fig. 3 and Fig 4) and these compounds and possibly the 5,5'-dimethoxy derivative further extend the number of C-DIMs that activate PPAR γ . It is apparent that the PPAR γ agonist activities of indole ring-substituted DIM-C-pPhC₆H₅ are structure-dependent; however, the cytotoxicities of these compounds are comparable, suggesting that this important aspect of their anticarcinogenic activity is PPAR γ -independent. This observation is consistent with a recent study showing that the antiproliferative activities of PPAR γ -active C-DIMs were not affected after cotreatment with PPAR γ antagonists [1].

Previous reports show that some PPAR γ agonists induce KLF4 in colon cancer cells; however, this response is structure and cell context-dependent. α - and β -CDODA-Me induced KLF4 expression in HT-29 and SW480 but not HCT-15 cells, whereas the 2-cyano derivative of betulinic induced KLF4 in HT-29 but not SW480 cells [16;29]. All three compounds are PPAR γ agonists but induction of KLF4 by α - and β -CDODA-Me was receptor-dependent and induction by the cyano betulinic acid derivative was PPAR γ -independent. Similar results were observed in this study (Fig. 5) where induction of KLF4 by DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs in SW480 and HT-29 cells was not inhibited after cotreatment with T007.

Receptor-dependent induction of caveolin-1 by CDODA-Me, PPAR γ -active C-DIMs and CN-BA has been reported in colon cancer cells; however, the induction responses were cell context-dependent [3;16;29]. For example, CN-BA induced caveolin-1 in HT-29 and HCT-15 cells but not SW480 cells, whereas β -CDODA-Me induces caveolin-1 in all three cell lines [16;29]. In this study, 5 μ M DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs induced caveolin-1 protein in HT-29 and SW480 cells (Fig. 6), and cotreatment with T007 inhibited the response in the latter cell line. In contrast, we did not observe induction of caveolin-1 in HCT-15 cells. The inhibitory response was not well-defined in HT-29 cells due to induction of caveolin-1 by T007 alone. Nevertheless, the pattern of caveolin-1 induction in the two cell lines by the indole ring-substituted compounds was similar to that previously described for PPAR γ -active C-DIMs in these same cell lines (3). However, in this study the failure to observe induction in HCT-15 cells was inconsistent with previous studies with C-DIMs in this cell line, and the differences observed are currently being reinvestigated.

In Panc28 cells, DIM-C-pPhCF₃ induced p21 expression that was PPAR γ -dependent [1], and Figure 7A confirms that both DIM-C-pPhC₆H₅ (#9) and DIM-C-pPhCF₃ (#1) induce p21 protein in this cell line. The 5,5'-dimethyl analog of DIM-C-pPhC₆H₅ also induced p21 protein but the corresponding 5,5'-dihydroxy analog was inactive at a concentration of 12.5 μ M. Moreover, the effects of DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs on p21 expression were unaffected after cotreatment with the PPAR γ antagonist T007. These results suggest that in Panc28 cells, DIM-C-pPhC₆H₅ and related compounds enhance PPAR γ -independent expression of p21, whereas previous studies reported induction of p21 by DIM-C-pPhCF₃ in Panc28 cells was PPAR γ -dependent [1].

These results observed in colon and pancreatic cancer cells further demonstrate that indole ring-substituted analogs of DIM-C-pPhC₆H₅ activate PPAR γ and some PPAR γ -mediated responses in colon cancer cells. However the role of the receptor in mediating the responses is cell context- and structure-dependent. It was also apparent that the PPAR γ -active and inactive C-DIM analogs exhibit comparable cytotoxicities suggesting that the antiproliferative activity of C-DIMs is primarily PPAR γ -independent. Currently, we are investigating structural modification of C-DIMs at various positions in the indole and phenyl rings in order to maximize their PPAR γ agonist activities and cytotoxic effects in cancer cell lines.

Acknowledgments

The financial assistance of the National Institutes of Health (ES09106, CA108718 and CA112337) and the Texas Agricultural Experiment Station is gratefully acknowledged.

REFERENCES

- Hong J, Samudio I, Liu S, Abdelrahim M, Safe S. Peroxisome proliferator-activated receptor γ -dependent activation of p21 in Panc-28 pancreatic cancer cells involves Sp1 and Sp4 proteins. *Endocrinology* 2004;vol. 145:5774–5785. [PubMed: 15345676]
- Vanderlaag K, Su Y, Frankel AE, Grage H, Smith R III, Khan S, Safe S. 1,1-Bis(3'-indolyl)-1-(p-substituted phenyl)methanes inhibit proliferation of estrogen receptor-negative breast cancer cells by activation of multiple pathways. *Breast Cancer Res.Treat.* 2007
- Chintharlapalli S, Smith R III, Samudio I, Zhang W, Safe S. 1,1-Bis(3'-indolyl)-1-(p-substitutedphenyl) methanes induce peroxisome proliferator-activated receptor γ -mediated growth inhibition, transactivation and differentiation markers in colon cancer cells. *Cancer Research* 2004;vol. 64:5994–6001. [PubMed: 15342379]
- Chintharlapalli S, Papineni S, Baek SJ, Liu S, Safe S. 1,1-Bis(3'-indolyl)-1-(p-substitutedphenyl) methanes are peroxisome proliferator-activated receptor gamma agonists but decrease HCT-116 colon cancer cell survival through receptor-independent activation of early growth response-1 and NAG-1. *Mol.Pharmacol* 2005;vol. 68:1782–1792. [PubMed: 16155208]
- Chintharlapalli S, Papineni S, Safe S. 1,1-Bis(3'-indolyl)-1-(p-substituted phenyl)methanes inhibit colon cancer cell and tumor growth through PPAR γ -dependent and PPAR γ -independent pathways. *Mol.Cancer Ther* 2006;vol. 5(no. 5):1362–1370. [PubMed: 16731770]
- Chintharlapalli S, Papineni S, Safe SH. 1,1-Bis(3'-indolyl)-1-(p-substitutedphenyl)methanes inhibit growth, induce apoptosis, and decrease the androgen receptor in LNCaP prostate cancer cells through PPAR γ -independent pathways. *Mol.Pharmacol* 2007;vol. 71(no. 2):558–569. [PubMed: 17093136]
- Contractor R, Samudio I, Estrov Z, Harris D, McCubrey JA, Safe S, Andreeff M, Konopleva M. A novel ring-substituted diindolylmethane 1,1-bis[3'-(5-methoxyindolyl)]-1-(p-t-butylphenyl)methane inhibits ERK activation and induces apoptosis in acute myeloid leukemia. *Cancer Research* 2005;vol. 65:2890–2898. [PubMed: 15805291]
- Qin C, Morrow D, Stewart J, Spencer K, Porter W, Smith R III, Phillips T, Abdelrahim M, Samudio I, Safe S. A new class of peroxisome proliferator-activated receptor γ (PPAR γ) agonists that inhibit growth of breast cancer cells: 1,1-bis(3'-indolyl)-1-(p-substitutedphenyl)methanes. *Mol.Cancer Therap* 2004;vol. 3:247–259. [PubMed: 15026545]
- Vanderlaag K, Samudio I, Burghardt R, Barhoumi R, Safe S. Inhibition of breast cancer cell growth and induction of cell death by 1,1-bis(3'-indolyl)methane (DIM) and 5,5'-dibromoDIM. *Cancer Lett* 2005;vol. 236(no. 2):198–212. [PubMed: 16051428]
- York M, Abdelrahim M, Chintharlapalli S, Lucero SD, Safe S. 1,1-Bis(3'-indolyl)-1-(p-substitutedphenyl)methanes induce apoptosis and inhibit renal cell carcinoma growth. *Clin.Cancer Res* 2007;vol. 13:6743–6752. [PubMed: 18006776]
- Abdelrahim M, Newman K, Vanderlaag K, Samudio I, Safe S. 3,3'-Diindolylmethane (DIM) and derivatives induce apoptosis in pancreatic cancer cells through endoplasmic reticulum stress-dependent upregulation of DR5. *Carcinogenesis* 2006;vol. 27:717–728. [PubMed: 16332727]

12. Calabro P, Samudio I, Safe SH, Willerson JT, Yeh ET. Inhibition of tumor necrosis factor- α -induced endothelial cell activation by a new class of PPAR γ agonists: an *in vitro* study. *J.Vascular Res* 2005;vol. 42:509–516.
13. Kassouf W, Chintharlapalli S, Abdelrahim M, Nelkin G, Safe S, Kamat AM. Inhibition of bladder tumor growth by 1,1-bis(3'-indolyl)-1-(*p*-substitutedphenyl)methanes: a new class of peroxisome proliferator-activated receptor γ agonists. *Cancer Research* 2006;vol. 66:412–418. [PubMed: 16397256]
14. Lei P, Abdelrahim M, Safe S. 1,1-Bis(3'-indolyl)-1-(*p*-substituted phenyl)methanes inhibit ovarian cancer cell growth through peroxisome proliferator-activated receptor-dependent and independent pathways. *Mol.Cancer Ther* 2006;vol. 5(no. 9):2324–2336. [PubMed: 16985067]
15. Cho SD, Yoon K, Chintharlapalli S, Abdelrahim M, Pei P, Hamilton S, Khan S, Ramaiah SK, Safe S. Nur77 agonists induce proapoptotic genes and responses in colon cancer cells through nuclear receptor-dependent and independent pathways. *Cancer Research* 2007;vol. 67:674–683. [PubMed: 17234778]
16. Chintharlapalli S, Papineni S, Jutooru I, McAlees A, Safe S. Structure-dependent activity of glycyrrhetic acid derivatives as peroxisome proliferator-activated receptor γ (PPAR γ) agonists in colon cancer cells. *Mol.Cancer Therap* 2007;vol. 6:1588–1598. [PubMed: 17513608]
17. Chen ZY, Tseng CC. 15-Deoxy- $\Delta^{12,14}$ prostaglandin J₂ up-regulates Krüppel-like factor 4 expression independently of peroxisome proliferator-activated receptor γ by activating the mitogen-activated protein kinase kinase/extracellular signal-regulated kinase signal transduction pathway in HT-29 colon cancer cells. *Mol.Pharmacol* 2005;vol. 68(no. 5):1203–1213. [PubMed: 16077033]
18. Escher P, Wahli W. Peroxisome proliferator-activated receptors: insight into multiple cellular functions. *Mutat.Res* 2000;vol. 448(no. 2):121–138. [PubMed: 10725467]
19. Fajas L, Debril MB, Auwerx J. Peroxisome proliferator-activated receptor- γ : from adipogenesis to carcinogenesis. *Journal of Molecular Endocrinology* 2001;vol. 27(no. 1):1–9. [PubMed: 11463572]
20. Suh N, Wang Y, Honda T, Gribble GW, Dmitrovsky E, Hickey WF, Maue RA, Place AE, Porter DM, Spinella MJ, Williams CR, Wu G, Dannenberg AJ, Flanders KC, Letterio JJ, Mangelsdorf DJ, Nathan CF, Nguyen L, Porter WW, Ren RF, Roberts AB, Roche NS, Subbaramaiah K, Sporn MB. A novel synthetic oleanane triterpenoid, 2-cyano-3,12-dioxolean-1,9-dien-28-oic acid, with potent differentiating, antiproliferative, and anti-inflammatory activity. *Cancer Research* 1999;vol. 59(no. 2):336–341. [PubMed: 9927043]
21. Rieusset J, Touri F, Michalik L, Escher P, Desvergne B, Niesor E, Wahli W. A new selective peroxisome proliferator-activated receptor γ antagonist with antiobesity and antidiabetic activity. *Molecular Endocrinology* 2002;vol. 16(no. 11):2628–2644. [PubMed: 12403851]
22. Berger JP, Petro AE, Macnaul KL, Kelly LJ, Zhang BB, Richards K, Elbrecht A, Johnson BA, Zhou G, Doebber TW, Biswas C, Parikh M, Sharma N, Tanen MR, Thompson GM, Ventre J, Adams AD, Mosley R, Surwit RS, Moller DE. Distinct properties and advantages of a novel peroxisome proliferator-activated protein γ selective modulator. *Molecular Endocrinology* 2003;vol. 17(no. 4):662–676. [PubMed: 12554792]
23. Koyama H, Miller DJ, Boueres JK, Desai RC, Jones AB, Berger JP, Macnaul KL, Kelly LJ, Doebber TW, Wu MS, Zhou G, Wang PR, Ippolito MC, Chao YS, Agrawal AK, Franklin R, Heck JV, Wright SD, Moller DE, Sahoo SP. (2*R*)-2-ethylchromane-2-carboxylic acids: discovery of novel PPAR α/γ dual agonists as antihyperglycemic and hypolipidemic agents. *J.Med.Chem* 2004;vol. 47(no. 12):3255–3263. [PubMed: 15163205]
24. Acton JJ III, Black RM, Jones AB, Moller DE, Colwell L, Doebber TW, Macnaul KL, Berger J, Wood HB. Benzoyl 2-methyl indoles as selective PPAR γ modulators. *Bioorg.Med.Chem.Lett* 2005;vol. 15(no. 2):357–362. [PubMed: 15603954]
25. Liu K, Black RM, Acton JJ III, Mosley R, Debenham S, Abola R, Yang M, Tschirret-Guth R, Colwell L, Liu C, Wu M, Wang CF, Macnaul KL, McCann ME, Moller DE, Berger JP, Meinke PT, Jones AB, Wood HB. Selective PPAR γ modulators with improved pharmacological profiles. *Bioorg.Med.Chem.Lett* 2005;vol. 15(no. 10):2437–2440. [PubMed: 15863293]
26. Schopfer FJ, Lin Y, Baker PR, Cui T, Garcia-Barrio M, Zhang J, Chen K, Chen YE, Freeman BA. Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor γ ligand. *Proc.Natl.Acad.Sci.U.S.A* 2005;vol. 102(no. 7):2340–2345. [PubMed: 15701701]

27. Baek SJ, Wilson LC, Hsi LC, Eling TE. Troglitazone, a peroxisome proliferator-activated receptor γ (PPAR γ) ligand, selectively induces the early growth response-1 gene independently of PPAR γ . A novel mechanism for its anti-tumorigenic activity. *Journal of Biological Chemistry* 2003;vol. 278 (no. 8):5845–5853. [PubMed: 12475986]
28. Baek SJ, Kim JS, Nixon JB, DiAugustine RP, Eling TE. Expression of NAG-1, a transforming growth factor- β superfamily member, by troglitazone requires the early growth response gene *EGR-1*. *Journal of Biological Chemistry* 2004;vol. 279(no. 8):6883–6892. [PubMed: 14662774]
29. Chintharlapalli S, Papineni S, Liu S, Jutooru I, Chadalapaka G, Cho SD, Murthy R, You YJ, Safe S. 2-Cyano-lup-1-en-3-oxo-20-oic acid, a cyano derivative of betulinic acid, activates peroxisome proliferator-activated receptor γ in colon and pancreatic cancer cells. *Carcinogenesis* 2007;vol. 28:2337–2346. [PubMed: 17724373]




Figure 1.
Structure of indole ring-substituted DIM-C-pPhC₆H₅.





Figure 2. Cell proliferation assays. Panc28 cells were treated with X-DIM-C-pPhC₆H₅ (X = H, X=NO₂, X=Br, X=OH X=Me and X=OMe), and cell numbers were determined after treatment for 24, 48, 72 and 96 hr as described in Materials and Methods. Columns = mean of three replicate determinations for each treatment group; bars = SE.

**Figure 3.**

X-DIM-C-pPhC₆H₅ activates PPAR γ in SW480 cells. (A) Activation of X-DIM-C-pPhC₆H₅. SW480 cells were transfected with PPAR γ -Gal4/pGal4 and treated with DMSO or different concentrations of X-DIM-C-pPhC₆H₅ (X=H, X=NO₂, X=Br, X=OH, X=Me and X=OMe), and luciferase activity was determined as described in Materials and Methods. (B) Activation of PPAR γ -active DIM-C-pPhCF₃, DIM-C-pPhtBu and DIM-C-pPhC₆H₅ and their 5,5'-dimethyl and 5,5'-dihydroxy derivatives. SW480 cells were treated as in (A), and luciferase activity was determined as described in Materials and Methods. Columns = mean of three replicate determinations for each treatment group; bars = SE. * = $P < 0.05$, significant induction. (C) Effects of PPAR γ antagonist T007 on induced transactivation in SW480 cells transfected

with PPAR γ -GAL4/pGAL4. Induction of luciferase activity by DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs was inhibited in cells cotreated with 10 μ M T007, a PPAR γ antagonist. Columns = mean of three replicate determinations for each treatment group; bars = SE; * = $P < 0.05$, significant induction; ** = $P < 0.05$, significant inhibition.





Figure 4. X-DIM-C-pPhC₆H₅ activates PPAR γ in Panc1 cells. (A) Activation of X-DIM-C-pPhC₆H₅. Panc1 cells were transfected with PPAR γ -Gal4/pGal4 and treated with DMSO or different concentrations of X-DIM-C-pPhC₆H₅ (X=H, X=NO₂, X=Br, X=OH, X=Me and X=OMe), and luciferase activity was determined as described in Materials and Methods. (B) Activation of PPAR γ -active DIM-C-pPhCF₃, DIM-C-pPhtBu and DIM-C-pPhC₆H₅ and their 5,5'-dimethyl and 5,5'-dihydroxy derivatives. Panc1 cells were treated as described in (A), and luciferase activity was determined as described in Materials and Methods. Columns = mean of three replicate determinations for each treatment group; bars = SE; * = $P < 0.05$, significant induction. (C) Effects of PPAR γ antagonist T007 on induced transactivation in Panc1 cells.

Cells were transfected with PPAR γ -GAL4/pGAL4, induction of luciferase activity by DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs was inhibited after cotreatment with 10 μ M T007, a PPAR γ antagonist. Columns = mean of three replicate determinations for each treatment group; bars = SE; * = $P < 0.05$, significant induction; ** = $P < 0.05$, significant inhibition.

A



B




Figure 5. Induction of KLF4 gene expression by DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs in SW480 and HT-29 cells. Induction of KLF-4 (A) and inhibition by T007 (B). Cells were treated with different concentrations of DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs or T007 alone or in combination for 24 hr, and KLF4 mRNA levels were determined by real-time polymerase chain reaction as described in the Materials and Methods. Each experiment was replicated (> 3X). T007 did not inhibit KLF4 mRNA induction by DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs.




Figure 6. Induction of caveolin-1 expression in colon cancer cells and effects of T007 on induction of caveolin-1. HCT-15 (A), HT-29 (B) or SW480 (C) cells were treated with DMSO or 5 μ M DIM-C-pPhC₆H₅ and the 5,5'-dihydroxy and 5,5'-dimethyl analogs alone or in combination with 10 μ M T007 and caveolin-1 expression was determined by western blot analysis as described in the Materials and Methods.




Figure 7. Induction of p21 protein. Panc-28 cells were treated with the different compounds alone (A) or in combination with T007 (B) as indicated for 24 hr, and whole-cell lysates were obtained and analyzed by immunoblots as described in the Materials and Methods.

Table 1

Growth inhibitory IC₅₀ values (μM) for X-IM-C-pPhC₆H₅ in Panc28 and SW480 cells treated for 24, 48, 72 or 96 hr.

	X=H	X=NO ₂	X=Br	X=OH	X=Me	X=OMe
<u>Panc28 cells</u>						
Day 1	5.17	4.08	4.45	5.68	4.80	5.08
Day 2	4.79	4.17	4.31	5.21	4.08	4.55
Day 3	4.42	3.04	3.90	4.52	2.90	3.22
Day 4	4.46	2.62	4.11	3.56	3.56	3.63
<u>SW480 cells</u>						
Day 1	4.77	2.25	3.31	2.14	3.62	3.06
Day 2	4.51	2.20	2.31	2.04	2.28	2.65
Day 3	3.99	2.03	1.94	2.15	2.28	2.82
Day 4	3.95	1.93	1.88	1.20	1.25	2.29