

Published in final edited form as:

Brain Behav Immun. 2010 May ; 24(4): 529–530. doi:10.1016/j.bbi.2010.02.002.

Psychological Stress, Telomeres, and Telomerase

Janice K. Kiecolt-Glaser^{a,b} and Ronald Glaser^{b,c}

^aDepartment of Psychiatry, The Ohio State University College of Medicine, USA

^bInstitute for Behavioral Medicine Research, Ohio State University College of Medicine, USA

^cDepartment of Molecular Virology, Immunology, and Medical Genetics, Ohio State University College of Medicine, USA

Telomeres and telomerase, hot topics in the aging literature in recent years, will undoubtedly receive even more attention following the award of the 2009 Nobel Prize in Medicine “for the discovery of how chromosomes are protected by telomeres and the enzyme telomerase” to Elizabeth Blackburn, her former student Carol Greider, and Jack Szostak. The paper by Epel and colleagues in this BBI issue (Epel et al., in press) represents the latest elegant behavioral work on this topic from the investigators who published the first paper linking stress and telomere length (Epel et al., 2004), an observation that has now been confirmed by several laboratories (Damjanovic et al., 2007; Parks et al., 2009).

In this report they replicate their original finding of lower basal telomerase activity associated with chronic stress, and they also show that an acute stressor boosts telomerase reactivity (Epel et al., in press). Furthermore, they provide evidence that the increased telomerase reactivity is independent of changes in numbers or percentages of monocytes, lymphocytes, and specific T-cell types, and they show that two facets of the stress response, the degree of threat and the magnitude of the cortisol response, are both associated with larger increases in telomerase activity.

These studies of stress and cell aging are consequential for several reasons. First, these cell aging measures have clinical significance for health: a growing literature has linked shorter telomeres and lower levels of the enzyme telomerase with adverse health behaviors, aging, age-related diseases, and mortality (Epel et al., in press).

The evidence linking psychological stress and cell aging is also important because it signposts new pathways on the road from stress to mortality—valuable new information about the ways that stress kills. Psychological stress can boost inflammation (Kiecolt-Glaser et al., 2003) as well as oxidative stress (Epel et al., 2004), and both inflammation and oxidative stress accelerate telomere attrition (Aviv, 2004). Inflammation triggers T-cell proliferation and enhances the leukocyte turnover rate, one known cause of telomere shortening, while oxidative stress accelerates telomere attrition by promoting telomere erosion during cellular replication (Aviv, 2004).

© 2010 Elsevier Inc. All rights reserved

Address correspondence to Janice K. Kiecolt-Glaser, Ph.D., Institute for Behavioral Medicine Research, The Ohio State University College of Medicine, IBMR Building, 460 Medical Center Drive, Room 130C, Columbus, OH 43210-1228, USA; 614-293-3499 (office), 614-366-3627 (fax), Janice.Kiecolt-Glaser@osumc.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Psychological stress impacts telomere length through another route as well. The enhanced replication of memory CD8+ T-cells induced by cytomegalovirus (CMV) and/or Epstein-Barr virus (EBV) antigens can exhaust virus-specific T-cells. These two herpesviruses latently infect their host; their periodic reactivation provides a source of viral antigens that stimulates a cellular immune response to viral proteins. Chronic exposure to high CMV and/or EBV antigen levels can promote progressive telomere shortening in antigen-specific T-cells; sustained stimulation of cell division in virus-specific T-cells has been associated with reductions in telomere length and telomerase activity in the CD8+ T cells among even normal healthy subjects (Pawelec et al., 2009; van Baarle et al., 2008).

Psychological stress clearly drives replication of several herpesviruses including CMV and EBV by impairing the ability of the cellular immune response to control viral latency (Glaser and Kiecolt-Glaser, 1994). EBV and CMV infections are ubiquitous in adults, e.g., greater than 90% of adults are seropositive for EBV (Glaser and Kiecolt-Glaser, 1994), and thus the risk associated with chronic stress-related reactivation is not trivial. Considerable circumstantial evidence has implicated CMV as a player in immunosenescence as well as mortality (Pawelec et al., 2009). Chronic psychological stress clearly enhances CMV and EBV replication (Glaser and Kiecolt-Glaser, 1994), and through this route could simultaneously promote reductions in telomere length and telomerase activity.

The original observation from Epel et al. linking chronic stress to telomere length and telomerase activity will be a central touchstone for psychoneuroimmunology researchers for the foreseeable future (Epel et al., 2004). Their newest work expands the mechanistic base and tells us useful things about the links between psychological and biological facets of the stress response and telomerase activity.

Acknowledgments

Work on this paper was supported in part by NIH grants AG029562, CA126857, CA131029, AT003912, Ohio State Comprehensive Cancer Center Core Grant CA16058, and NCRN Grant UL1RR025755.

References

- Aviv A. Telomeres and human aging: facts and fables. *Science of Aging, Knowledge, and Environment* 2004;51:pe43.
- Damjanovic AK, Yang Y, Glaser R, Kiecolt-Glaser JK, Nguyen H, Laskowski B, Zou Y, Beversdorf DQ, Weng N.-p. Accelerated telomere erosion is associated with a declining immune function of caregivers of Alzheimer's disease patients. *J. Immunol* 2007;179:4249–4254. [PubMed: 17785865]
- Epel ES, Blackburn EH, Lin J, Dhabhar FS, Adler NE, Morrow JD, Cawthon RM. Accelerated telomere shortening in response to life stress. *Proc. Natl. Acad. Sci. U. S. A* 2004;101:17312–17315. [PubMed: 15574496]
- Epel ES, Lin J, Dhabhar FS, Wolkowitz OM, Puterman E, Karan L, Blackburn EH. Dynamics of telomerase activity in response to acute psychological stress. *Brain. Behav. Immun.* in press.
- Glaser, R.; Kiecolt-Glaser, JK. Stress-associated immune modulation and its implications for reactivation of latent herpesviruses. In: Glaser, R.; Jones, J., editors. *Human herpesvirus infections*. Dekker; New York: 1994. p. 245-270.
- Kiecolt-Glaser JK, Preacher KJ, MacCallum RC, Atkinson C, Malarkey WB, Glaser R. Chronic stress and age-related increases in the proinflammatory cytokine IL-6. *Proc. Natl. Acad. Sci. U. S. A* 2003;100:9090–9095. [PubMed: 12840146]
- Parks CG, Miller DB, McCanlies EC, Cawthon RM, Andrew ME, DeRoo LA, Sandler DP. Telomere length, current perceived stress, and urinary stress hormones in women. *Cancer Epidemiol. Biomarkers Prev* 2009;18:551–560. [PubMed: 19190150]
- Pawelec G, Derhovanessian E, Larbi A, Strindhall J, Wikby A. Cytomegalovirus and human immunosenescence. *Rev Med Virol* 2009;19:47–56. [PubMed: 19035529]

van Baarle D, Nanlohy NM, Otto S, Plunkett FJ, Fletcher JM, Akbar AN. Progressive telomere shortening of Epstein-Barr virus-specific memory T cells during HIV infection: Contributor to exhaustion? *The Journal of Infectious Diseases* 2008;198:1353–1357. [PubMed: 18816191]