



Published in final edited form as:

Circ Res. 2010 March 19; 106(5): 815–817. doi:10.1161/CIRCRESAHA.109.216226.

M Cells in the Human Heart

Charles Antzelevitch, PhD, FAHA

Masonic Medical Research Laboratory, Utica, NY

Keywords

Ventricle; Electrophysiology; Anatomy; Cardiac arrhythmias

Dispersion of repolarization across the ventricular wall has been suggested to underlie the inscription of the normal electrocardiographic T wave and when amplified to contribute prominently to the development of cardiac arrhythmias.^{1–3}

The magnitude of transmural dispersion of repolarization (TDR) is due to intrinsic differences in the action potential duration (APD) of the three principal cell types that comprise the ventricular myocardium and the extent to which these repolarization differences are damped by electrotonic forces. An increase in tissue resistivity in the deep subepicardium,⁴ due to a sharp transition in cell orientation,³ reduced expression of connexin 43 (Cx43)^{5, 6} and increased density of collagen⁷ in this region, contributes to the expression of repolarization heterogeneities across the ventricular wall by limiting the degree of electrotonic interaction between the myocardial layers. Thus, the degree of electrotonic coupling, together with the intrinsic differences in APD, determine the extent to which TDR is expressed, its impact on arrhythmogenesis as well as on the morphology of the T wave. It is noteworthy that even in the absence of any difference in final repolarization time, electrotonic forces generated by transmural differences in the shape of the action potential can inscribe an upright T wave in the electrocardiogram (ECG). Theoretical studies have also been helpful in our understanding of the role of electrical coupling in the expression of TDR.⁸

TDR is in large part attributable to the presence of M cells between the endocardial and epicardial layers of the heart. The M cell, discovered in the early 1990's, and named in memory of Gordon K. Moe,^{9, 10} has as its hallmark the ability to prolong its action potential more than that of normal epicardium or endocardium in response to a slowing of rate or exposure to agents that prolong APD.^{9, 11–13}

Histologically, M cells are similar to epicardial and endocardial cells. Electrophysiologically and pharmacologically, they appear to be a hybrid between Purkinje and ventricular cells.¹⁴ Like Purkinje fibers, M cells show a prominent APD prolongation and develop early afterdepolarizations (EAD) in response to rapidly activating delayed rectifier potassium

Address for correspondence: Charles Antzelevitch, PhD, FACC, FAHA, FHRS, Executive Director and Director of Research, Gordon K. Moe Scholar, Masonic Medical Research Laboratory, 2150 Bleeker Street, Utica, N.Y. 13501, Phone: (315)735-2217, FAX: (315)735-5648, ca@mmrl.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosures: none

current (I_{Kr}) blockers, whereas epicardium and endocardium generally are less likely to do so. Like Purkinje fibers, M cells develop delayed afterdepolarizations (DAD) more readily in response to agents that calcium load or overload the cardiac cell. Thus, they may contribute to the development of both substrate and triggers associated with arrhythmogenesis.

In the dog, the ionic basis for these features of the M cell include the presence of a smaller slowly activating delayed rectifier current (I_{Ks}),¹⁵ a larger late sodium current (late I_{Na})¹⁶ and a larger Na-Ca exchange current (I_{Na-Ca}).¹⁷

Distribution of M cells within the ventricular wall has been investigated in greatest detail in the left ventricle of the canine heart. Although transitional cells are found throughout the wall in the canine left ventricle, M cells displaying the longest action potentials (at BCLs \geq 2000 msec) are often localized in the deep subendocardium to midmyocardium in the anterior wall⁴ and throughout the wall in the region of the right ventricular outflow tract (RVOT).¹⁰ M cells are also present in the deep cell layers of endocardial structures, including papillary muscles, trabeculae and the inter-ventricular septum^{18, 19}. Sicouri et al.,¹⁹ recently reported a greater TDR across the canine inter-ventricular septum than across the left ventricular (LV) free wall, a finding that is at odds with that of Morita and co-workers.²⁰ Cells with characteristics of M cells have been described in a wide variety of animal species including canine, guinea pig, rabbit, and pig ventricles.^{1, 4, 9, 12, 15, 18, 21–38}

There is a paucity of data dealing with the presence of M cells in the human heart. Drouin and co-workers were the first to report the presence of M cells in humans based on microelectrode recordings from tissue slices isolated from normal human hearts.³⁹ Li and co-workers²⁸ studied the characteristics of cells enzymatically dissociated from normal and failing human hearts and thus identified the presence of cells with the characteristics of M cells. Taggart and co-workers⁴⁰ attempted to map transmural distinctions of repolarization in the human heart *in vivo* using unipolar plunge electrodes, but failed to detect the presence of M cells or any type of TDR.

The extent to which transmural dispersion of repolarization (TDR) exists within the normal heart of animals *in vivo* has been a matter of considerable debate.^{10, 13, 26, 41–44} The controversy derives in large part from the fact that quantitation of TDR in animals *in vivo* is hampered by 1) the inability to record local repolarization accurately, 2) the unavoidable use of anesthesia, which reduces TDR and 3) less than optimal recording conditions. These same constraints apply to the measurement of TDR in the human heart^{40, 45}. Indirect evidence for the presence of a prominent TDR in the human heart in the absence of general anesthesia was provided in a recent study.⁴⁶

Detailed maps of repolarization characteristics in the human heart similar to those available for the dog heart were not available until recently. In this issue of *Circulation Research*, Glukhov et al.⁴⁷ present the results of optical recording of action potentials from coronary-perfused left ventricular wedge preparations isolated from normal and failing human hearts.

In nominally normal hearts, they recorded an average TDR of 104 ± 17 ms, which increased in some regions to values as high as 147 ms. M cells were found to cluster in islands located in the deep subendocardium with action potential duration (APD) averaging 527 ± 43 ms. Outside of the islands, subepicardial, midmyocardial, and subendocardial APD measured at 80% repolarization (APD_{80}) values were 380 ± 17 ms, 450 ± 26 ms, and 479 ± 23 ms. Although these TDR values are much greater than those reported in canine LV wedge preparations, the clustering of M cells in islands was observed in the canine heart by Akar et al. using optical recording techniques.⁴⁸ Interestingly, in failing hearts, TDR was reduced to 47 ± 6 ms and the authors report the absence of M cells. In failing hearts APD_{80} increased to 492 ± 38

ms, 512 ± 29 ms and 470 ± 28 ms in subendocardial, midmyocardial and subepicardial regions, respectively, approximating the APD values of the M cells in the normal heart, raising the interesting possibility that the ion channel remodeling associated with heart failure (reduced outward and augmented inward ion channel currents)⁴⁹, 50 transforms epicardial and endocardial cells into those with properties similar to those of M cells.

The hallmark of the M cell is its ability to prolong more than the other normal ventricular cell types in response to a slowing of heart rate and/or in response to APD prolonging drugs. 11 This has been the traditional definition of M cell for the past 20 years. Glukhov et al.⁴⁷ propose a new definition, which is based on the local APD gradient between the islands of delayed repolarization (M cells) and the neighboring myocardium. Based on their definition there is no evidence of M cells in failing hearts. Based on the traditional definition of the M cell, which distinguishes a M cell based on the ability of its APD to prolong prominently at slower rates, all of the cells in all three layers of the failing hearts are M cells.

The authors are to be applauded for making the effort to record a transmural ECG. This is the most important parameter to monitor the quality of the preparation, which is often circumvented by other groups claiming “high quality” wedge preparations. The normal convention for recording a transmural pseudo-ECG is to place the + electrode of the ECG leads facing the epicardial side of the preparation and the – terminal facing the endocardial surface, following the convention of precordial leads on the body surface. This appears to be reversed in their experiments, resulting an inverted pseudo-ECG. Thus the apparent ST segment elevation shown in Figure 1 of their paper is actually a depression and the negative T wave is actually a concordant positive T wave, which is expected based on the repolarization sequence. The ST segment depression suggests the presence of mild ischemia on the endocardial side of the preparation. It is difficult to determine the extent to which the ischemia/hypoxia may have been heterogeneous and it is possible that this non-homogeneity may be responsible for creating islands of abbreviated responses, which could give rise to the appearance of islands of M cells.

Congruent with the studies of Poelzing et al.⁵ in the dog and Yamada et al. in the rat⁶ showing reduced Cx43 in the epicardium, Glukhov et al., using immunostaining techniques, showed reduced expression of Cx43 in epicardium in normal, but particularly in failing, human hearts.

Also of interest are the results of Glukhov et al. showing reduced TDR in human failing hearts. This result is opposite to the that obtained in animal models, in which TDR has been shown to be markedly augmented with the development of congestive heart failure secondary to tachypacing.⁵¹

As the authors point out, these conclusions are based on limited data derived from relatively few (5) non-failing hearts that were not entirely normal and few (5) failing hearts of various etiologies. Accordingly, additional studies are needed to confirm and expand these findings, but the authors are to be congratulated on this significant advance in our understanding of human electrophysiology.

Non-standard Abbreviations and Acronyms

APD	action potential duration
APD ₈₀	APD measured at 80% repolarization
Cx43	connexin 43

DAD	develop delayed afterdepolarizations
EAD	early afterdepolarization
ECG	electrocardiogram
LV	left ventricle
I_{Kr}	rapidly activating delayed rectifier potassium current
I_{Ks}	slowly activating delayed rectifier potassium current
I_{Na-Ca}	Na-Ca exchange current
Late I_{Na}	late sodium channel current
RVOT	right ventricular outflow tract
TDR	transmural dispersion of repolarization

Acknowledgments

Sources of Funding:

CA is supported by grant HL47678 (CA) from NHLBI, and New York State and Florida Grand Lodges of Free and Accepted Masons.

Reference List

1. Yan GX, Antzelevitch C. Cellular basis for the normal T wave and the electrocardiographic manifestations of the long QT syndrome. *Circulation* 1998;98:1928–1936. [PubMed: 9799215]
2. Patel C, Burke JF, Patel H, Gupta S, Kowey PR, Antzelevitch C, Yan GX. Is there a significant transmural gradient in repolarization time in the intact heart? Cellular basis of the T wave: a century of controversy. *Circ Arrhythm Electrophysiol* 2009;2:80–88. [PubMed: 19808446]
3. Antzelevitch C. The role of spatial dispersion of repolarization in inherited and acquired sudden cardiac death syndromes. *Am J Physiol Heart Circ Physiol* 2007;293:H2024–H2038. [PubMed: 17586620]
4. Yan GX, Shimizu W, Antzelevitch C. Characteristics and distribution of M cells in arterially-perfused canine left ventricular wedge preparations. *Circulation* 1998;98:1921–1927. [PubMed: 9799214]
5. Poelzing S, Akar FG, Baron E, Rosenbaum DS. Heterogeneous connexin43 expression produces electrophysiological heterogeneities across ventricular wall. *Am J Physiol Heart Circ Physiol* 2004;286:H2001–H2009. [PubMed: 14704225]
6. Yamada KA, Kanter EM, Green KG, Saffitz JE. Transmural distribution of connexins in rodent hearts. *J Cardiovasc Electrophysiol* 2004;15:710–715. [PubMed: 15175068]
7. LeGrice IJ, Smaill BH, Chai LZ, Edgar SG, Gavin JB, Hunter PJ. Laminar structure of the heart: Laminar structure of the heart: ventricular myocytes arrangement and connective tissue architecture in the dog. *Am J Physiol* 1996;269:H571–H582. [PubMed: 7653621]
8. Viswanathan PC, Shaw RM, Rudy Y. Effects of I_{Kr} and I_{Ks} heterogeneity on action potential duration and its rate-dependence: A simulation study. *Circulation* 1999;99:2466–2474. [PubMed: 10318671]
9. Sicouri S, Antzelevitch C. A subpopulation of cells with unique electrophysiological properties in the deep subepicardium of the canine ventricle. The M cell. *Circ Res* 1991;68:1729–1741. [PubMed: 2036721]
10. Antzelevitch C, Shimizu W, Yan GX, Sicouri S, Weissenburger J, Nesterenko VV, Burashnikov A, Di Diego JM, Saffitz J, Thomas GP. The M cell: its contribution to the ECG and to normal and abnormal electrical function of the heart. *J Cardiovasc Electrophysiol* 1999;10:1124–1152. [PubMed: 10466495]

11. Antzelevitch C, Sicouri S, Litovsky SH, Lukas A, Krishnan SC, Di Diego JM, Gintant GA, Liu DW. Heterogeneity within the ventricular wall. Electrophysiology and pharmacology of epicardial, endocardial, and M cells. *Circ Res* 1991;69:1427–1449. [PubMed: 1659499]
12. Anyukhovsky EP, Sosunov EA, Rosen MR. Regional differences in electrophysiologic properties of epicardium, midmyocardium and endocardium: *In vitro* and *in vivo* correlations. *Circulation* 1996;94:1981–1988. [PubMed: 8873677]
13. Anyukhovsky EP, Sosunov EA, Gainullin RZ, Rosen MR. The controversial M cell. *J Cardiovasc Electrophysiol* 1999;10:244–260. [PubMed: 10090229]
14. Antzelevitch, C.; Dumaine, R. Electrical heterogeneity in the heart: physiological, pharmacological and clinical implications. In: Page, E.; Fozzard, HA.; Solaro, RJ., editors. *The Cardiovascular System. Volume 1, The Heart*. New York: The American Physiological Society by Oxford University Press; 2002. p. 654-692.
15. Liu DW, Antzelevitch C. Characteristics of the delayed rectifier current (I_{Kr} and I_{Ks}) in canine ventricular epicardial, midmyocardial, and endocardial myocytes. A weaker I_{Ks} contributes to the longer action potential of the M cell. *Circ Res* 1995;76:351–365. [PubMed: 7859382]
16. Zygmunt AC, Eddlestone GT, Thomas GP, Nesterenko VV, Antzelevitch C. Larger late sodium conductance in M cells contributes to electrical heterogeneity in canine ventricle. *Am J Physiol* 2001;281:H689–H697.
17. Zygmunt AC, Goodrow RJ, Antzelevitch C. I_{NaCa} contributes to electrical heterogeneity within the canine ventricle. *Am J Physiol Heart Circ Physiol* 2000;278:H1671–H1678. [PubMed: 10775148]
18. Sicouri S, Antzelevitch C. Electrophysiologic characteristics of M cells in the canine left ventricular free wall. *J Cardiovasc Electrophysiol* 1995;6:591–603. [PubMed: 8535556]
19. Sicouri S, Glass A, Ferreiro M, Antzelevitch C. Transseptal dispersion of repolarization and its role in the development of torsade de pointes arrhythmias. *J Cardiovasc Electrophysiol*. 2009 In press.
20. Morita ST, Zipes DP, Morita H, Wu J. Analysis of action potentials in the canine ventricular septum: no phenotypic expression of M cells. *Cardiovasc Res* 2007;74:96–103. [PubMed: 17266946]
21. Antzelevitch C, Sicouri S. Clinical relevance of cardiac arrhythmias generated by afterdepolarizations. Role of M cells in the generation of U waves, triggered activity and torsade de pointes. *J Am Coll Cardiol* 1994;23:259–277. [PubMed: 8277090]
22. Stankovicova T, Szilard M, De Scheerder I, Sipido KR. M cells and transmural heterogeneity of action potential configuration in myocytes from the left ventricular wall of the pig heart. *Cardiovasc Res* 2000;45:952–960. [PubMed: 10728421]
23. Liu DW, Gintant GA, Antzelevitch C. Ionic bases for electrophysiological distinctions among epicardial, midmyocardial, and endocardial myocytes from the free wall of the canine left ventricle. *Circ Res* 1993;72:671–687. [PubMed: 8431990]
24. Sicouri S, Antzelevitch C. Drug-induced afterdepolarizations and triggered activity occur in a discrete subpopulation of ventricular muscle cell (M cells) in the canine heart: quinidine and digitalis. *J Cardiovasc Electrophysiol* 1993;4:48–58. [PubMed: 8287236]
25. Sicouri S, Fish J, Antzelevitch C. Distribution of M cells in the canine ventricle. *J Cardiovasc Electrophysiol* 1994;5:824–837. [PubMed: 7874328]
26. Weissenburger J, Nesterenko VV, Antzelevitch C. Transmural heterogeneity of ventricular repolarization under baseline and long QT conditions in the canine heart *in vivo*: Torsades de Pointes develops with halothane but not pentobarbital anesthesia. *J Cardiovasc Electrophysiol* 2000;11:290–304. [PubMed: 10749352]
27. Sicouri S, Quist M, Antzelevitch C. Evidence for the presence of M cells in the guinea pig ventricle. *J Cardiovasc Electrophysiol* 1996;7:503–511. [PubMed: 8743756]
28. Li GR, Feng J, Yue L, Carrier M. Transmural heterogeneity of action potentials and I_{to1} in myocytes isolated from the human right ventricle. *Am J Physiol* 1998;275:H369–H377. [PubMed: 9683422]
29. Rodriguez-Sinovas A, Cinca J, Tapias A, Armadans L, Tresanchez M, Soler-Soler J. Lack of evidence of M-cells in porcine left ventricular myocardium. *Cardiovasc Res* 1997;33:307–313. [PubMed: 9074694]

30. Shimizu W, Antzelevitch C. Sodium channel block with mexiletine is effective in reducing dispersion of repolarization and preventing torsade de pointes in LQT2 and LQT3 models of the long-QT syndrome. *Circulation* 1997;96:2038–2047. [PubMed: 9323097]
31. El-Sherif N, Caref EB, Yin H, Restivo M. The electrophysiological mechanism of ventricular arrhythmias in the long QT syndrome: tridimensional mapping of activation and recovery patterns. *Circ Res* 1996;79:474–492. [PubMed: 8781481]
32. Weirich J, Bernhardt R, Loewen N, Wenzel W, Antoni H. Regional- and species-dependent effects of K⁺-channel blocking agents on subendocardium and mid-wall slices of human, rabbit, and guinea pig myocardium. *Pflugers Arch* 1996;431 R 130 Abstract.
33. Burashnikov A, Antzelevitch C. Acceleration-induced action potential prolongation and early afterdepolarizations. *J Cardiovasc Electrophysiol* 1998;9:934–948. [PubMed: 9786074]
34. Shimizu W, McMahon B, Antzelevitch C. Sodium pentobarbital reduces transmural dispersion of repolarization and prevents torsade de pointes in models of acquired and congenital long QT syndrome. *J Cardiovasc Electrophysiol* 1999;10:156–164.
35. Shimizu W, Antzelevitch C. Cellular basis for the ECG features of the LQT1 form of the long QT syndrome: effects of b-adrenergic agonists and antagonists and sodium channel blockers on transmural dispersion of repolarization and torsade de pointes. *Circulation* 1998;98:2314–2322. [PubMed: 9826320]
36. Shimizu W, Antzelevitch C. Cellular and ionic basis for T-wave alternans under Long QT conditions. *Circulation* 1999;99:1499–1507. [PubMed: 10086976]
37. Balati B, Varro A, Papp JG. Comparison of the cellular electrophysiological characteristics of canine left ventricular epicardium, M cells, endocardium and Purkinje fibres. *Acta Physiol Scand* 1998;164:181–190. [PubMed: 9805105]
38. McIntosh MA, Cobbe SM, Smith GL. Heterogeneous changes in action potential and intracellular Ca²⁺ in left ventricular myocyte sub-types from rabbits with heart failure. *Cardiovasc Res* 2000;45:397–409. [PubMed: 10728360]
39. Drouin E, Charpentier F, Gauthier C, Laurent K, Le Marec H. Electrophysiological characteristics of cells spanning the left ventricular wall of human heart: Evidence for the presence of M cells. *J Am Coll Cardiol* 1995;26:185–192. [PubMed: 7797750]
40. Taggart P, Sutton PM, Opthof T, Coronel R, Trimlett R, Pugsley W, Kallis P. Transmural repolarisation in the left ventricle in humans during normoxia and ischaemia. *Cardiovasc Res* 2001;50:454–462. [PubMed: 11376621]
41. Coronel R, de Bakker JM, Wilms-Schopman FJ, Opthof T, Linnenbank AC, Belterman CN, Janse MJ. Monophasic action potentials and activation recovery intervals as measures of ventricular action potential duration: Experimental evidence to resolve some controversies. *Heart Rhythm* 2006;3:1043–1050. [PubMed: 16945799]
42. Rosen MR, Anyukhovsky EP, Sosunov EA, Gainullin RZ. The M cell. *J Cardiovasc Electrophysiol* 1999;10:1297–1299. [PubMed: 10517665]
43. Janse MJ, Sosunov EA, Coronel R, Opthof T, Anyukhovsky EP, de Bakker JM, Plotnikov AN, Shlapakova IN, Danilo P Jr, Tijssen JG, Rosen MR. Repolarization gradients in the canine left ventricle before and after induction of short-term cardiac memory. *Circulation* 2005;112:1711–1718. [PubMed: 16157774]
44. Opthof T, Coronel R, Wilms-Schopman FJ, Plotnikov AN, Shlapakova IN, Danilo P Jr, Rosen MR, Janse MJ. Dispersion of repolarization in canine ventricle and the electrocardiographic T wave: T(p-e) interval does not reflect transmural dispersion. *Heart Rhythm* 2007;4:341–348. [PubMed: 17341400]
45. Antzelevitch C. Transmural dispersion of repolarization and the T wave. *Cardiovasc Res* 2001;50:426–431. [PubMed: 11376617]
46. Fish JM, Brugada J, Antzelevitch C. Potential proarrhythmic effects of biventricular pacing. *J Am Coll Cardiol* 2005;46:2340–2347. [PubMed: 16360069]
47. Glukhov AV, Fedorov VV, Lou Q, Ravikumar V, Kalish PW, Schuessler RB, Moazami N, Efimov IR. Transmural Dispersion Of Repolarization In Failing And Non Failing Human Ventricle. *Circ Res* 2010;106:XXX–XXX.

48. Akar FG, Yan GX, Antzelevitch C, Rosenbaum DS. Unique topographical distribution of M cells underlies reentrant mechanism of torsade de pointes in the long-QT syndrome. *Circulation* 2002;105:1247–1253. [PubMed: 11889021]
49. Akar FG, Tomaselli GF. Ion channels as novel therapeutic targets in heart failure. *Ann Med* 2005;37:44–54. [PubMed: 15902846]
50. Aiba T, Tomaselli GF. Electrical remodeling in the failing heart. *Curr Opin Cardiol* 2010;25:29–36. [PubMed: 19907317]
51. Akar FG, Rosenbaum DS. Transmural electrophysiological heterogeneities underlying arrhythmogenesis in heart failure. *Circ Res* 2003;93:638–645. [PubMed: 12933704]