



Published in final edited form as:

Trends Microbiol. 2010 April ; 18(4): 173–181. doi:10.1016/j.tim.2009.12.011.

Phage WO of *Wolbachia*: lambda of the endosymbiont world

Bethany N. Kent and Seth R. Bordenstein

Department of Biological Sciences, Vanderbilt University, Nashville, TN 37235, USA

Abstract

The discovery of an extraordinarily high level of mobile elements in the genome of *Wolbachia*, a widespread arthropod and nematode endosymbiont, suggests that this bacterium could be an excellent model for assessing the evolution and function of mobile DNA in specialized bacteria. Here, we discuss how studies on the temperate bacteriophage WO of *Wolbachia* have revealed unexpected levels of genomic flux and are challenging previously held views about the clonality of obligate intracellular bacteria. We also discuss the roles that this phage might play in the *Wolbachia*-arthropod symbiosis, and infer how this research can be translated to combating human diseases vectored by arthropods. We expect that this temperate phage will be a preeminent model system to understand phage genetics, evolution, and ecology in obligate intracellular bacteria. In this sense, phage WO might be likened to phage λ of the endosymbiont world.

Mobile elements in intracellular bacteria

The restrictive lifestyle of obligate intracellular bacteria can lead to a near minimal genome state that encodes only essential functions. This reduction is associated with a genome-wide deletion bias, population bottlenecks, and relaxed selection due to the ability of the bacteria to acquire nutrients from the host cell rather than synthesize them [1,2]. As a consequence of reductive evolution, mobile DNA elements have often been shown to be rare or absent from such streamlined bacteria [3-5]. However, genome sequence data shows that mobile elements are present at sometimes high frequency in obligate intracellular bacteria that switch hosts, including *Wolbachia*, *Rickettsia*, *Coxiella*, and *Phytoplasma* [4,6-10]. Thus, past findings suggesting that streamlined bacterial genomes lack mobile DNA are being revisited with new hypotheses on how these elements invade and survive in these reduced genomes.

The tripartite arthropod-*Wolbachia*-phage WO system is emerging as a model to study the role of mobile elements in obligate intracellular bacteria. In the last few years, the publication of several complete WO sequences, the discoveries of rampant horizontal transmission between coinfections, and the tritrophic interactions between phage, *Wolbachia*, and the arthropod host have propelled the field forward and will allow for rapid advancement in the study of WO evolution, function, and activity.

The biology of bacteriophage WO

Wolbachia species are members of the obligate intracellular *Rickettsiales* and forge parasitic relationships with arthropods and mutualistic relationships primarily with nematodes. During

© 2009 Elsevier Ltd. All rights reserved.

Corresponding author: Bordenstein, S. R. (s.bordenstein@vanderbilt.edu).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

their 100-million-year association with their hosts, the maternally-transmitted bacteria have evolved as “reproductive parasites” that cause cytoplasmic incompatibility (CI, see Glossary), feminization, parthenogenesis, and male killing in arthropods, while in nematodes and some arthropods, they are mutualistic and can be required for host oogenesis or larval development [11-13]. In addition to modifying reproduction, *Wolbachia* spp. have recently been shown to confer resistance against RNA viruses [14-16], influence locomotion in response to food cues [17] and increase egg laying of females reared on low- or high-iron diets in *Drosophila* [18]. In *Asobara tabida* insects and cell lines from *Drosophila simulans* flies and *Aedes* mosquitoes, *Wolbachia* is involved in iron metabolism of the host [19]. These bacteria can also be transmitted horizontally across species, which has led to a pandemic-level distribution in invertebrates: current estimates place *Wolbachia* in 66% of all arthropod species [20].

While identification of a bacteriophage in the *Wolbachia* infection of *Culex pipiens* mosquitoes was reported in the late 1970s [21], confirmation of a *Wolbachia* phage did not occur until twenty years later when a prophage region was identified in the genome of *Wolbachia* strain *w*Tai infecting *Teleogryllus taiwanemma* crickets [22]. Screening of *Wolbachia* infections from a variety of invertebrate hosts indicate that prophage WO (named after *Wolbachia*) is widespread in the genus [23-25]. PCR amplification of the minor capsid gene *orf7* showed that the phage infects 89% of the parasitic A and B *Wolbachia* supergroups (from arthropods) but is absent in the mutualistic C and D supergroups (from nematodes) [23,24]. However, vestiges of prophage DNA remain in the C and D supergroups, suggesting that at one point in evolutionary history they may have harbored phage too. Six prophage pseudogenes in the *w*Bm genome from the nematode *Brugia malayi* (Wbm5005, Wbm5030, Wbm5039, Wbm5040, Wbm5044, Wbm5080) are homologous to genes in A and B supergroup *Wolbachia*, as well as *Wolbachia*'s relatives *Rickettsia*, *Ehrlichia*, and *Anaplasma*. These genes are not part of phage WO. One additional phage pseudogene, Wbm5055, is homologous to a conserved hypothetical protein gene found in WO prophages from *w*Pip (WP1304), *w*Kue (gp17), and *w*Ri (WRi_007190), as well as in non-WO regions in *w*Mel and other locations in *w*Ri. Distribution of WO in arthropods might be greater than the current estimates, as the primers used to screen for presence or *orf7* were not degenerate enough to detect all of the *orf7* variants: for example, *Wolbachia* strain *w*Ri of *Drosophila simulans* was initially reported as having a single WO haplotype [24], but the genome sequence confirmed four prophage copies in the genome, three of which are unique [8].

Icosahedral WO phage particles have been purified from several arthropods harboring *Wolbachia* infections (Table 1). The virion heads range in size from 20 to 40 nm and, occasionally, a tail structure has been identified by transmission electron microscopy (TEM) [26,27]. Although initially reported to be a linear double-stranded DNA (dsDNA) molecule [28], the amplification of adjoined *att* sequences using inverse PCR demonstrated that the WO genome is circular [29] and replicates similarly to phage λ .

WO is a dynamic element that has a significant impact on the genetic diversity of *Wolbachia*. Complete sequences are known for prophages in strains *w*Mel [10], *w*Ri [8], *w*Kue [22], *w*CauB [28,29], and *w*Pip [30]. Analysis shows that WO molecular evolution is a complex process involving vertical transmission and horizontal transfer, recombination between phages, and hitchhiking of other mobile elements on WO. Although the details of phage transfer have not been discerned, evidence supports the potential movement of active WO particles between both related and divergent *Wolbachia* cells: electron microscopy shows particles aggregating outside of the *Wolbachia* cells after lysis and adjacent to developing spermatids in wasp testes [27]. Horizontal transfer of WO could occur either between multiple *Wolbachia* infections in a single host (Figure 1a) or, hypothetically, by paternal transmission of phage particles to a fertilized egg that harbors a phage-free *Wolbachia* strain (Figure 1b). Polymerase chain reaction (PCR) studies suggest that phage DNA might be transferred paternally from infected

males to uninfected females (S.R. Bordenstein, unpublished). WO horizontal gene transfer is strongly suggested by the fact that some divergent *Wolbachia* strains that coinfect the same host have identical *orf7* sequences [22,23,25,31]. This mechanism of transfer can be likened to an “intracellular arena” in which the host cell acts as a chemostat for phage transfers between *Wolbachia* coinfections [4,23]. Divergent phage haplotypes can also emerge from duplication of one initial viral infection. The two phage copies in *wCauB* are more similar to each other than to any other known WO sequence [29] and similarity for each of the five types in *wPip* is highest between another WO prophage in the same genome [30]. However, it appears that some genes in the phage haplotypes inhabiting the same genome have been acquired from WO phages from other *Wolbachia* strains.

The flux of prophage WO genomes is also supported by intragenic recombination between different phage haplotypes. The nucleotide sequence of the minor capsid gene *orf7* from strain *wKueA1* is chimeric, and population genetic analysis confirms the recombinogenic nature of WO [23]. The recombination rate of *orf7* is twelvefold greater than that of the rapidly evolving *Wolbachia* surface protein gene *wsp*, and fifteen-fold greater than the WO terminase-encoding gene *orf2*. Why these phage genes (*orf7* and *orf2*), which are located less than five kilobases apart and required for lytic phage production, are recombining at different rates remains to be determined.

The contribution of prophage WO to *Wolbachia* genetic diversity is not limited to phage-associated genes. Insertion sequences (IS) are frequently found in WO genomes, including IS3, IS4, IS5, IS6, IS110, and IS630 family elements, [8,10,30] and might be a major factor driving phage recombination [32,33]. Genes encoding transposases are present in nearly all sequenced WO prophage genomes and can laterally transfer between *Wolbachia* strains [34]. If the presence of these IS elements on WO does not hinder its lytic ability, they could hitchhike within the phage genome as it spreads to new cells and move to new locations in the newly infected host genome [35]. IS elements are responsible for a significant amount of genetic diversity between many *Wolbachia* strains; for example, in *wPip*, they truncate 44 genes [30].

Evolution of the WO core genome

In dsDNA phages of bacteria with a free-living replicative stage, evolution is categorized by the Modular Theory [32,36,37]. According to this theory, a phage genome can be divided into functional units or modules (each one responsible for head or tail formation, lysis, lysogeny, etc.), which can be mixed by recombination with other phages. Each module is often comprised of genes that have a shared evolutionary history owing to their physical linkage and functional coadaptation. Generalizing the principles of the Modular Theory to all dsDNA phages will require an expanded analysis in diverse ecological ranges [37]. In this regard, it seems opportune to test the theory on phages from obligate intracellular bacteria, as the intracellular niche may pose natural restraints on exposure to novel phage gene pools. While extensive recombination would be expected in phages that are exposed to a multitude of other phages, recombination between unrelated phages that have a limited niche environment would confirm that the Modular Theory holds even in the obligate intracellular bacteria.

The identification of phage termini for two WO phages (*WOCauB2* and *WOCauB3*) [29] and the complete sequences of 12 other active phages and prophages allow for an assessment of the WO core genome and an understanding of the molecular mechanisms by which these phages evolve. Modules for the assembly of head, baseplate and tail are readily identifiable based on gene homologies to the related lambdoid and P2 phages [10,22] (Figure 2), although tail module genes are present in only about half of the known WO sequences. Interspersed among the modules are genes of unknown function and others encoding putative virulence factors, transposases, and ankyrin repeat (ANK) proteins [8,10,29,30,38]. It is likely that the conserved

genes coding for hypothetical proteins located within specific modules are functionally required.

WO gene homologs occur in a diverse set of bacteria including *Alpha*-, *Beta*-, and *Gammaproteobacteria*, *Firmicutes*, *Cyanobacteria*, and *Bacteroidetes*, which could indicate that these phage genes have been transferred among multiple phyla [29]. Evidence that WO can acquire genes from, or transmit genes to, bacteria other than *Wolbachia* is apparent by the 70% nucleotide identity between prophage genes in *wMel* and a segment of a *Rickettsial* plasmid [38]. WO homologs also occur in the prophage regions of the facultative intracellular parasite *Bartonella henselae* [39]. Current data indicates that while phage WO genomes are modular, the functional gene modules do not readily exchange DNA with unrelated phages, as WO genes often have the highest sequence similarity to genes in other WO phages. The rarity of modular exchange with unrelated phages is likely due to the unique intracellular niche that phage WO occupies. Instead, evolutionary forces such as point mutation, deletions, recombination, and inversions tend to be the dominant modes of diversification. Notably, these modes of phage diversification still cause some of the largest fractions of absent or divergent genes between closely related *Wolbachia* genomes [38].

Lifecycle of phage WO

While the lytic and lysogenic nature of temperate phage WO has been demonstrated, the genetic mechanisms that drive prophage induction and lytic activity are currently unknown. Phage particles have been visualized for several *Wolbachia* strains (Table 1) but the precise identification of phage termini has occurred only from phages of the *Wolbachia* infection of *Cadra cautella* moths [29]. A serine-recombinase gene homolog that is likely to be responsible for integration of these phages is located at the termini of phages WOCauB2 (Figure 2) and WOCauB3, but the exact binding sites of the recombinase are unknown. These WO phages are not flanked by an inverted repeat, and comparison of the joined ends of the active phage genome (*attP*) with the integrated prophage terminal sequences (*attR* and *attL*) and flanking *Wolbachia* sequences (*attB*) found in common only a single nucleotide T in WOCauB2 and a trinucleotide TTG in WOCauB3 [29]. The serine recombinase gene found in the WOCauB phages is different to that of other sequenced WO prophages, with the exception of one *wRi* phage (*WRi_012450*), indicating that the other WO phages use a different site-specific recombination mechanism or are degenerate.

Expression of phage genes can be sex-specific and age-specific relative to the host arthropod. The expression of the minor capsid protein gene *orf7* was compared between developmental stages and sexes in *Wolbachia* from three different populations of *Culex pipiens* and one from *Culex quinquefasciatus* [26]. Adult females from all four *Culex* strains expressed *orf7*, while adult males of only two strains did the same. Expression also varied among developmental stages. In all four populations, eggs and early larval stages expressed *orf7*, but expression in later-stage larvae varied in the populations from no expression to strong expression. Strong expression returned by the pupal stage in three of the four strains. This evidence suggests that the biology of phage WO is closely linked to that of the arthropod host, hypothetically through direct interaction between host-encoded proteins and proteins encoded on the phage (ANK proteins, for example) or through changes that *Wolbachia* undergo because of the insect that have a downstream affect on phage WO. Studies using *Nasonia* parasitoid wasps have recently found that lytic phage production appears to be influenced by multiple abiotic and biotic factors including insect age, host species background, and temperature (S.R. Bordenstein, unpublished).

Effectors, toxins, and the tripartite relationship of WO, *Wolbachia*, and arthropods

In addition to the genomics and transmissibility of bacteriophage WO, there is considerable interest in the phage's function in the *Wolbachia*-arthropod symbiosis. Mobile elements were initially hypothesized to be responsible for causing reproductive parasitism because they can promiscuously transfer new functions between strains, potentially explaining the phylogenetic curiosity that different types of reproductive parasitism (CI, male killing, feminization, or parthenogenesis) do not cluster in groups in the *Wolbachia* tree. Analysis of WO genome sequences has identified several candidate proteins for interaction with eukaryotic cells (Table 2). Genes having homology to *vrlA* and *vrlC*, virulence-associated genes [40] on a pathogenicity island of the sheep pathogen *Dichelobacter nodosus*, are located in WO prophage sequences [28]. In *D. nodosus*, VrlC contains a conserved motif typically associated with sialidases [40]. A sialidase, which cleaves glycoconjugates on cell surfaces, might be involved in pathogenesis or *Wolbachia*'s ability to scavenge nutrients from the host cell as seen in a diverse range of bacteria such as *Pasteurellaceae* [41] and some *Mycoplasma* [42,43]. Notably, a weak correlation between sequence variability of VrlC and CI was observed in *Culex pipiens* mosquitoes [44].

Seven WO prophages also contain a homolog of a *Rickettsia* gene coding for a patatin-like phospholipase belonging to the phospholipase A₂ family. Proteins containing patatin-like domains have been linked to virulence in *Legionella pneumophila* [45] and *Pseudomonas aeruginosa* [46-48]. It is possible that the WO homolog might assist *Wolbachia* entry into host cells or be involved in other arthropod host cell interactions [29]. Further, the *wRi*, *wMel* and *wPip* prophages include a gene that might be part of a toxin/antitoxin system, by functioning as an 'addiction' module (i.e. killing those cells lacking the mobile element [49]). Homologs of this gene are also found elsewhere within the *Wolbachia* genome.

WOCauB3 contains a gene encoding a protein similar to *Salmonella enterica* ADP-ribosylating toxin, SpvB [50-52]. The WO homolog shows some similarity to bacterial proteins of the RHS and YD-repeat families, which have been implicated in interactions with eukaryotic host cells [53,54]. A protein with similar motifs is present in APSE, the bacteriophage of *Hamiltonella defensa* that confers pea aphid resistance to parasitoid attack [55,56]. Additionally, the SpvB homolog has sequence homology to a family of insecticidal toxins [29]. The diversity of putative effector proteins and toxins encoded among prophage genomes suggests that WO is involved in many important facets of *Wolbachia* biology. How the phage's effectors and toxins affect its host will be an important future topic of study.

While the identity of virulence factors varies between phage copies, genes coding for ankyrin-repeat (ANK) proteins are found in large number in *Wolbachia* genomes, and particularly in the vicinity of, or encoded on, WO [8,10,30,57]. ANKs are protein motifs that can mediate protein-protein interactions, act as transcription factors, and modify the activity of cell-cycle regulatory proteins in eukaryotes [58-60]. ANK proteins often contain transmembrane domains or signal peptides, offering two different mechanisms by which they can interact with the host cells: surface expression, or secretion into the insect cell. Because eukaryotic ANK proteins span functions involved in a variety of cell processes, *Wolbachia* ANK proteins have been hypothesized to have a role in reproductive parasitism.

The expression of ANK proteins was compared between *wMel*, which causes CI in *Drosophila melanogaster*, and the closely-related *wAu*, which infects *Drosophila simulans* but does not cause CI [61]. It was found that one ANK gene is absent in *wAu* but present in *wMel*, and the sequence of seven genes encoding ANK proteins in *wAu* differed significantly from that of the *wMel* homologs. Of these seven, one showed a difference in expression due to a gene truncation

caused by an insertion sequence within the open reading frame. Two of these ANK genes are located in the prophage region of *wMel* [62]. These differences in ANK protein expression and structure are candidates for the inability of *wAu* to cause sexual alteration. In *Culex* mosquitoes, two ANK genes associated with a prophage region are expressed sex-specifically and have sequence divergence associated with CI [63]. However, other genes such as *orf7* show sex specific expression even though there is no association of sequence variability with CI [26]. Although it is tempting to speculate that WO ANK genes could be involved in *Wolbachia*-host interactions, current evidence for this is complex and insufficient.

The Phage Density Model

Phage sequence analyses find no phylogenetic clustering of WO genotypes among the four major *Wolbachia*-induced sexual alterations. Further, some *Wolbachia* strains exist that induce CI, male killing, or parthenogenesis but lack the WO prophage [23,24]. These facts led to the development of the Phage Density Model as an alternative, but not mutually exclusive, explanation for the role of phage WO in sexual alterations (Figure 3) [27]. This theory proposes that variations in phage lysis are linked to the expression of sexual alterations through variations in *Wolbachia* densities. According to the model, lytic phages kill *Wolbachia* cells and thereby reduce bacterial densities in the tissues associated with reproductive modification. Because bacterial density is one of the most critical determinants of expression of *Wolbachia* functions [64-67], its variation affects the expression of sexual alterations. Evidence from TEM observations and quantitative studies of phage-*Wolbachia* interactions in *Nasonia vitripennis* wasps show particles exiting lysed cells into host reproductive tissues, and an inverse association of phage and *Wolbachia* titers [27]. A supergroup-B strain of *Nasonia vitripennis* with a mean phage density of less than two (estimated as the relative number of copies of the WO *orf7* gene in relation to those of *Wolbachia* gene *groEL*) exhibited 100% CI, whereas a supergroup-A strain with a mean phage density of six exhibited a decreased level of 67.7% CI. Among the supergroup-A infected males that showed variation in CI levels, males with complete CI had significantly lower phage densities than males expressing incomplete CI.

The model and evidence emphasize that the first tenet of phage function is that phages are parasites of bacteria, and that clarifying the separate roles of lytic and lysogenic phage development in *Wolbachia* biology will effectively structure inquiries into the function of phage WO. Currently, the weight of the experimental evidence suggests that the lytic phage is a mobile genetic parasite that can reduce *Wolbachia* densities, while the role of the lysogenic prophage remains a topic of future interest. Further, not all systems will harbor phage WO or show the same patterns as that observed in *Nasonia vitripennis*. For instance, the phage density model does not appear to be supported in *Culex* mosquitoes [68], but sample sizes in this study were too small to rule it out.

Biomedical applications for phage WO

Wolbachia has become increasingly important to human health and disease through two routes. First, vector control programs aimed at curbing the spread of insect-vector-borne diseases such as malaria and dengue fever will rely on the ability to release insect vectors transfected with *Wolbachia* infections that reduce the vectorial competence [69-74]. Second, the discoveries that river blindness, lymphatic filariasis [75] and heartworm [76] are associated with *Wolbachia*-induced pathologies raised the likelihood that antimicrobial therapies targeting *Wolbachia* may be effective in treatment of the systems [76-79]. For insect diseases, current strategies include (i) using *Wolbachia* to carry a transgene that would inhibit spread of the infectious agent [80,81], (ii) infecting mosquitoes with a life-shortening strain of *Wolbachia* so that the insect vector dies before it is capable of transmitting disease [71,82-84], or (iii)

releasing *Wolbachia*-infected males into the wild so that CI will inhibit mosquito reproduction and cause the native populations to crash to less-threatening levels [85,86]. In order for the first strategy to be effective, population replacement of natural vectors with transgenically modified ones must be rapid and overcome fitness costs associated with the transgene [81]. The speed with which *Wolbachia* can spread through a population due to its impact on host reproduction makes it an ideal method to transmit a transgene in wild populations of insects [81,87-89].

The ability for *Wolbachia* to be used in biological control of diseases is dependent on successful infection of mosquito vectors. In the first steps towards showing that a mosquito line for population replacement could be generated, the dengue vector *Aedes aegypti* was successfully infected with life-shortening *Wolbachia* strain *wMelPop* [73,90]. Under laboratory conditions, the lifespan of *Aedes aegypti* were halved after infection with *wMelPop* [71], and in older mosquitoes, a reduction in the ability to blood feed was noted [91], suggesting that a release of *wMelPop* into wild populations of mosquitoes could significantly reduce dengue transmission.

Strategies to combat malaria have shown less promise. While cell lines of the malaria vector *Anopheles gambiae* could be infected with *wMelPop* [92], the infection was avirulent and was not vertically transmitted to the next generation in adult mosquitoes [70]. In theory, mosquitoes infected with a transgenic *Wolbachia* expressing a protein that could hinder the transmission of malaria to humans would be an effective way to decrease the spread of the disease. Unfortunately, there are currently no tools available for the genetic manipulation of *Wolbachia* and no reports of successful transformation. Phage WO particles could be developed into the first transgenic tool of *Wolbachia*, as active phages might succeed in vectoring transgenes to recipient *Wolbachia* cells where traditional transformation strategies have failed. Additionally, the integrase and *att* sites in WO could be used to construct vector systems capable of integrating into the *Wolbachia* genome [29]. Once integration of mini-WO constructs is successful, phage WO could also be used as a knockout insertion element so that candidate genes underlying *Wolbachia* traits such as reproductive parasitism could be rapidly identified.

Concluding remarks and future directions

Studies of phage WO have shown its potential to have a substantial impact on the symbiosis between *Wolbachia* and host arthropods. Extrapolation of the phage infection frequency places WO in potentially millions of insect species, where it can contribute to *Wolbachia* genomic diversity and function in a number of ways, including horizontal gene transfer between different *Wolbachia* as well other endosymbionts, exchange of other mobile elements such as insertion sequences, intragenic recombination, gene loss, and an alteration of *Wolbachia* densities that can affect the penetrance of reproductive parasitism. The phenotypic impacts on the eukaryotic host cell could be equally diverse. WO encodes several different classes of proteins, such as virulence factor homologs and ANK proteins, which could influence *Wolbachia* or the arthropod host. Additionally, lytic WO decreases reproductive parasitism in *Nasonia* by lowering *Wolbachia* densities, but whether this relationship holds in other *Wolbachia*-arthropod relationships remains to be determined. Beyond *Wolbachia* lysis, the precise mechanisms by which the phage interacts with the invertebrate host is a topic of interest.

There is hope that phage WO particles may overcome current barriers to transform *Wolbachia* or that functional aspects of WO integration could be used to generate vector systems capable of supplying transgenes into *Wolbachia* genomes. Successful transformation of *Wolbachia* must first be demonstrated before a vector system would be of use.

Study of the biology of WO would not only expand knowledge of phage evolution and function, but could also lead to a role of this phage in the treatment of insect-vectored diseases. Therefore, future research on phage WO should encompass a wide variety of themes, including endosymbiosis, phage biology, and arthropod vector control (Box 1). As the study of new phages in diverse ecological niches necessitates new models akin to the well-studied phage λ , phage WO seems an adequate model for obligate intracellular bacteria.

Box 1. Questions for future research

Evolution

- Does the lack of modular exchange typify phage WO genome evolution?
- How common is phage WO across the *Wolbachia* genus?

Ecology

- What are the common mechanisms by which phage particles transfer?
- Do phage WO sequences cluster with geography or host range?
- What genetic factors regulate the temperate lifecycle of phage WO?

Host interaction

- Is WO retained due to a benefit to the *Wolbachia* bacterium or to the host arthropod?
- How applicable is the phage density model to other *Wolbachia*-arthropod systems?
- What role do the encoded effector proteins play in interactions with bacterial and eukaryotic cells?

Applications

- Can WO be used as a transgenic vector system either through active phage particles or through a mini-integration construct?

Acknowledgments

We thank Robert Brucker and Meghan Chafee for helpful feedback on the manuscript. This work was supported by grants NSF IOS-0852344 and NIH R01 GM085163-01 to S.R.B. This paper's contents are solely the responsibility of the authors and do not necessarily represent the official view of the NSF or NIH.

Glossary

Cytoplasmic incompatibility (CI):	a sperm-egg incompatibility that renders embryos inviable in crosses between infected males and uninfected females or females harboring a different strain of <i>Wolbachia</i> .
Feminization:	the process by which infected male embryos are converted to morphological and functional females.
Haplotype:	a distinct WO prophage type, based on nucleotide differences.
Male killing:	the process by which male embryos or larvae are preferentially killed relative to female ones.
Parthenogenesis:	a form of asexual reproduction where only female offspring are produced by infected females.

Phage termini: the left and right terminal ends of the phage at which points the phage is integrated into the host genome. Upon excision of phage WO, these ends will join such that the phage genome is circular.

References


1. Moran NA, et al. Genomics and evolution of heritable bacterial symbionts. *Annu Rev Genet* 2008;42:165–190. [PubMed: 18983256]
2. Wernegreen JJ. Genome evolution in bacterial endosymbionts of insects. *Nat Rev Genet* 2002;3:850–861. [PubMed: 12415315]
3. Moran NA, Plague GR. Genomic changes following host restriction in bacteria. *Curr Opin Genet Dev* 2004;14:627–633. [PubMed: 15531157]
4. Bordenstein SR, Reznikoff WS. Mobile DNA in obligate intracellular bacteria. *Nat Rev Microbiol* 2005;3:688–699. [PubMed: 16138097]
5. Touchon M, Rocha EP. Causes of insertion sequences abundance in prokaryotic genomes. *Mol Biol Evol* 2007;24:969–981. [PubMed: 17251179]
6. Ogata H, et al. The genome sequence of *Rickettsia felis* identifies the first putative conjugative plasmid in an obligate intracellular parasite. *PLoS Biol* 2005;3:e248. [PubMed: 15984913]
7. Wei W, et al. Ancient, recurrent phage attacks and recombination shaped dynamic sequence-variable mosaics at the root of phytoplasma genome evolution. *Proc Natl Acad Sci U S A* 2008;105:11827–11832. [PubMed: 18701718]
8. Klasson L, et al. The mosaic genome structure of the *Wolbachia* wRi strain infecting *Drosophila simulans*. *Proc Natl Acad Sci U S A* 2009;106:5725–5730. [PubMed: 19307581]
9. Wernegreen JJ. For better or worse: genomic consequences of intracellular mutualism and parasitism. *Curr Opin Genet Dev* 2005;15:572–583. [PubMed: 16230003]
10. Wu M, et al. Phylogenomics of the reproductive parasite *Wolbachia pipientis* wMel: a streamlined genome overrun by mobile genetic elements. *PLoS Biol* 2004;2:E69. [PubMed: 15024419]
11. Werren JH, et al. *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol* 2008;6:741–751. [PubMed: 18794912]
12. Dedeine F, et al. *Wolbachia* requirement for oogenesis: occurrence within the genus *Asobara* (Hymenoptera, Braconidae) and evidence for intraspecific variation in *A. tabida*. *Heredity* 2005;95:394–400. [PubMed: 16118660]
13. Dedeine F, et al. Intra-individual coexistence of a *Wolbachia* strain required for host oogenesis with two strains inducing cytoplasmic incompatibility in the wasp *Asobara tabida*. *Evolution* 2004;58:2167–2174. [PubMed: 15562683]
14. Hedges LM, et al. *Wolbachia* and virus protection in insects. *Science* 2008;322:702. [PubMed: 18974344]
15. Teixeira L, et al. The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol* 2008;6:e2. [PubMed: 19222304]
16. Osborne SE, et al. Variation in antiviral protection mediated by different *Wolbachia* strains in *Drosophila simulans*. *PLoS Pathog* 2009;5:e1000656. [PubMed: 19911047]
17. Peng Y, et al. *Wolbachia* infection alters olfactory-cued locomotion in *Drosophila* spp. *Appl Environ Microbiol* 2008;74:3943–3948. [PubMed: 18456851]
18. Brownlie JC, et al. Evidence for metabolic provisioning by a common invertebrate endosymbiont, *Wolbachia pipientis*, during periods of nutritional stress. *PLoS Pathog* 2009;5:e1000368. [PubMed: 19343208]
19. Kremer N, et al. *Wolbachia* interferes with ferritin expression and iron metabolism in insects. *PLoS Pathog* 2009;5:e1000630. [PubMed: 19851452]
20. Hilgenboecker K, et al. How many species are infected with *Wolbachia*? -- A statistical analysis of current data. *FEMS Microbiol Lett* 2008;281:215–220. [PubMed: 18312577]

21. Wright JD, et al. The ultrastructure of the rickettsia-like microorganism *Wolbachia pipientis* and associated virus-like bodies in the mosquito *Culex pipiens*. *J Ultrastruct Res* 1978;63:79–85. [PubMed: 671578]
22. Masui S, et al. Distribution and evolution of bacteriophage WO in *Wolbachia*, the endosymbiont causing sexual alterations in arthropods. *J Mol Evol* 2000;51:491–497. [PubMed: 11080372]
23. Bordenstein SR, Wernegreen JJ. Bacteriophage flux in endosymbionts (*Wolbachia*): infection frequency, lateral transfer, and recombination rates. *Mol Biol Evol* 2004;21:1981–1991. [PubMed: 15254259]
24. Gavotte L, et al. A survey of the bacteriophage WO in the endosymbiotic bacteria *Wolbachia*. *Mol Biol Evol* 2007;24:427–435. [PubMed: 17095536]
25. Gavotte L, et al. Diversity, distribution and specificity of WO phage infection in *Wolbachia* of four insect species. *Insect Mol Biol* 2004;13:147–153. [PubMed: 15056362]
26. Sanogo YO, Dobson SL. WO bacteriophage transcription in *Wolbachia*-infected *Culex pipiens*. *Insect Biochem Mol Biol* 2006;36:80–85. [PubMed: 16360953]
27. Bordenstein SR, et al. The tripartite associations between bacteriophage, *Wolbachia*, and arthropods. *PLoS Pathog* 2006;2:e43. [PubMed: 16710453]
28. Fujii Y, et al. Isolation and characterization of the bacteriophage WO from *Wolbachia*, an arthropod endosymbiont. *Biochem Biophys Res Commun* 2004;317:1183–1188. [PubMed: 15094394]
29. Tanaka K, et al. Complete WO phage sequences revealed their dynamic evolutionary trajectories and putative functional elements required for integration into *Wolbachia* genome. *Appl Environ Microbiol* 2009;75:5676–5686. [PubMed: 19592535]
30. Klasson L, et al. Genome evolution of *Wolbachia* strain wPip from the *Culex pipiens* group. *Mol Biol Evol* 2008;25:1877–1887. [PubMed: 18550617]
31. Chauvatcharin N, et al. Bacteriophage WO-B and *Wolbachia* in natural mosquito hosts: infection incidence, transmission mode and relative density. *Mol Ecol* 2006;15:2451–2461. [PubMed: 16842419]
32. Hatfull GF. Bacteriophage genomics. *Curr Opin Microbiol* 2008;11:447–453. [PubMed: 18824125]
33. Abedon ST. Phage evolution and ecology. *Adv Appl Microbiol* 2009;67:1–45. [PubMed: 19245935]
34. Cordaux R, et al. Intense transpositional activity of insertion sequences in an ancient obligate endosymbiont. *Mol Biol Evol* 2008;25:1889–1896. [PubMed: 18562339]
35. Reznikoff WS, et al. Comparative sequence analysis of IS50/Tn5 transposase. *J Bacteriol* 2004;186:8240–8247. [PubMed: 15576772]
36. Botstein D. A theory of modular evolution for bacteriophages. *Ann N Y Acad Sci* 1980;354:484–490. [PubMed: 6452848]
37. Hendrix RW. Bacteriophage genomics. *Curr Opin Microbiol* 2003;6:506–511. [PubMed: 14572544]
38. Ishmael N, et al. Extensive genomic diversity of closely related *Wolbachia* strains. *Microbiology* 2009;155:2211–2222. [PubMed: 19389774]
39. Alsmark CM, et al. The louse-borne human pathogen *Bartonella quintana* is a genomic derivative of the zoonotic agent *Bartonella henselae*. *Proc Natl Acad Sci U S A* 2004;101:9716–9721. [PubMed: 15210978]
40. Billington SJ, et al. Complete nucleotide sequence of the 27-kilobase virulence related locus (*vrl*) of *Dichelobacter nodosus*: evidence for extrachromosomal origin. *Infect Immun* 1999;67:1277–1286. [PubMed: 10024571]
41. Severi E, et al. Sialic acid utilization by bacterial pathogens. *Microbiology* 2007;153:2817–2822. [PubMed: 17768226]
42. May M, et al. Sialidase activity in *Mycoplasma synoviae*. *Avian Dis* 2007;51:829–833. [PubMed: 18251389]
43. Brown DR, et al. Spreading factors of *Mycoplasma alligatoris*, a flesh-eating mycoplasma. *J Bacteriol* 2004;186:3922–3927. [PubMed: 15175306]
44. Duron O, et al. Hypervariable prophage WO sequences describe an unexpected high number of *Wolbachia* variants in the mosquito *Culex pipiens*. *Proc Biol Sci* 2006;273:495–502. [PubMed: 16615218]


45. Aurass P, et al. *bdhA-patD* operon as a virulence determinant, revealed by a novel large-scale approach for identification of *Legionella pneumophila* mutants defective for amoeba infection. *Appl Environ Microbiol* 2009;75:4506–4515. [PubMed: 19411431]
46. Tam C, et al. Mutation of the phospholipase catalytic domain of the *Pseudomonas aeruginosa* cytotoxin ExoU abolishes colonization promoting activity and reduces corneal disease severity. *Exp Eye Res* 2007;85:799–805. [PubMed: 17905228]
47. Pankhaniya RR, et al. *Pseudomonas aeruginosa* causes acute lung injury via the catalytic activity of the patatin-like phospholipase domain of ExoU. *Crit Care Med* 2004;32:2293–2299. [PubMed: 15640644]
48. Sato H, et al. The mechanism of action of the *Pseudomonas aeruginosa*-encoded type III cytotoxin, ExoU. *EMBO J* 2003;22:2959–2969. [PubMed: 12805211]
49. Engelberg-Kulka H, Glaser G. Addiction modules and programmed cell death and antideath in bacterial cultures. *Annu Rev Microbiol* 1999;53:43–70. [PubMed: 10547685]
50. Fang FC, et al. Growth regulation of a *Salmonella* plasmid gene essential for virulence. *J Bacteriol* 1991;173:6783–6789. [PubMed: 1938884]
51. Grob P, Guiney DG. In vitro binding of the *Salmonella dublin* virulence plasmid regulatory protein SpvR to the promoter regions of *spvA* and *spvR*. *J Bacteriol* 1996;178:1813–1820. [PubMed: 8606153]
52. Guiney DG, et al. Growth-phase regulation of plasmid virulence genes in *Salmonella*. *Trends Microbiol* 1995;3:275–279. [PubMed: 7551642]
53. Hill CW, et al. Rhs elements of *Escherichia coli*: a family of genetic composites each encoding a large mosaic protein. *Mol Microbiol* 1994;12:865–871. [PubMed: 7934896]
54. Feulner G, et al. Structure of the *rhsA* locus from *Escherichia coli* K-12 and comparison of *rhsA* with other members of the *rhs* multigene family. *J Bacteriol* 1990;172:446–456. [PubMed: 2403547]
55. Degnan PH, Moran NA. Diverse phage-encoded toxins in a protective insect endosymbiont. *Appl Environ Microbiol* 2008;74:6782–6791. [PubMed: 18791000]
56. Oliver KM, et al. Bacteriophages encode factors required for protection in a symbiotic mutualism. *Science* 2009;325:992–994. [PubMed: 19696350]
57. Duron O, et al. Variability and expression of ankyrin domain genes in *Wolbachia* variants infecting the mosquito *Culex pipiens*. *J Bacteriol* 2007;189:4442–4448. [PubMed: 17449622]
58. Caturegli P, et al. *ankA*: an *Ehrlichia phagocytophila* group gene encoding a cytoplasmic protein antigen with ankyrin repeats. *Infect Immun* 2000;68:5277–5283. [PubMed: 10948155]
59. Elfring LK, et al. *Drosophila* PLUTONIUM protein is a specialized cell cycle regulator required at the onset of embryogenesis. *Mol Biol Cell* 1997;8:583–593. [PubMed: 9247640]
60. Li J, et al. Ankyrin repeat: a unique motif mediating protein-protein interactions. *Biochemistry* 2006;45:15168–15178. [PubMed: 17176038]
61. James AC, Ballard JW. Expression of cytoplasmic incompatibility in *Drosophila simulans* and its impact on infection frequencies and distribution of *Wolbachia pipientis*. *Evolution* 2000;54:1661–1672. [PubMed: 11108593]
62. Iturbe-Ormaetxe I, et al. Distribution, expression, and motif variability of ankyrin domain genes in *Wolbachia pipientis*. *J Bacteriol* 2005;187:5136–5145. [PubMed: 16030207]
63. Sinkins SP, et al. *Wolbachia* variability and host effects on crossing type in *Culex* mosquitoes. *Nature* 2005;436:257–260. [PubMed: 16015330]
64. Breeuwer JA, Werren JH. Cytoplasmic incompatibility and bacterial density in *Nasonia vitripennis*. *Genetics* 1993;135:565–574. [PubMed: 8244014]
65. Clark ME, et al. *Wolbachia* distribution and cytoplasmic incompatibility during sperm development: the cyst as the basic cellular unit of CI expression. *Mech Dev* 2003;120:185–198. [PubMed: 12559491]
66. Noda H, et al. Infection density of *Wolbachia* and incompatibility level in two planthopper species, *Laodelphax striatellus* and *Sogatella furcifera*. *Insect Biochem Mol Biol* 2001;31:727–737. [PubMed: 11267910]
67. Poinot D, et al. *Wolbachia* transfer from *Drosophila melanogaster* into *D. simulans*: Host effect and cytoplasmic incompatibility relationships. *Genetics* 1998;150:227–237. [PubMed: 9725842]

68. Walker T, et al. *Wolbachia* in the *Culex pipiens* group mosquitoes: introgression and superinfection. *J Hered* 2009;100:192–196. [PubMed: 18836146]
69. Brownstein JS, et al. The potential of virulent *Wolbachia* to modulate disease transmission by insects. *J Invertebr Pathol* 2003;84:24–29. [PubMed: 13678709]
70. Jin C, et al. The virulent *Wolbachia* strain wMelPop efficiently establishes somatic infections in the malaria vector *Anopheles gambiae*. *Appl Environ Microbiol* 2009;75:3373–3376. [PubMed: 19329661]
71. McMeniman CJ, et al. Stable introduction of a life-shortening *Wolbachia* infection into the mosquito *Aedes aegypti*. *Science* 2009;323:141–144. [PubMed: 19119237]
72. Rasgon J. Population replacement strategies for controlling vector populations and the use of *Wolbachia pipientis* for genetic drive. *J Vis Exp* 2007:225. [PubMed: 18979023]
73. Ruang-Areerate T, Kittayapong P. *Wolbachia* transinfection in *Aedes aegypti*: a potential gene driver of dengue vectors. *Proc Natl Acad Sci U S A* 2006;103:12534–12539. [PubMed: 16895998]
74. Kambris Z, et al. Immune activation by life-shortening *Wolbachia* and reduced filarial competence in mosquitoes. *Science* 2009;326:134–136. [PubMed: 19797660]
75. Taylor MJ. *Wolbachia* in the inflammatory pathogenesis of human filariasis. *Ann N Y Acad Sci* 2003;990:444–449. [PubMed: 12860672]
76. McCall JW, et al. Heartworm disease in animals and humans. *Adv Parasitol* 2008;66:193–285. [PubMed: 18486691]
77. Hoerauf A. Filariasis: new drugs and new opportunities for lymphatic filariasis and onchocerciasis. *Curr Opin Infect Dis* 2008;21:673–681. [PubMed: 18978537]
78. Enk CD. Onchocerciasis--river blindness. *Clin Dermatol* 2006;24:176–180. [PubMed: 16714198]
79. Shakya S, et al. Prior killing of intracellular bacteria *Wolbachia* reduces inflammatory reactions and improves antifilarial efficacy of diethylcarbamazine in rodent model of *Brugia malayi*. *Parasitol Res* 2008;102:963–972. [PubMed: 18180957]
80. Xi Z, et al. Generation of a novel *Wolbachia* infection in *Aedes albopictus* (Asian tiger mosquito) via embryonic microinjection. *Insect Biochem Mol Biol* 2005;35:903–910. [PubMed: 15944085]
81. Turelli M, Hoffmann AA. Microbe-induced cytoplasmic incompatibility as a mechanism for introducing transgenes into arthropod populations. *Insect Mol Biol* 1999;8:243–255. [PubMed: 10380108]
82. Rasgon JL, Scott TW. *Wolbachia* and cytoplasmic incompatibility in the California *Culex pipiens* mosquito species complex: parameter estimates and infection dynamics in natural populations. *Genetics* 2003;165:2029–2038. [PubMed: 14704183]
83. Min KT, Benzer S. *Wolbachia*, normally a symbiont of *Drosophila*, can be virulent, causing degeneration and early death. *Proc Natl Acad Sci U S A* 1997;94:10792–10796. [PubMed: 9380712]
84. McGraw EA, et al. *Wolbachia* density and virulence attenuation after transfer into a novel host. *Proc Natl Acad Sci U S A* 2002;99:2918–2923. [PubMed: 11880639]
85. Xi Z, et al. Interspecific transfer of *Wolbachia* into the mosquito disease vector *Aedes albopictus*. *Proc Biol Sci* 2006;273:1317–1322. [PubMed: 16777718]
86. Dobson SL, et al. The effect of *Wolbachia*-induced cytoplasmic incompatibility on host population size in natural and manipulated systems. *Proc Biol Sci* 2002;269:437–445. [PubMed: 11886634]
87. Sinkins SP. *Wolbachia* and cytoplasmic incompatibility in mosquitoes. *Insect Biochem Mol Biol* 2004;34:723–729. [PubMed: 15242714]
88. Rasgon JL, Scott TW. Impact of population age structure on *Wolbachia* transgene driver efficacy: ecologically complex factors and release of genetically modified mosquitoes. *Insect Biochem Mol Biol* 2004;34:707–713. [PubMed: 15242712]
89. Marshall JM. The effect of gene drive on containment of transgenic mosquitoes. *J Theor Biol* 2009;258:250–265. [PubMed: 19490857]
90. Xi Z, et al. *Wolbachia* establishment and invasion in an *Aedes aegypti* laboratory population. *Science* 2005;310:326–328. [PubMed: 16224027]
91. Turley AP, et al. *Wolbachia* infection reduces blood-feeding success in the dengue fever mosquito, *Aedes aegypti*. *PLoS Negl Trop Dis* 2009;3:e516. [PubMed: 19753103]

92. Rasgon JL, et al. Can *Anopheles gambiae* be infected with *Wolbachia pipientis*? Insights from an in vitro system. *Appl Environ Microbiol* 2006;72:7718–7722. [PubMed: 17028229]
93. Masui S, et al. Bacteriophage WO and virus-like particles in *Wolbachia*, an endosymbiont of arthropods. *Biochem Biophys Res Commun* 2001;283:1099–1104. [PubMed: 11355885]

(a) Phage Transfer Between Coinfecting *Wolbachia*

(b) Paternal Phage Transmission

**Figure 1.**

Horizontal transfer of bacteriophage WO. **(a)** Phage WO can transfer between two different *Wolbachia* strains that coinfect the same host cell [22,23,25,31]. The phage becomes lytic (i) and lyses its *Wolbachia* host cell (ii). An active phage particle then attaches to a phage-free *Wolbachia* that coinfects the same host cell (iii) and injects its DNA (iv), at which point the DNA integrates into the chromosome (v). **(b)** Phage WO might also, hypothetically, be transmitted paternally by sperm from an infected male to the egg of a female carrying a phage-free *Wolbachia*. Once the sperm fertilizes the egg (i) the transported phage is released (ii) and can infect *Wolbachia* as in steps (iii-v) above.




Figure 2.

The genome architecture of phage WO. WOCauB2 from *wCauB* is an active phage based on the detection of excised intermediates by inverse PCR and genome sequencing [29]. Its genome is 43 Kb in size and encodes 47 genes (numbered from gp1 to gp47). Functional gene homologs include a site-specific recombinase gene (teal), head region genes (purple), baseplate assembly genes (pink), tail protein genes (green), and a phage late control gene (red). Other interesting genes of note encode homologs of plasmid replication protein RepA and a sigma-70 transcription factor (grey). Several of the encoded proteins might interact with host proteins, including a patatin-like protein, VrlC.1 and VrlC.2 (orange), and ankyrin-repeat proteins (ANK, blue). Genes of unknown function are shown in white, and transposase (Tp) is shown in yellow.




Figure 3. The phage density model of cytoplasmic incompatibility (CI). **(a)** When phage WO is lysogenic and titers of *Wolbachia* are high in male reproductive tissues, high levels of CI prevent the production of viable offspring after mating with an uninfected female. **(b)** When phage WO in these *Wolbachia* becomes lytic, *Wolbachia* cell titers decrease due to cell lysis and cause the infected male and uninfected female to produce an increased number of offspring [27].

Table 1

Identification of WO prophages and active phage particles^a

Insect	WO prophages				Refs.	
	Common name	Species	Particle size by TEM ^b	Number of prophages ^c		Identification of phage DNA from active particles?
Mosquito		<i>Culex pipiens</i>	20 nm	5	No data available	[26,30]
		<i>Aedes albopictus</i>	~40 nm	1	No data available	[31]
Cricket		<i>Teleogryllus taiwanemima</i>	40 nm	1	PCR	[93]
Moth		<i>Cadra cautella</i>	40 nm	2	PCR, cloning, and sequencing	[28,29]
Wasp		<i>Nasonia vitripennis</i>	~25 nm	4	PCR	[27]
Fruit fly		<i>Drosophila melanogaster</i>	~25 nm	2	PCR	[25]
		<i>Drosophila simulans</i>	Not determined	4	PCR	[25]

^aPhage WO particles have been purified from several *Wolbachia*-insect systems using large-scale insect homogenization, density centrifugation, and purification through filters. In some cases, DNA has been amplified from phage particle isolations, confirming the presence of active phage particles.

^bTEM, transmission electron microscopy.

^cNumber of prophages in sequenced *Wolbachia* genomes

Table 2
Putative effectors and toxins encoded by the WO genomes of different *Wolbachia* strains

<i>Wolbachia</i> strains									
Effector or toxin	wCauB ^a	wRi ^b	wPip ^c	wMel ^d	wKue ^e	wWif ^f	wAna ^g	wMuni ^h	Putative function
SpvB	B3gp45 ⁱ								Interaction with host insect cells [29]
VHA/ VHC.1	B1gp15, B2gp29, B3gp30	WRi_007010 pseudogene	WPa_1322, WPa_0440 pseudogene	WD0580		Wendoof_ 01000548	WwAna1071	WUni_ 000700	Secretion into host insect cells [28]
VHC.2	B1gp16, B2gp30, B3gp31	WRi_007020	WPa_0441, WPa_1323	WD0579		Wendoof_ 01000385			Secretion into host insect cells [28]
Patain	B2gp45, B3gp44	WRi_06880	WPa_1340, WPa_272, WPa_0455	WD0565		Wendoof_ 01000457	WwAna0164		Entry into host cells
RhuM		WRi_005660, WRi_010320	WPa_431	WD0259	gp8				Unknown
Addition module toxin		WRi_005580	WPa_1330	WD0269, WD0600		Wendoof_ 01000378	WwAna0127	WUni_ 002995	Killing of cells lacking WO
DNA methylase		WRi_005640, WRi_010300	WPa_0258, WPa_0317, WPa_0429, WPa_1310	WD0263, WD0594		Wendoof_ 01000687, Wendoof_ 01000839	WwAna0008		Methylation of host DNA, protection of phage against restriction enzymes

^aPhages WOCauB1 (Accession No. AB161975.2), WOCauB2 (AB478515.1) and WOCauB3 (AB478516.1) from wCauB of *Cadra cautella* [28,29]

^bProphages from the wRi genome of *Drosophila simulans* Riverside (NC_012416) [8]

^cProphages from the wPip genome of *Culex quinquefasciatus* Pe1 (NC_010981) [30]

^dProphages from the wMel genome of *Drosophila melanogaster* (AE17196) [10]

^eProphage WO from wKue of *Ephesia kuehniella* (AB036666.1) [22]

^fProphage homologs from wWif of *Drosophila willistoni* (NZ_AAQP000000000)

^gProphage homologs from wAna of *Drosophila ananassae* (NZ_AAGB000000000)

^hProphage homologs from wMuni of *Muscidifurax uniraptor* (NZ_ACFF000000000)

ⁱAll gene numbers refer to the primary annotation of each genome in GenBank.