

Published in final edited form as:

Bioorg Med Chem Lett. 2010 May 1; 20(9): 2987–2990. doi:10.1016/j.bmcl.2010.02.102.

Peripheral Site Acetylcholinesterase Inhibitors Targeting Both Inflammation and Cholinergic Dysfunction

Sherri Young^a, Karine Fabio^a, Christophe Guillon^a, Pramod Mohanta^{a,†}, Timothy A. Halton^{a,‡}, Diane E. Heck^b, Robert A. Flowers II^a, Jeffrey D. Laskin^c, and Ned D. Heindel^{a,*}

^a Department of Chemistry, Lehigh University, Bethlehem, PA 18015, USA

^b Department of Environmental Health Science, New York Medical College School of Public Health, Valhalla, NY 10595, USA

^c CounterACT Center of Excellence, Rutgers University/UMDNJ, Piscataway, NJ 08854, USA

Abstract

The design and study of two classes of noncompetitive acetylcholinesterase inhibitors (AChEIs) which also function as NSAID prodrugs are reported. The most potent AChEIs disclosed contain an aromatic alkyl-aryl linker between an NSAID and a lipophilic choline mimic and they inhibit acetylcholinesterase (AChE) in the submicromolar range. These agents have the therapeutic potential to dually target inflammation by releasing an NSAID *in vivo* and activating the cholinergic anti-inflammatory pathway via cholinergic up-regulation.

Pro-inflammatory cytokine up-regulation plays a role in the pathogenesis of a wide range of disorders including osteoarthritis,¹ psoriasis,² multiple sclerosis,³ and other autoimmune disorders.⁴ Despite decades of research, non-steroidal anti-inflammatory drugs (NSAIDs) are still one of the most commonly used, highly effective treatments for such disorders. However, since chronic NSAID-use often leads to gastrointestinal (GI) side effects, NSAID ester prodrugs have been explored to mask the acidic GI-irritating portion of the NSAID.⁵ For conditions such as arthritis, topical NSAID prodrugs are particularly useful as their therapeutic action is localized, resulting in minimal systemic side effects.⁶

Alternately, acetylcholinesterase inhibitors (AChEIs) have implications in the treatment of severe inflammation resulting from sepsis,⁷ endotoxemia,⁸ and rheumatoid arthritis,⁹ as well as in the treatment of neuroinflammation associated with Alzheimer's disease^{10–12} and Myasthenia Gravis.¹³ The administration of CNS-active AChEIs such as galanthamine depletes systemic pro-inflammatory cytokines and ameliorates both central and peripheral inflammation.⁸ AChEIs seem to suppress inflammation via the cholinergic anti-inflammatory pathway, a mechanism by which the vagus nerve of the CNS regulates the production and release of tumor necrosis factor and other cytokines.^{14,15}

*Corresponding author. Tel.: 1-610-758-3464; fax.: 1-610-758-3461; ndh0@lehigh.edu.

†Current address: PTC Therapeutics, Inc., South Plainfield, NJ 07080-2449, USA

‡Current address: Digestive Care, Inc., Bethlehem, PA 18017-7059, USA

Supplementary data

Synthetic procedures, physical characterization of compounds, *in vitro* assay methods and Lineweaver-Burk plots can be found in the online version of this article.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

NSAID-AChEI conjugates have recently been explored in order to target neuroinflammation¹⁶ and vesicant-induced inflammation.¹⁷ These novel agents contain the AChEI, pyridostimine, linked via a hydrocarbon spacer to ibuprofen. When screened against sulfur mustard, the bifunctional agents demonstrated longer and more effective prevention of edema and inflammation than the AChEI alone with a post-treatment anti-inflammatory effect comparable to that of the parent NSAID.¹⁷ Furthermore, the same conjugates down-regulated nitric oxide and prostaglandin E₂ production and ameliorated autoimmune encephalomyelitis more than either component alone.¹⁶ NSAID-AChEIs seem to prove dually effective against inflammatory disorders by activating the cholinergic anti-inflammatory pathway and inhibiting cyclooxygenase. Based on previous findings, two novel classes of noncompetitive AChEIs containing common NSAIDs linked via an ester or an aromatic ester-carbonate backbone to choline bioisosteres have been developed (Figure 1).¹⁸ The synthesis and evaluation of these compounds as anti-inflammatory and anticholinesterase agents for topical or oral administration are presented herein.

Class 1 molecules have recently been designed as both up-regulators of choline and NSAID prodrugs.¹⁸ The cholinergic functions (-CO-O-CH₂CH₂X(CH₃)₃; X = C, Si, N⁺) are known bioisosteres of choline which inhibit AChE in both a competitive (as with the silicon derivative) and noncompetitive (as with the carbon derivative) manner.¹⁹ It is proposed that AChE binds to the choline mimic and activates the cholinergic anti-inflammatory pathway. Additionally, esterases in the plasma, skin and lipid membranes release the NSAID, thus targeting dual mechanisms of inflammation. Molecules containing NSAIDs directly linked to choline mimics via an ester bond (Figure 1, Class 2) have been included both as a control class as well as an additional class for therapeutic exploration.

Class 1 compounds were obtained by coupling either 3,3-dimethylbutyl carbonochloridate (**Ia**)²⁰ or 2-(trimethylsilyl)ethyl carbonochloridate (**Ib**)²¹ to the phenolic function of *p*-hydroxybenzyl alcohol. Coupling of **I** to an NSAID activated with 4-dimethylaminopyridine (DMAP) completed the synthesis (Scheme 1). Overall yields of this procedure ranged from 61 to 76 %, including synthesis of the chloroformate starting materials.

In the simple-ester series (Figure 1, Class 2), standard coupling conditions [1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC HCl)-DMAP] were employed for the carbon and silicon derivatives in a one step, moderate- to high-yielding procedure (55–97 %). Synthesis of the nitrogen-based NSAID-choline esters of Class 2 (**11**, **14**, and **17**) has been previously described by J. Young *et al.*²² and will not be discussed further. Using Young's method, products were isolated in moderate yields over three steps (30–40 %). All Class 2 agents were purified using non-chromatographic methods such as crystallization and extraction.

Following synthesis and characterization, all compounds were evaluated for anticholinesterase activity using Ellman's method²³ with tacrine,²⁴ one of the most potent commercial AChEIs, as a reference (Table 1). AChE contains a 20 Å deep gorge stretching from the enzyme surface to the active-site serine residue.²⁵ This structural moiety is often termed the aromatic or hydrophobic gorge since it is lined with Trp, Tyr and Phe residues.²⁵ A peripheral binding site termed the peripheral anionic site (PAS) is located at the top of the gorge, near the surface of the enzyme. Keeping in mind the nature of AChE, it was predicted that the highly lipophilic and aromatic Class 1 and 2 agents would interact with the Trp, Tyr and Phe residues at and near the PAS and inhibit enzyme activity.

Listed in Table 1 are IC₅₀ values for each compound against AChE. As a control, the parent NSAIDs were also screened for activity and were deemed inactive compared to most of the test compounds (IC₅₀ > 1000 μM, data not shown). A direct comparison of Class 1 and 2

activities reveals that Class 1 compounds are more potent than Class 2 by at least a factor of 4, in all cases. The most potent Class 1 inhibitor (**7**, Figure 2) has an IC₅₀ comparable to that of the AChEI, galanthamine, a natural alkaloid which has been approved for the treatment of Alzheimer's disease.²⁴ Since Class 2 compounds were found less active than Class 1, it is likely that the lipophilic aromatic linker is needed to maximize potential π - π stacking interactions²⁶ with the Trp, Tyr and Phe residues of the aromatic gorge.²⁵ The ester-carbonates (Class 1) are also at least 10 times more lipophilic than the corresponding simple esters, which may also increase hydrophobic interactions with AChE (refer to Table 1 for Clog P values). The more hydrophilic nitrogen derivatives of Class 2 (**11**, **14**, and **17**) were at least 20 times less potent than their carbon and silicon derivatives, further enhancing the argument that hydrophobic interactions at and near the active-site gorge may be responsible for the observed activities.

Overall, the most active AChEIs in both classes (**7**, **18**, and **19**) are all derivatives of the NSAID, diclofenac. This trend is likely due to the high lipophilicity of diclofenac which strengthens interactions with the hydrophobic residues of AChE. Furthermore, the chlorine substituents on one of the aromatic rings of diclofenac may enhance π - π stacking interactions with aromatic residues lining the gorge (likely Trp84, Phe330, or Trp279 in *Torpedo californica* AChE), as suggested by previous SAR and computational studies performed by Holzgrabe *et al.*²⁶ and Sussman *et al.*²⁷

The degree of reversibility of AChE-inhibitor interactions was investigated for all inhibitors. Following an incubation period of AChE with the test agents, gel filtration resulted in near-complete restoration of enzyme activity (80–100 %, see Table 1). It was important to demonstrate the reversible nature of our compounds because many of the known irreversible AChEIs are nerve agents, toxins, or pesticides and are not considered for therapeutic use.²⁸

The mechanism of AChE inhibition was further investigated by generating Lineweaver-Burk plots.²⁹ Briefly, the reciprocal of the reaction velocity (v^{-1}) was plotted against the reciprocal of the substrate concentration ($[\text{acetylthiocholine, ATCh}]^{-1}$) after assaying enzyme activity for varying inhibitor and ATCh concentrations. All compounds exhibited reversible, noncompetitive inhibition, with quadrant III interceptions (for detailed assay method and example Lineweaver-Burk plots, see Supporting Information). These analyses support the reversibility studies previously discussed. More in depth investigations which address whether compounds **1–19** are interacting with the PAS of AChE or an alternate site are ongoing.

To evaluate Class 1 and 2 agents as oral or topical prodrugs, hydrolytic release of the NSAID was followed by HPLC under various conditions. Gradual release of the parent NSAID was observed for both Class 1 and 2 agents in human plasma (80 % plasma in PBS), with half-lives not exceeding 8 h (Table 2). Negligible hydrolysis was observed in the PBS control during the time of the half-life determination. The ester-carbonate prodrugs released the parent drug more slowly ($t_{1/2}$ of 2–8 h) compared with the ester prodrugs ($t_{1/2}$ of 5–100 min). Compound **14** was completely hydrolyzed after 10 min, suggesting that the choline esters (**11**, **14**, and **17**) would not make suitable prodrugs.

Hydrolysis in acetate buffer was also determined at the pH on the surface of the skin (pH 5). No significant hydrolysis was observed for up to seven days under acidic conditions, suggesting that the ester and ester-carbonate linkages should be resistant to cleavage on the surface of the skin. Preliminary hydrolysis data suggest that Class 1 and 2 agents will act as NSAID prodrugs *in vivo*. Class 1 compounds demonstrate a more gradual NSAID-release when compared to Class 2. Additionally, they are more lipophilic than their Class 2 counterparts, making them more suitable as topical prodrugs.

Novel AChEIs have been developed with the potential to dually target inflammation by activating the cholinergic anti-inflammatory pathway and releasing an NSAID, *in vivo*. Lineweaver-Burk plots indicate that all test compounds are reversible, noncompetitive inhibitors of AChE. The most potent inhibitors (Class 1) contain a lipophilic aromatic spacer and are 4 times more active than those with solely an ester linkage between the anticholinergic and anti-inflammatory functionalities (Class 2). Of the compounds screened, diclofenac derivative **7** was most active likely due to its high lipophilicity, aromaticity and the chlorine-substituted ring.^{26,27} Hydrolysis data suggest that these agents should be effective as either oral or topical NSAID prodrugs.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.



Acknowledgments

This work was funded in part by the National Institutes of Health CounterACT Program through the National Institute of Arthritis and Musculoskeletal and Skin Diseases (award #U54AR055073). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the federal government. The authors would like to thank Professors Sam Niedbala and Jebrell Glover for use of their instrumentation.

References and notes

1. Özgüney I. *Expert Opin Pharmacother* 2008;9:1805–1816. [PubMed: 18570612]
2. Deeva I, Mariani S, De Luca C, Pacifico V, Leoni L, Raskovic D, Kharaeva Z, Korkina L, Pastore S. Wide-spectrum profile of inflammatory mediators in the plasma and scales of patients with psoriatic disease. *Cytokine*. 2009;10.1016/j.cyto.2009.09.014
3. Centonze D, Muzio L, Rossi S, Furlan R, Bernardi G, Martino G. The link between inflammation, synaptic transmission and neurodegeneration in multiple sclerosis. *Cell Death & Diff*. 2009;10.1038/cdd.2009.179
4. Kunz M, Ibrahim S. *Mediators Inflamm* 2009;2009:1–20.
5. Cannon, J. *Pharmacology for Chemists*. 2. Oxford University Press; New York: Concepts of Pharmacology; p. 13-17.
6. Trommer H, Neubert R. *Skin Pharmacol Physiol* 2006;19:106–121. [PubMed: 16685150]
7. Fodale V, Letterio B. *Crit Care Med* 2008;36:622–623. [PubMed: 18216614]
8. Pavlov V, Parrish W, Rosas-Ballina M, Ochani M, Puerta M, Ochani K, Chavan S, Al-Abed Y, Tracey K. *Brain Behav Immun* 2009;23:41–45. [PubMed: 18639629]
9. van Maanen M, Vervoordeldonk M, Tak P. *Nat Rev Rheumatol* 2009;5:229–232. [PubMed: 19337288]
10. Tsuno N. *Expert Rev Neurother* 2009;9:591–598. [PubMed: 19402770]
11. Guay D. *Consult Pharm* 2008;23:598–609. [PubMed: 19032006]
12. Marco L, do Carmo Carreiras M. *Recent Pat CNS Drug Discov* 2006;1:105–111. [PubMed: 18221196]
13. Brenner T, Nizri E, Irony-Tur-Sinai M, Hamra-Amitay Y, Wirguin I. *J Neuroimmunol* 2008;201–202:121–127.
14. Tracey K. *Nature* 2002;420:853–859. [PubMed: 12490958]
15. Tracey K. *J Clin Invest* 2007;117:289–296. [PubMed: 17273548]
16. Nizri E, Adani R, Meshulam H, Amitai G, Brenner T. *Neurosci Lett* 2005;376:46–50. [PubMed: 15694272]
17. Amitai G, Adani R, Fishbein E, Meshulam H, Laish I, Dahir S. *J Appl Toxicol* 2006;26:81–87. [PubMed: 16167317]
18. Laskin, JD.; Fabio, K.; Lacey, CJ.; Young, S.; Mohanta, P.; Guillon, C.; Heindel, ND.; Huang, M-T.; Heck, DE. *Unique Dual-Action Therapeutics*. U S Patent Pending. PCT/US2009/005961. Nov 3. 2009

19. Cohen S, Chishti S, Elkind J, Reese H, Cohen J. *J Med Chem* 1985;28:1309–1313. [PubMed: 4032433]
20. Hudson H, Koplick A, Poulton D. *J Chem Soc, Perkin Trans 2* 1979:57–66.
21. Berglund, A.; Akerblom, E.; Hedin, A. EU Patent. 0,209,586. 1986.
22. Venuti M, Young J, Maloney P, Johnson D, McGreevy K. *Pharm Res* 1989;6:867–873. [PubMed: 2608628]
23. Ellman G, Courtney K, Andres V, Feather-Stone R. *Biochem Pharmacol* 1961;7:88–95.
24. Guillou C, Mary A, Zafiarisoa Renko D, Gras E, Thal C. *Bioorg Med Chem Lett* 2000;10:637–639. [PubMed: 10762042]
25. Shen T, Tai K, Henchman R, McCammon A. *Acc Chem Res* 2002;35:332–340. [PubMed: 12069617]
26. Kapkova P, Stiefl N, Sürig U, Engels B, Baumann K, Holzgrabe U. *Arch Pharm Pharm Med Chem* 2003;336:523–540.
27. Dvir H, Wond D, Harel M, Barril X, Orozco M, Luque F, Muñoz-Torrero D, Camps P, Rosenberry T, Silman I, Sussman J. *Biochemistry* 2002;41:2970–2981. [PubMed: 11863435]
28. Taylor, P. *The Pharmacological Basis of Therapeutics*. Hardman, J.; Limbird, L.; Molinoff, P.; Ruddon, R.; Gilman, A., editors. McGraw-Hill; New York: 1996. p. 161-176.
29. (a) Lineweaver H, Burk D. *J Am Chem Soc* 1934;56:658–666. (b) Bergmeyer, H.; Gawehn, K. *Principles of Enzymatic Analysis*. Verlag Chemie; NY: 1978. p. 36-40.



Class 1 (<i>Ester-Carbonate Series</i>)			Class 2 (<i>Ester Series</i>)		
<i>Compound</i>	<i>NSAID</i>	<i>X</i>	<i>Compound</i>	<i>NSAID</i>	<i>X</i>
1	Ibuprofen	C	9	Ibuprofen	C
2	Ibuprofen	Si	10	Ibuprofen	Si
3	Naproxen	C	11	Ibuprofen	N ⁺
4	Naproxen	Si	12	Naproxen	C
5	Indomethacin	C	13	Naproxen	Si
6	Indomethacin	Si	14	Naproxen	N ⁺
7	Diclofenac	C	15	Indomethacin	C
8	Diclofenac	Si	16	Indomethacin	Si
			17	Indomethacin	N ⁺
			18	Diclofenac	C
			19	Diclofenac	Si

Figure 1.
Generic Structures of Class 1 (*top*) and Class 2 (*bottom*) agents.





Figure 2.
Structure of **7**, the most potent AChEI discussed herein ($IC_{50} = 0.51 \mu M$).

**Scheme 1.**

Synthesis of Class 1 agents. a) Et₃N, THF, -10 to 25°C; b) NSAID, EDC HCl, DMAP, CH₂Cl₂, 0 to 25°C.

Table 1

Anticholinesterase activities and Clog P values of Class 1 and 2 agents.

Class 1 (Ester-Carbonate Series)				Class 2 (Ester Series)			
Compound	IC ₅₀ (μM) ^a	% AChE Recovery ^b	Clog P ^c	Compound	IC ₅₀ (μM) ^a	% AChE Recovery ^b	Clog P ^c
1	1.93 ± 0.64	101 ± 8	7.37	9	24.6 ± 14.5	107 ± 17	6.47
2	1.19 ± 0.20	88 ± 11	7.53	10	25.7 ± 4.9	92 ± 11	6.64
3	1.74 ± 1.01	99 ± 4	6.50	11	>2000	99 ± 5	0.42
4	0.83 ± 0.15	98 ± 3	6.67	12	19.7 ± 1.7	117 ± 13	5.61
5	2.29 ± 0.94	78 ± 2	7.87	13	13.9 ± 0.3	100 ± 1	5.77
6	0.72 ± 0.13	86 ± 1	8.03	14	>1000	93 ± 13	-0.45
7	0.51 ± 0.02	87 ± 11	8.41	15	9.75 ± 0.89	80 ± 2	6.97
8	1.36 ± 0.13	104 ± 23	8.58	16	3.32 ± 0.36	73 ± 4	7.14
Tacrine HCl	0.055 ± 0.005	95		17	230 ± 32	101 ± 6	0.92
				18	2.69 ± 0.15	99 ± 4	7.52
				19	2.66 ± 0.25	90 ± 10	7.68

^aSource of AChE is electric eel (*Electrophorus electricus*). Each condition was run in duplicate.^bEeAChE was incubated for 30 min with 55 μM inhibitor at 25°C. Enzyme activity was assayed following gel filtration. Values reported are averages of at least two experiments.^cValues calculated using ChemBioDraw Ultra 11.0.

Table 2Hydrolysis of Class 1 and 2 Compounds in Human Plasma.^a

Class 1 (<i>Ester-Carbonate Series</i>)		Class 2 (<i>Ester Series</i>)	
Compound	$t_{1/2}$ (min) ^b	Compound	$t_{1/2}$ (min) ^b
1	204	12	38
3	135	13	63
4	253	14	< 5
5	468	18	111
7	357		

^aHydrolysis measured in 80 % human plasma in PBS at 37 °C.^bHalf-lives determined by plotting the semi-log of prodrug disappearance.