

Published in final edited form as:


Org Lett. 2010 May 7; 12(9): 2120–2122. doi:10.1021/ol100615j.

Asymmetric, Organocatalytic, Three-Step Synthesis of α -Hydroxy-(*E*)- β,γ -Unsaturated Esters

Lindsey C. Hess and Gary H. Posner*

Department of Chemistry, Johns Hopkins University, 3400 N Charles St., Baltimore, MD 21218

Abstract



An efficient and enantiocontrolled 3-step synthesis of α -hydroxy-(*E*)- β,γ -unsaturated esters is reported. Enantioenriched α -selenyl aldehydes, prepared in one step by asymmetric, organocatalytic α -selenylation of aldehydes, were directly subjected to a Wittig reaction followed by allylic selenide to selenoxide oxidation and final spontaneous [2,3]-sigmatropic rearrangement to yield the target compounds in 43–65 % overall yield and in 94–97 % ee.

Asymmetric synthesis utilizing organoselenium compounds has become increasingly popular in recent years.¹ Selenium incorporation can be done in either a nucleophilic or an electrophilic fashion.² Once selenium is introduced into the molecule, many different chemical transformations can occur including oxidations leading to either syn-elimination³ or, in the case of an allylic selenide, [2,3]-sigmatropic rearrangement to give an allylic alcohol.⁴ Pursuing our recently published work preparing chiral non-racemic γ -hydroxy-(*E*)- α,β -unsaturated sulfones and esters,⁵ we have developed a complementary asymmetric synthetic method producing α -hydroxy-(*E*)- β,γ -unsaturated esters.

α -Hydroxy esters and their corresponding acids are key structural units of valuable synthetic intermediates as well as natural products.^{1,6} Based on the utility of these structural units, we set out to design a simple, asymmetric general strategy for their synthesis. Surprisingly, most reports of selenium oxidation and [2,3]-sigmatropic rearrangements have been done utilizing chiral oxidants as opposed to installing chirality prior to oxidation.⁷ There have been a few reports involving diastereoselective oxidations of selenides containing chiral moieties,⁸ but the scope is quite limited, with low to moderate yields and only modest diastereomeric excess (de). Other methods have also been reported using conformationally locked ring systems to induce stereocontrol.⁹

ghp@jhu.edu.

 Supporting Information Available: Experimental procedures and full spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Recent reports on the organocatalytic, asymmetric α -selenenylation of aldehydes in high yields (> 85 %) and high enantiomeric excess (> 95 %) ^{10,11} gave an excellent method for controlling the absolute stereochemistry of our α -hydroxy-(*E*)- β,γ -unsaturated target systems.

We chose a variety of aldehydes (3-phenylpropanal, 2-cyclohexylethanal, N-Boc-4-piperidineethanal, hexadecanal, and 6-benzyloxyhexanal) as well as three different ester functional groups (OMe, *O*-*t*-butyl, and *O*-benzyl) to determine the generality of this reaction. α -Selenyl aldehydes **3a-e**, prepared according to an Italian protocol, ¹¹ were immediately subjected in situ to a Wittig reaction to give the enantiomerically enriched compounds **5-11** (Scheme 1, A) in 48–83 % yields. Oxidation of the γ -selenyl-(*E*)- α,β -unsaturated esters to the selenoxide using hydrogen peroxide caused spontaneous [2,3]-sigmatropic rearrangement to give the final enantioenriched α -hydroxy-(*E*)- β,γ -unsaturated esters (Scheme 1, B) in 63–90 % yields and excellent ee's (≥ 94 %). The exclusive (*E*)-geometry of the carbon-carbon double bond in intermediates **5-11** as well as the final products **12-18** was confirmed by ¹H NMR spectroscopy ($J = 14-16$ Hz). It is important to note that when α -hydroxy ester final products **12**, **13** and **14** were dissolved in a variety of solvents (e. g. THF, CHCl₃, CH₃CN, and hexanes) at room temperature, they were stable to racemization (as determined via chiral HPLC) for at least 24 hours. We also used mCPBA as an oxidant of selenium, and we obtained a comparable yield and ee when compared to hydrogen peroxide (system tested was hexadecanal). The preferred method of oxidation remained hydrogen peroxide due to the convenience of the reaction conditions (non-anhydrous).

To illustrate the utility of this method for natural product synthesis, we completed a formal total synthesis of (+)-symbioramide (**20**), a naturally occurring bioactive ceramide composed of *D*-erythro-dihydrosphingosine and (2*R*,3*E*)-2-hydroxy-3-octadecenoic acid. ¹² We prepared key intermediate (–)-**19** that had been used in a recent synthesis ¹³ of this biologically active ceramide (Scheme 2). Intermediate (–)-**19** had been prepared in 7 steps and 11.8 % overall yield from L-serine. ^{13a} Using the new method described here, we prepared this intermediate mono-protected diol (–)-**19** in only 5 steps and in 28 % overall yield and 95% ee (Scheme 3). It is important to note that, in order to achieve the natural stereochemistry at the chiral 2-position, we employed the opposite proline-derived catalyst in the asymmetric α -selenenylation of hexadecanal. This formal total synthesis of (+)-symbioramide (**20**) allowed us to confirm the absolute stereochemistry of our final α -hydroxy carbonyl products. Previously published characterization of mono-protected diol (–)-**19** indicates that the absolute stereochemistry at the 2 position is *R*, with $[\alpha]_D = -77^\circ$, matching our experimental data for intermediate (–)-**19** prepared via Scheme 3.

In one final example, we prepared also a vitamin D₃ chiron that incorporated a side chain with the structural motif generated via this synthetic method. An S_N2 displacement of iodide **22** ¹⁴ using allylmagnesium bromide yielded terminal olefin **23**. Oxidative cleavage afforded aldehyde **24**. Using the method described here (Scheme 4), we were able to prepare α -hydroxy-(*E*)- β,γ -unsaturated benzyl ester **26** as a single diastereomer. This fragment can be easily modified in order to prepare final C,D-ring side-chain modified vitamin D analogs.

In summary, we report an efficient, generalized, asymmetric, organocatalytic procedure for synthesis of α -hydroxy-(*E*)- β,γ -unsaturated esters in high enantiomeric purities. This three-step synthesis works with a variety of substrates, in good overall yields (46–65 %), and with excellent control of absolute stereochemistry (ee values ≥ 94 %). We have proven the utility of this method through a formal total synthesis of natural product (+)-symbioramide (**20**) as well as through the preparation of a 23-(*E*)-ene-25(*S*)-hydroxy building block useful for synthesis of new vitamin D side-chain analogs. ¹⁵

Supplementary Material






Refer to Web version on PubMed Central for supplementary material.

Acknowledgments


thank the NIH (CA 93547) for support.

References


1. Wirth T. *Tetrahedron* 1999;55:1.
2. (a) Sharpless KB, Lauer RF. *J Am Chem Soc* 1973;95:2697. (b) Sharpless KB, Young MW. *J Org Chem* 1975;40:947. (c) Grieco PA, Gilman S, Nishizawa M. *J Org Chem* 1976;41:1485. (d) Tomoda S, Nomura Y. *Chem Lett* 1981:1069.
3. (a) Clark RD, Heathcock CH. *J Org Chem* 1976;41:1396. [PubMed: 1263006] (b) Nicalaou KC, Lysenko Z. *J Am Chem Soc* 1977;99:3185. (c) Friedrich LE, Lam PYS. *J Org Chem* 1981;46:306. (d) Scarborough RM, Smith AB. *Tetrahedron Lett* 1977;18:4361. (e) Liotta D, Markiewicz W, Santiesteban H. *Tetrahedron Lett* 1977;18:4365.
4. (a) Reich HJ. *J Org Chem* 1975;40:2570. (b) Clive DL, Chittattu G, Curtis NJ, Menchen SM. *J Chem Soc, Chem Commun* 1978:770.
5. Petersen KS, Posner GH. *Org Lett* 2008;10:4685. [PubMed: 18811178]
6. (b) Tamura Y, Yakura T, Haruta JI, Kita Y. *Tetrahedron Lett* 1985;26:3837. (b) Tamura Y, Annoura H, Yamamoto H, Kondo H, Kita Y, Fujioka H. *Tetrahedron Lett* 1987;28:5709. (c) Takanami T, Tokoro H, Kato DI, Nishiyama S, Sugai T. *Tetrahedron Lett* 2005;46:3291. (d) Hong YT, Cho CW, Skucas E, Krische M. *Org Lett* 2007;9:3745. [PubMed: 17705502] (e) Faulkner DJ, Petersen MR. *J Am Chem Soc* 1973;95:553. [PubMed: 4687674] (f) Masamune TI, Ono M, Matsue H. *Bull Chem Soc Jpn.* 1975(g) Coppola, GM.; Schuster, HF. *α -Hydroxy Acids in Enantioselective Synthesis.* VCH; Weinheim: 1997.
7. (a) Davis FA, Stringer OD, McCauley JP. *Tetrahedron* 1985;41:4747. (b) Reich HJ, Yelm KE. *J Org Chem* 1992;57:5672. (c) Davis FA, Reddy RT. *J Org Chem* 1992;57:2599. (d) Komatsu N, Nishibayashi Y, Sugita T, Uemura SJ. *J Chem Soc, Chem Commun* 1992:46. (e) Komatsu N, Matsunaga S, Sugita T, Uemura S. *J Org Chem* 1993;58:3697. (f) Komatsu N, Nishibayashi Y, Sugita T, Uemura S. *J Am Chem Soc* 1993;115:5847. (g) Komatsu N, Nishibayashi Y, Uemura S. *Tetrahedron Lett* 1993;34:2339.
8. (a) Nishibayashi Y, Singh JD, Fukuzawa S, Uemura S. *J Org Chem* 1995;60:4114. (b) Davoren JE, Harcken C, Martin SF. *J Org Chem* 2008;73:391. [PubMed: 18088137]
9. (a) Zoretic PA, Chambers RJ, Marbury GD, Riebiro AA. *J Org Chem* 1985;50:2981. (b) Koprowski M, Krawczyk E, Skowronska A, McParlin M, Choi N, Radojevic S. *Tetrahedron* 2001;57:1105.
10. Sunden H, Rios R, Cordova A. *Tetrahedron Lett* 2007;48:7865.
11. Tiecco M, Carlone A, Sternativo S, Marini F, Bartoli G, Melchiorre P. *Angew Chem, Int Ed* 2007;46:6882.
12. Kobayashi J. *J Nat Prod* 1989;52:225.
13. (a) Azuma H, Takao R, Niuro H, Shikata K, Tamagaki S, Tachibana T, Ogino K. *J Org Chem* 2003;68:2790. [PubMed: 12662053] (b) Yoshida J, Masako N, Seki H, Hino J. *J Chem Soc, Perkin Trans 1* 1992;3:343.
14. Posner GH, Lee JK, White MC, Hutchings RH, Dai H, Kachinski JL, Dolan P, Kensler TW. *J Org Chem* 1997;62:3299. [PubMed: 11671717]




Scheme 1.
Preparation of α -hydroxy-(*E*)- β,γ -unsaturated esters



Scheme 2.
Previously Published Synthesis of (+)-symbioramide (20) Using Key Intermediate (-)-19



Scheme 3.
Formal Total Synthesis of (+)-symbioramide (**20**)



Scheme 4.
Synthesis of a 23-(*E*)-ene-25(*S*)-hydroxy Vitamin D₃ Chiron

Table 1

Reaction Results from Scheme 1.

compound (A,B)	R	R'	A yield, %	B yield, %	ee (%) ^a
5, 12	Bn (1a)	OMe (4a)	73	87	95
6, 13	Bn (1a)	O(<i>t</i> -Bu) (4b)	73	63	95
7, 14	Bn (1a)	OBn (4c)	81	70	95
8, 15	cyclohexyl (1b)	OBn (4c)	79	67	95
9, 16	N-Boc-piperidine (1c)	OBn (4c)	83	67	94
10, 17	BnO(CH ₂) ₄ (1d)	OBn (4c)	48	90	97
11, 18^b	CH ₃ (CH ₂) ₁₃ (1e)	OBn (4c)	77	83	95

^a ee values determined using chiral HPLC and racemic standards;^b catalyst (**2R**)-**2** was used.