

Molecular characterization of *Leptospira* spp. strains isolated from small rodents in Croatia

N. TURK^{1*}, Z. MILAS¹, J. MARGALETIC², V. STARESINA¹, A. SLAVICA³,
N. RIQUELME-SERTOUR⁴, E. BELLENGER⁴, G. BARANTON⁴ AND D. POSTIC⁴

¹ Department of Microbiology and Infectious Diseases, School of Veterinary Medicine University of Zagreb, Zagreb, Croatia

² Department of Forest Protection and Game, Faculty of Forestry University of Zagreb, Zagreb, Croatia

³ Chair of Game Biology & Pathology, School of Veterinary Medicine University of Zagreb, Zagreb, Croatia

⁴ Unité de Bactériologie Moléculaire et Médicale, Institut Pasteur, Paris, France

(Accepted 2 October 2002)

SUMMARY

We report the isolation and characterization of 16 *Leptospira* spp. strains isolated from small rodents captured in 11 different regions of inland Croatia. Large *NotI* and *SgrAI* restriction fragment allowed us to assign 10 isolates to the serovar *istrica*, 5 isolates to the serovar *tsaratsovo* and 1 isolate to the serovar *lora*. The phylogenetic analysis conducted from the sequences of the first 330 bp from the 16S rDNA gene revealed that the strains belonged to three different species, *L. borgpetersenii*, *L. kirschneri* and *L. interrogans*. Carrier rates in eight rodent species varied from 0 to 71·4%. *Mus musculus* showed the highest infection level and confirmed its role as a major reservoir of the serogroup Sejroë. For the first time we reported the occurrence of serovars *tsaratsovo* and *lora* in Croatia.

INTRODUCTION

Leptospirosis is a zoonosis widespread throughout the world, caused by pathogenic members of the genus *Leptospira* with great impact on both human and veterinary public health. Leptospire are immunologically and genetically heterogeneous spiral-shaped microorganisms that comprise 223 serovars organized into 24 serogroups [1] and several species [2–5]. Although leptospirosis was first described in Croatia in 1935 [6], our understanding of the circulating leptospiral serovars in circulation is still scarce. Various small rodents serve unambiguously as reservoir hosts for leptospire and they have potential to shed them in urine for extended periods. The evidence that small rodents carry leptospire and the presence

of antileptospiral agglutinins have been previously reported in certain parts of inland Croatia [7, 8]. To obtain a clear picture of the occurrence of the most common serovars, we started investigating the prevalence and identification of leptospire in small rodents in known natural foci of leptospirosis in Croatia.

There are various antigenic and genetic methods for the identification and characterization of leptospire. In the past years in Croatia new isolated serovars were originally classified on the basis of their antigenic traits following the conventional serotyping methods – the microscopic agglutination test and the cross-agglutinin absorption test (CAAT) with rabbit antisera [9]. Serovar identification by CAAT is fastidious, time-consuming, requires large volumes of culture and is hampered by its subjective interpretation (due to dark-field microscopy estimation of a 50% agglutinated leptospire cut-off). To overcome such difficulties,

* Author for correspondence: Department of Microbiology and Infectious Diseases, School of Veterinary Medicine University of Zagreb, Croatia, Heinzelova 55, 10000 Zagreb, Croatia.

Table 1. *Strains used in the study*

| Serogroup | Serovar | Strain (collection)* | Genomic species |
|-----------|-------------|----------------------|----------------------------|
| Australis | lora | Lora (IP) | <i>L. interrogans s.s.</i> |
| Australis | jalna | Jalna (IP) | <i>L. interrogans s.s.</i> |
| Australis | jalna | Jalna (KIT) | <i>L. interrogans s.s.</i> |
| Australis | bratislava | Jez-Bratislava (IP) | <i>L. interrogans s.s.</i> |
| Australis | bratislava | Jez-Bratislava (KIT) | <i>L. interrogans s.s.</i> |
| Australis | münchen | München C90 (IP) | <i>L. interrogans s.s.</i> |
| Sejroe | saxkoebing | Mus 24 (IP) | <i>L. interrogans s.s.</i> |
| Sejroe | hardjo | Hardjoprajitno (IP) | <i>L. interrogans s.s.</i> |
| Sejroe | roumanica | LM 294 (IP) | <i>L. interrogans s.s.</i> |
| Sejroe | wolffi | 3705 (IP) | <i>L. interrogans s.s.</i> |
| Sejroe | geyaweera | Geyaweera (IP) | <i>L. interrogans s.s.</i> |
| Sejroe | medanensis | Hond HC (IP) | <i>L. interrogans s.s.</i> |
| Sejroe | recreo | 380 (IP) | <i>L. interrogans s.s.</i> |
| Sejroe | polonica | 493 Poland (IP) | <i>L. borgpetersenii</i> |
| Sejroe | hardjobovis | Sponselee (IP) | <i>L. borgpetersenii</i> |
| Sejroe | sejroe | M 84 (IP) | <i>L. borgpetersenii</i> |
| Sejroe | istrica | Bratislava (IP) | <i>L. borgpetersenii</i> |
| Sejroe | nyanza | Kibos (IP) | <i>L. borgpetersenii</i> |
| Sejroe | dikkeni | Mannuthi (IP) | <i>L. borgpetersenii</i> |
| Sejroe | gorgas | 1413 U (IP) | <i>L. santarosai</i> |
| Sejroe | caribe | TRVL 61866 (IP) | <i>L. santarosai</i> |
| Sejroe | trinidad | TRVL 34056 (IP) | <i>L. santarosai</i> |
| Pomona | tsaratsovo | B 81/7 (IP) | <i>L. kirshneri</i> |
| Pomona | kunming | K 5 (IP) | <i>L. kirshneri</i> |

* IP, Collection of the WHO Collaboration Centre for Leptospire, Institut Pasteur, Paris, France; KIT, Collection of the Koninklijk Instituut voor de Tropen, Amsterdam, The Netherlands.

genetic methods have recently proved to be valuable for characterization of leptospira strains.

It had been shown previously that, although the sequences are very close to each other, 16S rDNA gene sequencing could allow the separation of *Leptospira* species and could be used for the comparison of different leptospira isolates [3, 10, 11]. Moreover, the partial sequencing of 16S rDNA gene amplicons (330-bp-long sequences corresponding to position 46–375 *Escherichia coli rrnB* numbering) has significant discriminatory power since almost 50% of the polymorphic positions recorded on the whole 16S rDNA gene of leptospira were localized within the first 330 bp [11].

Pulsed-field gel electrophoresis (PFGE) of large DNA fragments produced by rare-cutting restriction enzymes yield different electrophoretic patterns offering the advantage of a simple interpretation combined with a rapid result [12, 13].

In this paper, we report identification and characterization of leptospira isolates recently isolated from small rodents by phylogenetic analysis conducted from the partial 16S rDNA gene sequences and the

analysis of the polymorphism generated by PFGE. Using molecular tools, this study attempts to clarify through ecological approach, the occurrence and characters of the leptospiral serovars circulating in Croatia.

METHODS

Bacterial strains studied

Most of the leptospiral reference strains used in the study are from the Collection of the WHO Collaboration Centre for Leptospire, Institut Pasteur, Paris, France. Others were from collection kept at the Koninklijk Instituut voor de Tropen (KIT), Amsterdam, The Netherlands. They are listed in Table 1.

Study sites and investigation of small rodents

A total of 16 leptospiral strains (assigned as M1, M3, M10, M11, M12, M18, M27, M29, M30, M31, M33, M116, M218, M222, M261, M262) were isolated from

Table 2. Specification of captured small rodents, number of kidney cultures and leptospiral isolates obtained

| Species | Kidney cultures | Isolated serovars | | | Total | % |
|--------------------------------|-----------------|---|--------------------------------------|---------------------------------------|-------|------|
| | | <i>L. borgpetersenii</i> sejrøe isticra | <i>L. interrogans</i> australis lora | <i>L. kirshneri</i> pomona tsaratsovo | | |
| <i>Apodemus agrarius</i> | 74 | 0 | 0 | 4 | 4 | 5.4 |
| <i>Apodemus flavicollis</i> | 67 | 0 | 1 | 1 | 2 | 2.9 |
| <i>Apodemus sylvaticus</i> | 42 | 0 | 0 | 0 | 0 | 0 |
| <i>Auricola terrestris</i> | 1 | 0 | 0 | 0 | 0 | 0 |
| <i>Clethrionomys glareolus</i> | 23 | 0 | 0 | 0 | 0 | 0 |
| <i>Microtus agrestis</i> | 5 | 0 | 0 | 0 | 0 | 0 |
| <i>Microtus arvalis</i> | 3 | 0 | 0 | 0 | 0 | 0 |
| <i>Mus musculus</i> | 14 | 10 | 0 | 0 | 10 | 71.4 |
| Total | 227 | 10 | 1 | 5 | 16 | 7.0 |




Fig. 1. PFGE of *NotI* restriction fragments of serogroup Sejrøe. The digestion products were separated at 150 V for 36 h in 1% agarose 0.5 × Tris-borate-EDTA. Pulse time was 5–30 s. Lanes 1–10, serovar isticra isolates (M1, M3, M10, M11, M12, M18, M27, M30, M31, M33); 11, serovar saxkoebing Mus 24; 12, serovar hardjo Hardjoprajitno; 13, serovar polonica 493 Poland; 14, serovar romanica LM 294; 15, serovar wolffi 3705; 16, serovar hardjobovis Sponselee; 17, serovar sejrøe M 84; 18, serovar isticra Bratislava (signed with arrow); 19, serovar gorgas 1413 U; 20, serovar geyaweera Geyaweera; 21, serovar dikkeni Mannuthi; 22, serovar caribe TRVL 61866; 23, serovar medanensis Hond HC; 24, serovar recreo 380; 25, serovar nyanza Kibos; 26, serovar trinidad TRVL 34056, L, Lambda ladder PFG Marker (Biolabs).

the kidneys of 227 small rodents captured from March to October 2000 in 11 different geographical areas in inland Croatia where human and animal leptospirosis is a common disease (Karlovac, Velika Gorica, Popovaca, Kutina, Stara Gradiska, Nova Gradiska, Nova Kapela, Vinkovci, Otok, Vrbanja and Gunja). Rodents were captured using Sherman's live traps and the species were identified and separated. Animals were euthanized, aseptically dissected, and the kidney tissue was used for culture.

Culture conditions and serogroup identification

Isolates were grown in 5 ml Korthof's liquid medium [9] prior to the EMJH liquid medium [14, 15] at 30 °C to get a density suitable for use in agglutination reactions with 23 standard antisera (group sera) for the first typing according to serogroup affinities [16]. The growth was verified by counting in a Petroff-Hauser chamber. Agglutinations were performed by mixing equal quantities of a well-grown culture




Fig. 2. PFGE of *NotI* restriction fragments of serogroup Pomona. The digestion products were separated at 150 V for 36 h in 1% agarose 0.5 × Tris-borate-EDTA. Pulse time was 5–30 s. Lanes 1–5, serovar tsaratsovo isolates (M116, M218, M222, M261, M268); lane 6, serovar tsaratsovo B 81/7; lane 7, serovar kunming K 5. L, Lambda ladder PFG Marker (Biolabs).

(2×10^8 cells ml⁻¹) and serial dilutions of hyper-immune reference antisera produced in rabbits. The agglutination was read using dark-field microscopy.

Sequencing

The primers used to amplify the first 330 bp of the 16S rDNA gene were LEPTOA (5'-GGCGGCGGTCT-TAAACATG-3') with positions 38–57 and LEPTOB (5'-TTCCCCCATTGAGCAAGATT-3') with positions 367–347 [17]. The sequencing internal primers used were LEPTOC (5'-CAAGTCAAGCGGAG-TAGCA-3') with positions 58–76 and RS4 (5'-TCTT-AACTGCTGCCTCCCGT-3') [11]. Sequencing was performed by Genome Express (Montreuil, France). Access numbers of the sequences are: M1, AY149228; M3, AY149223; M10, AY149230; M11, AY149222; M12, AY149231; M18, AY149227; M27, AY149229; M29, AY149232; M30, AY149226; M31, AY149225; M33, AY149224; M116, AY149218; M218, AY149221; M222, AY149220; M261, AY149219; M262, AY149217.




Fig. 3. PFGE of *SgrAI* restriction fragments of serogroup Australis. The digestion products were separated at 150 V for 36 h in 1% agarose 0.5 × Tris-borate-EDTA. Pulse time was 5–30 s. Lane 1, serovar lora isolate (M29); lane 2, serovar lora Lora; lane 3, serovar jalna Jalna; lane 4, serovar jalna Jalna (KIT); lane 5, serovar bratislava Jez-Bratislava; lane 6, serovar bratislava Jez-Bratislava (KIT); lane 7, serovar münchen München C 90. L, Lambda ladder PFG Marker (Biolabs).

Phylogenetic analysis

The sequences obtained in this study and those available in data banks were aligned both automatically with Clustal V and manually with VSM [18]. They were analysed by distance methods, Unweighted Pair Group with Mathematical Average (UPGMA) [19] and Neighbour Joining (NJ) [20] with Mega Software [21]. The distances were calculated by the Jukes Cantor correction [20] and the tree construction was performed in a pairwise deletion procedure.

DNA extraction, enzyme digestion and serovar identification using PFGE

Preparation of the DNA was performed as previously described [13] and the plugs were stored in 0.5 M EDTA at 4 °C until used. Plugs were digested with *NotI* (Amersham) or *SgrAI* (Bio-Rad) in accordance with the manufacturer's instructions and then slices (1–2 mm thick) were transferred to a 1% agarose gel.




Fig. 4. Phylogenetic tree of partial 16S rDNA sequences (330 bp between positions 46 and 375 *E. coli* numbering) available from databanks and from strains determined in this study. The phylogenetic tree was drawn using the neighbour-joining method. The numbers at the branch nodes indicate the results of the bootstrap analysis.

PFGE was performed with a contour-clamped homogeneous electric field system (DRII apparatus, Bio-Rad Laboratories, Richmond, CA, USA). Separation

of DNA fragments was achieved with three different pulse times: 5–60 s for 35 h at 150 V; 5–30 s for 36 h at 150 V and 20–120 s for 36 h at 170 V in order to

separate small and large DNA fragments, respectively. The sizes of DNA fragments were estimated by comparison of bands with concatemerized λ bacteriophage genomes with a size range 50–1000 kb (Lambda ladder PFG Marker (Biolabs), Beverly, MA, USA). Gels were photographed under UV transillumination after staining with ethidium bromide. DNA fragment patterns were examined visually. Isolates with identical restriction fragment patterns with two enzymes were assigned as the same leptospiral serovar.

RESULTS

To obtain a better insight of the occurrence of the leptospiral serovars in areas of inland Croatia, a total of 227 small rodents scattered into 8 species were captured in 11 different locations and investigated for the presence of leptospire. Specification of captured animals as well as leptospira isolates are listed in Table 2. From 227 kidney tissue cultures, a total of 16 leptospiral isolates were obtained from 7.0% of the captured animals: 10 from *Mus musculus*, 2 from *Apodemus flavicollis* and 4 from *Apodemus agrarius* with carrier rates of 2.9–71.4% in different species (Table 2). After a first typing according to serogroup affinities the isolates appeared to belong to serogroups Sejrøe (10 isolates), Pomona (5 isolates) and Australis (1 isolate).

Intact genomic DNA from 16 leptospiral isolates was digested with restriction enzymes and submitted to PFGE. PFGE analysis of the *NotI* restriction patterns allowed us to differentiate serovars within the serogroups Sejrøe and Pomona while analysis of the *SgrAI* restriction pattern distinguished serovars into the serogroup Australis with similar *NotI* patterns. Results showed that 10 leptospiral isolates (all from *M. musculus*) appeared to have the same banding pattern which ranged in molecular size from approximately <50–450 kb similar to serovar *istrica* of the serogroup Sejrøe (Fig. 1). Five isolates (4 from *A. agrarius* and 1 from *A. flavicollis*) with the same banding pattern which ranged in molecular size from approximately <50–650 kb were similar to serovar *tsaratsovo* of the serogroup Pomona (Fig. 2) and finally, 1 isolate (from *A. flavicollis*) has banding pattern with the molecular size from approximately <50–500 kb similar to serovar *lora* of the serogroup Australis (Fig. 3). Reference strains of two serovars (*jalna* and *bratislava*) from serogroup Australis belonging to different reference collections (Table 1) yielded different fingerprinting profiles (Fig. 3). Similar findings of collection

heterogeneity between serovars belonging to the controversial species *L. meyeri* provided by three different collections was reported previously [11].

Partial sequencing of the 16S ribosomal gene revealed the belonging of strains to three different species, *L. borgpetersenii*, *L. kirschneri*, *L. interrogans* (Fig. 4). Otherwise, within each species the nucleotide sequences were remarkably conserved, since all sequences exhibited 100% identity. As previously observed [11], *L. interrogans* and *L. kirschneri* clustered very close together as expected from the alignment of sequences differing by only one nucleotide. However, the phylogenetic analysis allowed to clearly assign strains to different species by both Neighbour Joining (Fig. 4) or UPGMA (data not shown) methods.

DISCUSSION

In Croatia, no small rodent survey has been carried out since 1983 [8] although leptospirosis in humans and animals continued. Borcic et al. [7] found a leptospira carrier rate in 8.9% of examined small rodents in the Sava valley (northeast Croatia) and confirmed their previous studies [8] showing that the whole area of Posavina (northeast Croatia) to be a wide natural focus of leptospire of the pomona, grippotyphosa and sejrøe serovars, although it is believed that in the same area other leptospiral serovars may also find favourable conditions for their survival in nature.

Data on the leptospiral infection rates and the character of isolates from small rodents are rare in Europe. Stanko et al. [22] reported the prevalence of leptospiral antibodies in 5.0% of small rodents examined in Slovakia with the most frequently antibodies directed against the serovars grippotyphosa and sejrøe. The most frequent species of rodents were *A. flavicollis*, *A. agrarius* and *Clethrionomys glareolus*. Webster et al. [23] found antibodies to leptospira in 14% of wild brown rats (*Rattus norvegicus*) in British farms and Adler et al. [24] reported the leptospira carrier rate in 12.6% of the examined small rodents in an urban environment in Switzerland. We found *A. agrarius*, *A. flavicollis* and *A. sylvaticus* are most common among captured small rodents (Table 2) which confirms previous investigations [7, 8] in Croatia. It has to be mentioned that there was no proportional rate among different species of captured animals in all study sites (data not shown). So, the lack of some species in certain regions was more probably due to random sampling (trapping) of animals to be

examined rather than actual lack of species. The findings of the present investigations showed a surprisingly high prevalence of the serovar *istrica* (71.4%) among the population of *M. musculus*, indicating on possible epizootic state at the capture area. This high prevalence points out the risk of heavy soil contamination by leptospires excreted in the urine of these animals. Close interaction between humans, animals, soil and water in these regions defines a considerable hazard of leptospiral infection to the local population as well as for outdoor-reared domestic animals. Moreover, the majority of *M. musculus* were captured in a hunting area (Kutina) with a large population of wild animals and a high risk of leptospirosis spreading. Sebek et al. [25] already reported that *M. musculus* could be a reservoir for the serovar *sejroë*. The serovar *tsaratsovo* were isolated from 5.4% of *A. agrarius* captured. Although it is well known that maintaining host to the serovars of Pomona serogroup is *A. agrarius*, the finding of serovar *tsaratsovo* is surprising since it has never been reported in Croatia. The serovar *lora* is also newly reported in Croatia. The serovar *istrica* takes place in leptospirosis transmission chain in certain parts of east Croatia as described previously [8]. It is interesting that we did not found the serovar *icterochaemorragiae* among the small rodents examined despite its undoubted role in the epidemiology of leptospirosis in humans and animals in Croatia. This confirms previous studies [8, 26], suggesting that serovar *icterochaemorragiae* has no reservoir among 'wild' mice, and its transmission occurs exclusively through rats (*R. norvegicus*). Reference serovars within the serogroup *Sejroë* were distributed into three species. Serovars belonging to *L. borgpetersenii* were mostly isolated from Central Europe (Bulgaria, Czech Republic, Romania, Poland). Thus, isolation of a large number of strains belonging to a serovar from *L. borgpetersenii* is not surprising. Similarly, serovars within the serogroup Pomona isolated in the same geographic region mostly belonged to the species *L. kirschneri*. Thus, identification of *Leptospira* species isolated in Croatia was in accordance with previously known geographical distribution of different serovars.

Since 1990, it has been shown that PFGE [11–13, 27–29] is usually in agreement with serotyping when considering serovar identification. This method displays many advances, objectivity, rapidity, commercially available reagents and pictures. However, the routine use of this method is still in its infancy. Difference between the patterns provides a reliable identification for each isolate by comparing each reference

strain inside the serogroup. Restriction fingerprints can reveal degrees of relatedness among leptospiral strains; the more closely related the strains are genetically, the more similar their fingerprint patterns are.

In conclusion, using molecular tools we confirmed previous knowledge about occurrence of the serovar *istrica* in Croatia, but we reported for the first time the occurrence of the serovars *tsaratsovo* and *lora*. Results of this work are an initial step in the investigation of the molecular epidemiology of leptospirosis in Croatia. The role of different small rodents is still elusive and remains to be clarified. The relations of the reservoirs, humans and animals in the epidemiologic chain of leptospirosis in Croatia should be further studied, particularly on the molecular biology level.

ACKNOWLEDGEMENTS

We thank the Institut Pasteur which supported this work. The contribution of Dragica Zivkovic as a technical support is acknowledged.

REFERENCES

1. Kmety E, Dikken H. Classification of the species *Leptospira interrogans* and history of its serovars. Groningen: University Press, 1993.
2. Brenner DJ, Kaufmann AF, Sulzer KR, Steigerwalt AG, Rogers FC, Weyant RS. Further determination of DNA relatedness between serogroups and serovars in the family *Leptospiraceae* with a proposal for *Leptospira alexanderi* sp. nov. and four new *Leptospira* genomospecies. *Int J Syst Bacteriol* 1999; **49**: 839–58.
3. Perolat P, Chappel RJ, Adler B, et al. *Leptospira fainei* sp. nov., isolated from pigs in Australia. *Int J Syst Bacteriol* 1998; **48**: 851–8.
4. Ramadass P, Jarvis BDW, Corner CJ, Penny D, Marshall RB. Genetic characterization of pathogenic *Leptospira* species by DNA hybridization. *Int J Syst Bacteriol* 1992; **42**: 215–9.
5. Yasuda PH, Steigerwalt AG, Sulzer KR, Kaufmann AF, Rogers F, Brenner DJ. Deoxyribonucleic acid relatedness between serogroups and serovars in the family *Leptospiraceae* with proposals for seven new *Leptospira* species. *Int J Syst Bacteriol* 1987; **37**: 407–15.
6. Antunovic-Mikacic S. O prvom slucaju Weilove bolesti na nasem Primorju. [The first case of Weil's disease on our coastal region.] *Lij Vjes* 1935; **57**: 377.
7. Borcic B, Kovacic H, Sebek Z, Aleraj B, Tvrtkovic N. Small terrestrial mammals as reservoir of leptospires in the Sava Valley (Croatia). *Folia Parasitol* 1982; **29**: 177–82.
8. Borcic B, Kovacic H, Sebek Z, Aleraj B, Tvrtkovic N. Small terrestrial mammals as reservoirs of leptospires in the Drava Valley. *Vet Arhiv* 1983; **53**: 41–9.
9. Babudieri B. Laboratory diagnosis of leptospirosis. *Bull WHO* 1961; **24**: 45.

10. Hookey JV, Bryden J, Gatehouse L. The use of 16S rDNA sequence analysis to investigate the phylogeny of *Leptospiraceae* and related spirochaetes. *J Gen Microbiol* 1993; **139**: 2585–90.
11. Postic D, Riquelme-Sertour N, Merien F, Perolat P, Baranton G. Interest of partial 16S rDNA gene sequences to resolve heterogeneties between *Leptospira* collections: application to *L. meyeri*. *Res Microbiol* 2000; **151**: 333–41.
12. Herman JL, Baril C, Bellenger E, Perolat P, Baranton G, Saint Girons I. Genome conservation in isolates of *Leptospira interrogans*. *J Bacteriol* 1991; **173**: 7582–8.
13. Herman JL, Bellenger E, Perolat P, Baranton G, Saint Girons I. Pulsed-field gel electrophoresis of *NotI* digests of leptospiral DNA; a new rapid method of serovar identification. *J Clin Microbiol* 1992; **30**: 1696–702.
14. Ellinghausen HC, McCullough WG. Nutrition of *Leptospira pomona* and growth of 13 other serotypes: fractionation of oleic albumin complex and a medium of bovine albumin and polysorbate 80. *Am J Vet Res* 1965; **26**: 45–51.
15. Johnson RC, Harris WG. Differentiation of pathogenic and saprophytic leptospires. Growth at low temperatures. *J Bacteriol* 1967; **94**: 27–31.
16. Dikken H, Kmety E. Serological typing methods of leptospires. In: Bergan T, Norris JR, eds. *Methods in microbiology*, Vol. 11. New York: Academic Press, 1978: 259–307.
17. Merien F, Amouriaux P, Perolat P, Baranton G, Saint Girons I. Polymerase chain reaction for detection of *Leptospira* spp. in clinical samples. *J Clin Microbiol* 1992; **30**: 2219–24.
18. Ruimy R, Breittmayer V, Elbaze P, Lafay B, Boussemart O, Gauthier M, Christen R. Phylogenetic analysis and assessment of the genera *Vibrio*, *Photobacterium*, *Aeromonas* and *Plesiomonas* deduced from small-subunit rRNA sequences. *Int J Syst Bacteriol* 1994; **44**: 416–26.
19. Sneath PHA, Sokal RR. *Numerical taxonomy*. San Francisco, CA: Freeman, 1973.
20. Saitou N, Nei M. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987; **4**: 406–25.
21. Kumar S, Tamura K, Masatoshi N. MEGA: Molecular evolutionary genetics analysis, version 1.01. The Pennsylvania State University, University Park, Pa 16802, 1993.
22. Stanko M, Prokopcakova H, Fricova J, Petko B. Occurrence of antibodies to *Leptospira* in small mammals in eastern Slovakia. *Vet Med* 1996; **41**: 373–7.
23. Webster JP, Ellis WA, Macdonald DW. Prevalence of *Leptospira* spp. in wild brown rats (*Rattus norvegicus*) in UK farms. *Epidemiol Infect* 1995; **144**: 195–201.
24. Adler H, Vonstein S, Deplazes P, Stieger C, Frei R. Prevalence of *Leptospira* spp. in various species of small mammals caught in an inner-city area in Switzerland. *Epidemiol Infect* 2002; **128**: 107–9.
25. Sebek Z. A characterisation of leptospirosis foci on the territory of Czechoslovakia. In: Rosicky B, Heyberger K, eds. *Theoretical questions on natural foci diseases*. Prague: Publ House Czech Acad Sci, 1965.
26. Zaharija I. Istrazivanja leptospiroze u zivotinja u Hrvatskoj. VIII. Stakor (*E. norvegicus*) kao kliconosa leptospira u gradovima Osijeku, Puli i Porecu. [The investigation of leptospirosis in animals in Croatia. VIII. The rat (*E. norvegicus*) as a carrier of leptospires in the cities of Osijek, Pula and Porec.] *Vet Arhiv* 1969; **39**: 141–3.
27. Brenot A, Werts C, Ottone C, et al. First evidence for a restriction-modification system in *Leptospira* sp. *FEMS Microbiol Lett* 2001; **201**: 139–43.
28. Ciceroni L, Ciarrocchi S, Ciervo A, et al. Differentiation of leptospires of the serogroup Pomona by monoclonal antibodies, pulsed-field gel electrophoresis and arbitrarily primed polymerase chain reaction. *Res Microbiol* 2002; **153**: 37–44.
29. Collares-Pereira M, Korver H, Cao Thi BV, et al. Analysis of *Leptospira* isolates from mainland Portugal and the Azores islands. *FEMS Microbiol Lett* 2000; **185**: 181–7.