
Characterization of virulence factors in the newly described *Salmonella enterica* serotype Keurmassar emerging in Senegal (sub-Saharan Africa)

A. GASSAMA-SOW^{1*}, A. A. WANE¹, N. A. CANU², S. UZZAU²,
A. AIDARA-KANE¹ AND S. RUBINO²

¹ Laboratoire de Bactériologie Expérimentale, Institut Pasteur, Dakar, Sénégal

² Department of Biomedical Sciences, Center for Biotechnology, Development and Biodiversity Research, University of Sassari, Viale San Pietro, Sassari, Italy

(Accepted 10 November 2005, first published online 18 January 2006)

SUMMARY

From 2000 to 2001, nine strains of *Salmonella enterica* belonging to the new serotype Keurmassar have been isolated from human and poultry samples at the Senegalese National *Salmonella* and *Shigella* Reference Laboratory at the Pasteur Institute, in Dakar. All strains carried virulence factors including *Salmonella* Pathogenicity Islands (SPI)-1, -2, -3 and -5 encoded genes. Strains did not harbour virulence plasmid. Ribotyping analysis revealed a single clone identical to *Salmonella* Decatur isolated in Zimbabwe. These data suggest that strains are closely related, and may have been spread clonally. In this new serotype, insertion sequence IS200 is not present.

INTRODUCTION

Salmonella enterica is one of the most common causes of foodborne enteric infection in humans and the main source of infection is contaminated food of animal origin. According to estimations of the World Health Organization, there are annually 16·6 million cases of typhoid fever, with approximately 600 000 fatal outcomes, and 1·3 billion cases of acute gastroenteritis due to non-typhoidal salmonellosis, with 3 million deaths [1]. In developing countries, few data are available on the incidence of infections due to *Salmonella*. In these areas, only 1–10% of cases are reported, and where the disease is more severe, it is associated with 20–30% mortality [1].

In 2000–2001, the Senegalese National *Salmonella* and *Shigella* Reference Laboratory at the Pasteur

Institute in Dakar received 604 *Salmonella* samples; among them, a new serotype of *Salmonella* (*Salmonella enterica* serotype 35:c:1,2) named Keurmassar was identified. This new serotype was isolated from human and poultry samples [2]. Multiple virulence factors are involved in the pathogenesis of *Salmonella*. These virulence genes act together in a complex virulence function, and have been found on horizontally acquired DNA regions, such as pathogenicity islands, plasmids, transposons, and bacteriophages [3, 4].

Herein, we report for the first time in Africa, the characterization of virulence factors in the newly described *S. enterica* serotype Keurmassar.

MATERIALS AND METHODS

Nine *S. enterica* serotype Keurmassar strains were isolated in Senegal from human and poultry samples between 2000 to 2001 in the Senegalese National *Salmonella* and *Shigella* Reference Laboratory at the Pasteur Institute, Dakar.

* Author for correspondence: Dr Amy Gassama Sow, Laboratoire de Bactériologie Expérimentale, Institut Pasteur, 220 Dakar, Sénégal.
(Email: gassama@pasteur.sn)

Total genomic DNA was extracted as described previously [5]. Crude extracts for polymerase chain reaction (PCR) assays were obtained by the boiling method as described by Holmes [6]. The following molecular methods were used to type strains:

For ribotyping analysis, $\sim 2 \mu\text{g}$ of extracted DNA were digested overnight at 37°C with *Pvu*II. Restriction fragments were separated by electrophoresis in a 0.8% (w/v) agarose gel in Tris-acetate buffer for 16 h at 1.5 V/cm and transferred to Hybond N membranes by the method of Southern [7]. Restriction fragments of lambda DNA cleaved with *Hind*III were used as fragment size markers.

Analysis of IS200 content was performed by PCR using *S. enterica* serovar Typhimurium strain ATCC 14028s and *S. enterica* serotype Abortusovis strain SS44 as positive controls. The following primers (5'-CAG ATG CGC CTA TAA GGC T-3' and 5'-CTA GGC TGG GGG TTC CGG GAA-3') were used to amplify a fragment of 690 bp [8].

Virulence factors encoded within *Salmonella* Pathogenicity Islands (SPIs) were detected by PCR using the following primers: *invA* (5'-TGC CTA CAA GCA TGA AAT GG-3' and 5'-AAA CTG GAC CAC GGT TGA CAA-3'), *spiC* (5'-CCT GGA TAA TGA CTA TTG AT-3' and 5'-AGT TTA TGG TGA TTG CGT AT-3'), *misL* (5'-GTC GGC GAA TGC CGC GAA TA-3' and 5'-GCG CTG TTA ACG CTA ATA GT-3'), *orfL* (5'-GGA GTA TCG ATA AAG ATG TT-3' and 5'-GCG CGT AAC GTC AGA ATC AA-3'), *pipD* (5'-CGG CGA TTC ATG ACT TTG AT-3' and 5'-CGT TAT CAT TCG GAT CGT AA-3'), *spvR* (5'-CCC CGG GAA TTC GCT GCA TAA GGT AGA AGG-3' and 5'-CCC CGG GTA CCA TGG ATT TCT TGA TTA ATA AA-3'). Detection of *Gifsy*-1 and *Gifsy*-2 phages was performed as previously described [3]. *S. Typhimurium* strain ATCC 14028s was used as positive control.

RESULTS AND DISCUSSION

The nine strains of *S. Keurmassar* isolated from human and poultry specimens showed identical ribotype patterns suggesting that all strains isolated between 2000 to 2001 may belong to the same clone (Fig.). Unexpectedly, this ribotype was identical to the most frequently occurring profile observed in strains of *S. enterica* serotype Decatur [9]. Interestingly, strains of *S. Decatur* (6,7:c:1,5) have been rarely isolated worldwide, yet a number strains belonging to this serotype have recently been




Fig. Ribotype patterns obtained from *Salmonella enterica* subsp. *enterica* serotype Keurmassar. Lane 1, Lambda molecular weight ladder (kbp); lanes 2–11, *S. Keurmassar* (strains LO4020128, LO4020129, LO4020157, LO4020157, KO330119, KO412155, KO405195, KO329122, KO509194, KO508153); lane 12, *S. Decatur*, strain 1623.

isolated in Zimbabwe (Rubino *et al.*, unpublished observations).

The screening by PCR to detect IS200 was negative for all strains. This insertion sequence is present in most isolates of *Salmonella* and certain strains of *Shigella* and *Escherichia coli* [10, 11]. These data would suggest that *S. Keurmassar* belongs to a small group of non-typhoid *S. enterica* subspecies *enterica*, including serotypes Agona, Paratyphi C, Choleraesuis, and Typhisuis, that lack genomic copies of IS200 [9–11].

All isolates investigated for the presence of virulence factors revealed the presence of *invA*, *spiC*, *misL*, *pipD*, contained respectively in SPI-1, SPI-2, SPI-3, and SPI-5. These SPIs are required for bacterial penetration in the epithelial cells of intestine and for growth and survival of bacteria in the host [12]. Pathogenicity Islands (PAIs) have also been described in uropathogenic [13], and enteropathogenic *E. coli* [14].

SPI-4 gene *orfL* was absent in all strains; SPI-4 is involved in secretion of toxins that induce apoptosis in immune cells and it is required for survival in macrophages [15].

None of the *S. Keurmassar* strains harbour a virulence plasmid. A cluster of five genes, the *spv* locus, accounts for the overall contribution of virulence plasmids to pathogenicity and systemic infection in animal models: *spvR* a positive regulator gene and four effector genes, *spvA*, *spvB*, *spvC* and *spvD* [16]. These virulence genes are found in the most frequently isolated non-typhoid serotypes of *S. enterica* subsp. *enterica* including serotypes Typhimurium,

Enteritidis, Dublin, Choleraesuis, Gallinarum, and Abortusovis and in *S. enterica* subsp. *houtenae* [17].

All isolates of *S. Keurmassar* were non-lysogenic for Gifsy-1 and Gifsy-2, two prophages that carry several virulence genes. Gifsy-2 genome includes the *sodC-1* gene coding for a Cu Zn-dependent periplasmic superoxide dismutase; Gifsy-2 also encodes a virulence effector, *SseI*, secreted by the SPI-2 type III secretion system. The virulence of Gifsy-1 is undetectable in the presence of Gifsy-2; it becomes significant in cells that lack Gifsy-2, but carry the *sodC-1* gene [18]. In our study, the *sodC-1* gene was absent (data not shown).

Overall, genetic characterization of *S. Keurmassar* strains revealed the lack (i.e. Gifsy1, Gifsy2, *spv*) and/or the partial deletion (i.e. SPI-4) of several genetic loci previously known to play an important contribution to *S. enterica* pathogenicity. However, *S. Keurmassar* strains seem to be quite virulent to successfully establish *Salmonella*-induced enteritis in man.

In Africa, to our knowledge, this is the first report of the characterization of virulence factors in *Salmonella*.

ACKNOWLEDGEMENTS

This work was supported by a grant from the Sardinian Regional Government to S.U. (grant L.R. 19/96).

DECLARATION OF INTEREST

None.

REFERENCES

1. **Pang T, et al.** Typhoid fever and other salmonellosis: a continuing challenge. *Trends in Microbiology* 1995; **3**: 253–255.
2. **Cardinale E, et al.** Dual emergence in food and human of a novel multiresistant serotype of *Salmonella* in Senegal: *Salmonella enterica* subsp. *enterica* serotype 35:c:1,2. *Journal of Clinical Microbiology* 2001; **39**: S2373–S2374.
3. **Bacciu D, et al.** Transposition of heat-stable toxin *astA* gene into a Gifsy2-related prophage of *Salmonella enterica* serovar Abortusovis. *Journal of Bacteriology* 2004; **186**: 4568–4574.
4. **Häcker J, et al.** Pathogenicity islands of virulent bacteria: structure, function and impact on microbial evolution. *Molecular Microbiology* 1997; **23**: 1089–1097.
5. **Wilson K.** Preparation of genomic DNA from bacteria. In: Ausubel FA, et al., eds. *Current Protocols in Molecular Biology*, New York: John Wiley & Sons. 1994: pp. 2.4.1–2.4.5.
6. **Holmes DS, Quigley M.** A rapid boiling method for the preparation of bacterial plasmid. *Analytical Biochemistry* 1981; **114**: 193–197.
7. **Southern EM.** Detection of specific sequences among DNA fragments separated by gel electrophoresis. *Journal of Molecular Biology* 1975; **98**: 503–517.
8. **Bisercic M, Ochman N.** Natural populations of *Escherichia coli* and *Salmonella* Typhimurium harbor the same classes of insertion sequences. *Genetics* 1993; **133**: 449–453.
9. **Uzzau S, Hovi M, Stocker BA.** Application of ribotyping and IS200 fingerprinting to distinguish the five *Salmonella* serotype O:6,7:c:1,5 groups: *Choleraesuis sensu stricto*, *Choleraesuis* var. Kunzendorf, *Choleraesuis* var. Decatur, *Paratyphi C*, and *Typhisuis*. *Epidemiology and Infection* 1999; **123**: 37–46.
10. **Gibert I, Barbé J, Casadesús J.** Distribution of insertion sequence IS200 in *Salmonella* and *Shigella*. *Journal of General Microbiology* 1990; **136**: 2555–2560.
11. **Bisercic M, Ochman N.** The ancestry of insertion sequences common to *Escherichia coli* and *Salmonella* Typhimurium. *Journal of Bacteriology* 1993; **175**: 7863–7868.
12. **Marcus S, et al.** *Salmonella* pathogenicity islands: big virulence in small packages. *Microbes and Infection* 2000; **2**: 145–156.
13. **Blum G, et al.** Excision of large DNA regions termed pathogenicity islands from t-RNA-specific loci in the chromosome of an *Escherichia coli* wild-type pathogen. *Infection and Immunity* 1994; **62**: 606–614.
14. **McDaniel T, et al.** A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proceedings of the National Academy of Sciences USA* 1995; **92**: 1664–1668.
15. **Schmidt H, Hensel M.** Pathogenicity Islands in bacterial pathogenesis. *Clinical Microbiology Reviews* 2004; **17**: 14–56.
16. **Gulig P, et al.** Molecular analysis of *spv* virulence genes of the *Salmonella* virulence plasmids. *Molecular Microbiology* 1993; **7**: 825–830.
17. **Aabo S, Brown DJ, Olsen JE.** Virulence characterization of a strain of *Salmonella enterica* subspecies *houten* (subspecies IV) with chromosomal integrated *Salmonella* plasmid virulence (*spv*) genes. *Research in Microbiology* 2000; **151**: 183–189.
18. **Figuroa-Bossi N, Bossi L.** Inducible prophages contribute to *Salmonella* virulence in mice. *Molecular Microbiology* 1999; **33**: 167–176.