

## Prevalence and risk factors of *Helicobacter pylori* infection among healthy 3- to 5-year-old Israeli Arab children

KH. MUHSEN<sup>1,2</sup>, A. ATHAMNA<sup>2</sup>, M. ATHAMNA<sup>2</sup>, A. SPUNGIN-BIALIK<sup>1</sup>  
AND D. COHEN<sup>1\*</sup>

<sup>1</sup> Department of Epidemiology and Preventive Medicine, Sackler Faculty of Medicine, Tel Aviv University

<sup>2</sup> The Meshulash Research & Development Center, Kfar Qaraa, Israel

(Accepted 29 December 2005, first published online 2 March 2006)

### SUMMARY

We determined the prevalence and risk factors of *H. pylori* infection among 197 healthy 3- to 5-year-old Israeli Arab children, in a population under socioeconomic and environmental transition. Data on the socioeconomic and environmental characteristics were obtained by personal interviews. The presence of *H. pylori* infection was identified using an ELISA kit for detection of *H. pylori* antigens in stool specimens. The prevalence rate of *H. pylori* infection was 49·7% (95% CI 42·8–56·67). It varied significantly among the different villages. In the univariate analysis stratified by village, the risk of infection increased according to household crowding, number of siblings younger than 5 years and siblings' *H. pylori* positivity. In the multivariate analysis the village of residence and siblings' *H. pylori* positivity were the only variables that remained strongly associated with *H. pylori* infection. In a population such as that described in this study the socioeconomic and living conditions are major risk factors of *H. pylori* infection and the intra-familial transmission of *H. pylori* in early childhood has an important role.

### INTRODUCTION

*H. pylori* infection occurs worldwide and it is estimated that about 50% of the world's population is infected with the pathogen [1, 2]. The prevalence rates of *H. pylori* infection are higher in developing countries than in industrialized countries [1]. *H. pylori* is associated with gastric disorders, is responsible for the majority of duodenal and gastric ulcers, and is classified as grade one carcinogen in gastric cancer [1–4]. Acquiring *H. pylori* infection is most intensive during childhood [5], and clearly, it occurs earlier in life in developing countries than in developed countries [6, 7]. Having a low socioeconomic status,

living in overcrowded conditions, and having a low maternal education are associated with an increased risk of *H. pylori* infection [6, 8–11].

Reports from Israel indicate seroprevalence rates of *H. pylori* infection of 46% among young adults, and 72% among residents of rural communities aged  $\geq 30$  years [4, 12]. There are no published data on the epidemiology of *H. pylori* infection among healthy paediatric populations, either Jewish or Arab, in Israel.

We carried out the present study among the Israeli Arabs who form a unique population at an advanced stage of epidemiological transition with a significant decline in infant mortality rates and a clear shift of mortality patterns from infectious diseases to chronic diseases [13]. We assessed the prevalence and potential risk factors of *H. pylori* infection among healthy 3- to 5-year-old Israeli Arab children living in three villages in the Meshulash region of Israel. We tried to answer

\* Author for correspondence: Professor D. Cohen, Department of Epidemiology and Preventive Medicine, Sackler Faculty of Medicine, Tel-Aviv University, Ramat Aviv, Tel Aviv 69978, Israel. (Email: dancohen@post.tau.ac.il)

the question whether the epidemiological characteristics of *H. pylori* infection in populations under epidemiological and environmental transition such as the Israeli Arab population, are different from those identified in developed or developing countries.

## METHODS

### Study population

The Israeli Arabs live in four separate geographic areas: the Galilee in northern Israel, the Meshulash in central Israel, the Negev in southern Israel, and in East Jerusalem. The majority live in rural areas. The Israeli Arabs have much lower educational levels and are at a lower socioeconomic status compared with the Jewish population. Substantial differences between the two groups are also observed regarding the sanitation infrastructure. Nevertheless, the Israeli Arab population is in a positive transition process, with ongoing improvement of the educational level and medical system. Israeli citizens, including Arabs, have mandatory health insurance according to national health insurance law. The coverage of vaccination in this community is over 95%.

Healthy children, aged 3–5 years, living in the rural Meshulash region of Israel, who were participating in a cohort follow-up study of enteric infections were also asked to participate in a cross-sectional study of *H. pylori* infection. The study children attended kindergartens from the villages of Jeser El-Zarka, Faradis, and Kfar Qaraa.

The Institution Review Board of Tel Aviv University approved the study. Written informed consent was obtained from the parents of the participating children.

### Collection of data

Mothers of the study children were interviewed in Arabic using a structured questionnaire. The interviews were performed at the participants' homes, by interviewers who received standard training. The questionnaire was validated by a pilot study in the target population and included information about the following sociodemographic and environmental factors: age, gender, place of residence, monthly family income, mother's education, mother's age, father's education, father's age, number of siblings, number of siblings <5 years, number of persons living in the household, number of rooms in the

household, and reported contact between houseflies and foods. A crowding index was computed using the following formula: (no. of persons living in the house)/(no. of rooms in the house).

### Anthropometric measurements

Anthropometric measurements were performed by specially trained registered nurses. Body weight was measured to the nearest 0.1 kg using an analogue personal scale (calibrated before use), and height (to the nearest 0.1 cm) with a mobile stadiometer. *z* scores: height for age (HAZ), weight for age (WAZ), and weight for height (WHZ) were computed using Epi-Info software (CDC, Atlanta, GA, USA). The calculations were based on the 2000 CDC growth reference curves, which were primarily based on the US National Health Examination (NHES) and the National Health and Nutrition Examination Surveys (NHANES).

### Collection of stool specimens

Mothers of the study children were asked to collect a stool sample from their child. For a subset of study children living in Jeser El Zarka and Faradis a stool sample was also requested from their mothers as well as the sib that was nearest to the age to the study child. Fresh specimens were obtained from subjects using collection cups. After collection, the specimens were kept and transported on ice-packs in thermally isolated boxes and within 24 h of collection, aliquoted and frozen at the research laboratory at  $-20^{\circ}\text{C}$  until tested. The storage and transportation of specimens were conducted by similar means and according to the same protocol in the three villages. The whole process was supervised by a field coordinator.

### Detection of *H. pylori* infection

A commercial enzyme linked immunoassay kit (Premier HpSA, Meridian Bioscience Inc., Cincinnati, OH, USA) employing polyclonal anti-*H. pylori* antibody adsorbed to microwells was used to detect *H. pylori* antigen in stools according to the manufacturer's instructions. Optical density values of  $\geq 0.120$ , between 0.100 and 0.119, and  $< 0.100$  were considered positive, equivocal and negative respectively.

### Data management and analysis

The data obtained were managed and analysed using the following software: Excel (Microsoft), Access

Table 1. Socioeconomic and environmental characteristics of the three villages participating in the study: Jeser El Zarka, Faradis, and Kfar Qaraa, Israel, June–September 2004

	Jeser El Zarka n/total (%)	Faradis n/total (%)	Kfar Qaraa n/total (%)	P value‡
Mother's education				
Elementary	61/79 (77.2)	23/64 (35.9)	9/54 (16.7)	<0.001
Family income				
>4000 NIS*	11/79 (13.9)	31/64 (48.4)	25/49 (51)	<0.001
Crowding index†				
>2	56/79 (70.9)	21/64 (32.8)	4/54 (7.4)	<0.001
Contact of flies with food	46/79 (58.2)	8/64 (12.5)	6/54 (11.1)	<0.001

\* New Israeli Shekel.

† Number of people living in the house divided by the number of rooms in the house.

‡  $\chi^2$  test. Two tailed  $P < 0.05$  significant.

2000 (Microsoft), and SPSS version 12 (SPSS Inc., Chicago, IL, USA). Data entry was carried out by two trained persons and quality control was performed at the different stages of the data management.

Differences in socioeconomic variables characterizing the three villages were examined using the  $\chi^2$  test. Proportions and 95% confidence intervals (CI) were computed to estimate the prevalence rates of *H. pylori*. The analysis of data on the potential predictive factors was performed by stratification according to the place of residence, Jeser El Zarka vs. Faradis and Kfar Qaraa using the Mantel–Haenszel test. The data on all the tested variables were homogenous as documented by the Breslow–Day and Tarone's tests of homogeneity indicating the possibility to run this pooled analysis. For each variable pooled odds ratios (OR) of infection and 95% CI were computed. Variables that were significantly associated with *H. pylori* infection in the Mantel–Haenszel analysis were included in multivariate analysis using logistic regression models to study the independent effect of each variable on the risk of having *H. pylori* infection while controlling for other variables in the model. Adjusted OR and 95% CI were computed for each variable. Two-tailed  $P < 0.05$  was considered significant.

## RESULTS

Between June 2004 and October 2004, 197 children (105 males), mean age 3.7 years (s.d. = 0.55), were enrolled in the study. Fifty-four lived in Kfar Qaraa, 64 in Faradis and 79 in Jeser El Zarka. The environmental and socioeconomic characteristics of the three villages were significantly different as revealed by

analysis of variables such as the mother's education, monthly family income, crowding index and reported contact of flies with food (Table 1).

A total of 98 positive, three equivocal, and 96 negative stool samples for *H. pylori* were identified. In all further analyses, the three equivocal subjects were classified as negative.

The overall prevalence rate was 49.7% (95% CI 42.8–56.6). The prevalence rate in Jeser El Zarka was over double that of Kfar Qaraa and Faradis: 75.9% (95% CI 65.4–84) compared to 33.3% (95% CI 22.2–46.6) and 31.3% (95% CI 21.1–43.3) respectively ( $P < 0.001$ ).

Seventy-four mothers and 65 siblings with the mean age of 6 years (s.d. = 3 years), one for each study child were also tested for *H. pylori*. The rate of *H. pylori* positivity was 82.4% among the mothers and 58.5% among the siblings. There were no significant differences between families who participated in this part of the study compared with those who did not participate according to place of residence, monthly family income, mother's education and crowding index.

In view of the substantial differences among the three villages in socioeconomic and environmental factors and the significant association between the village of residence and *H. pylori* infection rates, the analysis of the potential risk factors was performed with stratification by the village of residence; Jeser El Zarka vs. Faradis and Kfar Qaraa, using the Mantel–Haenszel test. This analysis revealed a significant increased risk for *H. pylori* infection among the study children with an increase in the crowding index, number of siblings <5 years and presence in the household of a sib with *H. pylori* infection (Table 2). No statistically significant differences in the

Table 2. Stratified analysis of risk factors of *H. pylori* infection among 3- to 5-year old Israeli Arab children, from the Meshulash region, Israel, June–September 2004

	No. of samples tested		<i>H. pylori</i> positive [n (%)]		OR <sub>MH</sub> (95% CI)	P value*
	Jeser El Zarka	Faradis/ Kfar Qaraa	Jeser El Zarka	Faradis/ Kfar Qaraa		
Gender						
Males	35	70	26 (74.3)	21 (30)	1.2 (0.6–2.3)	0.49
Females	44	48	34 (77.3)	17 (35.4)		
Age						
≤4 yr	54	82	39 (72.2)	26 (31.7)	1.3 (0.6–2.6)	0.4
>4 yr	25	36	21 (84)	12 (33.3)		
Mother's age						
≤33 yr (median)	43	66	33 (76.7)	24 (36.4)	0.7 (0.4–1.3)	0.32
>33 yr	36	52	27 (75)	14 (26.9)		
Father's age						
≤36 yr (median)	39	64	30 (76.9)	22 (34.4)	0.8 (0.4–1.6)	0.62
>36 yr	35	52	26 (74.3)	16 (30.8)		
Mother's education						
Elementary	61	32	46 (75.4)	7 (21.9)	1.6 (0.7–3.5)	0.19
High school/higher education	18	86	14 (77.8)	31 (36)		
Father's education						
Elementary	49	32	35 (71.4)	10 (31.3)	1.5 (0.7–3.1)	0.27
High school/higher education	23	82	20 (87)	28 (34.1)		
Family income						
≤4000 NIS†	68	57	51 (75)	21 (36.8)	0.8 (0.4–1.6)	0.54
>4000 NIS	11	56	9 (81.8)	16 (28.6)		
Maternal <i>H. pylori</i> infection§						
Negative	7	6	2 (28.6)	2 (33.3)	2.3 (0.6–8.2)	0.19
Positive	32	29	24 (75)	7 (24.1)		
Siblings' <i>H. pylori</i> infection§						
Negative	7	20	3 (42.9)	4 (20)	4.3 (1.3–14.5)	0.016
Positive	28	10	22 (78.6)	5 (50)		
No. of siblings						
≤3 (median)	34	89	23 (67.6)	28 (31.5)	1.5 (0.7–2.9)	0.22
>3	45	29	37 (82.2)	10 (34.5)		
No. of siblings aged 0–5 yr						
≤1 (median)	40	97	27 (67.5)	28 (28.9)	2.3 (1.1–4.9)	0.018
>1	38	21	32 (84.2)	10 (47.6)		
Crowding index‡						
≤2 (median)	23	93	12 (52.2)	26 (28)	3.3 (1.6–6.6)	0.001
>2	56	25	48 (85.7)	12 (48)		
Contact of food by houseflies						
Yes	46	14	38 (82.6)	5 (35.7)	1.7 (0.8–3.7)	0.15
No	33	104	22 (66.7)	33 (31.7)		

OR<sub>MH</sub>, Mantel–Haenzel odds ratio.

\* Mantel–Haenzel test  $P < 0.05$  significant.

† New Israeli Shekel.

‡ Number of people living in the house divided by the number of rooms in the house.

§ Only children from Faradis and Jeser El Zarka were included in the analysis.

prevalence rates of *H. pylori* infection were detected between genders and according to children's age groups, mother's education, father's education,

parents' age groups, monthly family income, number of siblings, maternal *H. pylori* infection, and reported contact of food by houseflies (Table 2).

Table 3. *Multivariate analysis of the association between different risk factors and H. pylori positivity among 3- to 5-year old Israeli Arab children, from the Meshulash region, Israel, June–September 2004*

Variables	Adjusted OR (95% CI)*	P value
Village		
Faradis (ref.)	1	
Jeser El Zarka	3.3 (1.03–11)	0.043
Sibling's <i>H. pylori</i> infection		
Negative (ref.)	1	
Positive	4.4 (1.3–14.6)	0.016

\* Adjusted for the variables: village, sibling's *H. pylori* infection, crowding index and siblings aged 0–5 years. The associations between crowding index and siblings aged 0–5 years and *H. pylori* infection were not significant in this model.

An initial multivariate logistic regression analysis was carried out on data related to 196 subjects; it involved the variables that showed a statistically significant association with *H. pylori* positivity in the Mantel–Haenszel analysis except the variable 'sibling's *H. pylori* infection' for which information was available on 65 children.

All the variables that were included in the model retained their significant association with the risk of *H. pylori* infection. Children from Jeser El Zarka were at a three-fold higher risk of being infected with *H. pylori* than children from Faradis and Kfar Qaraa, adjusted OR 3.3 (95% CI 1.6–6.8). Children from households with a crowding index of >2 persons per room were at around a three-fold higher risk of *H. pylori* infection compared to children with a crowding index of <2 persons per room, adjusted OR 3.3 (95% CI 1.6, 6.7). When the number of siblings <5 years old was more than one, the risk of *H. pylori* infection increased to 2.4 compared with children having one or no siblings <5 years old, adjusted OR 2.4 (95% CI 1.1–5).

A second multivariate analysis was performed on the 65 subjects for whom all the variables significantly associated with *H. pylori* positivity in the univariate analysis were available, including the variable 'sibling's *H. pylori* infection'. This model revealed that the risk of *H. pylori* infection was 3.3 times higher among children from Jeser El Zarka compared with children from Faradis (Table 3). Children having a sibling positive for *H. pylori* infection had a 4.5-fold higher risk of *H. pylori* infection than children with no

*H. pylori* infection in the tested sibling (Table 3). The variables: crowding index, and number of siblings <5 years old did not retain a significant association with *H. pylori* infection.

### The association between *H. pylori* infection and anthropometric measurements

The anthropometric measurements were similar among *H. pylori*-positive and -negative subjects. The mean differences in height for age *z* scores (HAZ), weight for age *z* scores (WAZ), and weight for height *z* scores (WHZ) among *H. pylori*-positive and -negative subjects were 0.158, –0.013, and –0.166 respectively ( $P=0.2, 0.9, 0.2$ ).

## DISCUSSION

This community-based study was conducted among children in the Israeli Arab population, going through an ongoing process of socioeconomic and environmental transition.

To the best of our knowledge there are no published data regarding the epidemiology of *H. pylori* infection among healthy children, either Jewish or Arab in Israel. Using a recently developed ELISA system for direct detection of specific *H. pylori* antigen in stool specimens, we found high prevalence rates of *H. pylori* infection among healthy 3- to 5-year-old children. Data of a recent follow-up study [14] carried out among children in the United States reconfirmed the validity of the ELISA antigen detection system used in our study. Haggerty *et al.* [14] reported that the vast majority of the positive stool samples for *H. pylori* as detected by stool antigen ELISA system, were also positive by PCR. Only a few antigen-positive stools harboured *Helicobacter* of other species than *H. pylori*.

A previous article from Israel studying the seroprevalence of *H. pylori* antibodies reported a 46.5% *H. pylori* infection rate among Jewish young adults aged 18–19 years [4], similar to the overall rate found in our study in a much younger age group (preschool children). These findings may suggest an earlier acquisition of *H. pylori* infection among Arabs as compared to Jews in Israel.

The prevalence rate of *H. pylori* infection in our study is similar to rates described among children in developing countries [6, 9]. It varied significantly among the three villages, but was the highest in Jeser El Zarka. The substantial differences in the rates of

*H. pylori* infection in these three villages can be explained by their different socioeconomic and living conditions. Variations in *H. pylori* infection rates in a defined geographic area were also observed in Germany and Italy but in each of these studies the participants belonged to different subpopulations, according to their nationality or regarding rural vs. urban communities [15, 16]. Our study controls for these differences and suggests that in the same ethnic group, within a small geographic area, and in a rural environment, there are groups at substantially high risk of *H. pylori* infection.

Living in a village of low socioeconomic status, under conditions of crowding, with a high number of young siblings and with a sibling positive for *H. pylori* were factors that showed a significant and independent association with an increased risk of *H. pylori* infection among the preschool children. Two of these predictors, namely the village of residence and carrying *H. pylori* within the family, emerged as being the strongest variables associated with the risk of *H. pylori* infection.

The importance of socioeconomic status, household crowding and intra-familial transmission regarding the risk of acquiring *H. pylori* infection in early childhood has been reported in a series of articles [9, 17–20]. Each of these studies controlled, however, for a limited number of potential confounders. Our study was carried out in a paediatric population attending kindergartens, homogeneous in terms of ethnicity, age, and geographic place of residence and controls for a large number of potential confounders.

The relationship between sibling's *H. pylori* positivity and the increased risk of *H. pylori* infection among the study children probably supports the possibility of intra-familial transmission rather than exposure to a common source of infection since the analysis included children from different villages.

We assume that close personal contact in families where carriers of *H. pylori* are already present among siblings or parents can markedly increase the risk of oral–oral, gastro–oral or faecal–oral transmission through inadvertently spreading infected saliva, vomitus or having poor personal hygiene. Different modes or mechanisms of intra-familial transmission of *H. pylori* have been proposed, for example, sharing a cup [20] and sharing a bed or a bedroom with a *H. pylori*-infected sibling in early childhood [21].

The association between the child's *H. pylori* infection and maternal *H. pylori* infection was not

significant and could be explained by the small sample size and high positivity rate among mothers.

The issue of a possible association between nutritional status and *H. pylori* infection is controversial. This can be the result of differences in the methodology applied in the various studies including different sample sizes and the variety of the study populations among which stronger correlates of nutritional status could mask a potential specific association. In our study, anthropometric measurements were not associated with *H. pylori* infection. Other cross-sectional studies from Egypt and Bangladesh [9, 10, 22] reported similar findings. However, data which was obtained through a nested case control among Peruvian children, aged  $\geq 2$  years, showed less weight gain 2 months after *H. pylori* antibody seroconversion [23].

In summary, in the Israeli Arab population going through an ongoing process of socioeconomic and environmental transition, the prevalence and risk factors of *H. pylori* infection among healthy children at the age of 3–5 years are similar to those previously reported in developing countries.

#### ACKNOWLEDGEMENTS

The study was supported by grant no. 01-18-00437 from the Israel Ministry of Science. The authors thank the fieldwork team: Hiba Abu-Abed, Shiraz Muhsen, Ola Abu-Shehab, Manal Jurban, Roza Marai, for their contribution in the process of data collection, and Sophie Goren for her help in the data management.

#### DECLARATION OF INTEREST

None.

#### REFERENCES

1. **Suerbaum S, Michetti P.** *Helicobacter pylori* Infection. *New England Journal of Medicine* 2002; **347**: 1175–1187.
2. **Torres J, et al.** Comprehensive review of the natural history of *Helicobacter pylori* infection in children. *Archives of Medical Research* 2000; **31**: 431–469.
3. **Goodwin CS, Mendall MM, Nortfield TC.** *Helicobacter pylori* infection. *Lancet* 1997; **349**: 265–269.
4. **Gdalevich M, et al.** *Helicobacter pylori* infection and subsequent peptic duodenal disease among young adults. *International Journal of Epidemiology* 2000; **29**: 592–595.

5. **Mitchell A, et al.** Age-specific *Helicobacter pylori* seropositivity rates of children in an impoverished urban areas of northeast Brazil. *Journal of Clinical Microbiology* 2003; **41**: 1326–1328.
6. **Frenck RW, Clemens J.** *Helicobacter* in the developing world. *Microbes and Infection* 2003; **5**: 705–713.
7. **Malaty HM, et al.** Age at acquisition of *Helicobacter pylori* infection: a follow up study from infancy to adulthood. *Lancet* 2002; **16**: 931–935.
8. **Wizla-Derambure N, et al.** Familial and community environmental risk factors for *Helicobacter pylori* infection in children and adolescents. *Journal of Pediatrics and Gastroenterological Nutrition* 2001; **33**: 58–63.
9. **Clemens J, et al.** Sociodemographic, hygienic and nutritional correlates of *Helicobacter pylori* infection of young Bangladeshi children. *Pediatrics Infectious Disease Journal* 1996; **15**: 1113–1118.
10. **Mahalanabis D, et al.** *Helicobacter pylori* infection in the young in Bangladesh: prevalence, socioeconomic and nutritional aspects. *International Journal of Epidemiology* 1996; **25**: 894–898.
11. **Dominici P, et al.** Familial clustering of *Helicobacter pylori* infection: population based study. *British Medical Journal* 1999; **319**: 537–540.
12. **Gilboa S, et al.** *Helicobacter pylori* infection in rural settlements (Kibbutzim) in Israel. *International Journal of Epidemiology* 1995; **24**: 232–237.
13. **Israeli Center for Disease Control.** Ministry of Health, Health status in Israel 1999. Pitah Tikva: Ofset color, 1999; **37**: 45.
14. **Haggerty TD, et al.** Significance of transiently positive enzyme-linked immunosorbent assay results in detection of *Helicobacter pylori* in stool samples from children. *Journal of Clinical Microbiology* 2005; **43**: 2220–2223.
15. **Rothenbacher D, et al.** Prevalence and determinants of *Helicobacter pylori* infection in preschool children: a population based study from Germany. *International Journal of Epidemiology* 1998; **27**: 135–141.
16. **Dore MP, et al.** Risk factors associated with *Helicobacter pylori* infection among children in a defined geographic area. *Clinical Infectious Diseases* 2002; **35**: 240–245.
17. **Goodman K, Correa P.** Transmission of *Helicobacter pylori* among siblings. *Lancet* 2000; **355**: 358–362.
18. **Rothenbacher D, et al.** Role of infected parents in transmission of *Helicobacter pylori* to their children. *Pediatric Infectious Disease Journal* 2002; **21**: 674–679.
19. **Rocha GA, et al.** Transmission of *Helicobacter pylori* infection in families of preschool aged children from Minas Gerais, Brazil. *Tropical Medicine and International Health* 2003; **8**: 987–991.
20. **Rodrigues MN, et al.** Prevalence of *Helicobacter pylori* infection in children from urban community in north-east Brazil and risk factors of the infection. *European Journal of Gastroenterology and Hepatology* 2004; **16**: 201–205.
21. **Farrell S, et al.** Risk factors of *Helicobacter pylori* infection in children; an examination of the role played by intrafamilial bed sharing. *Pediatric Infectious Disease Journal* 2005; **24**: 149–152.
22. **Naficy AB, et al.** Seroepidemiology of *Helicobacter pylori* infection in a population of Egyptian children. *International Journal of Epidemiology* 2000; **29**: 928–932.
23. **Passaro DJ, et al.** Growth slowing after acute *Helicobacter pylori* infection is age dependent. *Journal of Pediatrics and Gastroenterological Nutrition* 2002; **35**: 522–526.