

# Mast Cell-derived Prostaglandin D<sub>2</sub> Controls Hyaluronan Synthesis in Human Orbital Fibroblasts via DP1 Activation

## IMPLICATIONS FOR THYROID EYE DISEASE\*

Received for publication, October 9, 2009, and in revised form, February 23, 2010. Published, JBC Papers in Press, March 22, 2010, DOI 10.1074/jbc.M109.074534

Naxin Guo<sup>†</sup>, Carolyn J. Baglole<sup>§¶1</sup>, Charles W. O'Loughlin<sup>†</sup>, Steven E. Feldon<sup>†</sup>, and Richard P. Phipps<sup>†§¶2</sup>

From the Departments of <sup>†</sup>Ophthalmology and <sup>§</sup>Environmental Medicine and the <sup>¶</sup>Lung Biology and Disease Program, University of Rochester School of Medicine and Dentistry, Rochester, New York 14642

Thyroid eye disease (TED) is a debilitating disorder characterized by the accumulation of adipocytes and hyaluronan (HA). Production of HA by fibroblasts leads to remarkable increases in tissue volume and to the anterior displacement of the eyes. Prostaglandin D<sub>2</sub> (PGD<sub>2</sub>), mainly produced by mast cells, promotes orbital fibroblast adipogenesis. The mechanism by which PGD<sub>2</sub> influences orbital fibroblasts and their synthesis of HA is poorly understood. We report here that mast cell-derived PGD<sub>2</sub> is a key factor that promotes HA biosynthesis by orbital fibroblasts. Primary orbital fibroblasts from TED patients were isolated and used to test the effects of PGD<sub>2</sub>, prostaglandin J<sub>2</sub>, as well as prostaglandin D receptor (DP) agonists and antagonists on HA synthesis. The expression of HA synthase (HAS), hyaluronidase, DP1, and DP2 mRNA levels was assessed by PCR. Small interfering RNAs against *HAS1* or *HAS2* were used to assess the importance of HAS isoforms on HA production. Treatment of human orbital fibroblasts with PGD<sub>2</sub> and PGJ<sub>2</sub> increased HA synthesis and HAS mRNA. *HAS2* was the dominant isoform responsible for HA production by PGD<sub>2</sub>. The effect of PGD<sub>2</sub> on HA production was mimicked by the selective DP1 agonist BW245C. The DP1 antagonist MK-0524 completely blocked PGD<sub>2</sub>-induced HA synthesis. Human mast cells (HMC-1) produced PGD<sub>2</sub>. Co-culture of HMC-1 cells with orbital fibroblasts induced HA production and inhibition of mast cell-derived PGD<sub>2</sub> prevented HA synthesis. Mast cell-derived PGD<sub>2</sub> increased HA production via activation of DP1. Selectively targeting the production of PGD<sub>2</sub> and/or activation of DP1 may prevent pathological changes associated with TED.

Graves disease is an autoimmune disorder in which 40–60% of patients develop Graves ophthalmopathy, also called thyroid eye disease (TED).<sup>3</sup> TED is characterized by expansion of the

orbital fat compartment and extraocular muscles (1). Intense inflammation and infiltration of immune cells, including T cells, macrophages, and mast cells, in the retrobulbar space of TED patients are key factors that drive the proliferation and differentiation of orbital fibroblasts to adipocytes (2, 3). In addition to the adipogenic potential of orbital fibroblasts, these cells are also key producers of extracellular matrix glycosaminoglycans. One of the key pathological findings in TED patients is the overproduction and accumulation of the glycosaminoglycan hyaluronan (HA). The extremely hydrophilic nature of HA leads to remarkable increases in tissue volume and to the anterior displacement of the eye, or exophthalmos (4), resulting in the disfigurement and vision impairment (5) characteristic of TED.

HA is synthesized as an acidic, negatively charged, high molecular weight polysaccharide via the actions of hyaluronan synthases (HAS) (6), of which there are three isoforms: *HAS1*, *HAS2*, and *HAS3*. Increased HA synthesis closely correlates with the expression levels of HAS (7), which are themselves induced by growth factors, cytokines (7–9), and prostaglandins (PGs) (10, 11). One PG that may have an important implication in TED is PGD<sub>2</sub>. PGD<sub>2</sub> is a metabolite of arachidonic acid that is formed by the actions of cyclooxygenases and PGD<sub>2</sub> synthases (PGDS) (12, 13). Many of the biological actions of PGD<sub>2</sub> are mediated through two G protein-coupled receptors: DP receptor 1 (DP1) and DP2 (also called chemoattractant receptor-homologous molecule (CRTH2)) (14–16). These receptors elicit divergent effects by the coupling to either G<sub>s</sub> (DP1) or G<sub>i</sub> (DP2) to elevate cyclic AMP (cAMP) or intracellular calcium (Ca<sup>2+</sup>), respectively. PGD<sub>2</sub> can also spontaneously undergo a series of dehydration reactions to form the PGJ family of prostaglandins, including 15d-PGJ<sub>2</sub>, an endogenous ligand for the peroxisome proliferator-activated receptor (PPAR $\gamma$ ) (17, 18).

Human orbital fibroblasts express PPAR $\gamma$  (3), and PPAR $\gamma$  is crucial for the differentiation of fibroblasts to adipocytes (19). We recently published that activated human T lymphocytes isolated from patients with TED produce much more PGD<sub>2</sub>-derived PGs compared with T cells from healthy individuals (3). Mast cells are also a key cellular source of PGs, with PGD<sub>2</sub> being the major prostanoid released (3, 20, 21). Mast cells are a central immune cell in the pathogenesis of TED. Not only is there

\* This research was supported by EY017123 and EY011708 and a Research to Prevent Blindness Foundation Grant.

<sup>1</sup> Supported by a Parker B. Francis Fellowship and an American Thoracic Society Research Grant.

<sup>2</sup> To whom correspondence should be addressed: University of Rochester School of Medicine and Dentistry, Dept. of Environmental Medicine, Box 850, 601 Elmwood Ave., Rochester, NY 14642. Tel.: 585-275-8326; Fax: 585-506-0239; E-mail: Richard\_Phipps@urmc.rochester.edu.

<sup>3</sup> The abbreviations used are: TED, thyroid eye disease; cAMP, cyclic AMP; DP, D prostanoid receptor; ELISA, enzyme-linked immunosorbent assay; FBS, fetal bovine serum; HA, hyaluronan; HAS, hyaluronan synthase(s); PGD and PGJ, prostaglandin D and J, respectively; PGDS, PGD<sub>2</sub> synthase; H-PGDS, hematopoietic type PGD<sub>2</sub> synthase; HYAL, hyaluronidase; RT, reverse transcription; qRT, quantitative RT; siRNA, small interfering RNA;

PPAR $\gamma$ , peroxisome proliferator-activated receptor  $\gamma$ ; PG, prostaglandin(s); IBMX, isobutylmethylxanthine; SC, scrambled control.

intense mast cell infiltration and degranulation (22) associated with adipocytes (23) in TED patients, but co-culture of mast cells with orbital fibroblasts up-regulates HA synthesis (24). Whether this increase in HA production by orbital fibroblasts is the result of PGD<sub>2</sub> acting via the direct modulation of DP receptors is not known. Herein, we report for the first time that PGD<sub>2</sub> increases HA synthesis in orbital fibroblasts via the induction of HAS2. Pharmacological inhibition of DP1, but not DP2, prevented the PGD<sub>2</sub>-induced up-regulation of HA. We also show that inhibition of PGD<sub>2</sub> synthesis by mast cells prevents HA synthesis by orbital fibroblasts. These results have important implications for future therapies directed against treating those afflicted with TED. Preventing PGD<sub>2</sub> synthesis by mast cells and/or DP1 activation on orbital fibroblasts may reduce the severity of the disease.

## EXPERIMENTAL PROCEDURES

### Reagents

PGD<sub>2</sub>, PGJ<sub>2</sub>, MK0524, ramatroban (BAY-u3405), BW 245C, 4-benzhydryloxy-1-[3-(1*H*-tetrazol-5-yl)-propyl]-piperidine (HQL-79), 13,14-dihydro-15-keto-PGD<sub>2</sub>, anti-DP1, anti-hematopoietic type PGDS (H-PGDS), anti-Cox-1 (cyclooxygenase-1), and anti-Cox-2 antibodies were purchased from Cayman Chemical (Ann Arbor, MI). Forskolin, A23187, and isobutylmethylxanthine (IBMX) were purchased from Sigma. Anti-DP2 antibody was purchased from R&D Systems (Minneapolis, MN). All of the drugs were dissolved in DMSO.

### Tissue Collection and Cell Culture

**Orbital Fibroblasts**—Primary orbital fibroblasts were isolated from TED patients undergoing orbital decompression surgery. The protocols for orbital biopsy and blood sample (see below) were approved by the internal review board, and informed, written consent was obtained from all patients. The primary fibroblasts were established by standard explant techniques (25, 26) and cultured in RPMI 1640 supplemented with 10% fetal bovine serum (FBS) (Hyclone, Logan, UT), 2-mercaptoethanol (Eastman Kodak Co.), L-glutamine (Invitrogen), HEPES (U.S. Biochemical Corp.), nonessential amino acids, sodium pyruvate, and gentamicin (Invitrogen). Fibroblasts were characterized by their adherent morphology; expression of vimentin and types I and III collagen; and absence of CD45, factor VIII, or cytokeratin. Fibroblasts were used at the earliest passage possible (between passages 4 and 10).

**Human T Cells**—Lymphocytes were isolated from 60 ml of peripheral blood obtained during orbital decompression surgery. Whole blood was separated over a Ficoll-Paque Plus gradient (Amersham Biosciences) to obtain peripheral blood mononuclear cells. T cells were enriched using CD3/CD28 T cell Expander beads (Invitrogen). Specifically,  $5 \times 10^6$  lymphocytes were incubated with CD3/CD28 beads at a 1:1 ratio in RPMI 1640 with 10% FBS medium at 37 °C for 2 days. After that, 50 units/ml recombinant interleukin-2 was added to the culture and incubated for another 2–3 days. On day 5, cells were diluted to a concentration of  $0.5 \times 10^6$  cells/ml in medium containing 50 units/ml recombinant interleukin-2 and incubated for an additional 3–7 days. Cellular purity was assessed using an anti-CD3-PE antibody (BD Biosciences, San Jose, CA)

and analyzed on a FACSCalibur flow cytometer (BD Biosciences). The T-cell purity was >95% (data not shown).

### Co-culture of Mast Cells with Orbital Fibroblasts

HMC-1 mast cells were allowed to proliferate in Iscove's modified Dulbecco's medium enriched with 5% FBS. Co-cultures were initiated by introducing HMC-1 cells to confluent cultures of fibroblasts as described previously (24). Mast cells were washed with RPMI 1640 medium (0.5% FBS) once before the addition to the fibroblasts (1:1 ratio unless otherwise specified). After 4 h, the medium and HMC-1 cells were removed by gently rinsing in phosphate-buffered saline, and fresh RPMI 1640 medium (0.5% FBS) was used to cover the cells. Alternatively, fibroblasts and mast cells were co-cultured in a transwell system (0.4  $\mu$ m; Greiner Bio-one, New York) with fibroblasts being cultured on the bottom and the mast cells in the top chamber. The ratio of mast cells to fibroblasts was varied from 5:1 to 20:1.

### Quantitation of HA

Aliquots of culture medium were removed at the indicated time points and centrifuged (5 min,  $8,000 \times g$  at 4 °C), and HA levels were quantified by enzyme-linked immunosorbent assay (ELISA) according to the manufacturer's protocol (R&D Systems). Briefly, samples were incubated in HA-binding protein-coated microwells, allowing HA in samples to react with the immobilized HA-binding protein. After the removal of unbound molecules by washing, biotinylated HA-binding protein was added to the microwells. Following a second wash, streptavidin conjugated to horseradish peroxidase was added. After the last wash, the chromogenic substrate was added, and HA levels were determined using a Varioskan Flash plate reader (Thermo Fisher Scientific, Milford, MA).

### Agarose Gel Electrophoresis of HA

Fibroblasts were cultured to confluence in 10-cm<sup>2</sup> dishes in RPMI containing 10% FBS, followed by RPMI with 2% FBS for 3 days. Fibroblasts were then treated with 5  $\mu$ M PGD<sub>2</sub> or control (DMSO) for 18 h. Cell culture supernatant was centrifuged (5 min at  $8,000 \times g$  at 4 °C), and half of the sample was digested with 1 turbidity-reducing unit/ml *Streptomyces* hyaluronidase (Sigma) at 37 °C for 4 h; the remainder of the sample was mock-digested. Following digestion with proteinase K (120  $\mu$ g/ml) (Qiagen, Valencia, CA) at 37 °C for 4 h, the samples were concentrated using a Vivaspin 10,000 Da cut-off ultrafiltration spin column (Sartorius Stedim Biotech, Goettingen, Germany), washed with phosphate-buffered saline, and analyzed by 1% agarose electrophoresis. Select-HA HiLadder (Hyalose LLC, Oklahoma City, OK) and blue dextran (Sigma; molecular mass  $2 \times 10^6$  Da) were also run as standards. The gel was stained with 0.005% Stains-All (Sigma) in 50% ethanol overnight at room temperature before being scanned on an HP scanner.

### Western Blotting

Following treatments, cells were lysed in 1 $\times$  SDS sample buffer, and protein concentrations were determined using the DC protein assay kit (Bio-Rad). Samples were separated on a 9 or 12% SDS-polyacrylamide gel, transferred to polyvinylidene

## PGD<sub>2</sub> Controls Hyaluronan Synthesis in Orbital Fibroblasts

difluoride membranes, and subjected to immunoblotting with antibodies against DP1 (1:1,000), DP2 (1:500), H-PGDS (1:500), Cox-1 (1:500), or Cox-2 (1:500). As a loading control, membranes were reprobed for glyceraldehyde-3-phosphate dehydrogenase (GAPDH; 1:3,000; Calbiochem).

### Measurement of cAMP

cAMP levels were quantified using a time-resolved fluorescence resonance energy transfer immunoassay from PerkinElmer Life Sciences according to the manufacturer's protocol. Briefly, orbital fibroblasts were detached with a non-enzymatic cell dissociation solution (Versene, Invitrogen) and resuspended in Stimulation Buffer (pH 7.4; 1× HBSS buffer containing 5 mmol/liter HEPES buffer and 0.01–0.1% bovine serum albumin) containing 0.5 mM IBMX and labeled with the Alexa Fluor® 647-labeled anti-cAMP antibody. Finally, the cells were incubated with drugs at the indicated times. Following the addition of the detection buffer, the amount of cAMP signal was measured.

### Measurement of PGD<sub>2</sub>

HMC-1 cells (1 × 10<sup>6</sup>/well) were seeded into 6-well plates and cultured in Iscove's modified Dulbecco's medium with 5% FBS in the absence or presence of HQL-79 (H-PGDS inhibitor, 100 μM) at 37 °C for 60 min. The cells were then washed and resuspended in RPMI 1640 medium with 0.5% FBS. After a 2-h incubation, the cells were stimulated with 10 μM calcium ionophore A23187 for 30 min. The culture media were collected, and PGD<sub>2</sub> in the supernatants was quantified by PGD<sub>2</sub>-MOX EIA kits (Cayman Chemical).

### Reverse Transcription (RT)-PCR and Quantitative RT (qRT)-PCR

RNA was isolated from orbital fibroblasts or T cells using an RNeasy minikit (Qiagen) and reverse transcribed to cDNA using an iScript cDNA synthesis kit (Bio-Rad) and diluted 5-fold in RNase-free H<sub>2</sub>O. Forward and reverse primers for each gene are shown in Table 1. The PCR was performed using the Qiagen Fast Cycling PCR kit for 30 cycles in a Bio-Rad thermal cycler. The PCR product was visualized on a 5% acrylamide gel stained with ethidium bromide. Amplifications without reverse transcriptase were carried out as negative controls. qRT-PCR was performed in a Bio-Rad iCycler containing B-R SYBR Green SuperMix for IQ (Quanta Biosciences, Gaithersburg, MD). Efficiency of the amplification was determined to be >90%.

Amplification of 7 S ribosomal RNA was carried out for each cDNA (in triplicate) for normalization. The threshold cycle number (C<sub>t</sub>) of amplification in each sample was determined by Bio-Rad software, and the relative mRNA abundance was calculated as the C<sub>t</sub> for amplification of a gene-specific cDNA minus average C<sub>t</sub> for 7 S, expressed as a power of 2 (*i.e.* 2<sup>ΔC<sub>t</sub></sup>).

### Gene Knockdown Using siRNA

Orbital fibroblasts were cultured to 80–90% confluence and transfected with 80 nM *HAS1*, *HAS2*, or scrambled control (SC) siRNA (Santa Cruz Biotechnology, Inc., Santa Cruz, CA) using Lipofectamine 2000 (Invitrogen). Forty-eight hours post-transfection, RPMI containing 0.5% FBS was added. Twenty-four

**TABLE 1**

**Gene primers**

Gene	Primer sequence	Size <i>bp</i>
<i>HAS1</i> (NM-001523)	Forward	5'-TGTGTATCCTGCATCAGCGGT-3'
	Reverse	5'-CTGGAGGTGTACTTGGTAGCATAAACC-3'
<i>HAS2</i> (NM-005328)	Forward	5'-GCCTCATCTGTGGAGATGGT-3'
	Reverse	5'-ATGCACTGAACACACCCAAA-3'
<i>HAS3</i> (NM-005329)	Forward	5'-GGCATTATCAAGGCCACCTA-3'
	Reverse	5'-AGGCCAATGAAGTTCACCAC-3'
<i>HYAL1</i> (NM-007312)	Forward	5'-GATGTCAGTGTCTTCGATGTGGTA-3'
	Reverse	5'-GGGAGCTATAGAAAATGTCATGTCA-3'
<i>HYAL2</i> (NM-003773)	Forward	5'-CTAATGAGGGTTTGTGAACCAGAATAT-3'
	Reverse	5'-GCAGAATCGAAGCGTGGATAC-3'
<i>HYAL3</i> (NM-003549)	Forward	5'-CCTCCAGTGCCCTCTTCC-3'
	Reverse	5'-ACAAGGTCATCCTGGGACAG-3'
<i>DPI</i> (NM-000953)	Forward	5'-TCTGCGCGTACCTTTCATG-3'
	Reverse	5'-TCCTCGTGGACCATCTGGATA-3'
<i>DP2</i> (NM-004778)	Forward	5'-TTTCTCAACATGTTCCGCCAG-3'
	Reverse	5'-AAGCACCAGGCAGACTTTGT-3'
7 S RNA (NR-002715)	Forward	5'-ACCACCAGGTTGCCTAAGGA-3'
	Reverse	5'-CACGGAGTTTGCCTGCT-3'

hours later, the cells were treated with or without 5 μM PGD<sub>2</sub> for 2 h (for qRT-PCR analysis) or 18 h (for analysis of HA secretion).

### Statistical Analysis


Statistical analysis was performed using GraphPad Prism (GraphPad Software, Inc., La Jolla, CA). For comparison between groups of three or more, an analysis of variance with a Newman-Keuls multiple comparison test was used to determine differences between treatments. *Error bars* represent the S.D. of triplicate samples. A *p* value of <0.05 is considered significant. All experiments were performed at least three times.

## RESULTS

*Prostaglandins of the D and J Series Induce HA Synthesis by Human Orbital Fibroblasts*—We previously reported that PGD<sub>2</sub> and PGJ<sub>2</sub> induce adipogenesis in human orbital fibroblasts (3). Here, we examined whether PGD<sub>2</sub> or PGJ<sub>2</sub> would also induce HA production. Two strains of human orbital fibroblasts (OF1 and OF2) from two different TED patients were treated with increasing concentrations of PGD<sub>2</sub> or PGJ<sub>2</sub> for 18 h, and HA levels were detected in the cell culture supernatant by a commercial HA ELISA. Fig. 1A shows that basal levels of HA differed significantly between the two orbital fibroblast strains (OF1 (1.9 ± 1.4 ng/μg protein) *versus* OF2 (0.5 ± 0.6 ng/μg protein); #, *p* < 0.05). Treatment of the fibroblast strain designated OF1 with either PGJ<sub>2</sub> or PGD<sub>2</sub> resulted in significantly more HA production compared with OF2. PGD<sub>2</sub> and PGJ<sub>2</sub> also significantly increased HA synthesis by both fibroblast strains in a dose-dependent manner, with 5 μM PGD<sub>2</sub> and 2 μM PGJ<sub>2</sub> yielding the largest induction (Fig. 1A). Higher concentrations of PGD<sub>2</sub> or PGJ<sub>2</sub> did not further increase HA syn-



## PGD<sub>2</sub> Controls Hyaluronan Synthesis in Orbital Fibroblasts



**FIGURE 1. PGD<sub>2</sub> and PGJ<sub>2</sub> induce HA synthesis in human orbital fibroblasts.** *A*, confluent strains of human orbital fibroblasts (OF1 and OF2) were cultured in RPMI 1640 with 0.5% FBS for 3 days prior to treatment with PGJ<sub>2</sub>, PGD<sub>2</sub>, or vehicle (DMSO) for 18 h. The cell culture medium was assayed for HA by an HA ELISA as described under "Experimental Procedures." There was a significant increase in HA production by two strains of orbital fibroblasts following treatment with PGD<sub>2</sub> (1–5 μM) and PGJ<sub>2</sub> (2 μM). The experiment was performed in triplicate. \*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$  compared with vehicle control. #,  $p < 0.05$ ; ###,  $p < 0.001$ , OF1 versus OF2. Results are expressed as the mean  $\pm$  S.D. (*error bars*). *B*, agarose gel HA analysis: Orbital fibroblasts (OF1 and OF2) were cultured in RPMI 1640 with 2% FBS for 3 days and then treated with 5 μM PGD<sub>2</sub> (D<sub>2</sub>) or vehicle (V) for 18 h, and the conditioned medium was analyzed by agarose gel electrophoresis. Both OF1 and OF2 exhibited basal HA (*lanes 3 and 5, respectively; blue*). When treated with PGD<sub>2</sub>, there was an increase in color intensity, indicating increased HA production (*lanes 4 and 6; compare with lanes 3 and 5*). A *Streptomyces* hyaluronidase-digested sample (HA'ase) (from PGD<sub>2</sub>-treated OF2) was included as a negative control (*lane 7*).

thesis in either of the two fibroblast strains (data not shown). Therefore, most of the remaining experiments were conducted using 5 μM PGD<sub>2</sub> or 2 μM PGJ<sub>2</sub>.

We also performed agarose gel electrophoresis to analyze the molecular size(s) of HA produced by the orbital fibroblasts. HA is an extremely high molecular weight ( $M_r$ ) polysaccharide ( $\sim 10^5$  to  $4 \times 10^6$  Da) and can be separated from other glycosaminoglycans by ultrafiltration combined with agarose gel electrophoresis (27). Fig. 1*B* confirms the presence of HA in the cell culture media of two fibroblast strains (OF1 and OF2) (*lanes 3–6, blue*). Treatment with PGD<sub>2</sub> (D<sub>2</sub>) increased HA compared with vehicle (V) (*lanes 4 and 6; compare with lanes 3 and 5*). OF1 also produces relatively more low  $M_r$  HA compared with OF2. We also used *Streptomyces* hyaluronidase to digest the HA in the samples. Unlike other hyaluronidases, this enzyme is specific for hyaluronic acid and is inactive with chondroitin and chondroitin sulfate (28). The *Streptomyces* hyaluronidase-digested sample (HA'ase, *lane 7*) did not show any blue staining, confirming the HA specificity of the assay.

Together, these results demonstrate that orbital fibroblasts increase the production of HA when exposed to PGD<sub>2</sub>.

*Differential Expression of PGD<sub>2</sub>-induced Hyaluronidase (HYAL) in Two Orbital Fibroblast Strains—*HYALs (HYAL1 to -3) depolymerize HA into low  $M_r$  polymers (29). To determine if the apparent increase in low  $M_r$  HA in PGD<sub>2</sub>-treated OF1 fibroblast strain might be the result of increased HYAL expression, we treated OF1 and OF2 with 5 μM PGD<sub>2</sub> and assessed mRNA. There was a significant induction in HYAL1, HYAL2, and HYAL3 mRNA following treatment of OF1 with PGD<sub>2</sub> (Fig. 2). In contrast, OF2 failed to significantly increase HYAL mRNA expression. The induction in HYAL in OF1 caused by PGD<sub>2</sub> was significantly greater than PGD<sub>2</sub>-treated OF2 HYAL mRNA levels (Fig. 2). These results are the first to demonstrate regulation of HYAL mRNA by PGD<sub>2</sub> in human orbital fibroblasts.


*PGD<sub>2</sub> Induces HAS mRNA Expression in Human Orbital Fibroblasts—*We next evaluated the effect of PGD<sub>2</sub> on HAS mRNA expression. Fig. 3*A* shows that all three HAS genes are expressed in orbital fibroblasts. In untreated cells (vehicle), HAS1 mRNA is very low and scarcely detectable. In contrast, HAS2 and HAS3 mRNA are relatively greater. Following 2 h of PGD<sub>2</sub> treatment, the mRNA for all three

HAS isoforms increased significantly (Fig. 3, *A* and *B*), with HAS1 exhibiting the greatest induction. The increase in HAS1 at 2 h was significantly greater compared with either HAS2 or HAS3 (Fig. 3*B*; ###,  $p < 0.001$ ). The mRNA for all three HAS isoforms declined rapidly, where by 16 h, there was no significant difference compared with untreated (0 h). 7 S ribosomal mRNA expression remained the same, regardless of treatment (Fig. 3*A*, compare *lanes 5* and *9*) and was used to normalize the qRT-PCR results (Fig. 3*B*).

*PGD<sub>2</sub>-induced HA Production by Orbital Fibroblasts Is Dependent on HAS2 Expression—*To verify whether or not HA synthesis was directly related to HAS expression, siRNA was used to selectively knock down HAS1 or HAS2 mRNA levels in fibroblasts that were treated with PGD<sub>2</sub>. Fig. 4*A* shows that siRNA directed against HAS1 or HAS2 selectively and significantly reduced (by  $\sim 75$ –80% of mRNA levels) PGD<sub>2</sub>-induced HAS1 (*open bars*) and HAS2 (*black bars*), compared with the SC siRNA.

We then analyzed for HA in orbital fibroblasts that were untreated (*open bars*) or treated with PGD<sub>2</sub> (*black bars*) after

## PGD<sub>2</sub> Controls Hyaluronan Synthesis in Orbital Fibroblasts



**FIGURE 2. Differential induction in HYAL mRNA between two orbital fibroblast strains.** cDNA from orbital fibroblast strains OF1 and OF2 treated with 5  $\mu$ M PGD<sub>2</sub> or vehicle (Untreated) for 4 h was assessed by qRT-PCR. There was a significant increase in mRNA expression for all three HYAL isoforms (HYAL1, HYAL2, and HYAL3) when OF1 was treated with 5  $\mu$ M PGD<sub>2</sub> (\*\*\*,  $p < 0.0001$  compared with untreated). There was no significant increase in HYAL1 to -3 in PGD<sub>2</sub>-treated OF2 cells (ns, compared with untreated). The induction of HYAL1, HYAL2, and HYAL3 mRNA in PGD<sub>2</sub>-treated OF1 was significantly higher than PGD<sub>2</sub>-exposed OF2 HYAL expression (\$\$\$,  $p < 0.0001$  for each respective HYAL). Results are expressed as the mean  $\pm$  S.D. (error bars) ( $n = 3$ ).

transfection with siRNA against *HAS1* and/or *HAS2* (Fig. 4B). Those fibroblasts transfected with the SC siRNA exhibited a significant increase in HA synthesis in response to PGD<sub>2</sub> (Fig. 4B; compare with Fig. 1A). Despite the significant increase in *HAS1* mRNA induced by PGD<sub>2</sub> (Fig. 3), attenuation of *HAS1* expression did not block PGD<sub>2</sub>-induced HA production (Fig. 4B). In contrast, reduction in *HAS2* mRNA expression significantly reduced both untreated and PGD<sub>2</sub>-induced HA production when compared with the SC siRNA control (Fig. 4B). This reduction in HA synthesis in the *HAS2* siRNA-transfected, PGD<sub>2</sub>-treated fibroblasts was not significantly different compared with untreated SC siRNA-transfected cells (ns,  $p = 0.13$ ). These data support that *HAS2* is the dominant isoform responsible for the increased HA synthesis by orbital fibroblasts in response to PGD<sub>2</sub>.

**PGD<sub>2</sub> and PGJ<sub>2</sub> Induce HA Production in Human Orbital Fibroblasts via DP1, but Not DP2, Activation**—Both PGD<sub>2</sub> and PGJ<sub>2</sub> can exert their biological functions via activation of DP1 or DP2 (30). We therefore examined whether or not human orbital fibroblasts express DP receptors. RT-PCR and Western blot analysis revealed that orbital fibroblasts (OF1 and OF2) express both *DP1* and *DP2* mRNA and protein, respectively (Fig. 5, A and B). Protein expression of DP1 and DP2 in orbital fibroblasts was comparable with that of human T cells, which are well known to express DP receptors, particularly DP2 (15, 31). Both orbital fibroblast strains (OF1 and OF2) expressed considerably more DP1 mRNA and protein compared with DP2 (Fig. 5, A and B).

To determine which DP was involved in the PGD<sub>2</sub>-mediated induction of HA, we first utilized well described selective pharmacological antagonists directed against either DP1 (MK-0524) (32, 33) or DP2 (ramatroban) (30, 34). Fibroblasts were untreated or were pretreated with either MK-0524 or ramatroban for 1 h with or without PGD<sub>2</sub> or PGJ<sub>2</sub>. Fig. 5C shows that neither MK-0524 (MK) nor ramatroban (RAM) influenced basal HA levels (no significance (ns) compared with untreated;


open bar). Fibroblasts treated with either PGD<sub>2</sub> (5  $\mu$ M) or PGJ<sub>2</sub> (2  $\mu$ M) significantly increased HA synthesis (Fig. 5C; compare with Figs. 1A and 4B). Pretreatment with the DP1 antagonist MK-0524 completely blocked the ability of both PGD<sub>2</sub> and PGJ<sub>2</sub> to induce HA production in orbital fibroblasts. By contrast, pretreatment with the DP2 antagonist ramatroban was unable to prevent the induction of HA by PGD<sub>2</sub> or PGJ<sub>2</sub>. This suggests that DP1 is the dominant receptor subtype involved in the induction of HA by PG of the D and J series.

To confirm the importance of DP1 in the induction of HA in human orbital fibroblasts, we utilized BW245C, a selective DP1 agonist (35), and assessed *HAS* mRNA expression and HA production. The results presented in Fig. 5D demonstrate that treatment with BW245C (10  $\mu$ M) significantly induced *HAS* mRNA expression. Here, there was a significant increase in *HAS1* and *HAS2* mRNA expression as early as 2 h. *HAS3* mRNA increased by 6 h of exposure to BW245C. In addition, BW245C dose-dependently increased HA levels (Fig. 5E). At concentrations of 5 and 10  $\mu$ M, there was a significant increase in HA production, compared with untreated fibroblasts. In contrast, the high affinity selective DP2 agonist 13,14-dihydro-15-keto-PGD<sub>2</sub> (36–38) (in the low micromolar concentration range) did not significantly increase HA levels (data not shown). Taken together, these data strongly suggest that PGD<sub>2</sub> and PGJ<sub>2</sub> induced HA production in orbital fibroblasts through activation of the DP1.

**DP1 Activation Boosts Intracellular cAMP to Increase HA by Orbital Fibroblasts**—Activation of DP1, a G<sub>s</sub> protein-coupled receptor, leads to an increase in intracellular cAMP. We hypothesized that activation of DP1 by PGD<sub>2</sub> and subsequent production of HA would require the generation of cAMP. We therefore measured intracellular cAMP following treatment with PGD<sub>2</sub> and the DP1 agonist BW245C. Treatment with PGD<sub>2</sub> (5  $\mu$ M) and BW245C (10  $\mu$ M) significantly increased intracellular cAMP production within 15–30 min (Fig. 6A). cAMP levels remained elevated through 60 min (Fig. 6A, black bars).

To investigate whether or not the elevation in intracellular levels of cAMP was directly attributable to the increased HA production by orbital fibroblasts, cells were treated with forskolin, which activates adenylyl cyclase to augment intracellular levels of cAMP (39, 40). As we predicted, forskolin alone significantly increased HA production compared with vehicle-treated fibroblasts (Fig. 6B). We then treated the cells with PGD<sub>2</sub> or PGJ<sub>2</sub> together with IBMX, a nonselective phosphodiesterase inhibitor that boosts cAMP levels by preventing its degradation (41). IBMX, in conjunction with either PGD<sub>2</sub> or PGJ<sub>2</sub>, significantly enhanced the ability of PGD<sub>2</sub> or PGJ<sub>2</sub> to generate HA (Fig. 6B). Collectively, these data support the idea that PGD<sub>2</sub> and PGJ<sub>2</sub>, via activation of DP1 and generation of cAMP, increase HA production in human orbital fibroblasts.

**Mast Cell-derived PGD<sub>2</sub> Activates Orbital Fibroblasts to Produce HA via *HAS2***—Mast cells are a key immune cell proposed to be involved in the pathogenesis of TED via their ability to activate fibroblasts. Mast cells are also a key cell type that produces PGD<sub>2</sub> *in vivo*. We therefore hypothesized that mast cell-derived PGD<sub>2</sub> would increase HA production by orbital fibroblasts. We first sought to determine if human mast cells



**FIGURE 3. PGD<sub>2</sub> induces HAS mRNA in human orbital fibroblasts.** *A*, RT-PCR analysis. cDNA from orbital fibroblasts treated with 5 μM PGD<sub>2</sub> or vehicle for 2 h was amplified by RT-PCR and separated on 5% acrylamide gels. All three HAS isoforms are expressed in human orbital fibroblasts. Following treatment with PGD<sub>2</sub>, there was a relative increase in the abundance of *HAS1*, *HAS2*, and *HAS3* mRNA. 7 S was used as a control. *B*, qRT-PCR. Total RNA from orbital fibroblasts treated with 5 μM PGD<sub>2</sub> for 1–24 h was analyzed by qRT-PCR as described under “Experimental Procedures.” There was a significant increase in HAS mRNA levels (*HAS1*, *HAS2*, and *HAS3*) beginning at 2 h (\*\*\*, *p* < 0.001 compared with time 0 for each HAS isoform). Of the three, *HAS1* yielded the greatest increase (2807 ± 213-fold); this increase in *HAS1* was significantly greater than *HAS2* or *HAS3* (###, *p* < 0.001). Expression of *HAS1* and *HAS2* remained significantly elevated through 4 h (\*, *p* < 0.05). By 16 h, mRNA for *HAS1*, *HAS2*, and *HAS3* was not different from control. Results are expressed as the mean ± S.D. (error bars) of triplicate samples performed on duplicate cultures.

produce PGD<sub>2</sub> *in vitro*. HMC-1 cells, a commonly used human mast cell line (42, 43), were used to first examine whether mast cells express H-PGDS. Western blot analysis of unstimulated HMC-1 cells revealed that these cells basally express both Cox-1 and H-PGDS (Fig. 7A). We then activated HMC-1 cells with A23187, a calcium ionophore (42), in the presence or absence of HQL-79, a specific inhibitor of H-PGDS (44, 45), and determined PGD<sub>2</sub> production. Fig. 7B demonstrates that activated HMC-1 cells significantly increase PGD<sub>2</sub> compared with unstimulated cells. Inhibition of H-PGDS with HQL-79 significantly reduced the level of PGD<sub>2</sub> in activated HMC-1 cells (Fig. 7B).

Using co-culture, we further investigated the association among mast cells, orbital fibroblasts, and HA production. First,

we cultured orbital fibroblasts with HMC-1 cells (at a ratio of 1:1) for 4 h. Following the removal of the mast cells and the addition of fresh medium, the supernatant was collected for HA analysis. When fibroblasts and mast cells were cultured together, there was a significant increase in HA production compared with fibroblasts alone (Fig. 8A, *open versus black bar*; \*\*, *p* < 0.001).


We then utilized a transwell system, whereby orbital fibroblasts were physically separated from HMC-1 cells. Following 24 h of co-culture, conditioned medium in both chambers was collected for HA ELISA. HA levels were negligible in the chamber (upper) containing only the HMC-1 cells. HA production also did not increase as a consequence of increasing cell number (Fig. 8B, *open bars*), suggesting that HMC-1 cells do not produce HA in significant quantities. In contrast, HA levels in the lower chamber (orbital fibroblasts) were significantly higher than in the HMC-1 (upper) chamber (Fig. 8B; ###, *p* < 0.001, upper chamber compared with lower chamber). Co-culture of HMC-1 cells with orbital fibroblasts significantly increased HA levels with increasing ratios of HMC-1/orbital fibroblasts. Thus, even when HMC-1 cells were physically separated from fibroblasts, HA levels increased, suggesting that a factor(s) secreted by the HMC-1 cells drives HA synthesis by fibroblasts.

To determine if PGD<sub>2</sub> was the factor secreted by HMC-1 cells that was responsible for driving HA synthesis by orbital fibroblasts, we pre-treated HMC-1 cells with HQL-79 (to block PGD<sub>2</sub> synthesis) and then added these cells to the upper chamber of the transwell at a ratio of 20:1 (HMC-1/orbital fibroblast). Fig. 8C shows that treatment of HMC-1 cells with HQL-79 significantly attenuated the production of HA by orbital fibroblasts compared with treatment of HMC-1 cells with vehicle alone. These data suggest that PGD<sub>2</sub> secreted by mast cells plays a dominant role in HA production by orbital fibroblasts during TED pathogenesis.

Finally, to determine whether HA production by co-culture of fibroblasts with mast cells was via *HAS2*, we transfected orbital fibroblasts with control, *HAS1*, or *HAS2* siRNA and co-cultured the transfected fibroblasts with the mast cells. As expected, co-culture with untransfected fibro-



## PGD<sub>2</sub> Controls Hyaluronan Synthesis in Orbital Fibroblasts



**FIGURE 4. PGD<sub>2</sub>-induced HA production by orbital fibroblasts is dependent on HAS2 expression.** *A*, orbital fibroblasts were transfected with siRNA for *HAS1*, *HAS2*, or a combination as described under "Experimental Procedures." The cells were cultured for 24 h in RPMI 1640 with 0.5% FBS and treated with 5 μM PGD<sub>2</sub> for 2 h, and HAS gene levels were analyzed by qRT-PCR. The mRNA for *HAS1* and *HAS2* in the PGD<sub>2</sub>-treated SC siRNA samples was standardized to 100 for comparison of gene expression. Each siRNA reduced its target mRNA expression selectively and significantly (up to 80%). \*\*,  $p < 0.01$ . *B*, orbital fibroblasts transfected with siRNA for *HAS1*, *HAS2*, or a combination were cultured in RPMI 1640 with 0.5% FBS for 24 h and then exposed to 5 μM PGD<sub>2</sub> for 18 h, and HA was analyzed by HA ELISA. Fibroblasts treated with PGD<sub>2</sub> (black bars) increased HA synthesis. Knockdown of *HAS2*, but not *HAS1*, significantly reduced the ability of PGD<sub>2</sub> to induce HA in human orbital fibroblasts. \*,  $p < 0.05$ ; \*\*,  $p < 0.01$  compared with PGD<sub>2</sub>-treated SC siRNA-transfected; ##,  $p < 0.01$ , untreated SC siRNA versus untreated *HAS2* siRNA; ns, no significance, *HAS2* siRNA-transfected, PGD<sub>2</sub>-treated fibroblasts versus untreated SC siRNA-transfected cells.

blasts with HMC-1 cells significantly increased HA production (Fig. 9). Transfection of fibroblasts with either control siRNA or *HAS1* siRNA failed to attenuate HA production elicited by co-culture with HMC-1 mast cells. In contrast, transfection of orbital fibroblasts with *HAS2* siRNA significantly decreased HA production by orbital fibroblasts. Here, there was a significant decrease in HA basal levels (untransfected, no mast cells) as well as by fibroblasts co-cultured with HMC-1 cells (Fig. 9). Collectively, these data highlight the dominant role of *HAS2* in HA production elicited by mast cell-derived PGD<sub>2</sub>.


## DISCUSSION

TED is a debilitating disorder that causes disfigurement and vision impairment. TED afflicts ~40% of patients with Graves hyperthyroidism, a thyroid-specific autoimmune disease characterized by the presence of autoantibodies against the thyroid-stimulating hormone receptor. Treatment of mod-

erate to severe TED often involves invasive procedures, including orbital radiotherapy and orbital decompression surgery (46). There are few effective pharmacological treatments for TED due, in part, to a poor understanding the pathogenic mechanisms leading to clinical manifestations of TED. Current evidence, including data from our laboratories (3, 47, 48), suggests that activation of orbital fibroblasts by infiltrating inflammatory cells, particularly T cells and mast cells, plays an important role in TED pathogenesis (49). Orbital fibroblast proliferation and ECM production, particularly HA (50, 51), are key events that contribute to manifestations of TED, such as periorbital edema, exophthalmos, and extraocular motility dysfunction (1, 52). Here, we have provided data indicating that mast cell-derived PGD<sub>2</sub> is a key factor that regulates the production of HA by orbital fibroblasts via activation of DP1 (Figs. 1, 5, and 7). Importantly, prevention of DP1 signaling (Fig. 5) or PGD<sub>2</sub> production by mast cells (Fig. 8) attenuated HA synthesis by the fibroblasts.

HA is a large negatively charged polysaccharide that is overproduced in the retroocular space of patients with TED (53). HA has remarkable viscosity and ability to retain water, which leads to increases in orbital tissue volume and anterior displacement of the eye, culminating in exophthalmos (4). Fibroblasts are the major source of HA in the orbit. Activation of orbital fibroblasts by immunoglobulins from TED patients (4) and cytokines, such as interleukin-1β (51), interferon-γ (54), and transforming growth factor-β (55), increase HA production. Our study is the first to show that PGD<sub>2</sub>, a non-cytokine mediator, contributes to HA synthesis by orbital fibroblasts (Fig. 1). The increase in HA production by PGD<sub>2</sub> is directly related to increased expression of *HAS2* (Figs. 3, 4, and 9). Despite the fact that all three HAS isoforms (*HAS1*, *HAS2*, and *HAS3*) are significantly increased, only the attenuation of *HAS2* expression completely abrogated PGD<sub>2</sub>-induced HA. This is in agreement with observations made in *HAS2*<sup>-/-</sup> mouse embryos, which are virtually devoid of HA (56). Of interest, *HAS1* mRNA exhibited the most dramatic increase in response to PGD<sub>2</sub> yet does not contribute significantly to HA biosynthesis (Fig. 4). This observation may be explained by the fact that *HAS1* protein rapidly loses its enzymatic activity (57). Thus, our data suggest that *HAS2* is responsible for most, if not all, HA production by orbital fibroblasts in response to PGD<sub>2</sub>.


The induction of HAS mRNA and subsequent production of HA required the generation of the second messenger cAMP via activation of DP1. We are the first to show that DP receptors are expressed on orbital fibroblasts (Fig. 5). Pharmacological activation of DP1 significantly increased HAS mRNA and HA biosynthesis (Fig. 5). Moreover, the DP1 antagonist MK-0524 completely inhibited PGD<sub>2</sub>-induced HA production, whereas DP2 antagonist ramatroban had no effect. These findings suggest that PGD<sub>2</sub>-induced HA production is mediated solely by DP1. MK-0524 (also known as Laropiprant and CORDAPTIVE™) is an orally acting drug that is currently being developed in conjunction with niacin as a cholesterol-lowering drug (58). Our results indicate that selectively targeting the DP1 system (via MK-0524) in the eye



**FIGURE 5. PGD<sub>2</sub> and PGJ<sub>2</sub> induce HA production in human orbital fibroblasts via DP1, but not DP2, activation.** *A*, RT-PCR. Human orbital fibroblasts (OF1 and OF2) and T cells express *DP1* and *DP2* mRNA. Note the variability in the expression of *DP2* between the two fibroblast strains. *B*, Western blot analysis revealed that orbital fibroblasts and T cells express both *DP1* and *DP2* protein. Membranes were reprobed for glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) to ensure equal protein loading. *C*, pharmacological inhibition of *DP1*, but not *DP2*, blocks PGD<sub>2</sub>- and PGJ<sub>2</sub>-induced HA production. Orbital fibroblasts were left untreated (*Vehicle*) or were pretreated with 100 nM of the *DP1* antagonist MK-0524 (*MK*) or *DP2* antagonist ramatroban (*RAM*) for 1 h with or without PGD<sub>2</sub> or PGJ<sub>2</sub> for 18 h, and HA ELISA was performed. Both PGD<sub>2</sub> and PGJ<sub>2</sub> significantly increased HA production compared with untreated ( $p < 0.001$  (###) and  $p < 0.01$  (##), respectively). Pretreatment of orbital fibroblasts with *MK* significantly decreased the ability of PGD<sub>2</sub> (\*\*,  $p < 0.01$ ) and PGJ<sub>2</sub> (\*,  $p < 0.05$ ) to induce HA levels. *RAM* was not able to prevent PGD<sub>2</sub>- and PGJ<sub>2</sub>-increased HA levels ( $p < 0.05$  (#) and  $p < 0.01$  (##) compared with vehicle control, respectively). *ns*, not significant compared with untreated (vehicle). *D*, treatment with the *DP1* agonist BW245C significantly increased the expression of HAS mRNA. Orbital fibroblasts were cultured in reduced serum for 3 days and exposed to BW245C (10 μM) for the indicated times. There was a significant increase in *HAS1* (329 ± 128-fold increase; \*\*,  $p < 0.01$ ) and *HAS2* (17 ± 0.14-fold increase; \*\*\*,  $p < 0.001$ ) at 2 h compared with vehicle control. *HAS3* mRNA increased by 6 h (4.8 ± 0.06; \*\*\*,  $p < 0.01$ ). *E*, activation of *DP1* by the selective agonist BW245C induces HA. There was a significant increase in HA when fibroblasts were treated with 5 and 10 μM BW245C ( $p < 0.05$  (\*) and  $p < 0.01$  (\*\*), respectively, compared with vehicle control. Samples were run in duplicate, utilizing three separate human orbital fibroblast strains (representative results are shown).




## PGD<sub>2</sub> Controls Hyaluronan Synthesis in Orbital Fibroblasts



**FIGURE 6. PGD<sub>2</sub>-induced HA synthesis is through the DP1-cAMP signal pathway.** *A*, DP1 activation by PGD<sub>2</sub> or BW245C increases intracellular cAMP level. Orbital fibroblasts were treated with PGD<sub>2</sub> or BW245C for up to 60 min, and intracellular cAMP was detected as described under "Experimental Procedures." There was a significant increase in cAMP within 15 min of treatment with PGD<sub>2</sub> compared with vehicle control (\*,  $p < 0.05$ ). cAMP further increased by 30 and 60 min (\*\*\*,  $p < 0.001$ ). *B*, cultures of confluent orbital fibroblasts were treated with 5 μM forskolin or 200 μM IBMX, with or without 5 μM PGD<sub>2</sub> or 2 μM PGJ<sub>2</sub>, for 18 h, and the cell culture supernatant was collected for HA ELISA. There was a significant increase in HA when cells were treated with forskolin, PGD<sub>2</sub>, or PGJ<sub>2</sub> compared with vehicle-treated (open bar) (\*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ). Augmenting cAMP (via IBMX) in conjunction with PGD<sub>2</sub> or PGJ<sub>2</sub> significantly increased HA when compared with PGD<sub>2</sub> and PGJ<sub>2</sub> alone (#,  $p < 0.05$ ). Results are expressed as the mean ± S.D. (error bars).

may prevent the severity and/or occurrence of proptosis in TED patients by reducing HA production.

PGD<sub>2</sub> also regulates the migration of inflammatory cells by facilitating vasodilatation and increasing vascular permeability (59). Within the eye, this would facilitate transendothelial migration of infiltrating mast cells and lymphocytes, thereby contributing to the increase in immune cells characteristic of TED. Association of mast cells with fibroblasts/adipocytes in the orbital tissue is commonly observed in biopsies of TED patients as well as animal models of TED (22, 23, 60–63). Mast cell degranulation and close proximity to adipocytes are suggestive of their participation in the TED process (23). Our data presented here strongly support the hypothesis that mast cell-fibroblast interactions participate in TED. We first demonstrated that fibroblasts robustly respond to PGD<sub>2</sub> to increase HA synthesis (Fig. 1). Using both co-culture and transwell systems, we then showed that




**FIGURE 7. PGD<sub>2</sub> production by mast cells is dependent on H-PGDS activity.** *A*, Western blot of HMC-1 cells indicates the expression of Cox-1 and H-PGDS. Unactivated HMC-1 cells do not express Cox-2. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used as a housekeeping protein. *B*, inhibition of H-PGDS activity ameliorates the production of PGD<sub>2</sub> by activated mast cells. HMC-1 cells were treated with HQL-79 for 1 h, followed by activation with A23187, and cell culture supernatant was assessed for PGD<sub>2</sub> levels by commercial EIA kit. Activation of HMC-1 cells with A23187 significantly increased PGD<sub>2</sub> production (###,  $p < 0.001$  compared with untreated). Inclusion of HQL-79 significantly reduced PGD<sub>2</sub> production (\*\*\*,  $p < 0.001$ ).

mast cells increase orbital fibroblast production of HA via PGD<sub>2</sub> produced by the mast cell (Fig. 8). This accumulated HA within the orbit can, in turn, facilitate inflammatory cell infiltration (64) to the orbit, thereby perpetuating clinical symptoms associated with TED. These data, in conjunction with our previously published data showing that PGD<sub>2</sub> promotes adipogenesis (48), highlights PGD<sub>2</sub> as a key regulator of orbital fibroblast function.


It has been well described that fibroblasts are heterogeneous, differing not only between organ systems but also within a given organ (25, 65–68). We previously published that fibroblasts from the lung exhibit significant interindividual variability in their apoptotic response to cigarette smoke (69), a finding that may explain why only a fraction of smokers develop lung diseases, such as chronic obstructive pulmonary disease. Therefore, it is noteworthy that there was significant variability in HA production, HA size, and HYAL expression between orbital fibroblasts derived from two different individuals (Figs. 1 and 2). The increase in low  $M_r$  HA by OF1 (Fig. 1B) correlates with increased expression of *HYAL1* to -3 (Fig. 2). These enzymes are responsible for the depolymerization of HA, thereby generating HA fragments (29). These low  $M_r$  HA fragments are known to promote inflammation (70) by activating immune cells and increasing inflammatory gene expression (71). Thus, increased HA production, coupled with the generation of inflammation-promoting low  $M_r$  HA, by orbital fibroblasts from an individual could be a decisive factor in predicting which Graves patients ultimately develop symptomatic TED.

Using a combination of molecular and pharmacological approaches, we have provided data on the role of mast cell-derived PGD<sub>2</sub> in promoting HA biosynthesis by orbital fibroblasts, key effector cells in the pathogenesis of TED. We also show that the DP1/cAMP signaling pathway rapidly activates HAS mRNA induction to promote HA production. PGD<sub>2</sub>, in addition to causing HA accumulation, may also



**FIGURE 8. Mast cell-derived PGD<sub>2</sub> activates orbital fibroblasts to produce HA.** *A*, contact co-culture. Confluent orbital fibroblasts were seeded with HMC-1 cells at a cell ratio of 1:1 for 4 h. The mast cells were then removed, the fibroblasts were washed, and fresh medium was added for another 18 h. The medium was collected for HA ELISA. Co-culture of orbital fibroblasts with HMC-1 cells significantly increased HA synthesis (\*\*,  $p < 0.001$ ). *B*, transwell co-culture. HMC-1 cells and confluent fibroblasts were co-cultured in a transwell system, where the fibroblasts and HMC-1 cells were separated by a 0.4-μm membrane; the HMC-1/orbital fibroblast (OF) ratio was 5:1, 10:1, or 20:1. The conditioned medium from both chambers was collected for HA ELISA. There was a significant difference in HA levels between HMC-1 (upper chamber, open bars) and orbital fibroblasts (lower chamber, black bars) (#,  $p < 0.05$ ; ###,  $p < 0.001$ ). Co-culture of orbital fibroblasts with HMC-1 cells significantly increased HA production only by the fibroblasts (lower chamber) (\*\*\*,  $p < 0.001$  compared with no HMC-1 cells). *C*, inhibition of PGD<sub>2</sub> secretion by HMC-1 cells prevents HA production by orbital fibroblasts. HMC-1 cells were treated with the H-PGDS inhibitor HQL-79 prior to co-culturing (ratio 20:1) with orbital fibroblasts. There was a significant increase in HA synthesis when fibroblasts were cultured with HMC-1 cells (\*\*,  $p < 0.01$  compared with no HMC). This increase in HA was attenuated when PGD<sub>2</sub> production in mast cells was prevented by HQL-79 (##,  $p < 0.001$ , HQL-79 compared with vehicle).

provide an environment conducive to lymphocyte infiltration into orbital tissue via DP1. The lymphocytic accumulation within the orbit may, in turn, lead to additional release of inflammatory mediators, including PGD<sub>2</sub>. Therefore,



**FIGURE 9. siRNA against orbital fibroblast HAS2 prevents mast cell-derived PGD<sub>2</sub> induction of HA.** Orbital fibroblasts were transfected with HAS1, HAS2, or SC siRNA using Lipofectamine 2000 (Invitrogen). Following this, the fibroblasts were co-cultured for 24 h with HMC-1 cells in a transwell system at a ratio of 20:1 (mast cells/orbital fibroblasts), and the media were collected for HA detection. Untransfected orbital fibroblast co-cultured with HMC-1 cells significantly increased HA production (\*\*\*,  $p < 0.0001$ ; compared with untransfected without co-culture). Co-culture of fibroblasts with HMC-1 cells also induced a significant increase in HA in SC siRNA-transfected (\*\*,  $p < 0.001$ ) and HAS1 siRNA-transfected (\*,  $p < 0.05$ ) orbital fibroblasts. HAS2 siRNA-transfected orbital fibroblasts failed to increase HA synthesis when co-cultured with HMC-1 cells (###,  $p < 0.0001$  compared with untransfected, control siRNA, and HAS1 siRNA). Results are expressed as mean  $\pm$  S.D. ( $n = 4-6$ ).

selectively targeting the production of lipid mediators (*i.e.* PGD<sub>2</sub>) and/or activation of receptor systems (DP1) may reduce or prevent symptoms associated with TED.

REFERENCES

- Prabhakar, B. S., Bahn, R. S., and Smith, T. J. (2003) *Endocr. Rev.* **24**, 802–835
- Lehmann, G. M., Garcia-Bates, T. M., Smith, T. J., Feldon, S. E., and Phipps, R. P. (2008) *PPAR Res.* **2008**, 895901
- Feldon, S. E., O’loughlin, C. W., Ray, D. M., Landskroner-Eiger, S., Seweryniak, K. E., and Phipps, R. P. (2006) *Am. J. Pathol.* **169**, 1183–1193
- Smith, T. J., and Hoa, N. (2004) *J. Clin. Endocrinol. Metab.* **89**, 5076–5080
- Kuriyan, A. E., Phipps, R. P., and Feldon, S. E. (2008) *Curr. Opin. Ophthalmol.* **19**, 499–506
- Jiang, D., Liang, J., and Noble, P. W. (2007) *Annu. Rev. Cell Dev. Biol.* **23**, 435–461
- Makkonen, K. M., Pasonen-Seppänen, S., Törrönen, K., Tammi, M. I., and Carlberg, C. (2009) *J. Biol. Chem.* **284**, 18270–18281
- Campo, G. M., Avenoso, A., Campo, S., D’Ascola, A., Traina, P., and Calatroni, A. (2009) *Br. J. Biomed. Sci.* **66**, 28–36
- Guo, N., Kanter, D., Funderburgh, M. L., Mann, M. M., Du, Y., and Funderburgh, J. L. (2007) *J. Biol. Chem.* **282**, 12475–12483
- Marzoll, A., Nagy, N., Wördehoff, L., Dai, G., Fries, S., Lindner, V., Grosser, T., and Fischer, J. W. (2009) *J. Cell Mol. Med.* **13**, 3713–3719
- Fischer, J. W., and Schrör, K. (2007) *Thromb. Haemost.* **98**, 287–295
- Goetzl, E. J., An, S., and Smith, W. L. (1995) *FASEB J.* **9**, 1051–1058
- Herlong, J. L., and Scott, T. R. (2006) *Immunol. Lett.* **102**, 121–131
- Boie, Y., Sawyer, N., Slipetz, D. M., Metters, K. M., and Abramovitz, M. (1995) *J. Biol. Chem.* **270**, 18910–18916
- Nagata, K., Hirai, H., Tanaka, K., Ogawa, K., Aso, T., Sugamura, K., Nakamura, M., and Takano, S. (1999) *FEBS Lett.* **459**, 195–199

16. Kostenis, E., and Ulven, T. (2006) *Trends Mol. Med.* **12**, 148–158
17. Forman, B. M., Tontonoz, P., Chen, J., Brun, R. P., Spiegelman, B. M., and Evans, R. M. (1995) *Cell* **83**, 803–812
18. Kliewer, S. A., Lenhard, J. M., Willson, T. M., Patel, I., Morris, D. C., and Lehmann, J. M. (1995) *Cell* **83**, 813–819
19. Tamori, Y., Masugi, J., Nishino, N., and Kasuga, M. (2002) *Diabetes* **51**, 2045–2055
20. Xue, L., Barrow, A., and Pettipher, R. (2009) *Clin. Exp. Immunol.* **156**, 126–133
21. Lewis, R. A., Soter, N. A., Diamond, P. T., Austen, K. F., Oates, J. A., and Roberts, L. J., 2nd (1982) *J. Immunol.* **129**, 1627–1631
22. Lauer, S. A., Silkiss, R. Z., and McCormick, S. A. (2008) *Ophthalmol. Plast. Reconstr. Surg.* **24**, 257–261
23. Boschi, A., Daumerie, C., Spiritus, M., Beguin, C., Senou, M., Yuksel, D., Duplidy, M., Costagliola, S., Ludgate, M., and Many, M. C. (2005) *Br. J. Ophthalmol.* **89**, 724–729
24. Smith, T. J., and Parikh, S. J. (1999) *Endocrinology* **140**, 3518–3525
25. Smith, T. J., Sempowski, G. D., Wang, H. S., Del Vecchio, P. J., Lippe, S. D., and Phipps, R. P. (1995) *J. Clin. Endocrinol. Metab.* **80**, 2620–2625
26. Baglole, C. J., Reddy, S. Y., Pollock, S. J., Feldon, S. E., Sime, P. J., Smith, T. J., and Phipps, R. P. (2005) *Methods Mol. Med.* **117**, 115–127
27. Lee, H. G., and Cowman, M. K. (1994) *Anal. Biochem.* **219**, 278–287
28. Ohya, T., and Kaneko, Y. (1970) *Biochim. Biophys. Acta* **198**, 607–609
29. Noble, P. W. (2002) *Matrix Biol.* **21**, 25–29
30. Pettipher, R., Hansel, T. T., and Armer, R. (2007) *Nat. Rev. Drug Discov.* **6**, 313–325
31. Nagata, K., and Hirai, H. (2003) *Prostaglandins Leukot. Essent. Fatty Acids* **69**, 169–177
32. Sturino, C. F., O'Neill, G., Lachance, N., Boyd, M., Berthelette, C., Labelle, M., Li, L., Roy, B., Scheiget, J., Tsou, N., Aubin, Y., Bateman, K. P., Chaurret, N., Day, S. H., Lévesque, J. F., Seto, C., Silva, J. H., Trimble, L. A., Carriere, M. C., Denis, D., Greig, G., Kargman, S., Lamontagne, S., Mathieu, M. C., Sawyer, N., Slipetz, D., Abraham, W. M., Jones, T., McAuliffe, M., Piechuta, H., Nicoll-Griffith, D. A., Wang, Z., Zamboni, R., Young, R. N., and Metters, K. M. (2007) *J. Med. Chem.* **50**, 794–806
33. Leblanc, Y., Roy, P., Dufresne, C., Lachance, N., Wang, Z., O'Neill, G., Greig, G., Denis, D., Mathieu, M. C., Slipetz, D., Sawyer, N., and Tsou, N. (2009) *Bioorg. Med. Chem. Lett.* **19**, 2125–2128
34. Royer, J. F., Schratl, P., Carrillo, J. J., Jupp, R., Barker, J., Weyman-Jones, C., Beri, R., Sargent, C., Schmidt, J. A., Lang-Loidolt, D., and Heinemann, A. (2008) *Eur. J. Clin. Invest.* **38**, 663–671
35. Kabashima, K., and Narumiyi, S. (2003) *Prostaglandins Leukot. Essent. Fatty Acids* **69**, 187–194
36. Cheng, K., Wu, T. J., Wu, K. K., Sturino, C., Metters, K., Gottesdiener, K., Wright, S. D., Wang, Z., O'Neill, G., Lai, E., and Waters, M. G. (2006) *Proc. Natl. Acad. Sci. U.S.A.* **103**, 6682–6687
37. Yoshimura-Uchiyama, C., Iikura, M., Yamaguchi, M., Nagase, H., Ishii, A., Matsushima, K., Yamamoto, K., Shichijo, M., Bacon, K. B., and Hirai, K. (2004) *Clin. Exp. Allergy* **34**, 1283–1290
38. Hirai, H., Tanaka, K., Yoshie, O., Ogawa, K., Kenmotsu, K., Takamori, Y., Ichimasa, M., Sugamura, K., Nakamura, M., Takano, S., and Nagata, K. (2001) *J. Exp. Med.* **193**, 255–261
39. Morris, B. J., Adams, D. J., Beveridge, D. J., van der Weyden, L., Mangs, H., and Leedman, P. J. (2004) *Acta Physiol. Scand.* **181**, 369–373
40. Metzger, H., and Lindner, E. (1981) *Arzneimittelforschung* **31**, 1248–1250
41. Weinberg, E., Zeldich, E., Weinreb, M. M., Moses, O., Nemcovsky, C., and Weinreb, M. (2009) *J. Cell. Biochem.* **108**, 207–215
42. Kim, S. H., and Shin, T. Y. (2009) *Toxicol. In Vitro* **23**, 1215–1219
43. Margulis, A., Nocka, K. H., Brennan, A. M., Deng, B., Fleming, M., Goldman, S. J., and Kasaian, M. T. (2009) *J. Immunol.* **183**, 1739–1750
44. Kanaoka, Y., and Urade, Y. (2003) *Prostaglandins Leukot. Essent. Fatty Acids* **69**, 163–167
45. Aritake, K., Kado, Y., Inoue, T., Miyano, M., and Urade, Y. (2006) *J. Biol. Chem.* **281**, 15277–15286
46. Bartalena, L., and Tanda, M. L. (2009) *N. Engl. J. Med.* **360**, 994–1001
47. Hwang, C. J., Afifyan, N., Sand, D., Naik, V., Said, J., Pollock, S. J., Chen, B., Phipps, R. P., Goldberg, R. A., Smith, T. J., and Douglas, R. S. (2009) *Invest. Ophthalmol. Vis. Sci.* **50**, 2262–2268
48. Feldon, S. E., Park, D. J., O'Loughlin, C. W., Nguyen, V. T., Landskroner-Eiger, S., Chang, D., Thatcher, T. H., and Phipps, R. P. (2005) *Invest. Ophthalmol. Vis. Sci.* **46**, 3913–3921
49. Lehmann, G. M., Feldon, S. E., Smith, T. J., and Phipps, R. P. (2008) *Thyroid* **18**, 959–965
50. Gianoukakis, A. G., Jennings, T. A., King, C. S., Sheehan, C. E., Hoa, N., Heldin, P., and Smith, T. J. (2007) *Endocrinology* **148**, 54–62
51. Kaback, L. A., and Smith, T. J. (1999) *J. Clin. Endocrinol. Metab.* **84**, 4079–4084
52. van Steensel, L., Paridaens, D., Schrijver, B., Dingjan, G. M., van Daele, P. L., van Hagen, P. M., van den Bosch, W. A., Drexhage, H. A., Hooijkaas, H., and Dik, W. A. (2009) *Invest. Ophthalmol. Vis. Sci.* **50**, 3091–3098
53. Kahaly, G., Förster, G., and Hansen, C. (1998) *Thyroid* **8**, 429–432
54. Smith, T. J., Bahn, R. S., Gorman, C. A., and Cheavens, M. (1991) *J. Clin. Endocrinol. Metab.* **72**, 1169–1171
55. Wang, H. S., Tung, W. H., Tang, K. T., Wong, Y. K., Huang, G. J., Wu, J. C., Guo, Y. J., and Chen, C. C. (2005) *J. Cell. Biochem.* **95**, 256–267
56. Camenisch, T. D., Spicer, A. P., Brehm-Gibson, T., Biesterfeldt, J., Augustine, M. L., Calabro, A., Jr., Kubalak, S., Klewer, S. E., and McDonald, J. A. (2000) *J. Clin. Invest.* **106**, 349–360
57. Itano, N., Sawai, T., Yoshida, M., Lenas, P., Yamada, Y., Imagawa, M., Shinomura, T., Hamaguchi, M., Yoshida, Y., Ohnuki, Y., Miyauchi, S., Spicer, A. P., McDonald, J. A., and Kimata, K. (1999) *J. Biol. Chem.* **274**, 25085–25092
58. Schwartz, J. I., Liu, F., Stroh, M., Gipson, A., Johnson-Levonas, A. O., Lasseter, K. C., Lai, E., and Wagner, J. A. (2009) *Am. J. Ther.* **16**, 215–223
59. Ulven, T., and Kostenis, E. (2006) *Curr. Top. Med. Chem.* **6**, 1427–1444
60. Hufnagel, T. J., Hickey, W. F., Cobbs, W. H., Jakobiec, F. A., Iwamoto, T., and Eagle, R. C. (1984) *Ophthalmology* **91**, 1411–1419
61. Many, M. C., Costagliola, S., Detrait, M., Deneff, F., Vassart, G., and Ludgate, M. C. (1999) *J. Immunol.* **162**, 4966–4974
62. Costagliola, S., Many, M. C., Deneff, J. F., Pohlenz, J., Refetoff, S., and Vassart, G. (2000) *J. Clin. Invest.* **105**, 803–811
63. Yamada, M., Li, A. W., West, K. A., Chang, C. H., and Wall, J. R. (2002) *Autoimmunity* **35**, 403–413
64. Toole, B. P. (2004) *Nat. Rev. Cancer* **4**, 528–539
65. Phipps, R. P., Borrello, M. A., and Blieden, T. M. (1997) *J. Periodontol. Res.* **32**, 159–165
66. Borrello, M. A., and Phipps, R. P. (1996) *Cell. Immunol.* **173**, 198–206
67. Hagood, J. S., Lasky, J. A., Nesbitt, J. E., and Segarini, P. (2001) *Chest* **120**, Suppl. 1, 64S–66S
68. Khoo, T. K., Coenen, M. J., Schiefer, A. R., Kumar, S., and Bahn, R. S. (2008) *Thyroid* **18**, 1291–1296
69. Baglole, C. J., Bushinsky, S. M., Garcia, T. M., Kode, A., Rahman, I., Sime, P. J., and Phipps, R. P. (2006) *Am. J. Physiol. Lung Cell Mol. Physiol.* **291**, L19–L29
70. de la Motte, C., Nigro, J., Vasanji, A., Rho, H., Kessler, S., Bandyopadhyay, S., Danese, S., Flocchi, C., and Stern, R. (2009) *Am. J. Pathol.* **174**, 2254–2264
71. Gao, F., Koenitzer, J. R., Tobilewski, J. M., Jiang, D., Liang, J., Noble, P. W., and Oury, T. D. (2008) *J. Biol. Chem.* **283**, 6058–6066