

# Highlights on the capacities of "Gel-based" proteomics

François Chevalier

## Abstract

Gel-based proteomic is the most popular and versatile method of global protein separation and quantification. This is a mature approach to screen the protein expression at the large scale, and a cheaper approach as compared with gel-free proteomics. Based on two independent biochemical characteristics of proteins, two-dimensional electrophoresis combines isoelectric focusing, which separates proteins according to their isoelectric point, and SDS-PAGE, which separates them further according to their molecular mass. The next typical steps of the flow of gel-based proteomics are spots visualization and evaluation, expression analysis and finally protein identification by mass spectrometry. For the study of differentially expressed proteins, two-dimensional electrophoresis allows simultaneously to detect, quantify and compare up to thousand protein spots isoforms, including post-translational modifications, in the same gel and in a wide range of biological systems. In this review article, the limits, benefits, and perspectives of gel-based proteomic approaches are discussed using concrete examples.

## Introduction

Proteomics, one of the most important areas of research in the post-genomic era, is not new in terms of its experimental foundations [1]. It is a natural consequence of the huge advances in genome sequencing, bioinformatics and the development of robust, sensitive, reliable and reproducible analytical techniques [2-12]. Genomics projects have produced a large number of DNA sequences from a wide range of organisms, including humans and mammals. This "genomics revolution" has changed the concept of the comprehensive analysis of biological processes and systems. It is now hypothesized that biological processes and systems can be described based on the comparison of global, quantitative gene expression patterns from cells or tissues representing different states. The discovery of posttranscriptional mechanisms that control rate of synthesis and half-life of proteins and the ensuing nonpredictive correlation between mRNA and protein levels expressed by a particular gene indicate that direct measurement of protein expression also is essential for the analysis of biological processes and systems. Global analysis of gene expression at the protein level is now also termed proteomics. The standard method for quantitative proteome analysis combines protein separation by

high resolution (isoelectric focusing/SDS-PAGE) two-dimensional gel electrophoresis (2DE) with mass spectrometric (MS) or tandem MS (MS/MS) identification of selected protein spots [5,9,11,13-16]. Important technical advances related to 2DE and protein MS have increased sensitivity, reproducibility, and throughput of proteome analysis while creating an integrated technology. Quantitation of protein expression in a proteome provides the first clue into how the cell responds to changes in its surrounding environments. The resulting over- or under-expressed proteins are deemed to play important roles in the precise regulation of cellular activities that are directly related to a given exogenous stimulus. Conventional 2DE, in combination with advanced mass spectrometric techniques, has facilitated the rapid characterization of thousands of proteins in a single polyacrylamide gel. The uniqueness of 2DE for easy visualisation of protein isoforms, using two physical parameters such as isoelectric point and molecular weight, renders this technology itself extremely informative. The method routinely analyzes more than 1000 different protein spots separated on a single two-dimensional gel and, thus, is well suited for the global analysis of protein expression in an organism. However, high-throughput quantitation of proteins from different cell lysates remains a challenging issue, owing to the poor reproducibility of 2DE, as well as low sensitivity and narrow linear dynamic ranges in the

\* Correspondence: [francois.chevalier@cea.fr](mailto:francois.chevalier@cea.fr)

<sup>1</sup> Proteomic Laboratory, IRCM, CEA, Fontenay aux Roses, France  
Full list of author information is available at the end of the article

detection methods [17-21] Recent developments of fluorescent dyes, such as the different commercially available SYPRO dyes, partially addressed some of these problems [22-30]. These dyes detect as little as 1 ng of proteins, and at the same time they offer more than 1000-fold linear dynamic range. The more critical issue, however, is the reproducibility problem of 2DE. Even the identical protein samples that are run on two separate two-dimensional gels will normally produce very similar but not identical 2DE protein maps, owing to the gel-to-gel and operator-to-operator variations. This can be circumvented using multiplexing methods such as fluorescent two-dimensional "Difference Gel Electrophoresis" (2-D DIGE), which substantially reduces variability by displaying two or more complex protein mixtures labeled with different fluorescent dyes in a single 2D gel [21,31-38].

In this review, we focus on the latest developments in 2DE within the context of large-scale proteomics to reveal the advantages, limits and perspectives of the 2DE-based proteomic approach.

## **A - Gel-based proteomic: from sample preparation to protein separation**

### **1 - Protein solubilisation**

In order to take advantage of the high resolution capacity of 2DE, proteins have to be completely denatured, disassembled, reduced and solubilised to disrupt molecular interactions and to ensure that each spot represents an individual polypeptide.

Although a large number of standard protocols has been published, these protocols have to be adapted and further optimized for the type of sample (bacteria/yeast/mammalian cells; cells/tissue; animal/vegetal material; etc...) to be analyzed, as well as for the proteins of interest (cytosolic/nuclear; total "soluble" or membrane "insoluble" proteins; etc...).

After cell disruption, native proteins must be denatured and reduced to disrupt intra- and intermolecular interactions, and solubilized while maintaining the inherent charge properties. Sample solubilization is carried out using a buffer containing chaotropes (urea and/or thiourea), nonionic (Triton X-100) and/or zwitterionic detergents (CHAPS), reducing agents (DTT), carrier ampholytes and most of the time protease and phosphatase inhibitor cocktails are mandatory.

### **2 - The first dimension: isoelectric-focusing with immobilized pH gradients (IPGs)**

Proteins are amphoteric molecules; they carry positive, negative or zero net charge, depending on their amino acid composition. The net charge of a protein is the sum of all the negative and positive charges. The isoelectric point (pI) of a protein is the specific pH at which the net charge of the protein is zero. Proteins are positively

charged at pH values below their pI and are negatively charged at pH values above their pI. IEF is an electrophoretic separation based on this specific biochemical characteristic of proteins.

Basically, the first dimension of the 2DE is achieved with a "strip". It is a dry gel that is formed by the polymerization of acrylamide monomers, linked by bis-acrylamide with molecules of covalently linked immobilin. Immobilins are chemical components that are derived from acrylamide and have additional ionizable non-amphoteric functions. Immobilins of various pKa can create an immobilized pH gradient inside the acrylamide gel. Immobilin was developed by Professors Righetti and Görg at the beginning of the 1990s and is now widely used in 2DE because the IEF gradient is very stable over time and in a high electric field, and shows good reproducibility and a large capacity for separation [9,39-46].

The strip acrylamide gels are dried and cast on a plastic backing. Prior to use, they are rehydrated in a solution containing a pI-corresponding cocktail of carrier ampholytes and with the correct amount of proteins in the solubilization buffer. The carrier ampholytes are amphoteric molecules with a high buffering capacity near their pI. Commercial carrier ampholyte mixtures, which comprise species with pIs spanning a specific pH range, help the proteins to move.

When an electric field is applied, the negatively charged molecules (proteins and ampholytes) move towards the anode (positive/red electrode) and the positively charged molecules move towards the cathode (negative/black electrode). When the proteins are aligned according to their pI, the global net charge is zero and the protein is unable to move and is then focused. Focusing is achieved with a dedicated apparatus that is able to deliver up to 8000 or 10,000 V, but with a limitation in current intensity (50  $\mu$ A maximum/strip) to reduce heat. The strips are usually first rehydrated without current for at least 5 h (passive rehydration), rehydrated with 50 V for 5 h (active rehydration) and then focused until at least 30 to 80 kV/h.

The equilibration step is critical for 2DE. In this step, the strips are saturated with sodium dodecyl sulfate (SDS), an anionic detergent that can denature proteins and form a negatively charged protein/SDS complex. The amount of SDS bound to a protein is directly proportional to the mass of the protein. Thus, proteins that are completely covered by negative charges are separated on the basis of molecular mass.

The equilibration solution also contains buffer, with urea and glycerol. Equilibration of the strips is achieved in two steps: (1) with an equilibration solution containing DTT, to maintain a reducing environment; and (2) with an equilibration solution containing iodoacetamide, to alkylate reduced thiol groups, preventing their re-oxidation during electrophoresis.

### 3 - The second dimension: SDS-PAGE

In SDS polyacrylamide gel electrophoresis (SDS-PAGE), migration is determined not by the intrinsic electric charge of polypeptides but by their molecular weight. The SDS-denatured and reduced proteins are separated according to an apparent molecular weight, in comparison with a molecular weight marker. A linear relationship between the logarithm of the molecular weight and the distance of migration of the proteins can be used; it depends essentially on the percentage of polyacrylamide.

Equilibrated strips are embedded with 1% (w/v) low-melting-point agarose in TRIS/Glycine/SDS running buffer and with 0.01% bromophenol blue on the top of the second dimension acrylamide gel. Gels are usually run with 1 or 2 W of current in the first hour, followed by 15 mA/gel overnight with a temperature regulation (10°C to 18°C). When the bromophenol blue migration front reaches the bottom of the gel, the second dimension is finished and the acrylamide gel can be removed from the glass plates.

### 4 - Gel staining

The gel must firstly be immersed in a fixation solution containing acid (phosphoric acid or acetic acid) and alcohol (ethanol or methanol) as a function of the staining protocol selected. Numerous stains can be used, but with very different costs [17]. Conventional "visible" dyes are Coomassie Blue, colloidal Coomassie Blue and silver nitrate, with quite different sensitivities: 50, 10 and 0.5 ng of detectable protein/spot respectively [17,20,25,47-51]. Commercially available fluorescent dyes, such as Sypro Ruby, Flamingo and Deep Purple, have sensitivities of about 1 ng of detectable protein/spot [21,23,26,28,52-55]. Fluorescent dyes have the advantage of a 4 log dynamic linear range but the disadvantage of being more expensive. In comparison with fluorescent dyes, silver nitrate stain has a dynamic linear range of only 1.5 log, and is not recommended for a gel comparison study.

### 5 - Bioinformatics analysis of 2-D images

Stained gels are scanned on a "visible" or "fluorescent" scanner as a function of the staining protocol selected. The image can then be imported to specific software to be analysed and compared. For a comparison study, at least three repetitions of the same sample should be run; many migration artifacts can occur during 2DE and, to reduce such variability, a mean of several gels is essential. Software, such as Image Master, Progenesis, PDQuest and Samespots, can be used to detect spots and to compare the spot intensity between samples [53-60]. Spots of interest, *i.e.* spots specific to a sample or spots over-expressed on a condition/treatment, can be selected for further MS analysis. Several "computer-based" comparisons can be performed with a 2DE map. As a proteomic

map is specific of a given cell, tissue or organism in a specific physiological condition, it is possible to compare not only one spot to one spot, but a set of spots to a set of spots, for example between two closed organisms. In a precedent study, we investigated the natural variation in the proteome among 8 *Arabidopsis thaliana* ecotypes, of which 3 were previously shown to display atypical responses to environmental stress [61]. The 2DE proteomic maps revealed important variations in terms of function between ecotypes [62]. Hierarchical clustering of proteomes according to either the amount of all anonymous spots, that of the 25 major spots or the functions of these major spots identified the same classes of ecotypes, and grouped the three atypical ecotypes (Fig. 1).


### 6 - Protein identification

To identify the proteins within the spots of interest (according to image analysis), a gel with a greater amount of protein is prepared. In this case, IEF step must be performed at least until 100 kV/h. The other steps of the 2DE are very similar to the previously described protocol. Colloidal Coomassie Blue or fluorescent dyes are recommended for the staining of the preparative gel, because they have good compatibility with MS [22,23,28,63]. In contrast, silver nitrate will give poor results, even if MS-compatible protocols are available [21,49,50]. It should be noted that a specific spot picker robot, able to work with fluorescence, is essential when working with fluorescent dyes. On a precedent study, we analyzed the total protein maps visualized when using classical visible stains and different fluorescent dyes [49]. For this purpose, a soluble extract from *Arabidopsis thaliana* was taken as a model of sequenced eukaryotic genome and resolved by 2-DE. Besides specificities in background quality, propensity to saturation, and staining reproducibility, large differences were observed between dyes in terms of sensitivity, especially for low abundance spots. The effects of the staining procedure on MALDI-TOF MS characterization were analyzed too on a set of 48 protein spots that were selected for their contrasting abundance, pI, and Mr. Gels were stained with either classical visible stains colloidal (Coomassie blue and silver nitrate), and different fluorescent dyes (Sypro Ruby and Deep Purple). It appeared that Sypro Ruby combined several favorable features: no dependence of the identification rate upon the physico-chemical properties of proteins, no impact on frequency of missed cleavages, and a higher predicted identification rate (Fig. 2).

## B - Benefits of Gel-based proteomic

### 1 - A reduced gel to gel variation using "Difference in Gel Electrophoresis" (2D-DIGE)

Difference in gel electrophoresis (DIGE), first conceived by Unlu et al. in 1997, takes advantages of structurally




**Figure 1 Hierarchical clustering of *Arabidopsis thaliana* ecotypes based on 2DE spot amounts.** Soluble proteins from ecotype Col-0 were separated using pH 4-7 IPG in first dimension and 11% SDS-PAGE in second dimension. Proteins spots were visualized by colloidal Coomassie blue staining. The amount of each spot was estimated by its normalized volume as obtained by image analysis [62]. Euclidian distances were then computed for all spots to build the similarity matrix for ecotypes, and clustering was performed using the Ward's method to link the variables.

similar cyanine-based dyes to label different pools of protein samples, which are then co-separated on a single 2DE gel [34].

The biggest advantage of DIGE over other two-dimensional-based technologies is that it enables the analysis of two or more protein samples simultaneously on a single 2DE [31,32,35,36]. Because the same proteins present in two different samples were pre-labeled with two different dyes (i.e., Cy3 and Cy5, respectively), they could be combined and separated on the same 2DE without the loss of the relative protein abundance in the original samples. At the end of protein separation, the relative ratio of proteins in the two original samples could be readily obtained by comparing the fluorescence intensity of the same protein spots under different detection channels (e.g., Cy3 and Cy5) using a commercial fluorescence gel scanner. Because only one gel is used in DIGE, and the same proteins from two different protein samples comigrate as single spots, there is no need for the generation of "averaged" gels, as well as superimposition of different gels, making spot comparison and protein quantitation much more convenient and reliable. This makes DIGE potentially amendable for high-throughput proteomics applications.

DIGE has shown significant advantages over conventional 2DE in a number of applications. Up to three kinds of fluorescent cyanine dyes have been employed in DIGE, namely, Cy2, Cy3, and Cy5, which allows for simultaneous analysis of up to three different protein samples in a single gel. DIGE is a valuable method for high-throughput studies of protein expression profiles, providing opportunities to detect and quantify accurately "difficult" proteins, such as low-abundance proteins.

A problem in DIGE lies in the hydrophobicity of the cyanine dyes, which label the protein by reacting covalently, to a large extent, with surface-exposed lysines in the protein, and lead to removal of multiple charges from the protein. Consequently, this decreases the solubility of the labeled protein, and in some cases may lead to protein precipitation prior to gel electrophoresis. To address this problem, minimal labeling is generally employed in DIGE. Typically the labelling reaction is optimized such that only 1-5% of total lysines in a given protein are labeled. Alternatively, Shaw et al. have developed a new batch of DIGE Cy3 and Cy5 dyes, which label only free cysteines in a protein by "saturation" labeling [37]. This strategy offers greater sensitivity than the conventional



DIGE method. The biggest drawback, however, is that it only labels proteins that contain free cysteines, meaning that a certain percentage of proteins in a proteome will not be labeled with this strategy, let alone downstream detection and characterization of these proteins.

## 2 - Characterization of protein isoforms


An area of increasing interest in proteomics is the identification of post-translational modifications and/or spliced forms of a same gene or protein [64-67]. The process of determining whether a protein is expressed in a particular proteome has become a relatively simple task with the automation of the 'in-gel' digest and subsequent identification of the resulting peptides with mass spectrometers. Today, most proteins are identified by either assigning them definitive protein attributes, such as peptide masses generated by MALDI-TOF mass spectrometry and the short amino acid sequences generated by tandem MS. It is clear that when several spliced variants are present in a proteome, such approach for protein identification mostly characterizes peptides common to all spliced variants. In a precedent study, we used the advantages of 2DE separation to analyze alpha-amylase diversity in human saliva [68]. Because each alpha-amylase isoforms exist as a discrete purified protein, any

information obtained from the analysis of this protein is unique to its original proteome (Fig. 3). 2DE was combined with systematic MALDI-TOF MS analysis and more than 140 protein spots identifying the alpha-amylase were shown to constitute a stable but very complex pattern. Careful analysis of mass spectra and simultaneous hierarchical clustering of the observed peptides and of the electrophoretic features of spots defined several groups of isoforms (Fig. 3 right part) with specific sequence characteristics, potentially related with special biological activities. In a recent study, 2DE separation was successfully used to analyze isoforms and polymers forms of bovine milk proteins [69]. A combination of reducing and non-reducing steps was used to reveal proteins polymers occurring before or after heat treatment of milk (Fig. 4). This original 2DE strategy revealed numerous disulfide-mediated interactions and was proposed to analyze reduction/oxidation of milk and dairy product proteins.

## C - Limits of "Gel-based proteomic"

### 1 - Membrane proteins

The resolution of membrane proteins remains an area of considerable concern in gel-based proteomics [70-75]. There remains an attitude that it is difficult or impossible




**Figure 3 Characterization of isoforms of alpha-amylase in human saliva.** A 2DE map of human salivary proteins was performed [68]. Proteins were resolved using pH 3-10NL IPG and 12% SDS-PAGE, and stained with colloidal Coomassie blue. Alpha-amylase spots subsequently identified by MALDI-TOF MS are indicated by respective spot number (left part). According to the mass features of alpha-amylase (right part) identified spots, (A) coverage of the alpha-amylase sequence by the total population of peptides identified (black boxes) in the different alpha-amylase spots, (B) simultaneous clustering of the 67 alpha-amylase spots according to the MW range measured on gels, the MS identification of peptides in the N-terminal and C-terminal and central regions of the sequence, (C) individual spot coverage of the alpha-amylase sequence by peptides identified (black boxes) by MALDI-TOF MS.

to effectively resolve membrane proteins using 2DE. Indeed, few membrane proteins are seen on 2D gels when conventional sample-preparation methods are used. Membrane proteins are poorly soluble in the detergent/chaotropic conditions available for IEF, and are inherently insoluble in gel matrices under these conditions and thus are poorly resolved by IEF and subsequent 2DE. Fractionation, in combination with the correct solubilizing reagents, produces sample extracts that are highly enriched for membrane proteins. Sequential extraction of proteins from a sample by increasing protein solubility at each step is an effective strategy for first removing the more abundant soluble proteins and then for concentrating the less abundant and less soluble membrane proteins. A specific 2DE strategy, using the cationic detergent benzyltrimethyl-n-hexadecylammonium chloride in the first and the anionic detergent SDS in the second dimension, was successfully used to analyze membranes proteins in various systems [76-78].

## 2 - Low-abundance proteins

Low-abundance proteins are rarely seen on traditional 2D maps because large quantities of abundant soluble proteins obscure their detection [21,79-81]. Most 2DE-based proteomic studies employ a 'one-extract-one-gel' approach and the majority of proteins identified in these

studies are in high abundance. These low-abundance proteins are considered to be some of the most important, including regulatory proteins, signal transduction proteins and receptors. Consequently, the analysis of low-abundance proteins is becoming increasingly common in cellular proteomics. The dynamic range of protein concentration can differ considerably between biological samples. For yeast, the most abundant proteins are present at around 2 000 000 copies per cell, whereas the least abundant proteins are present at around 100 copies per cell, a dynamic range of only 4 orders of magnitude. However, in plasma, the predicted dynamic range of proteins is up to 12 orders of magnitude. Analysis of individual compartments not only provides information on protein localization, but also allows detection of protein populations otherwise not detectable in whole cell proteomes. Detection of the low-abundance proteins requires most of the time removal of abundant proteins from the sample. For example, the complexity of the serum and plasma proteome presents extreme analytical challenges in comprehensive analysis due to the wide dynamic range of protein concentrations. Therefore, robust sample preparation methods remain one of the important steps in the proteome characterization workflow. A specific depletion of high-abundant proteins from human serum and plasma using affinity columns is of particular interest to



improve dynamic range for proteomic analysis and enable the identification of low-abundant plasma proteins [80,82]. On another hand, IPG technology can be used with narrow (2-3 pH units) and very narrow (~1 pH unit) gradients that enable many more proteins to be resolved. Indeed, the advent of immobilized pH gradients has greatly improved the reproducibility of 2D gels and has made it easier for new users to implement this technology. The loading capacity of narrow-range IPGs is considerably higher than broad-range IPGs, thus enabling the visualization and identification of previously unseen proteins. Sub-fractionation can be used to improve the recovery of low-abundance proteins too. For example, membrane preparation methods are commercially available and allow a specific separation between abundant/soluble proteins and membrane/low-abundance proteins. More recently, a system is available to perform a specific depletion of high-abundant proteins and a reduction of protein concentration differences [82,83]. The protein population is "equalized", by sharply reducing the concentration of the most abundant components, while simultaneously enhancing the concentration of the most dilute species.

### 3 - Alkaline proteins

Alkaline proteins were particularly difficult to resolve on 2D gels. First, the most common commercially available pH gradients, until recently, were pH 4-7 and pH 3-10 and these do not provide significant alkaline-protein res-

olution. As more alkaline pH range immobilized pH gradients become commercially available, resolution of proteins in IPGs up to pH 12 has been demonstrated. Strongly alkaline proteins such as ribosomal and nuclear proteins with closely related pIs between 10.5 and 11.8 were focused to the steady state by using 3-12, 6-12 and 9- 12 pI ranges [84-86]. For highly resolved 2-D patterns, different optimization steps with respect to pH engineering and gel composition were necessary, such as the substitution of dimethylacrylamide for acrylamide, the addition near the cathode of a paper strip soaked with DTT providing a continuous influx of DTT to compensate for the loss of DTT [41,45], and the addition of isopropanol to the IPG rehydration solution in order to suppress the reverse electroendosmotic flow which causes highly streaky 2-D patterns in narrow pH range IPGs 9-12 and 10-12 [86].

### 4 - High molecular weight proteins

Conventional 2DE, using SDS-PAGE as second dimension is not very useful to visualize high-molecular-weight (HMW) proteins. Even with gradient acrylamide gels, it is very difficult to obtain a good separation of proteins up to 250 kDa. HMW proteins and protein-complexes can be analysed using agarose gel IEF [87]. Agarose 2-DE is sufficiently good at separating HMM proteins with molecular masses as large as 500 kDa from various diseased tissues and cells [88]. Indeed, this method was successfully used to analyze HMW complexes from yeast [87], HMW proteins from human plasma [89], hepatocellular carcinoma, prostate and colorectal cancer [88,90-92].

### Conclusion

Thanks to its high resolving power and its large sample loading capacity, 2DE allows several hundred proteins to be displayed simultaneously on a single gel, producing a direct and global view of a sample proteome at a given time point. Reference maps of numerous distinct samples have now been published, providing, to researchers worldwide, standardized libraries of proteins known to be present in these samples. But 2DE has some limitations that must be taken into account. Despite maximal precautions, there will be some degree of gel-to-gel and run-to-run variability in the expression of the same protein set, which could be overcome by maintaining a variability coefficient of reference spots as low as possible. It can be largely circumvented using a DIGE strategy. Additionally, some proteins may escape the capabilities of conventional 2DE for several reasons, including the poor solubility of membrane proteins and out of range characteristics of extreme proteins such as high or low pI and molecular weight. Despite all these drawbacks, 2DE can demonstrate changes in relative abundance of visualized proteins and can detect protein isoforms, variants, polymer

complexes and posttranslational modifications. Quantitative proteomics can be achieved by assessing differences in protein expression across gels using 2DE dedicated software and proteins in varying spots can be identified by MS. The uniqueness of 2DE for easy visualization of protein isoforms renders this technology itself extremely informative and it is currently the most rapid method for direct targeting of protein expression differences.

#### Competing interests

The author declares that they have no competing interests.

#### Authors' contributions

FC carried out all of this work. All authors read and approved the final manuscript.

#### Author Details

Proteomic Laboratory, iRCM, CEA, Fontenay aux Roses, France

Received: 5 March 2010 Accepted: 28 April 2010

Published: 28 April 2010

#### References

- O'Farrell PH: **High resolution two-dimensional electrophoresis of proteins.** *Journal of Biological Chemistry* 1975, **250**:4007-4021.
- Wan JH, He FC: **Technical development of proteomics.** *Chinese Science Bulletin* 1999, **44**:1441-1447.
- Gromov PS, Celis JE: **From genomics to proteomics.** *Molecular Biology* 2000, **34**:508-520.
- Govorun VM, Archakov AI: **Proteomic technologies in modern biomedical science.** *Biochemistry-Moscow* 2002, **67**:1109-1123.
- Garfin DE: **Two-dimensional gel electrophoresis: an overview.** *Trends in Analytical Chemistry* 2003, **22**:263-272.
- Collinsova M, Jiracek J: **Current development in proteomics.** *Chemické Listy* 2004, **98**:1112-1118.
- Bradshaw RA, Burlingame AL: **From proteins to proteomics.** *Lab Invest* 2005, **85**:267-272.
- Bergh G Van den, Arckens L: **Recent advances in 2D electrophoresis: an array of possibilities.** *Expert Review of Proteomics* 2005, **2**:243-252.
- Carrette O, Burkhard PR, Sanchez JC, Hochstrasser DF: **State-of-the-art two-dimensional gel electrophoresis: a key tool of proteomics research.** *Nature Protocols* 2006, **1**:812-823.
- Bergeron JJM, Bradshaw RA: **What has proteomics accomplished?** *Molecular & Cellular Proteomics* 2007, **6**:1824-1826.
- Penque D: **Two-dimensional gel electrophoresis and mass spectrometry for biomarker discovery.** *Proteomics Clinical Applications* 2009, **3**:155-172.
- Bergh G Van den, Arckens L: **High Resolution Protein Display by Two-Dimensional Electrophoresis.** *Current Analytical Chemistry* 2009, **5**:106-115.
- HumpherySmith I, Cordwell SJ, Blackstock WP: **Proteome research: Complementarity and limitations with respect to the RNA and DNA worlds.** *Electrophoresis* 1997, **18**:1217-1242.
- Celis JE, Gromov P: **2D protein electrophoresis: can it be perfected?** *Current Opinion in Biotechnology* 1999, **10**:16-21.
- Ong SE, Pandey A: **An evaluation of the use of two-dimensional gel electrophoresis in proteomics.** *Biomolecular Engineering* 2001, **18**:195-205.
- Lopez JL: **Two-dimensional electrophoresis in proteome expression analysis.** *Journal of Chromatography B-Analytical Technologies in the Biomedical and Life Sciences* 2007, **849**:190-202.
- Chevalier F, Rofidal V, Rossignol M: **Visible and fluorescent staining of two-dimensional gels.** *Methods in Molecular Biology* 2007, **355**:145-156.
- Harris LR, Churchward MA, Butt RH, Coorsen JR: **Assessing detection methods for gel-based proteomic analyses.** *Journal of Proteome Research* 2007, **6**:1418-1425.
- Volkova KD, Kovalska VB, Yarmoluk SM: **Modern techniques for protein detection on polyacrylamide gels: problems arising from the use of dyes of undisclosed structures for scientific purposes.** *Biotechnic & Histochemistry* 2007, **82**:201-208.
- Smejkal GB: **The Coomassie chronicles: past, present and future perspectives in polyacrylamide gel staining.** *Expert Review of Proteomics* 2004, **1**:381-387.
- Patton WF: **Detection technologies in proteome analysis.** *Journal of Chromatography B-Analytical Technologies in the Biomedical and Life Sciences* 2002, **771**:3-31.
- Cong WT, Hwang SY, Jin LT, Choi JK: **Sensitive fluorescent staining for proteomic analysis of proteins in 1-D and 2-D SDS-PAGE and its comparison with SYPRO Ruby by PMF.** *Electrophoresis* 2008, **29**:4304-4315.
- Ball MS, Karuso P: **Mass spectral compatibility of four proteomics stains.** *Journal of Proteome Research* 2007, **6**:4313-4320.
- Chevalier F, Centeno D, Rofidal V, Tauzin M, Martin O, Sommerer N, Rossignol M: **Different impact of staining procedures using visible stains and fluorescent dyes for large-scale investigation of proteomes by MALDI-TOF mass spectrometry.** *Journal of Proteome Research* 2006, **5**:512-520.
- Lanne B, Panfilov O: **Protein staining influences the quality of mass spectra obtained by peptide mass fingerprinting after separation on 2-D gels. A comparison of staining with coomassie brilliant blue and SYPRO Ruby.** *Journal of Proteome Research* 2005, **4**:175-179.
- Lamanda A, Zahn A, Roder D, Langen H: **Improved Ruthenium II tris (bathophenanthroline disulfonate) staining and destaining protocol for a better signal-to-background ratio and improved baseline resolution.** *Proteomics* 2004, **4**:599-608.
- White IR, Pickford R, Wood J, Skehel JM, Gangadharan B, Cutler P: **A statistical comparison of silver and SYPRO Ruby staining for proteomic analysis.** *Electrophoresis* 2004, **25**:3048-3054.
- Berggren KN, Schulenberg B, Lopez MF, Steinberg TH, Bogdanova A, Smejkal G, Wang A, Patton WF: **An improved formulation of SYPRO Ruby protein gel stain: Comparison with the original formulation and with a ruthenium II tris (bathophenanthroline disulfonate) formulation.** *Proteomics* 2002, **2**:486-498.
- Rabilloud T, Strub JM, Luche S, van Dorsselaer A, Lunardi J: **Comparison between Sypro Ruby and ruthenium II tris (bathophenanthroline disulfonate) as fluorescent stains for protein detection in gels.** *Proteomics* 2001, **1**:699-704.
- Berggren K, Steinberg TH, Lauber WM, Carroll JA, Lopez MF, Chernokalskaya E, Zieske L, Diwu ZJ, Haugland RP, Patton WF: **A luminescent ruthenium complex for ultrasensitive detection of proteins immobilized on membrane supports.** *Analytical Biochemistry* 1999, **276**:129-143.
- Karp NA, Feret R, Rubtsov DV, Lilley KS: **Comparison of DIGE and post-stained gel electrophoresis with both traditional and SameSpots analysis for quantitative proteomics.** *Proteomics* 2008, **8**:948-960.
- Hrebicek T, Duerschmid K, Auer N, Bayer K, Rizzi A: **Effect of CyDye minimum labeling in differential gel electrophoresis on the reliability of protein identification.** *Electrophoresis* 2007, **28**:1161-1169.
- Karp NA, McCormick PS, Russell MR, Lilley KS: **Experimental and statistical considerations to avoid false conclusions in proteomics studies using differential in-gel electrophoresis.** *Molecular & Cellular Proteomics* 2007, **6**:1354-1364.
- Viswanathan S, Unlu M, Minden JS: **Two-dimensional difference gel electrophoresis.** *Nature Protocols* 2006, **1**:1351-1358.
- Wheelock AM, Morin D, Bartosiewicz M, Buckpitt AR: **Use of a fluorescent internal protein standard to achieve quantitative two-dimensional gel electrophoresis.** *Proteomics* 2006, **6**:1385-1398.
- Gade D, Thiermann J, Markowsky D, Rabus R: **Evaluation of two-dimensional difference gel electrophoresis for protein profiling.** *Journal of Molecular Microbiology and Biotechnology* 2003, **5**:240-251.
- Shaw J, Rowlinson R, Nickson J, Stone T, Sweet A, Williams K, Tonge R: **Evaluation of saturation labelling two-dimensional difference gel electrophoresis fluorescent dyes.** *Proteomics* 2003, **3**:1181-1195.
- Westermeier R, Loyland S, Asbury R: **Proteomics technology.** *Journal of Clinical Ligand Assay* 2002, **25**:242-252.
- Bjellqvist B, Ek K, Righetti PG, Gianazza E, Gorg A, Westermeier R, Postel W: **Isoelectric-focusing in immobilized pH gradients - principle, methodology and some applications.** *Journal of Biochemical and Biophysical Methods* 1982, **6**:317-339.



40. Righetti PG, Castagna A, Hamdan M: **Recent trends in proteome analysis.** *Advances in Chromatography* 2003, **42**:269-321. *Advances in Chromatography*
41. Altland K, Becher P, Rossmann U, Bjellqvist B: **Isoelectric-focusing of basic-proteins - the problem of oxidation of cysteines.** *Electrophoresis* 1988, **9**:474-485.
42. Bouchal P, Kucera I: **Two-dimensional electrophoresis in proteomics: Principles and applications.** *Chemische Listy* 2002, **97**:29-36.
43. Friedman DB, Hoving S, Westermeyer R: **Isoelectric focusing and two-dimensional gel electrophoresis.** *Guide to Protein Purification* Second edition. 2009, **466**:515-540. *Methods in Enzymology*
44. Gorg A, Postel W, Gunther S: **The current state of two-dimensional electrophoresis with immobilized pH gradients.** *Electrophoresis* 1988, **9**:531-546.
45. Gorg A, Boguth G, Obermaier C, Posch A, Weiss W: **2-dimensional polyacrylamide-gel electrophoresis with immobilized pH gradients in the first dimension (IPG-DALT) - the state-of-the-art and the controversy of vertical versus horizontal systems.** *Electrophoresis* 1995, **16**:1079-1086.
46. Gorg A, Weiss W, Dunn MJ: **Current two-dimensional electrophoresis technology for proteomics.** *Proteomics* 2004, **4**:3665-3685.
47. Rabilloud T: **A comparison between low background silver diamine and silver-nitrate protein stains.** *Electrophoresis* 1992, **13**:429-439.
48. Neuheff V, Stamm R, Pardowitz I, Arold N, Ehrhardt W, Taube D: **Essential problems in quantification of proteins following colloidal staining with coomassie brilliant blue dyes in polyacrylamide gels, and their solution.** *Electrophoresis* 1990, **11**:1101-1117.
49. Chevalier F, Rofidal V, Vanova P, Bergoin A, Rossignol M: **Proteomic capacity of recent fluorescent dyes for protein staining.** *Phytochemistry* 2004, **65**:1499-1506.
50. Mortz E, Krogh TN, Vorum H, Gorg A: **Improved silver staining protocols for high sensitivity protein identification using matrix-assisted laser desorption/ionization-time of flight analysis.** *Proteomics* 2001, **1**:1359-1363.
51. Jin LT, Li XK, Cong WT, Hwang SY, Choi JK: **Previsible silver staining of protein in electrophoresis gels with mass spectrometry compatibility.** *Analytical Biochemistry* 2008, **383**:137-143.
52. Nebrich G, Liegmann H, Wacker M, Herrmann M, Sagi D, Landowsky A, Klose J: **Proteomer, a novel software application for management of proteomic 2DE-gel data-II application.** *Molecular & Cellular Proteomics* 2005, **4**:S296-S296.
53. Wheelock AM, Buckpitt AR: **Software-induced variance in two-dimensional gel electrophoresis image analysis.** *Electrophoresis* 2005, **26**:4508-4520.
54. Maurer MH: **Software analysis of two-dimensional electrophoretic gels in proteomic experiments.** *Current Bioinformatics* 2006, **1**:255-262.
55. Wheelock AM, Goto S: **Effects of post-electrophoretic analysis on variance in gel-based proteomics.** *Expert Review of Proteomics* 2006, **3**:129-142.
56. Rosengren AT, Salmi JM, Aittokallio T, Westerholm J, Laheesmaa R, Nyman TA, Nevalainen OS: **Comparison of PDQuest and Progenesis software packages in the analysis of two-dimensional electrophoresis gels.** *Proteomics* 2003, **3**:1936-1946.
57. Nebrich G, Liegmann H, Wacker M, Herrmann M, Sagi D, Landowsky A, Klose J: **Proteomer, a novel software application for management of proteomic 2DE-gel data-II application.** *Molecular & Cellular Proteomics* 2005, **4**:S296-S296.
58. Clark BN, Gutstein HB: **The myth of automated, high-throughput two-dimensional gel analysis.** *Proteomics* 2008, **8**:1197-1203.
59. Pancaud A, Affolter M, Moreillon P, Kussmann M: **Experimental and computational approaches to quantitative proteomics: Status quo and outlook.** *Journal of Proteomics* 2008, **71**:19-33.
60. Kang YY, Techanukul T, Mantalaris A, Nagy JM: **Comparison of Three Commercially Available DIGE Analysis Software Packages: Minimal User Intervention in Gel-Based Proteomics.** *Journal of Proteome Research* 2009, **8**:1077-1084.
61. Chevalier F, Pata M, Nacry P, Doumas P, Rossignol M: **Effects of phosphate availability on the root system architecture: large-scale analysis of the natural variation between Arabidopsis accessions.** *Plant Cell and Environment* 2003, **26**:1839-1850.
62. Chevalier F, Martin O, Rofidal V, Devauchelle AD, Barteau S, Sommerer N, Rossignol M: **Proteomic investigation of natural variation between Arabidopsis ecotypes.** *Proteomics* 2004, **4**:1372-1381.
63. Nock CM, Ball MS, White IR, Skehel JM, Bill L, Karuso P: **Mass spectrometric compatibility of Deep Purple and SYPRO Ruby total protein stains for high-throughput proteomics using large-format two-dimensional gel electrophoresis.** *Rapid Communications in Mass Spectrometry* 2008, **22**:881-886.
64. Holland JW, Deeth HC, Alewood PF: **Proteomic analysis of K-casein micro-heterogeneity.** *Proteomics* 2004, **4**:743-752.
65. Schulenberg B, Goodman TN, Aggeler R, Capaldi RA, Patton WF: **Characterization of dynamic and steady-state protein phosphorylation using a fluorescent phosphoprotein gel stain and mass spectrometry.** *Electrophoresis* 2004, **25**:2526-2532.
66. Ahler K, Jungbauer A: **Chromatographic and electrophoretic characterization of protein variants.** *Journal of Chromatography B-Analytical Technologies in the Biomedical and Life Sciences* 2006, **841**:110-122.
67. Poth AG, Deeth HC, Alewood PF, Holland JW: **Analysis of the Human Casein Phosphoproteome by 2-D Electrophoresis and MALDI-TOF/TOF MS Reveals New Phosphoforms.** *Journal of Proteome Research* 2008, **7**:5017-5027.
68. Hirtz C, Chevalier F, Centeno D, Rofidal V, Egea JC, Rossignol M, Sommerer N, de Periee DD: **MS characterization of multiple forms of alpha-amylase in human saliva.** *Proteomics* 2005, **5**:4597-4607.
69. Chevalier F, Hirtz C, Sommerer N, Kelly AL: **Use of Reducing/Nonreducing Two-Dimensional Electrophoresis for the Study of Disulfide-Mediated Interactions between Proteins in Raw and Heated Bovine Milk.** *Journal of Agricultural and Food Chemistry* 2009, **57**:5948-5955.
70. Luche S, Santoni V, Rabilloud T: **Evaluation of nonionic and zwitterionic detergents as membrane protein solubilizers in two-dimensional electrophoresis.** *Proteomics* 2003, **3**:249-253.
71. Santoni V, Kieffer S, Desclaux D, Masson F, Rabilloud T: **Membrane proteomics: Use of additive main effects with multiplicative interaction model to classify plasma membrane proteins according to their solubility and electrophoretic properties.** *Electrophoresis* 2000, **21**:3329-3344.
72. Santoni V, Molloy M, Rabilloud T: **Membrane proteins and proteomics: Un amour impossible?** *Electrophoresis* 2000, **21**:1054-1070.
73. Santoni V, Doumas P, Rouquie D, Mansion M, Rabilloud T, Rossignol M: **Large scale characterization of plant plasma membrane proteins.** *Biochimie* 1999, **81**:655-661.
74. Santoni V, Rabilloud T, Doumas P, Rouquie D, Mansion M, Kieffer S, Garin J, Rossignol M: **Towards the recovery of hydrophobic proteins on two-dimensional electrophoresis gels.** *Electrophoresis* 1999, **20**:705-711.
75. Chevalier M, Santoni V, Poinas A, Rouquie D, Fuchs A, Kieffer S, Rossignol M, Lunardi J, Garin J, Rabilloud T: **New zwitterionic detergents improve the analysis of membrane proteins by two-dimensional electrophoresis.** *Electrophoresis* 1998, **19**:1901-1909.
76. Moebius J, Zahedi RP, Lewandrowski U, Berger C, Walter U, Sickmann A: **The human platelet membrane proteome reveals several new potential membrane proteins.** *Molecular & Cellular Proteomics* 2005, **4**:1754-1761.
77. Zahedi RP, Meisinger C, Sickmann A: **Two-dimensional benzyldimethyl-n-hexadecylammonium chloride/SDS-PAGE for membrane proteomics.** *Proteomics* 2005, **5**:3581-3588.
78. Dreger M, Bengtsson L, Schoneberg T, Otto H, Hucho F: **Nuclear envelope proteomics: Novel integral membrane proteins of the inner nuclear membrane.** *Proceedings of the National Academy of Sciences of the United States of America* 2001, **98**:11943-11948.
79. Yamada M, Murakami K, Wallingford JC, Yuki Y: **Identification of low-abundance proteins of bovine colostrum and mature milk using two-dimensional electrophoresis followed by microsequencing and mass spectrometry.** *Electrophoresis* 2002, **23**:1153-1160.
80. Greenough C, Jenkins RE, Kitteringham NR, Pirmohamed M, Park BK, Pennington SR: **A method for the rapid depletion of albumin and immunoglobulin from human plasma.** *Proteomics* 2004, **4**:3107-3111.
81. Ahmed N, Rice GE: **Strategies for revealing lower abundance proteins in two-dimensional protein maps.** *Journal of Chromatography B-Analytical Technologies in the Biomedical and Life Sciences* 2005, **815**:39-50.

82. Righetti PG, Castagna A, Boschetti E, Lomas L: **Equalizer beads; The quest for a democratic proteome.** *Molecular & Cellular Proteomics* 2005, **4**:S12-S12.
83. Righetti PG, Boschetti E, Lomas L, Citterio A: **Protein Equalizer (TM) Technology: The quest for a democratic proteome.** *Proteomics* 2006, **6**:3980-3992.
84. Drews O, Reil G, Parlar H, Gorg A: **Setting up standards and a reference map for the alkaline proteome of the Gram-positive bacterium *Lactococcus lactis*.** *Proteomics* 2004, **4**:1293-1304.
85. Wildgruber R, Reil G, Drews O, Parlar H, Gorg A: **Web-based two-dimensional database of *Saccharomyces cerevisiae* proteins using immobilized pH gradients from pH 6 to pH 12 and matrix-assisted laser desorption/ionization-time of flight mass spectrometry.** *Proteomics* 2002, **2**:727-732.
86. Gorg A, Obermaier C, Boguth G, Csordas A, Diaz JJ, Madjar JJ: **Very alkaline immobilized pH gradients for two-dimensional electrophoresis of ribosomal and nuclear proteins.** *Electrophoresis* 1997, **18**:328-337.
87. Yokoyama R, Iwafune Y, Kawasaki H, Hirano H: **Isoelectric focusing of high-molecular-weight protein complex under native conditions using agarose gel.** *Analytical Biochemistry* 2009, **387**:60-63.
88. Oh-Ishi M, Maeda T: **Disease proteomics of high-molecular-mass proteins by two-dimensional gel electrophoresis with agarose gels in the first dimension (Agarose 2-DE).** *Journal of Chromatography B-Analytical Technologies in the Biomedical and Life Sciences* 2007, **849**:211-222.
89. Jin Y, Manabe T: **Performance of agarose IEF gels as the first dimension support for non-denaturing micro-2-DE in the separation of high-molecular-mass plasma proteins and protein complexes.** *Electrophoresis* 2009, **30**:939-948.
90. Seimiya M, Tomonaga T, Matsushita K, Sunaga M, Oh-Ishi M, Kodera Y, Maeda T, Takano S, Togawa A, Yoshitomi H, et al.: **Identification of novel immunohistochemical tumor markers for primary hepatocellular carcinoma; Clathrin heavy chain and formiminotransferase cyclodeaminase.** *Hepatology* 2008, **48**:519-530.
91. Kuruma H, Egawa S, Oh-Ishi M, Kodera Y, Satoh M, Chen WG, Okusa H, Matsumoto K, Maeda T, Baba S: **High molecular mass proteome of androgen-independent prostate cancer.** *Proteomics* 2005, **5**:1097-1112.
92. Tomonaga T, Matsushita K, Yamaguchi S, Oh-Ishi M, Kodera Y, Maeda T, Shimada H, Ochiai T, Nomura F: **Identification of altered protein expression and post-translational modifications in primary colorectal cancer by using agarose two-dimensional gel electrophoresis.** *Clinical Cancer Research* 2004, **10**:2007-2014.

doi: 10.1186/1477-5956-8-23

**Cite this article as:** Chevalier, Highlights on the capacities of "Gel-based" proteomics *Proteome Science* 2010, **8**:23

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

