



Published in final edited form as:

Rheum Dis Clin North Am. 2010 May ; 36(2): 367–383. doi:10.1016/j.rdc.2010.02.005.

A multitude of kinases—Which are the best targets in treating rheumatoid arthritis?

Tamsin M. Lindstrom, PhD^{1,2} and William H. Robinson, MD PhD^{1,2}

¹Geriatric Research Education and Clinical Center (GRECC), VA Palo Alto Health Care System, Palo Alto, CA, 94304, USA

²Division of Immunology and Rheumatology, Stanford University School of Medicine, Stanford, CA, 94305, USA

Synopsis

Small-molecule kinase inhibitors are increasingly taking center stage in the quest for new drugs for the treatment of rheumatoid arthritis (RA). By targeting kinases, many of which orchestrate multiple signaling pathways, small-molecule inhibitors can exert potent anti-inflammatory and immunomodulatory effects. The success of small-molecule kinase inhibitors in the treatment of cancer, coupled with greater insight into inflammatory and immune signaling, has spurred efforts to identify kinases that could be targeted for the treatment of chronic inflammatory disorders such as RA. Several kinases have been convincingly implicated in the pathogenesis of RA, and many kinase inhibitors have proven efficacious in the treatment of inflammatory arthritis in animals; but few kinase inhibitors have so far been tested in RA clinical trials. Here we discuss the challenges and progress in the pursuit of small-molecule kinase inhibitors for RA, including the lessons learnt from the failure of erstwhile front-runner inhibitors and the tentative promise of inhibitors making their debut on the RA stage.

Keywords

MAPKs; tyrosine kinases; IKK; JAK; Syk; small-molecule inhibitors

Introduction

The advent of biologic therapeutics, most notably anti-tumor necrosis factor (TNF) agents, has dramatically improved the treatment of rheumatoid arthritis (RA). Nevertheless, the available biologics rarely result in disease remission and provide clinical benefit only in subsets of RA patients. In addition, biologics can be administered only by injection and are expensive. Alternative therapies for RA are needed, and small-molecule kinase inhibitors may fit the bill. Small molecules have several features that give them the edge over other therapeutics: they are orally bioavailable, cell permeable, and inexpensive to manufacture. Insight into intracellular signaling pathways involved in inflammation and immunity has allowed the

© 2009 Elsevier Inc. All rights reserved.

Correspondence should be addressed to T.M.L. or W.H.R., MC154R, VA Palo Alto Health Care System, 3801 Miranda Ave, Palo Alto, CA 94304, USA; (650) 849-1207 phone; (650) 849-1208 fax; tamsinlindstrom@gmail.com; wrobins@stanford.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

rational design of small molecules that can counteract aberrant immune responses. Many of these small molecules inhibit kinases, which often lie at the nexus of multiple proinflammatory pathways, and may thus have potent anti-inflammatory properties. The therapeutic potential of kinase inhibitors is showcased by their success in the treatment of cancer.

Both adaptive and innate immune responses are involved in the pathogenesis of RA, a systemic autoimmune disease characterized by destruction of the synovial joints. Initiation of the disease involves systemic dysregulation of T- and B-cell responses, which leads to a breach in self-tolerance and eventually to the mounting of an immune response against the synovial joints. During the chronic inflammatory stage of the disease, mast cells, macrophages, neutrophils, T cells, and B cells all infiltrate the synovium, where they release proinflammatory cytokines and matrix metalloproteinases (MMPs) that erode the synovial cartilage. Inflammation in the joints also triggers the development of apoptosis-resistant, hyperproliferative fibroblast-like synoviocytes (FLS), which produce further proinflammatory cytokines. The synovial hyperplasia, in turn, leads to the formation of a destructive pannus that invades surrounding cartilage and bone. Finally, inflammation suppresses the formation of bone-forming osteoblasts and augments the formation of bone-resorbing osteoclasts, leading to the erosion of bone. Several kinases have been shown to play important roles in one or more of these pathogenic processes. Here we discuss the therapeutic potential of small molecules targeting specific protein kinases in the treatment of RA, and provide an overview of the progress to date. Lipid kinases—in particular the phosphatidylinositol 3 kinases (PI3Ks)—are also emerging as attractive drug targets in the treatment of inflammation. The therapeutic potential of blocking PI3Ks in RA has been recently reviewed⁷⁷ and will not be discussed further here.

Mitogen-activated protein kinases (MAPK): advances and setbacks

MAPK signaling comprises three interrelated pathways mediated by the MAPKs p38, extracellular signal-regulated kinase (ERK), or c-Jun terminal kinase (JNK). Each of these pathways involves the sequential activation of multiple kinases, such that the MAPKs are activated by MAPK kinases (MKKs), which are themselves activated by MAPKK kinases (MKKKs). Thus, the p38 kinases (α , β , γ , and δ) are activated by MKK3 and MKK6; the ERKs (1 and 2) by MEK1 and MEK2; and the JNKs (1, 2, and 3) by MKK4 and MKK7.⁷⁵ JNK, ERK, and p38 are the terminal kinases of these pathways and serve to regulate an array of cellular responses through the phosphorylation of serine/threonine residues in discrete sets of transcription factors. All three of these MAPKs are activated in RA synovium⁸² and have been proposed as therapeutic targets in the treatment of RA.

p38

Enthusiasm for inhibitors of p38—until recently heralded as one of the most promising class of oral therapeutics for RA—has finally subsided. Numerous p38 inhibitors have been developed and tested in preclinical and clinical studies. While the preclinical data were encouraging, with p38 inhibition shown to suppress inflammation and joint destruction in multiple different models of RA,³⁰ these initial successes did not extend to the treatment of RA. The first generation of small-molecule p38 inhibitors, which targeted all four isoforms of p38, failed in clinical trials owing to liver, brain, and skin toxicities. Nevertheless, the discovery that p38 α is the important isoform in RA, acting to drive the expression of proinflammatory cytokines and the formation of osteoclasts,^{6,83} engendered hope that selective inhibition of p38 α would avoid the adverse effects of the pan-38 inhibitors. Unfortunately, p38 α -specific inhibitors did not perform much better (Table 1). For instance, clinical development of Scio-323 and AMG-548 was terminated because of skin toxicity and liver toxicity, respectively,³² while the p38 α inhibitors that did advance to phase II clinical trials proved to be ineffective.^{12,16}

Both the toxicity and the inefficacy of p38 inhibitors are most likely target based, rendering the systemic targeting of p38 unviable. Multiple structurally unrelated p38 inhibitors have been shown to be toxic to the liver and skin and to induce only transient reductions in markers of inflammation.³⁰⁻³² p38 α 's pivotal position in the regulation of inflammation is thought to underlie these phenomena. Although its proinflammatory role has long been recognized, p38 α has more recently been found to play an anti-inflammatory role, too. Not only does it drive the expression of important anti-inflammatory genes, but also it mediates intracellular feedback loops that constrain the activity of other proinflammatory pathways. For instance, p38 α activates mitogen- and stress-activated protein kinase (MSK) 1 and MSK2, which contribute to the resolution of inflammation through the transcriptional activation of anti-inflammatory genes such as interleukin (IL)-10, IL-1 receptor antagonist, and protein phosphatase dual specificity.²⁻¹⁷⁻⁵¹ p38 α also reigns in inflammation by phosphorylating TAK-associated kinase 1 (TAB1) and thereby inhibiting TAK1, which regulates the proinflammatory JNK and I κ B kinase (IKK) pathways, as well as p38 α itself.³⁰ Thus, blockade of p38 α would allow inflammation to proceed unchecked. Genetic evidence supports the idea that p38 α inhibition underlies the toxicity and inefficacy of p38 inhibitors: Myeloid cell-specific ablation of p38 α in mice results in increased ERK and JNK activity, and in vascular permeability and edema;⁵¹ double deficiency in MSK1 and MSK2 leads to prolonged inflammation in a model of toxic contact eczema;² and hepatocyte-specific ablation of p38 α in mice results in excessive activation of the pro-apoptotic JNK in the liver following LPS challenge.⁴²

While the death knell may have sounded for inhibitors of p38, components downstream of p38 α may yet constitute viable therapeutic targets. MAPK-activated protein kinase 2 (MK2), a kinase downstream of p38 α that posttranscriptionally promotes the expression of proinflammatory genes, has been proposed as one such candidate.³⁰ Targeting of MK2 should spare p38 α -mediated anti-inflammatory mechanisms, including the p38 α -TAB1 feedback loop and expression of anti-inflammatory genes. Support for such an approach comes from the finding that MK2-deficient mice are protected against collagen-induced arthritis (CIA).⁴⁰ One small-molecule MK2 inhibitor has already been shown to reduce lipopolysaccharide (LPS)-induced TNF production in rats, and many more are being synthesized.³⁻⁴⁹

MEKs and ERKs

The critical role of MEK-ERK signaling in cell proliferation has led to MEK1/2 and ERK1/2 being investigated as candidate targets in clinical trials in cancer.⁷⁵ MEK-ERK signaling is upregulated in synovial tissues in RA and in CIA,⁹⁴ and promotes proliferation of RA FLS *in vitro*.⁶⁵ MEK-ERK signaling may thus contribute to the pathogenesis of RA by driving formation of the tumor-like pannus that is characteristic of RA. But the MEK-ERK cascade is not solely a proliferative one—it is also proinflammatory, inducing the production of IL-1 β , IL-6, TNF, and MMPs and itself being activated by proinflammatory cytokines. In addition to promoting inflammation and tissue destruction in the synovial joints, ERK signaling is important in lymphocyte activation and differentiation. ERK mediates B-cell receptor (BCR) and CD40 receptor signaling in B cells, and T-cell receptor (TCR) signaling in T cells.^{64,87} Interestingly, recent data suggest that ERK dysregulation in both CD4⁺ and CD8⁺ T cells may even contribute to the breakdown of T-cell tolerance in RA, by lowering the threshold for T-cell activation.⁸⁷

Several small-molecule inhibitors of MEK 1/2 have shown efficacy in animal models of RA. Oral administration of PD184352 to mice with CIA suppressed synovitis, pannus formation, and cartilage and bone erosion; it also prevented proteoglycan loss in articular cartilage in a rabbit model of IL-1 β -induced arthritis.⁹⁴ These effects correlated closely with the inhibition of ERK phosphorylation in mouse joints. Prophylactic, intraperitoneal administration of

subtherapeutic doses of U0126 to SKG mice, which spontaneously develop autoimmune arthritis owing to a mutation in *zap70*.⁷⁹ delayed disease onset and reduced disease severity, supporting the concept that ERK dysregulation may contribute to the development of RA.⁸⁷ A third MEK 1/2 inhibitor, ARRY-162, inhibited inflammation and bone resorption in mice with CIA and in rats with adjuvant-induced arthritis (AIA), and exhibited additive efficacy when combined with standard-of-care agents such as anti-TNFs and methotrexate.¹⁰² These promising findings saw ARRY-162 enter clinical development; however, despite being well tolerated, ARRY-162 did not fare any better than placebo in a recent phase II, 12-week trial in patients with active RA on a background of methotrexate treatment (Table 1).

In addition to inhibitors of MEK 1/2, an inhibitor of the downstream ERKs has been assessed in a mouse model of RA. Intraperitoneal administration of the ERK 1/2 inhibitor FR180204 to mice before the induction of CIA reduced the clinical signs of arthritis, the production of anti-collagen type II (CII) antibodies, and CII-specific proliferation of T cells.⁶⁸ Conversely, recent studies suggest that targeting components upstream of MEK may also provide efficacy in RA. Tumor progression locus 2 (Tpl2) is the MKKK that activates MEK 1 and 2 and hence the ERKs.²⁵⁻²⁸ Studies using *tpl2*^{-/-} mice have shown that Tpl2 is required for LPS-induced production of circulating TNF *in vivo* and for LPS-induced production of TNF by macrophages *in vitro*.²⁵⁻⁹⁰ Furthermore, Tpl2 deficiency has been shown to protect mice from TNF-induced inflammatory bowel disease⁵⁴ and arthritis (unpublished data and 18). Several small-molecule inhibitors of Tpl2, developed by Wyeth Research, have been assessed for their ability to suppress TPL2-MEK-ERK-induced inflammation. Compound 1 suppressed LPS- and IL-1 β -induced production of TNF by human monocytes, as well as IL-1 β -induced production of IL-6, IL-8, prostaglandin E2, and MMPs by RA FLS.³⁷ Compound 44 inhibited the production of TNF in an LPS-induced model of inflammation in rats.³⁵ Results from the testing of Tpl2 inhibitors in animal models of RA have not been described to date.

Thus, small-molecule inhibitors exist for the targeting of the TPL2-MEK-ERK pathway at three different levels. However, the inefficacy of the MEK 1/2 inhibitor ARRY-162 in a phase II RA trial, together with concerns that MEK/ERK inhibition could result in the development of lupus-like disease,^{20,21}⁸¹ raise doubts over the potential of MEK/ERK inhibitors for the treatment of RA. Safety might also be an issue with Tpl2 inhibitors, but these could potentially provide greater therapeutic efficacy than MEK/ERK inhibitors. Although the signaling defect in Tpl2-deficient macrophages and B cells appears to be restricted to activation of the MEK/ERK pathway,^{25,28} Tpl2 regulates the activation of JNK and nuclear factor kappa B (NF- κ B), in addition to ERK, in mouse embryonic fibroblasts.¹⁸ Because synovial-fibroblast production of proinflammatory and degradative mediators is important in the pathogenesis of RA, inhibition of Tpl2 might provide added benefit by suppressing both ERK-driven activation of lymphocytes and ERK-, JNK-, and NF- κ B-driven activation of synovial fibroblasts.

JNK

Activated by stress signals and cytokines, JNKs play important roles in apoptosis, inflammation, and matrix degradation.^{56,97} JNKs exist as three isoforms: JNK1, JNK2, and JNK3. JNK1 and JNK2 are ubiquitously expressed, and phosphorylation of these isoforms is detected in RA synovium but not in osteoarthritic synovium;⁹¹ JNK3 expression is largely restricted to the brain, heart, and testes, and therefore not thought to be involved in RA.^{36,59} As discussed below, some of the efficacy of spleen tyrosine kinase (Syk) inhibitors in RA could potentially be attributed to the inhibition of JNKs, because the tyrosine kinase Syk lies upstream of JNK in the MAPK signaling cascade. Notably, Syk-activated JNKs drive the expression of IL-6 and MMP-3 in RA FLS.¹¹ Induction of MMP expression is defective in JNK1- or JNK2-deficient murine FLS, and pharmacologic inhibition of JNK blocks induction of MMP

expression in RA FLS.³⁹ In addition to promoting synoviocyte production of proinflammatory mediators, JNK1 regulates the differentiation of T cells into Th1 cells.²²

The JNK-driven expression of MMPs appears to be critical in the destruction of joints in inflammatory arthritis. Subcutaneous administration of SP600125, a small-molecule inhibitor that targets all three JNK isoforms, suppressed cartilage and bone erosion in rat AIA, effects associated with inhibition of both JNK activity and MMP expression in the joints.³⁹ Oral administration of another pan-JNK inhibitor, AS601245, attenuated CIA in mice, reducing synovial inflammation and cartilage degradation.³¹ JNK1 deficiency does not confer resistance to destructive arthritis in JNK1-deficient, TNF-transgenic mice, nor does it reduce the activity of JNK-mediated signaling.⁵³ In addition, JNK2 deficiency confers only modest protection against the development of CAIA.³⁹ Together, these findings suggest that inhibition of both JNK1 and JNK2 is required for the effective attenuation of inflammatory arthritis.

Although developed as a JNK inhibitor, SP600125 has been shown to inhibit 13 other protein kinases with similar or greater potency and to have an unfavorable pharmacokinetic profile.⁴⁻⁹¹ Likewise, AS601245 exhibits only moderate selectivity for JNK.³¹ More specific inhibition of the JNK signaling cascade can be achieved by targeting the physical interaction between JNK and other components of the cascade. JNK-interacting protein-1 (JIP1) is a scaffolding protein that promotes JNK activity by facilitating the interaction between JNK and upstream kinases.¹⁰¹ Overexpression of JIP1, however, suppresses JNK activity (presumably owing to the sequestration of JNK-interacting components), and a peptide corresponding to the minimal region of JIP1 (pepJIP1) has been developed as an inhibitor of JNK.⁴³ While peptide therapeutics are associated with disadvantages such as their rapid degradation *in vivo* and the need for administration via injection, a small-molecule mimic of pepJIP1, BI-78D3, was recently developed and shown to exert anti-inflammatory effects *in vivo*, restoring insulin sensitivity in a mouse model of type 2 diabetes.⁸⁸ In addition, a small-molecule inhibitor that selectively blocks the DNA-binding activity of AP-1, an important JNK-activated transcription-factor complex, was recently shown to be efficacious in a mouse model of arthritis. Oral administration of the AP-1 inhibitor T-5224 both prevented and treated CIA in mice, abrogating joint destruction and suppressing MMP and IL-1 β expression.¹

Although toxicity in animal models treated with inhibitors of the JNK pathway has not been reported, long-term suppression of JNK could potentially have adverse effects due to JNK's role in regulating apoptosis.⁹⁷ JNK1-deficient mice spontaneously develop intestinal tumors and are more susceptible to the development of TPA-induced skin tumors.^{86,96} Thus, increased tumorigenicity may limit the value of JNK inhibitors for the treatment of chronic inflammatory disorders such as RA.

Tyrosine kinases: the frontrunners

Tyrosine kinases targeted in RA clinical trials

JAKs—Janus kinases (JAKs) play important roles in innate and adaptive immune responses, serving to transduce signals from cytokine receptors that lack intrinsic kinase activity. Cytokine receptors containing the common γ -chain subunit signal through JAK1 and JAK3, while receptors for hematopoietic growth factors or gp40-containing cytokines signal through JAK2. JAK1 and JAK2 are ubiquitously expressed and are essential for lymphopoiesis and hematopoiesis, respectively.³³ JAK3 is expressed primarily in cells of the immune system and is critical in lymphocyte activation, function, and proliferation;⁴⁸ accordingly, the defect in JAK3-deficient mice appears to be restricted to T cells, B cells, and natural killer cells.⁶⁶⁻⁹⁵

Given their multifarious roles in innate and adaptive immunity, one might well expect JAKs to be involved in the pathogenesis of RA. It was not until recently, however, that JAKs began

to be explored as candidate therapeutic targets in RA. Progress has since been rapid. The finding that inhibition of JAK3 ameliorates clinical signs of inflammatory arthritis by >90% and protects against joint damage in rodent models of RA⁶³ was swiftly followed by assessment of the therapeutic efficacy of two small-molecule JAK inhibitors—CP690550 and INCB18424—in patients with RA. CP690550 was developed as a JAK3 inhibitor but also inhibits JAK2, albeit less potently; its selectivity for the JAKs has been confirmed by testing against a panel of 317 kinases.⁴⁷ INCB18424 is an inhibitor primarily of JAKs 1 and 2. High hopes are now pinned on these JAK inhibitors. They are arguably the best-performing investigational small molecule drugs in RA at present, with both CP690550 and INCB18424 proving efficacious and well tolerated in initial phase II clinical trials (CP690550, 6-week trial; INCB18424, 28-day trial) (Table 1). Which of these two JAK inhibitors will prove to be safer in the long term remains to be seen. The restriction of JAK3 expression to hematopoietic cells might mean that a JAK3 inhibitor will have fewer target-based adverse effects than a JAK1/2 inhibitor; on the other hand, *JAK3* mutations in humans are known to cause severe immunodeficiency syndrome.^{58,78} In addition, the nature of the adverse effects seen with CP690550 suggest that therapeutically efficacious doses of this compound result in inhibition of JAK2 in addition to JAK3.⁵⁵ Conversely, JAK3 signaling may be indirectly affected by inhibitors of JAK1, since JAK1 and JAK3 cooperate in the transduction of multiple signals.⁹⁹ The outcomes of phase IIb trials of CP690550 and INCB18424 are eagerly awaited.

Syk—Another prime therapeutic contender is R788, the prodrug for the R406 small-molecule inhibitor of Syk. Syk is expressed in all hematopoietic cells, mediating immunoreceptor signaling such as BCR signaling in B cells and FcγR signaling in mast cells, macrophages, neutrophils, and basophils.⁵ It is also expressed in nonhematopoietic cells, in which it transduces signals from receptors for TNF, IL-1, and LPS. Syk activity is upregulated in RA synovium compared to control osteoarthritic synovium and mediates the production of IL-6 and MMP-3—major culprits in joint destruction—in TNF-stimulated RA FLS.¹¹ Syk also promotes osteoclast activity.⁵ Thus, Syk may promote both the adaptive immune responses and the destructive effector processes that underlie RA, making it an attractive therapeutic target. Indeed, the R406 Syk inhibitor suppressed inflammation and joint destruction in two antibody-mediated models of RA in mice,⁷ as well as in a T-cell-mediated model of RA in rats.⁷³ In a preliminary 12-week phase II trial in RA, R788 (which is rapidly converted to R406 following oral administration) proved efficacious and generally well tolerated.¹⁰⁰ Notably, R788 administration resulted in a rapid and sustained decrease in serum IL-6 and MMP-3 levels, an indication that Syk inhibition may be able to halt joint damage. The long-term efficacy and safety of R788 is the focus of an ongoing open-label study of the RA patients who completed the initial R788 phase II trial. Although relatively specific for Syk,⁷ R788 did cause hypertension in a limited number of RA patients, which may reflect off-target inhibition of the vascular-endothelial growth factor receptor (VEGFR).¹⁰⁰ This observation has raised some concern about the safety of R788 in RA, a disease associated with increased cardiovascular complications.⁴⁴ As for target-mediated adverse effects, the ubiquity of Syk may be an issue, but its non-redundant functions in adulthood may not be as widespread as its expression.⁵ Interestingly, Syk has been shown to signal upstream of JNK in mast cells⁶⁰ and in RA FLS;¹¹ therefore, Syk inhibition could potentially share some of the advantages and disadvantages of JNK inhibition (see section on JNK).

Tyrosine kinases targeted in animal models of RA

Several other tyrosine kinases have been implicated in RA, partly on the basis of observations in cancer patients treated with imatinib mesylate (imatinib). Imatinib, the first kinase inhibitor introduced into clinical practice, targets several tyrosine kinases, including Bcr-Abl, PDGFR, c-Fms, c-Kit, Syk, and Lck. Case studies documented the alleviation of RA symptoms in patients administered imatinib for the treatment of chronic myelogenous leukemias or c-Kit-

expressing gastrointestinal stromal tumors,^{19,23} suggesting that one or more of the imatinib-targeted kinases are important in the pathogenesis of RA. Prompted by these findings, Eklund and colleagues administered imatinib to three patients with treatment-refractory RA. All three patients showed some degree of clinical improvement;²⁶ one patient continued treatment for 24 months and showed marked and long-lasting clinical improvement.²⁷ However, two of the three patients in this study discontinued imatinib treatment at two and at four months, owing to adverse events. Furthermore, the outcomes of a double-blind, placebo-controlled, 3-month, phase II trial conducted by Novartis, in which imatinib was administered to patients with active RA despite methotrexate treatment, were never reported. Although toxicities—including cardiotoxicity due to inhibition of Abl⁵⁰—may limit the use of imatinib in non-oncologic chronic diseases, selectively inhibiting the imatinib-targeted kinases that are important in RA may provide a more favorable risk-to-benefit ratio. In mouse studies, imatinib-induced attenuation of CIA was associated with suppression of c-Fms activation in synovial macrophages, of PDGFR activation in FLS, and of c-Kit activation in mast cells.⁷² The involvement of each of these tyrosine kinases in RA has been independently investigated.

Accumulating evidence suggests that c-Fms and its ligand macrophage colony-stimulating factor (M-CSF) are involved in the pathogenesis of RA. M-CSF-c-Fms signaling is integral to macrophage and osteoclast formation, as evidenced by the osteopetrosis and the reduction in tissue macrophages in both M-CSF- and c-Fms-deficient mice.¹⁵ M-CSF levels are elevated in the synovial fluid and serum of RA patients,^{71,103} and administration of exogenous M-CSF to mice exacerbates submaximal CIA.⁹ Conversely, M-CSF-deficient mice are resistant to the development of CIA, and neutralizing antibodies against M-CSF or c-Fms attenuate mouse CIA.^{9,52} Several small-molecule inhibitors of c-Fms have been developed and tested in models of RA. In parallel experiments, the c-Fms-specific inhibitor GW2580 was shown to be as efficacious as imatinib in attenuating inflammatory arthritis in antibody-mediated and T-cell-mediated mouse models of RA.⁷¹ In these models, prophylactic, oral administration of GW2580 reduced synovitis, pannus formation, and cartilage and bone erosion; GW2580 was also able to treat established arthritis. The amelioration of arthritis was associated with reduced macrophage infiltration and c-Fms expression in the synovial joints. *In vitro*, GW2580 inhibited the differentiation of monocytes into macrophages and osteoclasts; the resorption of bone by osteoclasts; and the priming of TNF production in FcR-stimulated macrophages.⁷¹ Thus, c-Fms inhibitors may have potential in the treatment of RA through the mitigation of the non-antigen-specific processes that underpin the chronic inflammatory stage of RA. GW2580 has also been shown to attenuate tissue and bone destruction in the joints of rats with AIA, though no effects on joint inflammation were detected in this model.¹³ Two other orally bioavailable c-Fms inhibitors, Ki20027 and cyanopyrrole 8, have been shown to reduce joint inflammation and bone destruction in rodent models of RA, but these compounds are less selective than GW2580.^{45,67} Tested against a panel of 179 kinases, GW2580 proved relatively selective, inhibiting only c-Fms (IC₅₀ of 0.03 μM) and TrkA (IC₅₀ of 0.88 μM).¹³ The restriction of c-Fms expression to monocyte-lineage cells might mean that c-Fms inhibitors would be relatively safe and well tolerated. Nevertheless, elevations in levels of liver enzymes in arthritic mice treated with GW2580, though not associated with histological evidence of pathology, could indicate potential toxicities of GW2580.¹³

Although PDGFR and c-Kit have been implicated in RA, small-molecule inhibitors that selectively inhibit either one of these kinases are not currently available. PDGFR is a ubiquitous tyrosine kinase that plays a key role in fibroblast proliferation, and imatinib has been shown to inhibit PDGFR-mediated proliferation of FLS derived from arthritic mice or from RA patients.^{72,80} Therefore, PDGFR is thought to contribute to RA pathogenesis by promoting synovial hyperplasia and thus pannus formation. c-Kit, on the other hand, has been proposed to contribute by mediating the aberrant activation of mast cells. c-Kit is essential for the survival and activation of mast cells, and release of proinflammatory mediators from synovial mast

cells precedes the onset of clinical signs of inflammation in certain antibody-mediated models of RA.⁵⁷ However, the importance of mast cells in autoimmune arthritis is contentious. In initial studies, mouse strains deficient in mast cells—owing to either a loss-of-function mutation in the gene encoding the c-Kit ligand (*Kit^{Sl}/Kit^{Sl-d}* mice) or a mutation in *c-Kit* (*Kit^WKit^{W-v}* mice)—were shown to be resistant to arthritis induced by K/BxN serum transfer; moreover, engraftment of mast cells restored susceptibility to arthritis in these mice.⁵⁷ These findings cast mast cells as the cellular link between autoantibodies and arthritis. Subsequent studies, however, showed that *Kit^{W-sh}* mice, which are mast-cell-deficient owing to a mutation that abrogates *c-Kit* expression specifically in mast cells, develop full-blown CAIA.¹⁰⁴ Thus, c-Kit may contribute to RA through effects in a cell type other than the mast cell. To date, the most potent and specific small-molecule inhibitor of c-Kit is masitinib (AB1010), with an IC₅₀ of 200 nM for inhibition of recombinant c-Kit.²⁴ However, masitinib also inhibits PDGFR and LynB at nanomolar concentrations—though, unlike imatinib, it is a weak inhibitor of c-Fms and Abl. In a small, open-label, dose-ranging, 12-week, phase II trial in RA patients, masitinib exhibited only moderate efficacy (in the absence of a placebo control).⁹³ Furthermore, patient withdrawal rate was high, owing to adverse effects. Thus, whether inhibiting c-Kit or PDGFR would be of therapeutic value in RA is currently unclear.

Another interesting kinase is Bruton's tyrosine kinase (BTK). It is expressed primarily in B cells, mast cells, platelets, and myeloid cells.⁷⁶ Mutations in the *BTK* gene result in X-linked agammaglobulinaemia (XLA), a disease characterized by marked reduction in numbers of mature B cells and by severe immunodeficiency. BTK transduces BCR signaling in B cells, FcεR1 signaling in mast cells, and toll-like receptor (TLR) signaling in monocytes. Monocytes from XLA patients exhibit defective TNF production in response to TLR stimulation, while BTK-deficient mast cells exhibit impairment of degranulation, histamine release, and cytokine production.⁷⁶ A relatively selective BTK inhibitor, compound 4 was shown to be efficacious in an LPS-induced mouse model of RA—but its therapeutic use may be limited because it is an irreversible inhibitor.⁷⁰⁻⁷⁶ Cgi1746, a reversible orally bioavailable BTK inhibitor with good selectivity, showed efficacy in mouse CIA.⁷⁶ In addition, the rationally designed BTK inhibitor LFM-A13—an analog of a metabolite of the drug leflunomide that is used to treat RA—has been shown to suppress FcεRI-induced release of histamine from rat mast cells.⁴¹ Encouragingly, preclinical studies have demonstrated favorable pharmacokinetic and toxicity profiles of LFM-A13 in mice, rats, and dogs.⁹⁸

The tyrosine kinase VEGFR has also been implicated in RA and is reviewed elsewhere.¹⁴ However, therapeutic targeting of VEGFR may be associated with cardiotoxicity and hypertension,²⁹ which may be of particular concern in a disease such as RA that is often accompanied by cardiovascular dysfunction.

Inhibitor of κB kinase 2 (IKK2): resurgence of an old favorite

The NF-κB pathway is considered the master regulator of inflammation and immunity. It plays a pivotal role in inflammatory and autoimmune diseases—and no less so in RA. Interestingly, several drugs used in the treatment of RA, including sulfasalazine, glucocorticoids, leflunomide, and gold compounds, can inhibit NF-κB. NF-κB is intimately involved in the autoimmune, inflammatory, and destructive processes that underlie RA.⁸⁹ It promotes (i) proliferation of T cells, by inducing the expression of IL-2; (ii) antibody production and class switching in B cells; (iii) recruitment of inflammatory cells, by inducing the expression of adhesion molecules and chemokines; (iv) production of proinflammatory cytokines by multiple cell types; and (v) synovial hyperplasia, by driving angiogenesis and FLS proliferation and survival. In addition, NF-κB directly promotes erosion of cartilage and bone by three different mechanisms: it induces the expression of the matrix-degrading MMPs, it mediates the survival and differentiation of bone-resorbing osteoclasts, and it inhibits the formation of bone-forming

osteoblasts. Underscoring the importance of NF- κ B in inflammatory arthritis, mice deficient in the p50 or c-rel NF- κ B subunits are resistant to the development of CIA,⁸ as are transgenic mice overexpressing a super-repressor form of the NF- κ B inhibitor I κ B α in the T-cell lineage. 85

The NF- κ B transcription factor is regulated by the upstream IKK complex, consisting of the kinases IKK1 and IKK2 and the regulatory component NF- κ B essential modulator (NEMO). IKK2 is the kinase that plays the dominant role in activation of the canonical, proinflammatory NF- κ B pathway, and thus selective inhibition of IKK2 has been explored as an anti-inflammatory therapeutic approach. Numerous orally bioavailable, small-molecule inhibitors of IKK2 have been shown to profoundly suppress both the development and the progression of inflammatory arthritis in rodent models of RA.^{34,38,61,62,69,74,84} Confirming that targeting of IKK2 underlies these effects, intra-articular gene transfer of a dominant-negative form of IKK2 was shown to attenuate rat AIA.⁹² While the importance of NF- κ B in inflammation and immunity has long been recognized, NF- κ B inhibitors have yet to make it into the clinic. The reason for this is that NF- κ B is also important in normal physiology—chronically shutting down NF- κ B is expected to incur a number of serious adverse effects, including tissue injury due to generalized apoptosis and increased susceptibility to infection. Nevertheless, recent findings suggest that IKK/NF- κ B inhibitors may have better prospects than once thought. For instance, although NF- κ B is indispensable for liver development in the fetus, it appears that inhibition of NF- κ B in the developed liver is not hepatotoxic, and may even be hepatoprotective.⁸⁹ Moreover, approaches allowing partial suppression of NF- κ B activity are starting to yield promising results.

One such approach is the use of a cell-permeable peptide corresponding to the NEMO binding domain (NBD) of IKK2 to disrupt the interaction of IKK2 with NEMO, thereby blocking the formation of the IKK complex. The NBD peptide has been shown to inhibit LPS-induced osteoclastogenesis both *in vitro* and *in vivo* and to suppress inflammation and bone destruction in the joints of mice with CIA—without inducing any overt toxicity.⁴⁶ This favorable therapeutic index has been ascribed to the abrogation of inflammation-induced, but not basal, NF- κ B activity. Consistent with this, the NBD peptide reduced serum and joint levels of TNF, IL-1 β , and MMP-9 in CIA mice to those seen in naïve mice. Because peptide therapeutics are beset by a number of disadvantages, the full potential of this approach may not be realized until compounds that mimic the effect of the NBD peptide are developed. Another approach to the partial inhibition of NF- κ B involves simply administering submaximal doses of a small-molecule inhibitor of IKK2 activity. Oral, prophylactic administration of the IKK2 inhibitor BMS-066 was recently shown to protect against the development of rat AIA and mouse CIA at doses that only partially and transiently inhibit NF- κ B activity.³⁴ The observed protection was attributed to the cumulative effect of partial inhibition of multiple NF- κ B-dependent pathogenic processes. Thus, the scope of NF- κ B's role in immunity and inflammation, once thought to preclude the therapeutic targeting of the NF- κ B pathway, may be turned to advantage. Dampening, rather than completely blocking, IKK-NF- κ B signaling seems to be the way to go.

Conclusions

The success of small-molecule kinase inhibitors in the treatment of cancer has spurred efforts to identify kinase targets for the treatment of RA. Many kinases have been convincingly implicated in the pathogenesis of RA, and many kinase inhibitors have proven efficacious in the treatment of inflammatory arthritis in animals. However, few kinase inhibitors have so far made it into clinical development, let alone survived the rigors of a phase II RA clinical trial. This is partly because the therapeutic index of a therapy needs to be higher for a chronic inflammatory disorder such as RA than for cancer. The kinase inhibitors approved for the

treatment of cancer are not very selective, and inhibition of multiple kinases heightens the risk of adverse effects. In selecting suitable therapeutic targets for chronic diseases, not only must potential off-target effects be minimized, but also target-based toxicities must be rigorously scrutinized. For instance, the experience with p38 α inhibitors highlights the importance of appraising the potential effects of kinase inhibition on feedback loops. Furthermore, the identification of several commonly targeted kinases as important regulators of cardiac function underscores the need for careful selection of kinase targets to preclude cardiotoxicity.²⁹ Finally, caution should be exerted in assigning culpability to a specific kinase on the basis of the effects of small-molecule inhibitors, most of which lack specificity.

Despite these hurdles, the treatment of RA with oral kinase inhibitors seems within reach. Attaining the fine line between therapeutic efficacy and toxicity is crucial and tricky; but it may be possible. Unlike cancer, which is frequently driven by mutations in kinases and thus requires treatment with high doses of kinase inhibitors, inflammatory diseases are driven by aberrant activation of wild-type kinases, against which low doses of inhibitor may be effective. Lower doses of kinase inhibitors should result in greater selectivity and reduced toxicity. Moreover, as recently illustrated for IKK, inhibition of an essential kinase—if not absolute—may be tolerated. Such partial sparing of target kinase activity may well underlie the tolerability of many of the kinase inhibitors tested and should perhaps be an overt goal in the development of new kinase inhibitors. Emergent kinome-profiling technologies are expected to facilitate both the discovery of additional kinases involved in RA and the development of more-selective kinase inhibitors. Greater specificity may also be achieved by targeting substrate-specific docking sites on kinases, rather than the highly conserved ATP-binding sites, as illustrated by the pepJIP1 and its small-molecule mimic, T-5224. Finally, the burgeoning efforts at biomarker discovery in RA may one day mean that even those kinase inhibitors now relegated to the scrap heap can be used as effective and safe therapy in specific patient subsets.

Acknowledgments

This work was supported by NIH NIAMS R01 AR-054822, an American College of Rheumatology Within-Our-Reach grant, and Veterans Affairs Health Care System funding awarded to W.H.R.

References

1. Aikawa Y, Morimoto K, Yamamoto T, et al. Treatment of arthritis with a selective inhibitor of c-Fos/activator protein-1. *Nat Biotechnol* 2008;26:817. [PubMed: 18587386]
2. Ananieva O, Darragh J, Johansen C, et al. The kinases MSK1 and MSK2 act as negative regulators of Toll-like receptor signaling. *Nat Immunol* 2008;9:1028. [PubMed: 18690222]
3. Anderson DR, Hegde S, Reinhard E, et al. Aminocyanopyridine inhibitors of mitogen activated protein kinase-activated protein kinase 2 (MK-2). *Bioorg Med Chem Lett* 2005;15:1587. [PubMed: 15745802]
4. Bain J, McLauchlan H, Elliott M, et al. The specificities of protein kinase inhibitors: an update. *Biochem J* 2003;371:199. [PubMed: 12534346]
5. Bajpai M, Chopra P, Dastidar SG, et al. Spleen tyrosine kinase: a novel target for therapeutic intervention of rheumatoid arthritis. *Expert Opin Investig Drugs* 2008;17:641.
6. Bohm C, Hayer S, Kilian A, et al. The alpha-isoform of p38 MAPK specifically regulates arthritic bone loss. *J Immunol* 2009;183:5938. [PubMed: 19828631]
7. Braselmann S, Taylor V, Zhao H, et al. R406, an orally available spleen tyrosine kinase inhibitor blocks fc receptor signaling and reduces immune complex-mediated inflammation. *J Pharmacol Exp Ther* 2006;319:998. [PubMed: 16946104]
8. Campbell IK, Gerondakis S, O'Donnell K, et al. Distinct roles for the NF-kappaB1 (p50) and c-Rel transcription factors in inflammatory arthritis. *J Clin Invest* 2000;105:1799. [PubMed: 10862795]
9. Campbell IK, Rich MJ, Bischof RJ, et al. The colony-stimulating factors and collagen-induced arthritis: exacerbation of disease by M-CSF and G-CSF and requirement for endogenous M-CSF. *J Leukoc Biol* 2000;68:144. [PubMed: 10914502]

10. Camps M, Ruckle T, Ji H, et al. Blockade of PI3Kgamma suppresses joint inflammation and damage in mouse models of rheumatoid arthritis. *Nat Med* 2005;11:936. [PubMed: 16127437]
11. Cha HS, Boyle DL, Inoue T, et al. A novel spleen tyrosine kinase inhibitor blocks c-Jun N-terminal kinase-mediated gene expression in synoviocytes. *J Pharmacol Exp Ther* 2006;317:571. [PubMed: 16452391]
12. Cohen SB, Cheng TT, Chindalore V, et al. Evaluation of the efficacy and safety of pamapimod, a p38 MAP kinase inhibitor, in a double-blind, methotrexate-controlled study of patients with active rheumatoid arthritis. *Arthritis Rheum* 2009;60:335. [PubMed: 19180516]
13. Conway JG, Pink H, Bergquist ML, et al. Effects of the cFMS kinase inhibitor 5-(3-methoxy-4-((4-methoxybenzyl)oxy)benzyl)pyrimidine-2,4-diamine (GW2580) in normal and arthritic rats. *J Pharmacol Exp Ther* 2008;326:41. [PubMed: 18434589]
14. D'Aura, Swanson C.; Paniagua, RT.; Lindstrom, TM., et al. Tyrosine kinases as targets for the treatment of rheumatoid arthritis. *Nat Rev Rheumatol* 2009;5:317. [PubMed: 19491913]
15. Dai XM, Ryan GR, Hapel AJ, et al. Targeted disruption of the mouse colony-stimulating factor 1 receptor gene results in osteopetrosis, mononuclear phagocyte deficiency, increased primitive progenitor cell frequencies, and reproductive defects. *Blood* 2002;99:111. [PubMed: 11756160]
16. Damjanov N, Kauffman RS, Spencer-Green GT. Efficacy, pharmacodynamics, and safety of VX-702, a novel p38 MAPK inhibitor, in rheumatoid arthritis: results of two randomized, double-blind, placebo-controlled clinical studies. *Arthritis Rheum* 2009;60:1232. [PubMed: 19404957]
17. Darragh J, Ananieva O, Courtney A, et al. MSK1 regulates the transcription of IL-1ra in response to TLR activation in macrophages. *Biochem J* 425:595. [PubMed: 19922413]
18. Das S, Cho J, Lambert I, et al. Tpl2/cot signals activate ERK, JNK, and NF-kappaB in a cell-type and stimulus-specific manner. *J Biol Chem* 2005;280:23748. [PubMed: 15833743]
19. Demetri GD, von Mehren M, Blanke CD, et al. Efficacy and safety of imatinib mesylate in advanced gastrointestinal stromal tumors. *N Engl J Med* 2002;347:472. [PubMed: 12181401]
20. Deng C, Kaplan MJ, Yang J, et al. Decreased Ras-mitogen-activated protein kinase signaling may cause DNA hypomethylation in T lymphocytes from lupus patients. *Arthritis Rheum* 2001;44:397. [PubMed: 11229472]
21. Deng C, Lu Q, Zhang Z, et al. Hydralazine may induce autoimmunity by inhibiting extracellular signal-regulated kinase pathway signaling. *Arthritis Rheum* 2003;48:746. [PubMed: 12632429]
22. Dong C, Yang DD, Wysk M, et al. Defective T cell differentiation in the absence of Jnk1. *Science* 1998;282:2092. [PubMed: 9851932]
23. Druker BJ, Talpaz M, Resta DJ, et al. Efficacy and safety of a specific inhibitor of the BCR-ABL tyrosine kinase in chronic myeloid leukemia. *N Engl J Med* 2001;344:1031. [PubMed: 11287972]
24. Dubreuil P, Letard S, Ciufolini M, et al. Masitinib (AB1010), a potent and selective tyrosine kinase inhibitor targeting KIT. *PLoS One* 2009;4:e7258. [PubMed: 19789626]
25. Dumitru CD, Ceci JD, Tsatsanis C, et al. TNF-alpha induction by LPS is regulated posttranscriptionally via a Tpl2/ERK-dependent pathway. *Cell* 2000;103:1071. [PubMed: 11163183]
26. Eklund KK, Joensuu H. Treatment of rheumatoid arthritis with imatinib mesylate: clinical improvement in three refractory cases. *Ann Med* 2003;35:362. [PubMed: 12952023]
27. Eklund KK, Lindstedt K, Sandler C, et al. Maintained efficacy of the tyrosine kinase inhibitor imatinib mesylate in a patient with rheumatoid arthritis. *J Clin Rheumatol* 2008;14:294. [PubMed: 18824923]
28. Eliopoulos AG, Wang CC, Dumitru CD, et al. Tpl2 transduces CD40 and TNF signals that activate ERK and regulates IgE induction by CD40. *EMBO J* 2003;22:3855. [PubMed: 12881420]
29. Force T, Krause DS, Van Etten RA. Molecular mechanisms of cardiotoxicity of tyrosine kinase inhibition. *Nat Rev Cancer* 2007;7:332. [PubMed: 17457301]
30. Gaestel M, Kotlyarov A, Kracht M. Targeting innate immunity protein kinase signalling in inflammation. *Nat Rev Drug Discov* 2009;8:480. [PubMed: 19483709]
31. Gaillard P, Jeanclaude-Etter I, Ardisson V, et al. Design and synthesis of the first generation of novel potent, selective, and in vivo active (benzothiazol-2-yl)acetonitrile inhibitors of the c-Jun N-terminal kinase. *J Med Chem* 2005;48:4596. [PubMed: 15999997]
32. Genovese MC. Inhibition of p38: has the fat lady sung? *Arthritis Rheum* 2009;60:317. [PubMed: 19180514]

33. Ghoreschi K, Laurence A, O'Shea JJ. Janus kinases in immune cell signaling. *Immunol Rev* 2009;228:273. [PubMed: 19290934]
34. Gillooly KM, Pattoli MA, Taylor TL, et al. Periodic, partial inhibition of IkappaB Kinase beta-mediated signaling yields therapeutic benefit in preclinical models of rheumatoid arthritis. *J Pharmacol Exp Ther* 2009;331:349. [PubMed: 19652024]
35. Green N, Hu Y, Janz K, et al. Inhibitors of tumor progression loci-2 (Tpl2) kinase and tumor necrosis factor alpha (TNF-alpha) production: selectivity and in vivo antiinflammatory activity of novel 8-substituted-4-anilino-6-aminoquinoline-3-carbonitriles. *J Med Chem* 2007;50:4728. [PubMed: 17715908]
36. Gupta S, Barrett T, Whitmarsh AJ, et al. Selective interaction of JNK protein kinase isoforms with transcription factors. *EMBO J* 1996;15:2760. [PubMed: 8654373]
37. Hall JP, Kurdi Y, Hsu S, et al. Pharmacologic inhibition of tpl2 blocks inflammatory responses in primary human monocytes, synoviocytes, and blood. *J Biol Chem* 2007;282:33295. [PubMed: 17848581]
38. Hammaker D, Sweeney S, Firestein GS. Signal transduction networks in rheumatoid arthritis. *Ann Rheum Dis* 2003;62:ii86. [PubMed: 14532158]
39. Han Z, Boyle DL, Chang L, et al. c-Jun N-terminal kinase is required for metalloproteinase expression and joint destruction in inflammatory arthritis. *J Clin Invest* 2001;108:73. [PubMed: 11435459]
40. Hegen M, Gaestel M, Nickerson-Nutter CL, et al. MAPKAP kinase 2-deficient mice are resistant to collagen-induced arthritis. *J Immunol* 2006;177:1913. [PubMed: 16849504]
41. Heinonen JE, Smith CI, Nore BF. Silencing of Bruton's tyrosine kinase (Btk) using short interfering RNA duplexes (siRNA). *FEBS Lett* 2002;527:274. [PubMed: 12220673]
42. Heinrichsdorff J, Luedde T, Perdiguero E, et al. p38 alpha MAPK inhibits JNK activation and collaborates with IkappaB kinase 2 to prevent endotoxin-induced liver failure. *EMBO Rep* 2008;9:1048. [PubMed: 18704119]
43. Heo YS, Kim SK, Seo CI, et al. Structural basis for the selective inhibition of JNK1 by the scaffolding protein JIP1 and SP600125. *EMBO J* 2004;23:2185. [PubMed: 15141161]
44. Hueber AJ, McInnes IB. Is spleen tyrosine kinase inhibition an effective therapy for patients with RA? *Nat Clin Pract Rheumatol* 2009;5:130. [PubMed: 19252517]
45. Illig CR, Chen J, Wall MJ, et al. Discovery of novel FMS kinase inhibitors as anti-inflammatory agents. *Bioorg Med Chem Lett* 2008;18:1642. [PubMed: 18242992]
46. Jimi E, Aoki K, Saito H, et al. Selective inhibition of NF-kappa B blocks osteoclastogenesis and prevents inflammatory bone destruction in vivo. *Nat Med* 2004;10:617. [PubMed: 15156202]
47. Karaman MW, Herrgard S, Treiber DK, et al. A quantitative analysis of kinase inhibitor selectivity. *Nat Biotechnol* 2008;26:127. [PubMed: 18183025]
48. Kawamura M, McVicar DW, Johnston JA, et al. Molecular cloning of L-JAK, a Janus family protein-tyrosine kinase expressed in natural killer cells and activated leukocytes. *Proc Natl Acad Sci U S A* 1994;91:6374. [PubMed: 8022790]
49. Keminer O, Kraemer J, Kahmann J, et al. Novel MK2 inhibitors by fragment screening. *Comb Chem High Throughput Screen* 2009;12:697. [PubMed: 19531016]
50. Kerkela R, Grazette L, Yacobi R, et al. Cardiotoxicity of the cancer therapeutic agent imatinib mesylate. *Nat Med* 2006;12:908. [PubMed: 16862153]
51. Kim C, Sano Y, Todorova K, et al. The kinase p38 alpha serves cell type-specific inflammatory functions in skin injury and coordinates pro- and anti-inflammatory gene expression. *Nat Immunol* 2008;9:1019. [PubMed: 18677317]
52. Kitaura H, Zhou P, Kim HJ, et al. M-CSF mediates TNF-induced inflammatory osteolysis. *J Clin Invest* 2005;115:3418. [PubMed: 16294221]
53. Koller M, Hayer S, Redlich K, et al. JNK1 is not essential for TNF-mediated joint disease. *Arthritis Res Ther* 2005;7:R166. [PubMed: 15642137]
54. Kontoyiannis D, Boulougouris G, Manoloukos M, et al. Genetic dissection of the cellular pathways and signaling mechanisms in modeled tumor necrosis factor-induced Crohn's-like inflammatory bowel disease. *J Exp Med* 2002;196:1563. [PubMed: 12486099]

55. Kremer JM, Bloom BJ, Breedveld FC, et al. The safety and efficacy of a JAK inhibitor in patients with active rheumatoid arthritis: Results of a double-blind, placebo-controlled phase IIa trial of three dosage levels of CP-690,550 versus placebo. *Arthritis Rheum* 2009;60:1895. [PubMed: 19565475]
56. Kyriakis JM, Avruch J. Sounding the alarm: protein kinase cascades activated by stress and inflammation. *J Biol Chem* 1996;271:24313. [PubMed: 8798679]
57. Lee DM, Friend DS, Gurish MF, et al. Mast cells: a cellular link between autoantibodies and inflammatory arthritis. *Science* 2002;297:1689. [PubMed: 12215644]
58. Macchi P, Villa A, Giliani S, et al. Mutations of Jak-3 gene in patients with autosomal severe combined immune deficiency (SCID). *Nature* 1995;377:65. [PubMed: 7659163]
59. Martin JH, Mohit AA, Miller CA. Developmental expression in the mouse nervous system of the p493F12 SAP kinase. *Brain Res Mol Brain Res* 1996;35:47. [PubMed: 8717339]
60. Matsubara S, Li G, Takeda K, et al. Inhibition of spleen tyrosine kinase prevents mast cell activation and airway hyperresponsiveness. *Am J Respir Crit Care Med* 2006;173:56. [PubMed: 16192454]
61. Mbalaviele G, Sommers CD, Bonar SL, et al. A novel, highly selective, tight binding I κ B kinase-2 (IKK-2) inhibitor: a tool to correlate IKK-2 activity to the fate and functions of the components of the nuclear factor- κ B pathway in arthritis-relevant cells and animal models. *J Pharmacol Exp Ther* 2009;329:14. [PubMed: 19168710]
62. McIntyre KW, Shuster DJ, Gillooly KM, et al. A highly selective inhibitor of I κ B kinase, BMS-345541, blocks both joint inflammation and destruction in collagen-induced arthritis in mice. *Arthritis Rheum* 2003;48:2652. [PubMed: 13130486]
63. Milici AJ, Kudlacz EM, Audoly L, et al. Cartilage preservation by inhibition of Janus kinase 3 in two rodent models of rheumatoid arthritis. *Arthritis Res Ther* 2008;10:R14. [PubMed: 18234077]
64. Mizuno T, Rothstein TL. B cell receptor (BCR) cross-talk: CD40 engagement enhances BCR-induced ERK activation. *J Immunol* 2005;174:3369. [PubMed: 15749869]
65. Morel J, Audo R, Hahne M, et al. Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) induces rheumatoid arthritis synovial fibroblast proliferation through mitogen-activated protein kinases and phosphatidylinositol 3-kinase/Akt. *J Biol Chem* 2005;280:15709. [PubMed: 15684417]
66. Nosaka T, van Deursen JM, Tripp RA, et al. Defective lymphoid development in mice lacking Jak3. *Science* 1995;270:800. [PubMed: 7481769]
67. Ohno H, Uemura Y, Murooka H, et al. The orally-active and selective c-Fms tyrosine kinase inhibitor Ki20227 inhibits disease progression in a collagen-induced arthritis mouse model. *Eur J Immunol* 2008;38:283. [PubMed: 18085662]
68. Ohori M, Takeuchi M, Maruki R, et al. FR180204, a novel and selective inhibitor of extracellular signal-regulated kinase, ameliorates collagen-induced arthritis in mice. *Naunyn Schmiedebergs Arch Pharmacol* 2007;374:311. [PubMed: 17123065]
69. Okazaki Y, Sawada T, Nagatani K, et al. Effect of nuclear factor- κ B inhibition on rheumatoid fibroblast-like synoviocytes and collagen induced arthritis. *J Rheumatol* 2005;32:1440. [PubMed: 16078317]
70. Pan Z, Scheerens H, Li SJ, et al. Discovery of selective irreversible inhibitors for Bruton's tyrosine kinase. *ChemMedChem* 2007;2:58. [PubMed: 17154430]
71. Paniagua RT, Chang A, Mariano M, et al. c-Fms-mediated differentiation and priming of monocyte lineage cells plays a central role in autoimmune arthritis. *Arthritis Res Ther*. in press.
72. Paniagua RT, Sharpe O, Ho PP, et al. Selective tyrosine kinase inhibition by imatinib mesylate for the treatment of autoimmune arthritis. *J Clin Invest* 2006;116:2633. [PubMed: 16981009]
73. Pine PR, Chang B, Schoettler N, et al. Inflammation and bone erosion are suppressed in models of rheumatoid arthritis following treatment with a novel Syk inhibitor. *Clin Immunol* 2007;124:244. [PubMed: 17537677]
74. Podolin PL, Callahan JF, Bolognese BJ, et al. Attenuation of murine collagen-induced arthritis by a novel, potent, selective small molecule inhibitor of I κ B Kinase 2, TPCA-1 (2-[(aminocarbonyl) amino]-5-(4-fluorophenyl)-3-thiophenecarboxamide), occurs via reduction of proinflammatory cytokines and antigen-induced T cell Proliferation. *J Pharmacol Exp Ther* 2005;312:373. [PubMed: 15316093]
75. Roberts PJ, Der CJ. Targeting the Raf-MEK-ERK mitogen-activated protein kinase cascade for the treatment of cancer. *Oncogene* 2007;26:3291. [PubMed: 17496923]

76. Rokosz LL, Beasley JR, Carroll CD, et al. Kinase inhibitors as drugs for chronic inflammatory and immunological diseases: progress and challenges. *Expert Opin Ther Targets* 2008;12:883. [PubMed: 18554156]
77. Rommel C, Camps M, Ji H. PI3K delta and PI3K gamma: partners in crime in inflammation in rheumatoid arthritis and beyond? *Nat Rev Immunol* 2007;7:191. [PubMed: 17290298]
78. Russell SM, Tayebi N, Nakajima H, et al. Mutation of Jak3 in a patient with SCID: essential role of Jak3 in lymphoid development. *Science* 1995;270:797. [PubMed: 7481768]
79. Sakaguchi N, Takahashi T, Hata H, et al. Altered thymic T-cell selection due to a mutation of the ZAP-70 gene causes autoimmune arthritis in mice. *Nature* 2003;426:454. [PubMed: 14647385]
80. Sandler C, Joutsiniemi S, Lindstedt KA, et al. Imatinib mesylate inhibits platelet derived growth factor stimulated proliferation of rheumatoid synovial fibroblasts. *Biochem Biophys Res Commun* 2006;347:31. [PubMed: 16806061]
81. Sawalha AH, Richardson B. MEK/ERK pathway inhibitors as a treatment for inflammatory arthritis might result in the development of lupus: comment on the article by Thiel et al. *Arthritis Rheum* 2008;58:1203. [PubMed: 18383376]
82. Schett G, Tohidast-Akrad M, Smolen JS, et al. Activation, differential localization, and regulation of the stress-activated protein kinases, extracellular signal-regulated kinase, c-JUN N-terminal kinase, and p38 mitogen-activated protein kinase, in synovial tissue and cells in rheumatoid arthritis. *Arthritis Rheum* 2000;43:2501. [PubMed: 11083274]
83. Schett G, Zwerina J, Firestein G. The p38 mitogen-activated protein kinase (MAPK) pathway in rheumatoid arthritis. *Ann Rheum Dis* 2008;67:909. [PubMed: 17827184]
84. Schopf L, Savinainen A, Anderson K, et al. IKKbeta inhibition protects against bone and cartilage destruction in a rat model of rheumatoid arthritis. *Arthritis Rheum* 2006;54:3163. [PubMed: 17009244]
85. Seetharaman R, Mora AL, Nabozny G, et al. Essential role of T cell NF-kappa B activation in collagen-induced arthritis. *J Immunol* 1999;163:1577. [PubMed: 10415062]
86. She QB, Chen N, Bode AM, et al. Deficiency of c-Jun-NH(2)-terminal kinase-1 in mice enhances skin tumor development by 12-O-tetradecanoylphorbol-13-acetate. *Cancer Res* 2002;62:1343. [PubMed: 11888903]
87. Singh K, Deshpande P, Pryshchep S, et al. ERK-dependent T cell receptor threshold calibration in rheumatoid arthritis. *J Immunol* 2009;183:8258. [PubMed: 20007589]
88. Stebbins JL, De SK, Machleidt T, et al. Identification of a new JNK inhibitor targeting the JNK-JIP interaction site. *Proc Natl Acad Sci U S A* 2008;105:16809. [PubMed: 18922779]
89. Strnad J, Burke JR. IkappaB kinase inhibitors for treating autoimmune and inflammatory disorders: potential and challenges. *Trends Pharmacol Sci* 2007;28:142. [PubMed: 17287032]
90. Sugimoto K, Ohata M, Miyoshi J, et al. A serine/threonine kinase, Cot/Tpl2, modulates bacterial DNA-induced IL-12 production and Th cell differentiation. *J Clin Invest* 2004;114:857. [PubMed: 15372110]
91. Sweeney SE, Firestein GS. Mitogen activated protein kinase inhibitors: where are we now and where are we going? *Ann Rheum Dis* 2006;65:iii83. [PubMed: 17038480]
92. Tak PP, Gerlag DM, Aupperle KR, et al. Inhibitor of nuclear factor kappaB kinase beta is a key regulator of synovial inflammation. *Arthritis Rheum* 2001;44:1897. [PubMed: 11508443]
93. Tebib J, Mariette X, Bourgeois P, et al. Masitinib in the treatment of active rheumatoid arthritis: results of a multicentre, open-label, dose-ranging, phase 2a study. *Arthritis Res Ther* 2009;11:R95. [PubMed: 19549290]
94. Thiel MJ, Schaefer CJ, Lesch ME, et al. Central role of the MEK/ERK MAP kinase pathway in a mouse model of rheumatoid arthritis: potential proinflammatory mechanisms. *Arthritis Rheum* 2007;56:3347. [PubMed: 17907188]
95. Thomis DC, Gurniak CB, Tivol E, et al. Defects in B lymphocyte maturation and T lymphocyte activation in mice lacking Jak3. *Science* 1995;270:794. [PubMed: 7481767]
96. Tong C, Yin Z, Song Z, et al. c-Jun NH2-terminal kinase 1 plays a critical role in intestinal homeostasis and tumor suppression. *Am J Pathol* 2007;171:297. [PubMed: 17591974]
97. Tournier C, Hess P, Yang DD, et al. Requirement of JNK for stress-induced activation of the cytochrome c-mediated death pathway. *Science* 2000;288:870. [PubMed: 10797012]

98. Uckun FM, Tibbles H, Venkatachalam T, et al. Preclinical toxicity and pharmacokinetics of the Bruton's tyrosine kinase-targeting anti-leukemic drug candidate, alpha-cyano-beta-hydroxy-beta-methyl-N- (2,5-dibromophenyl) propenamide (LFM-A13). *Arzneimittelforschung* 2007;57:31. [PubMed: 17341007]
99. Waldburger JM, Firestein GS. Garden of therapeutic delights: new targets in rheumatic diseases. *Arthritis Res Ther* 2009;11:206. [PubMed: 19232066]
100. Weinblatt ME, Kavanaugh A, Burgos-Vargas R, et al. Treatment of rheumatoid arthritis with a Syk kinase inhibitor: a twelve-week, randomized, placebo-controlled trial. *Arthritis Rheum* 2008;58:3309. [PubMed: 18975322]
101. Whitmarsh AJ, Cavanagh J, Tournier C, et al. A mammalian scaffold complex that selectively mediates MAP kinase activation. *Science* 1998;281:1671. [PubMed: 9733513]
102. Winkler, J.; Wright, D.; Pheneger, J., et al. ARRY-162, a potent and selective inhibitor of Mek 1/2. Preclinical and clinical evidence of activity in arthritis; 9th World Congress on Inflammation; 2009.
103. Yang PT, Kasai H, Xiao WG, et al. Increased expression of macrophage colony-stimulating factor in ankylosing spondylitis and rheumatoid arthritis. *Ann Rheum Dis* 2006;65:1671. [PubMed: 17105859]
104. Zhou JS, Xing W, Friend DS, et al. Mast cell deficiency in Kit(W-sh) mice does not impair antibody-mediated arthritis. *J Exp Med* 2007;204:2797. [PubMed: 17998392]

Table 1

Small-molecule kinase inhibitors in RA clinical trials

Kinase target	Inhibitor (company)	Outcome in RA clinical trials
p38α	SCIO-323 (Scios)	clinical development terminated owing to skin toxicity ³²
	AMG-548 (Amgen)	clinical development terminated owing to liver toxicity ³²
	Pamapimod (Roche)	phase II trial comparing efficacy and safety with MTX treatment ¹² ➤ ACR20 response rate was higher in the MTX-treated group than in the pamapimod-treated group
	VX-702 (Vertex)	phase II trial in patients with RA on a background of MTX ¹⁶ ➤ Tolerated but no significant increase over placebo in ACR20 response rate
JAKs	INCB18424 (Incyte)	phase II trial in patients with RA on a background of MTX, SSZ, antimalarials, and/or prednisone* ➤ ACR20, ACR50, ACR70, and ACR90 response rates were 83%, 50%, 25%, and 17%, respectively, in the 15 mg group, compared with 33%, 11%, 0%, and 0% in the placebo group
	CP690550 (Pfizer)	phase II trial in patients with RA refractory to MTX, etanercept, infliximab, or adalimumab ⁵⁵ ➤ ACR20 response rates were 70.5%, 81.2%, and 76.8% in the 5 mg, 15 mg, and 30 mg groups, respectively, compared with 29.2% in the placebo group ➤ ACR50 and ACR70 response rates were significantly greater in all treatment groups compared to the placebo group
Syk	R788 (Rigel)	phase II trial in patients with RA on a background of MTX, SSZ, antimalarials, prednisone and/or NSAIDs ¹⁰⁰ ➤ ACR20, ACR50, and ACR70 response rates were 72%, 57%, and 40%, respectively, in the 150 mg group compared with 38%, 19%, and 4% in the placebo group
MEK 1/2	ARRY-162 (Array BioPharma)	phase II trial in patients with RA refractory to MTX [†] ➤ Tolerated but no significant increase over placebo in ACR20 response
c-Kit/PDGFR	AB1010 (AB Science)	open-label, uncontrolled, phase II trial in patients with RA refractory to MTX or anti-TNF agents ⁹³ ➤ ACR20, ACR50, and ACR70 response rates were 54%, 26%, and 8%, respectively ➤ High rate (37%) of patient withdrawal owing to adverse effects
	Imatinib mesylate (Novartis)	phase II trial in patients with active RA on a background of MTX [§] ➤ Outcomes not reported

* Incyte press release. Incyte's JAK Inhibitor Demonstrates Marked Clinical Benefits in Phase IIa Rheumatoid Arthritis Study. <http://www.incyte.com/Eular%20PR%206%2012%2008%20-%20Final.pdf>. Released June 12, 2008. Accessed January 25, 2010.

† Array BioPharma. Array BioPharma Announces Top-Line Results from Rheumatoid Arthritis Phase 2 Trial. http://www.drugs.com/clinical_trials/array-biopharma-announces-top-line-results-rheumatoid-arthritis-phase-2-trial-7991.html. Announced September 3, 2009. Accessed January 25, 2010.

§ Novartis Pharmaceuticals. A Study of Imatinib 400 mg Once Daily in Combination With Methotrexate in the Treatment of Rheumatoid Arthritis. <http://www.clinicaltrials.gov/ct2/show/NCT00154336?term=imatinib+and+rheumatoid&rank=1>. Last updated November 19, 2009. Accessed January 25, 2010.

JNK, c-Jun terminal kinase; Syk, spleen tyrosine kinase; PDGFR, platelet-derived growth factor; MEK, MAPK-ERK kinase; MTX, methotrexate; TNF, tumor necrosis factor; SSZ, sulfasalazine

Table 2

Small-molecule kinase inhibitors showing treatment benefit in animal models of RA.

Kinase targeted	Small-molecule inhibitor (company)	RA model
MEK 1/2	PD184352 (Pfizer)	mouse CIA; rabbit cytokine-induced cartilage degradation ⁹⁴
	U0126 (DuPont Pharmaceuticals)	mouse SKG ⁸⁷
	ARRY-162* (Array BioPharma)	rat CIA and AIA ¹⁰²
ERK 1/2	FR180204 (Astellas)	mouse CIA ⁶⁸
JNK 1/2/3	SP600125 (Celgene)	rat AIA ³⁹
	AS601245 (EMD Serono)	mouse CIA ³¹
IKK2	BMS-345541 (Bristol-Myers Squibb)	mouse CIA ⁶²
	BMS-066 (Bristol-Myers Squibb)	mouse CIA and rat AIA ³⁴
	TPCA-1 (GlaxoSmithKline)	mouse CIA ⁷⁴
	PHA-408 (Pfizer)	rat SCWA ⁶¹
	ML120B (Millenium Pharmaceuticals)	rat AIA ⁸⁴
	IMD-0560 (IMDD)	mouse CIA ⁶⁹
c-Fms	SPC839 (Celgene)	rat AIA ³⁸
	GW2580 (GlaxoSmithKline)	rat AIA ¹³ ; mouse CIA, CAIA, K/BxN ⁷¹
	Ki20227 (Kirin)	mouse CIA ⁶⁷
BTK	Cyanopyrrole 8 (Johnson & Johnson)	mouse CIA ⁴⁵
	Compound 4 (Celera Genomics)	mouse LPS-induced arthritis ⁷⁰
PI3Kγ	Cgi1746 (CGI Pharmaceuticals)	mouse CIA ⁷⁶
	AS-604850, AS-605240 (EMD Serono)	mouse CAIA ¹⁰

RA, rheumatoid arthritis; CIA, collagen-induced arthritis; CAIA, anti-collagen antibody-induced arthritis; AIA, adjuvant-induced arthritis; SCWA, streptococcal cell wall-induced arthritis; MEK, MAPK-ERK kinase; ERK, extracellular signal-regulated kinase; JNK, c-Jun N-terminal kinase; IKK2, inhibitor of κ B kinase 2; PI3K γ , phosphoinositide-3 kinase γ ; BTK, Bruton's tyrosine kinase; LPS, lipopolysaccharide; IMDD, Institute of Medicinal and Molecular Design.

* In clinical trials in RA patients (see Table 1)