

Exogenous insulin enhances humoral immune responses in short-day, but not long-day, Siberian hamsters (*Phodopus sungorus*)

Nicholas W. Garcia^{1,†}, Timothy J. Greives^{1,‡}, Devin A. Zysling¹,
Susannah S. French², Emily M. Chester¹ and Gregory E. Demas^{1,*}

¹Department of Biology, Program in Neuroscience and Center for the Integrative Study of Animal Behavior, Indiana University, Bloomington, IN 47405, USA

²Department of Biology, Utah State University, Logan, UT 84322, USA

Many animals experience marked seasonal fluctuations in environmental conditions. In response, animals display adaptive alterations in physiology and behaviour, including seasonal changes in immune function. During winter, animals must reallocate finite energy stores from relatively costly, less exigent systems (e.g. reproduction and immunity) to systems critical for immediate survival (e.g. thermoregulation). Seasonal changes in immunity are probably mediated by neuroendocrine factors signalling current energetic state. One potential hormonal candidate is insulin, a metabolic hormone released in response to elevated blood glucose levels. The aim of the present study was to explore the potential role of insulin in signalling energy status to the immune system in a seasonally breeding animal, the Siberian hamster (*Phodopus sungorus*). Specifically, exogenous insulin was administered to male hamsters housed in either long ‘summer-like’ or short ‘winter-like’ days. Animals were then challenged with an innocuous antigen and immune responses were measured. Insulin treatment significantly enhanced humoral immune responses in short, but not long days. In addition, insulin treatment increased food intake and decreased blood glucose levels across photoperiodic treatments. Collectively, these data support the hypothesis that insulin acts as an endocrine signal integrating seasonal energetic changes and immune responses in seasonally breeding rodents.

Keywords: immunity; energy balance; insulin; antibody response

1. INTRODUCTION

Most non-tropical animals experience marked seasonal fluctuations in environmental conditions, which can act as seasonal stressors. During winter, animals typically experience reduced food availability and low ambient temperatures that can limit total energy budgets by decreasing available energy or increasing energy output, respectively. To cope with these potential seasonal stressors, many small mammals have evolved mechanisms to use photoperiodic (day length) cues to appropriately alter energy investment into physiological systems to match current environmental conditions. For example, long-day, reproductively active rodents downregulate their reproductive neuroendocrine axes and cease breeding activity during the short days of winter; reproductive activity is restored following the return of long day lengths associated with spring and summer (Bronson & Heideman 1994). In addition, marked seasonal changes in foraging, food intake and body mass occur in a wide range of vertebrate species (Bartness & Demas 2004).

A growing body of evidence suggests that immunity, like reproduction, is energetically costly and plays a key role in physiological trade-offs (Lochmiller & Deerenberg 2000; Demas 2004; Martin *et al.* 2008). If immune

function is costly, we would predict seasonal changes in immune function in response to changing photoperiods. Empirical support for this hypothesis exists; long-day breeding mammals in seasonal environments modulate immune function during the short days of winter by altering circulating antibody responses, lymphocyte proliferation, delayed-type hypersensitivity, wound healing and numbers of circulating leucocytes and lymphocytes (Yellon *et al.* 1999; Drazen *et al.* 2000; Bilbo *et al.* 2002; Kinsey *et al.* 2003; Weil *et al.* 2007). Despite these observations, considerably less is known regarding the physiological mechanisms mediating photoperiodic changes in immune function. Although changes in gonadal steroid hormones (e.g. testosterone and oestradiol) and pineal melatonin can explain some of the photoperiodic effects on immunity (Demas & Nelson 1998a,b; Bilbo & Nelson 2001; Drazen *et al.* 2001), additional endocrine factors are probably involved, including hormones that regulate energy balance (Demas 2004). For example, leptin, a hormone secreted predominantly by adipose tissue, modulates immune responses in several rodent species (Lord *et al.* 1998; Demas & Sakaria 2005). Furthermore, seasonal changes in leptin concentrations appear to mediate, at least in part, seasonal changes in immunity (Drazen *et al.* 2001). Although leptin increases immune responses in short-day hamsters, antibody responses are not fully restored to normal long-day levels, suggesting that additional endocrine factors are probably involved in mediating this response.

* Author for correspondence (gdemas@indiana.edu).

† Present address: Department of Biology, University of North Carolina, Chapel Hill, NC 27599, USA.

‡ Present address: Max Planck Institute for Ornithology, Radolfzell 78315, Germany.

The pancreatic peptide insulin is secreted in response to increased energy intake and regulates the storage of excess energy. It is well known that insulin signals energy availability to peripheral tissues and the central nervous system (Schwartz & Kahn 1999; Benoit *et al.* 2004). The role of insulin in signalling energy status to the immune system and modulating subsequent responses, however, remains unknown. Because insulin is secreted in response to changes in blood glucose levels, it serves as an important indicator of current energy availability within an animal (Benoit *et al.* 2004). Furthermore, insulin levels are correlated with adiposity in most mammals studied thus far, including Siberian hamsters (*Phodopus sungorus*; Bartness *et al.* 1995). Therefore, insulin may serve as both an indicator of short-term available energy (i.e. glucose) as well as long-term energy stores (i.e. adiposity). Although insulin appears to play an important role in signalling peripheral energy stores, little is known regarding the potential role of this hormone in regulating immune responses. Insulin receptors are expressed by immune cells (e.g. macrophages, B and T lymphocytes), and stimulation of these receptors with exogenous hormone affects glucose transport and can alter immune responses (reviewed in Wolowczuk *et al.* 2008). It has also been suggested that the insulin receptor plays a critical role in the proliferation and survival of immune cells (Knutson 1991). To our knowledge, however, virtually nothing is known regarding the role of insulin in regulating seasonal changes in immune responses.

The aim of the current investigation was to test the hypothesis that insulin plays a role in regulating seasonal changes in immunity. Specifically, if insulin acts as a link between energy balance and immunity, then experimental increases in the insulin signal, indicating an excess of available energy, should enhance immune function and attenuate short-day decreases in immunity. To test this hypothesis, we housed Siberian hamsters in either long or short days, manipulated insulin levels via treatment with exogenous hormone and examined the effects of these manipulations on both innate and acquired immune responses.

2. MATERIAL AND METHODS

(a) *Animals and housing conditions*

Adult (greater than 60 days old) male Siberian hamsters ($n = 50$) were obtained from our breeding colony maintained on long days (L:D 16:8) at Indiana University. Hamsters were individually housed in polypropylene cages ($23.3 \times 15.3 \times 15.9$ cm) for one week prior to the start of the experiment. Food (Purina rat chow, St Louis, MO, USA) and tap water were provided ad libitum throughout the experiment. Room temperature was maintained at $21 \pm 2^\circ\text{C}$ and relative humidity at $50 \pm 10\%$. Animals were then randomly assigned to either long (L:D 16:8) ($n = 18$) or short days (L:D 8:16) ($n = 32$) and initial body mass was measured. All procedures were approved by the Bloomington Institutional Animal Care and Use Committee (BIACUC).

(b) *Responsiveness to photoperiod*

Frequently, a subset of hamsters fails to show photo-responsiveness (i.e. do not display typical gonadal regression, reductions in fat stores or changes in pelage density and coloration) despite prolonged maintenance in short days;

these animals are known as photoperiodic non-responders (Puchalski & Lynch 1986). After eight weeks of exposure to short-day photoperiods, 15 animals were determined to be non-responders (defined by a less than or equal to 10% reduction in body mass and maintenance of summer pelage coloration) and were removed from all subsequent analyses (Greives *et al.* 2008). At the conclusion of the experiment, necropsies were performed on the remaining animals and paired testes were weighed to confirm responsiveness (defined as a paired gonadal mass < 0.15 g; Greives *et al.* 2007). One additional hamster was determined to be a non-responder based on paired testes mass (paired testes mass of 0.63 g) and was removed from subsequent analyses.

(c) *Food intake and insulin administration*

Weekly body masses and daily food intake were measured for all hamsters prior to insulin treatment. Following one week of baseline food and body mass measurements, insulin injections were initiated. Long- and short-day hamsters were divided into one of two injection groups: control animals received 0.1 ml of 0.9 per cent saline vehicle (long day, $n = 9$; short day, $n = 8$), whereas experimental animals received 0.1 ml of 40 U kg^{-1} protamine zinc insulin (PZI) (IDEXX Pharmaceuticals, Greensboro, NC, USA) (long day, $n = 9$; short day, $n = 8$) dissolved in saline. PZI was used because it is a long-lasting form of insulin that has been shown to affect energetic state in rodents, including hamsters. The 40 U kg^{-1} dose was chosen based on a pilot study from our laboratory that revealed lower doses (20 U kg^{-1}) had no effect on blood glucose levels or subsequent immune measures (D. A. Zysling & G. E. Demas 2008, unpublished data), as well as previously published research in this species (Bartness *et al.* 1991). The dose of insulin was gradually increased until over a period of 14 days to 40 U kg^{-1} to avoid severe insulin-induced hypoglycaemia that could result in death (Bartness *et al.* 1991; Bartness & Clein 1994; Demas & Bartness 1999). To this end, daily subcutaneous (s.c.) injections of PZI were administered for 24 days according to the following injection schedule: day 1: 0.625 U kg^{-1} ; days 2–3: 1.25 U kg^{-1} ; day 4–6: 5 U kg^{-1} ; days 7–10: 10 U kg^{-1} ; days 11–14: 20 U kg^{-1} and days 15–24: 40 U kg^{-1} .

(d) *Immunizations*

Fifteen days after the first injection (i.e. once experimental animals had reached the desired 40 U kg^{-1} dose of insulin), all hamsters received a single s.c. injection of 100 μg of the antigen keyhole limpet haemocyanin (KLH) suspended in 0.1 ml sterile saline (day 0; Zysling *et al.* 2006). The animals were previously naive and were returned to the colony room following injections. KLH is an innocuous respiratory protein derived from the giant keyhole limpet (*Megathura crenulata*). KLH was used because it generates a robust antigenic response in rodents, but does not make the animals sick (e.g. prolonged inflammation or fever; Dixon *et al.* 1966). Blood was drawn from the retro-orbital sinus at two different sampling periods (days 5 and 10 post-immunization). These sampling periods were chosen to capture basal (day 5) and peak (day 10) immunoglobulin G (IgG) production (the predominant Ig class present in blood) during the course of the immune response to KLH (Demas *et al.* 1997a; Drazen *et al.* 2000). Day 5 also corresponds to the peak of immunoglobulin M (IgM) production, which is the initial antibody class produced in response to an infection (Demas *et al.*

1997a,b; Drazen *et al.* 2000). On each sampling day, animals were brought into the surgery room, lightly anaesthetized with isoflurane vapours (Aerrane, Henry Schein, Melville, NY, USA), and blood samples were drawn from the retro-orbital sinus between 1000 h and 1200 h EST. Samples were allowed to clot for 1 h, the clots were removed and the samples were centrifuged (at 4°C) for 30 min at 2500 rpm. Serum aliquots were aspirated and stored in sealable polypropylene microcentrifuge tubes at -80°C until assayed for anti-KLH immunoglobulins and leptin.

(e) Blood glucose measurement

Blood glucose levels were measured 10 days after KLH injection (i.e. after 25 days of daily insulin treatment). Blood samples were collected 1 h after insulin or vehicle administration and blood glucose levels were determined using a digital glucose meter (Roche Diagnostics, Indianapolis, IN, USA). Specifically, 5 µl of whole blood was transferred onto the meter's check strips and the readout was recorded. The meter was previously calibrated using an internal standard provided by the manufacturer.

(f) Antibody response to KLH

Serum samples were analysed by enzyme-linked immunosorbent assay (ELISA) to determine IgM and IgG concentrations (Klein-Schneegans *et al.* 1989). Microtiter plates were coated with KLH in sodium bicarbonate buffer (pH 9.6), incubated overnight at 4°C and washed with phosphate buffered saline (PBS) with 0.05 per cent Tween 20 (PBS-T; pH 7.4). Plates were then blocked with 5 per cent non-fat dry milk in PBS and washed again with PBS-T. Serum samples were diluted 1 : 20 with PBS and added to the plate wells in duplicate. Negative control samples (i.e. serum from KLH-naive hamsters) and positive control samples (i.e. serum pooled from hamsters previously shown to have significantly high anti-KLH antibody response) were also diluted 1 : 20 with PBS and added to the plate wells in duplicate. Plates were incubated at 37°C for 3 h and then washed with PBS-T. Secondary antibody was then added to the wells and the plates were incubated for 1.5 h for day 5 (IgM) samples and 1 h for day 10 (IgG) samples. Alkaline phosphatase-conjugated anti-mouse IgM (MP Biomedicals, Salon, OH, USA) diluted 1 : 500 in PBS was used for day 5 samples and alkaline phosphatase-conjugated anti-Syrian hamster IgG (Rockland, Gilbertsville, PA, USA) was diluted 1 : 500 in PBS and used for day 10 samples.

Following incubation, plates were again washed with PBS-T, and 150 µl of the enzyme substrate *p*-nitro-phenyl phosphate (Sigma, St Louis, MO, USA; 1 mg ml⁻¹ in diethanolamine substrate buffer) was added to each well. Absorbance was measured (Bio-Rad Benchmark Plate Reader, Hercules, CA, USA) at 405 nm. The mean for each sample was calculated and expressed as a percentage of the positive control mean (% plate positive).

(g) Serum leptin

Insulin is known to influence serum leptin levels (Saladin *et al.* 1995); therefore, we measured circulating leptin concentrations in the present study. Serum leptin concentrations were determined in day 10 serum samples (1 : 2 dilution with assay buffer) using an ELISA from a commercially prepared murine kit (EZML-82K, Millipore, St Charles, MO, USA). This assay was previously validated for use with Siberian hamsters (French *et al.* 2009). All procedures were followed as directed by the manufacturer's

instructions. The antiserum used was highly specific for leptin; cross-reactivity with related hormones is less than 0.05 per cent and the sensitivity of the assay is 0.05 ng ml. Intra-assay variability was consistently low and coefficients of variance were less than 10 per cent for all samples.

(h) Bactericidal assay

As a functional assessment of an animal's ability to eliminate a bacterial infection, we used an *ex vivo* bactericidal assay, based on a modification (French *et al.* 2009) of a previously published protocol (Matson *et al.* 2006). This assay quantifies the relative number of *Escherichia coli* colony forming units (CFUs) that grow after incubation with serum. Differences in CFU presumably represent differences in the ability of serum proteins (i.e. complement) to kill bacterial colonies. Briefly, *E. coli* (ATCC #8739, Microbiologics, St Cloud, MN, USA) (1 pellet = 10⁷ CFU) was added to 40 ml 1 M sterile PBS warmed to 35–37°C, which was vortexed to create a bacterial stock solution and activated by incubation for 30 min at 37°C. Serum samples were diluted 1 : 40 in glutamine-enriched CO₂-independent media (Invitrogen Corp., Carlsbad, CA, USA). This dilution was validated for serum with a dose-response curve prior to the experiment. The stock bacteria solution (500 000 CFU ml⁻¹) was diluted with sterile 1 M PBS to create a 50 000 CFU ml⁻¹ working solution. To obtain estimates of bacterial numbers (i.e. positive control), the working solution was diluted 1 : 10 with glutamine-enriched CO₂-independent media. For each sample, the working solution was added at a 1 : 10 ratio to the diluted serum sample. The bacteria/serum cocktails were incubated for 30 min at 37°C. All samples were vortexed and 50 µl was added to Petri plates in duplicate and spread with a flame-sterilized spreader. All plates were stored upside down overnight at 37°C. Following incubation, bacteria colonies were counted on each plate, and duplicates were averaged. The mean value for each sample was expressed as a per cent of bacteria killed relative to the control plates, in which no killing occurred.

(i) Necropsies and tissue collection

At the conclusion of the study, hamsters were euthanized via an overdose of ketamine cocktail (20 mg ml⁻¹ ketamine and 4 mg ml⁻¹ xylazine in 0.9% saline solution; Nelson & Demas 1996). Inguinal white adipose tissue (IWAT), retroperitoneal WAT (RWAT), epididymal WAT (EWAT) and paired testes were collected, cleaned of connective tissues and weighed to the nearest 0.01 g.

(j) Statistical analyses

All data were analysed with Minitab 15 (Mintab Inc., State College, PA, USA) using a two-way (photoperiod × treatment) analysis of variance (ANOVA), except for food consumption data, which were analysed by a repeated measures ANOVA using JMP 7.0 (SAS, Cary, NC, USA). Data for repeated measures analysis were taken from the 9-day period that began following the first administration of full-dose insulin. A composite adipose tissue score was calculated for each hamster by summing the individual WAT pad masses (Zysling & Demas 2007).

All pair-wise comparisons of means were conducted using Tukey's honestly significant difference post hoc comparisons. Differences between group means were considered significant at *p* < 0.05. Datasets were analysed for normality using the Anderson-Darling test. Bactericidal assay data did not meet requirements for normality and were




Figure 1. Mean (\pm s.e.m.) final body masses (a) and food intake (b) in Siberian hamsters housed in either long or short days and administered exogenous insulin (striped bar) or vehicle (filled bar) control. Significant differences in food intake are indicated by different letters, whereas significant differences in final body masses are indicated by an asterisk (*) if $p < 0.05$.

transformed by taking the arcsine of the square root of each data point, a standard transformation for percentages.

3. RESULTS

(a) Body and tissue masses and food intake

Short-day hamsters had significantly lower final body masses than long-day animals ($F_{1,32} = 65.24$, $p < 0.001$; figure 1a); however, insulin treatment had no significant effect on body mass ($F_{1,32} = 0.63$, $p > 0.05$). In addition, short-day animals had significantly smaller IWAT ($F_{1,32} = 65.99$, $p < 0.001$), RWAT ($F_{1,32} = 52.87$, $p < 0.001$) and EWAT pad masses ($F_{1,32} = 65.81$, $p < 0.001$), as well as lower composite adipose tissue scores ($F_{1,32} = 77.46$, $p < 0.001$; figure 2a) compared with long-day animals. Insulin treatment had no effect on any IWAT pad masses or composite adipose tissue scores ($p > 0.05$ in all cases). Short-day hamsters had significantly regressed paired testes when compared with long-days animals ($F_{1,32} = 494.05$, $p < 0.001$; figure 2b). Insulin treatment had no effect on paired testes mass ($F_{1,32} = 0.01$, $p > 0.05$).

(b) Food intake

During the initial baseline (i.e. pre-insulin treatment) period, short-day hamsters consumed significantly less food than long-day animals ($F_{1,34} = 6.34$; $p < 0.001$; figure 1b). During the experimental treatment period, both photoperiod ($F_{1,30} = 21.43$, $p < 0.001$; figure 1b)




Figure 2. Mean (\pm s.e.m.) composite white adipose tissue masses (a) and paired testes masses (b) in Siberian hamsters housed in either long or short days and administered exogenous insulin (striped bar) or vehicle (filled bar) control. Significant differences between pair-wise means are indicated by an asterisk (*) if $p < 0.05$.

and insulin treatment ($F_{1,30} = 9.89$, $p < 0.001$) significantly affected food consumption following KLH exposure; short-day animals consumed less food than long-day animals, and animals receiving insulin injections consumed more when compared with vehicle-injected animals. There was no significant interaction between the two treatments ($F_{1,30} = 0.31$, $p > 0.05$).

(c) Blood glucose levels

Blood glucose levels were significantly affected by photoperiod treatment ($F_{1,32} = 8.08$, $p = 0.008$; figure 3a); blood glucose levels were significantly lower in short-day hamsters compared with long-day hamsters. Insulin-treated short-day hamsters had significantly lower blood glucose levels when compared with insulin-treated long-day hamsters ($p = 0.029$); however, there was no difference between long- and short-day hamsters treated with saline ($p > 0.05$). There was also a significant main effect of hormone treatment on glucose with lower blood glucose levels in hamsters receiving insulin when compared with hamsters injected with saline ($F_{1,32} = 29.82$, $p < 0.001$; figure 3a). Specifically, blood glucose levels were lower in long-day hamsters treated with insulin when compared with saline-injected long-day hamsters ($p = 0.027$).

(d) Immune responses

Insulin-treated hamsters displayed significantly elevated IgG levels when compared with animals injected with




Figure 3. Mean (\pm s.e.m.) blood glucose (a) and serum leptin concentrations (b) in Siberian hamsters housed in either long or short days and administered exogenous insulin (striped bar) or vehicle (filled bar) control. Significant differences in blood glucose are indicated by different letters, whereas significant differences in serum leptin concentration are indicated by an asterisk (*) if $p < 0.05$.

saline ($F_{1,32} = 7.67$, $p = 0.010$; figure 4b). Specifically, short-day animals treated with insulin showed significantly greater IgG responses with compared with short-day saline-treated animals ($p = 0.044$). A similar, but non-significant trend, was found for serum anti-KLH IgM levels ($F_{1,31} = 3.28$, $p = 0.081$; figure 4a). Blood glucose levels were negatively correlated with anti-KLH IgG (figure 5a) and anti-KLH IgM per cent plate positive values ($R^2 = 0.187$, $p = 0.012$; $R^2 = 0.140$, $p = 0.035$). There was no effect of photoperiod on either serum anti-KLH IgG levels ($p > 0.05$), or serum anti-KLH IgM levels ($p > 0.05$). There was no significant interaction between photoperiod and insulin treatment on either serum anti-KLH IgG levels ($p > 0.05$), or serum anti-KLH IgM levels ($p > 0.05$). Neither photoperiod ($F_{1,65} = 1.30$, $p > 0.05$) nor insulin treatment ($F_{1,65} = 2.77$, $p > 0.05$) had a significant effect on serum bacterial killing ability (figure 5).

(e) Serum leptin concentrations

Serum leptin concentrations were significantly lower in short-day animals when compared with long-day animals ($F_{1,26} = 23.47$, $p < 0.001$; figure 3b); however, there were no significant effects of insulin treatment ($F_{1,26} = 1.12$, $p > 0.05$), or an interaction between insulin and photoperiod treatments ($F_{1,26} = 0.10$, $p > 0.05$; figure 3b). All other pair-wise comparisons were not statistically




Figure 4. Mean (\pm s.e.m.) serum anti-KLH IgM (a) and IgG (b) levels in Siberian hamsters housed in either long or short days and administered exogenous insulin (striped bar) or vehicle (filled bar) control. Significant differences between pair-wise means are indicated by an asterisk (*) if $p < 0.05$.

significant ($p > 0.05$ in all cases). Serum leptin levels were correlated with adipose tissue composite scores ($R^2 = 0.435$, $p < 0.001$), but not day 10 blood glucose levels ($R^2 = 0.009$, $p > 0.05$) or anti-KLH IgG plate positive values ($R^2 = 0.001$, $p > 0.05$).

4. DISCUSSION

In the current study, we investigated a potential role for the pancreatic peptide insulin in mediating seasonal energetic trade-offs with immunity in Siberian hamsters. Specifically, we asked: (i) whether exogenous administration of insulin alters humoral and innate immune responses and (ii) whether this effect is photoperiod-dependent. As expected, short-day animals displayed gonadal regression and marked reductions in total body fat relative to long-day hamsters. Exogenous insulin increased KLH-specific IgG production, but only in short-day hamsters; insulin did not affect humoral immunity in long-day animals. Anti-KLH IgM followed a similar, non-significant, trend. In contrast, insulin had no effect on innate immunity (i.e. bacterial killing ability) regardless of photoperiodic condition. Serum leptin was reduced in short-day animals, but was not altered in response to insulin treatment. The findings from the present study support the idea that insulin can influence immune responses in hamsters and the effects of insulin on immunity can vary according to photoperiodic status. In addition, these results suggest that insulin acts as a peripheral signal of energy availability and probably mediates, at least in part, energetic trade-offs




Figure 5. (a) Pearson product-moment correlation between blood glucose levels (day 10 sample) and IgG ($R^2 = 0.187$, $p = 0.012$). Mean (\pm s.e.m.) bacterial killing ability (b) in Siberian hamsters housed in either long or short days and given exogenous administration of insulin (striped bar) or vehicle (filled bar) control.

between immunity and other physiological systems (e.g. reproduction).

Blood glucose levels were significantly lower 1 h after insulin injection, suggesting that hamsters are physiologically responsive to hormone. Short-day hamsters appear to be more sensitive to the insulin signal as blood glucose levels in short-day animals were significantly lower than long-day animals given insulin. In contrast, there was no significant difference in glucose levels between long-day and short-day control hamsters. A similar pattern was observed with food intake across experimental treatments. Consistent with previous findings in this species (Bartness *et al.* 1995), long-day insulin-treated hamsters consumed significantly more food than all other groups. In contrast, an increase in humoral immune response was observed in insulin-treated short-day hamsters, but not long-day hamsters. Thus, insulin-induced immunoenhancement was not probably a direct result of increased food intake. These findings suggest that insulin-induced immunoenhancement is not merely a product of abundant energy, but is also dependent on the photoperiodic state of the animal. Previous work in this and other species suggested that short-, but not long-day conditions can buffer immune suppression from an energetic stressor, suggesting long-day hamsters may be more sensitive to energy changes (Demas *et al.* 1997b; Zysling *et al.* 2006). However, where energy abundance is concerned, this may not be the case. Under stressful winter conditions, an animal may stand to

profit from upregulating immunity if it receives a signal of immediate energy.

Changes in insulin levels alter a wide range of metabolic processes, which may alter immune responses. For example, insulin can influence circulating levels of leptin (Saladin *et al.* 1995; Warne *et al.* 2009) and can alter the metabolic actions of adipocytes (Muller *et al.* 1997). Leptin, in turn, can affect humoral immune responses in a variety of mammalian species, including hamsters (Lord *et al.* 1998; Drazen *et al.* 2001). Thus, the effects of insulin on immunity may be indirect, via changes in leptin levels. Serum leptin concentrations were determined in the present study to assess whether the observed effects on immunity are caused by altered leptin levels. Leptin levels differed according to photoperiod treatment and, consistent with previous studies, were significantly lower in short-day hamsters (Drazen *et al.* 2000). Insulin treatment, however, had no effect on serum leptin levels. Because leptin concentrations were lower in short-day hamsters and insulin increased antibody responses in these animals, leptin is not probably playing a role in the photoperiod-dependent effects of insulin on humoral immunity.

Although metabolic factors other than leptin cannot be ruled out, insulin may act directly on peripheral immune tissues to enhance immune responses in short-day hamsters. Like most tissues, peripheral lymphoid tissue possesses insulin receptors (Gavin *et al.* 1973). Interestingly, the insulin receptor is expressed on resting neutrophils, monocytes and B lymphocytes, but not T lymphocytes; activation of T cells, however, results in marked upregulation of insulin receptors on these cells (Viardot *et al.* 2007). Consistent with this idea, lymphocyte proliferation and activation is altered by *in vitro* insulin administration (Viardot *et al.* 2007), suggesting that insulin can act directly on lymphocytes to modulate acquired immune responses.

Alternatively, insulin may act indirectly, via the central nervous system, to affect immunity. Such a mechanism has been suggested as part of the effects of leptin on humoral immunity (Demas 2002). Previous research has demonstrated that central insulin receptor expression is reduced under short-day conditions in the arcuate nucleus (ARC) and short-day animals are presumably less sensitive to the insulin signal (Tups *et al.* 2006). Similar findings of short-day decreases in insulin receptors have been reported in Japanese quail in the infundibular nucleus, a structure homologous to the mammalian ARC (Anraku *et al.* 2007). In contrast to previous findings, short-day hamsters in the present study appear more sensitive to exogenous insulin (e.g. display largest decrease in blood glucose levels and increase in immunoglobulin response), suggesting possible upregulation of insulin receptors, either centrally or in peripheral tissues (e.g. lymphoid tissues). According to a tissue-specific analysis of the insulin receptor expression in the mouse transcriptome, the insulin receptor is a ubiquitous housekeeping gene: there is no significant difference between expression in the hypothalamus compared with lymphoid organs (Su *et al.* 2002). To our knowledge, no studies have examined photoperiodic changes in insulin receptors on lymphoid tissues in this or other seasonally breeding species.

Previous studies in Siberian hamsters have reported a photoperiod-dependent decrease in immune responses. Specifically, short-day hamsters have a lower antibody

response than long-day animals (Drazen *et al.* 2001; Demas *et al.* 2002). In the current study, the expected decrease in immune response in short-day hamsters, however, was not observed; short-day levels of anti-KLH immunoglobulins did not differ between long- and short-day hamsters. This is somewhat surprising given that short-day hamsters exhibited all other typical photoperiod-induced characteristics including decreased body mass, food intake, adipose tissue mass and gonadal regression (Bartness 1996). Although it is not known why immunity was not altered in short-day hamsters, it is possible that daily injections and handling acted as repeated stressors, potentially altering immune responses and masking expected photoperiodic changes in immunity. Male hamsters typically have elevated glucocorticoid levels in short days and exhibited greater immune responses to restraint stress (Bilbo & Nelson 2003). Unlike the effects on antibody levels, exogenous insulin had no effect on innate immunity as assessed by bacterial killing ability. Innate immune responses tend to remain uniform. Innate responses use intrinsic microbial molecules, pathogen-associated molecular patterns (PAMPs), to bind foreign molecules (Medzhitov & Janeway 2002). Alteration in the ability to recognize PAMPs is commonly deleterious to pathogen recognition (Beutler 2004). Therefore, we might not expect to see a difference in innate immunity even under conditions where a signal of additional energy is involved.

Regardless of the precise mechanisms of action, the present results support the idea that insulin alters humoral immune responses, either directly or indirectly, in Siberian hamsters. Further research is required to determine the specific mechanisms mediating this effect. Taken together, these results suggest that insulin serves as an important peripheral signal linking energy availability and immunity. Exogenous insulin can enhance immune responses in short-day animals and, like leptin, can enhance energy availability owing to increased food intake. These findings support the idea that reduced energy availability can lead to changes in immunity which, in turn, can affect disease susceptibility. In addition, these results suggest an important role for insulin in regulating energetic trade-offs among competing physiological systems and provide an important step towards understanding the neuroendocrine mechanisms regulating seasonal changes in mammalian immunity.

This work was reviewed and approved by the Institutional Animal Care and Use Committee at Indiana University. This work was supported by a SICB Grant-in-Aid and NSF International Research Fellowship to T.J.G., NIH Training Grant HD 049336-04 Common Themes in Reproductive Diversity Training grant, NIH to S.S.F. and T.J.G., a fellowship from the Center for the Integrative Study of Animal Behaviour (CISAB) to E.M.C., and a Faculty Research Support (FRSP) grant and NSF grant IOB-0543798 to G.E.D.

REFERENCES

- Anraku, T. *et al.* 2007 Photoperiodic changes in hypothalamic insulin receptor gene expression are regulated by gonadal testosterone. *Brain Res.* **1163**, 86–90. (doi:10.1016/j.brainres.2007.06.028)
- Bartness, T. J. 1996 Photoperiod, sex, gonadal steroids, and housing density affect body fat in hamsters. *Physiol. Behav.* **60**, 517–529. (doi:10.1016/S0031-9384(96)80027-8)
- Bartness, T. J. & Clein, M. R. 1994 Effects of food deprivation and restriction, and metabolic blockers on food hoarding in Siberian hamsters. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **266**, R1111–R1117.
- Bartness, T. J. & Demas, G. E. 2004 Comparative studies of food intake. In *Handbook of neurobiology*, vol. 10, *Neurobiology of food and fluid intake* (eds E. M. Stricker & S. Woods), pp. 423–467. New York, NY: Kluwer Press.
- Bartness, T. J., McGriff, W. R. & Maharaj, M. P. 1991 Effects of diabetes and insulin on photoperiodic responses in Siberian hamsters. *Physiol. Behav.* **49**, 613–620. (doi:10.1016/0031-9384(91)90287-X)
- Bartness, T. J., Morley, J. E. & Levine, A. S. 1995 Effects of food deprivation and metabolic fuel utilization on the photoperiodic control of food intake in Siberian hamsters. *Physiol. Behav.* **57**, 61–68. (doi:10.1016/0031-9384(94)00203-H)
- Benoit, S. C., Clegg, D. J., Seeley, R. J. & Woods, S. C. 2004 Insulin and leptin as adiposity signals. *Recent Prog. Horm. Res.* **59**, 267–285. (doi:10.1210/rp.59.1.267)
- Beutler, B. 2004 Innate immunity: an overview. *Mol. Immunol.* **40**, 845–859. (doi:10.1016/j.molimm.2003.10.005)
- Bilbo, S. D. & Nelson, R. J. 2001 Sex steroid hormones enhance immune function in male and female Siberian hamsters. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **280**, R207–R213.
- Bilbo, S. D. & Nelson, R. J. 2003 Sex differences in photoperiodic and stress-induced enhancement of immune function in Siberian hamsters. *Brain Behav. Immun.* **17**, 462–472. (doi:10.1016/S0889-1591(03)00063-1)
- Bilbo, S. D., Drazen, D. L., Quan, N., He, L. & Nelson, R. J. 2002 Short day lengths attenuate the symptoms of infection in Siberian hamsters. *Proc. R. Soc. Lond. B* **269**, 447–454. (doi:10.1098/rspb.2001.1915)
- Bronson, F. H. & Heideman, P. E. 1994 Seasonal regulation of reproduction in mammals. In *The physiology of reproduction*, vol. 2, 2nd edn (eds E. Knobil & J. D. Neill), pp. 541–584. New York, NY: Raven Press.
- Demas, G. E. 2002 Splenic denervation blocks leptin-induced enhancement of humoral immunity in Siberian hamsters (*Phodopus sungorus*). *Neuroendocrinology* **76**, 178–184. (doi:10.1159/000064527)
- Demas, G. E. 2004 The energetics of immunity: a neuroendocrine link between energy balance and immune function. *Horm. Behav.* **45**, 173–180. (doi:10.1016/j.yhbeh.2003.11.002)
- Demas, G. E. & Bartness, T. J. 1999 Effects of food deprivation and metabolic fuel utilization on food hoarding by jirds (*Meriones shawi*). *Physiol. Behav.* **67**, 243–248. (doi:10.1016/S0031-9384(99)00066-9)
- Demas, G. E. & Nelson, R. J. 1998a Exogenous melatonin enhances cell-mediated, but not humoral, immune function in adult male deer mice (*Peromyscus maniculatus*). *J. Biol. Rhythms* **13**, 245–252. (doi:10.1177/074873098129000084)
- Demas, G. E. & Nelson, R. J. 1998b Short-day enhancement of immune function is independent of steroid hormones in deer mice (*Peromyscus maniculatus*). *J. Comp. Physiol. B* **168**, 419–426.
- Demas, G. E. & Sakaria, S. 2005 Leptin regulates energetic tradeoffs between body fat and humoral immunity. *Proc. R. Soc. B* **272**, 1845–1850. (doi:10.1098/rspb.2005.3126)
- Demas, G. E., Chefer, V., Talan, M. I. & Nelson, R. J. 1997a Metabolic costs of mounting an antigen-stimulated immune response in adult and aged C57BL/6J mice.

- Am. J. Physiol. Regul. Integr. Comp. Physiol.* **42**, R1631–R1637.
- Demas, G. E., DeVries, A. C. & Nelson, R. J. 1997b Effects of photoperiod and 2-deoxy-D-glucose-induced metabolic stress on immune function in female deer mice (*Peromyscus maniculatus*). *Am. J. Physiol.* **272**, R1762–R1767.
- Demas, G. E., Drazen, D. L., Jasnow, A. M., Bartness, T. J. & Nelson, R. J. 2002 Sympathoadrenal system differentially affects photoperiodic changes in humoral immunity of Siberian hamsters (*Phodopus sungorus*). *J. Neuroendocrinol.* **14**, 29–35. (doi:10.1046/j.0007-1331.2001.00736.x)
- Dixon, F. J., Jacot-Guillarmod, H. & McConahey, P. J. 1966 The antibody responses of rabbits and rats to hemocyanin. *J. Immunol.* **97**, 350–355.
- Drazen, D. L., Kriegsfeld, L. J., Schneider, J. E. & Nelson, R. J. 2000 Leptin, but not immune function, is linked to reproductive responsiveness to photoperiod. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **278**, R1401–R1407.
- Drazen, D. L., Demas, G. E. & Nelson, R. J. 2001 Leptin effects on immune function and energy balance are photoperiod dependent in Siberian hamsters (*Phodopus sungorus*). *Endocrinology* **142**, 2768–2775. (doi:10.1210/en.142.7.2768)
- French, S. S., Greives, T. J., Zysling, D. A., Chester, E. M. & Demas, G. E. 2009 Leptin increases maternal investment. *Proc. R. Soc. B* **276**, 4003–4011. (doi:10.1098/rspb.2009.1199)
- Gavin III, J. R., Gorden, P., Roth, J., Archer, J. A. & Buell, D. N. 1973 Characteristics of the human lymphocyte insulin receptor. *J. Biol. Chem.* **248**, 2202–2207.
- Greives, T. J., Mason, A. O., Scotti, M.-A. L., Levine, J., Ketterson, E. D., Kriegsfeld, L. J. & Demas, G. E. 2007 Environmental control of kisspeptin: implications for seasonal reproduction. *Endocrinology* **148**, 1158–1166. (doi:10.1210/en.2006-1249)
- Greives, T. J., Kriegsfeld, L. J. & Demas, G. E. 2008 Exogenous kisspeptin does not alter photoperiod-induced gonadal regression in Siberian hamsters (*Phodopus sungorus*). *Gen. Comp. Endocrinol.* **156**, 552–558. (doi:10.1016/j.ygcen.2008.02.017)
- Kinsey, S. G., Prendergast, B. J. & Nelson, R. J. 2003 Photoperiod and stress affect wound healing in Siberian hamsters. *Physiol. Behav.* **78**, 205–211. (doi:10.1016/S0031-9384(02)00967-8)
- Klein-Schneegans, A. S., Gaveriaux, C., Fonteneau, P. & Loor, F. 1989 Indirect double sandwich ELISA for the specific and quantitative measurement of mouse IgM, IgA and IgG subclasses. *J. Immunol. Methods* **119**, 117–125.
- Knutson, V. 1991 Cellular trafficking and processing of the insulin receptor. *FASEB J.* **5**, 2130–2138.
- Lochmiller, R. L. & Deerenberg, C. 2000 Trade-offs in evolutionary immunology: just what is the cost of immunity? *Oikos* **88**, 87–98. (doi:10.1034/j.1600-0706.2000.880110.x)
- Lord, G. M., Matarese, G., Howard, J. K., Baker, R. J., Bloom, S. R. & Lechler, R. I. 1998 Leptin modulates the T-cell immune response and reverses starvation-induced immunosuppression. *Nature* **394**, 897–901. (doi:10.1038/29795)
- Martin, L. B., Weil, Z. M. & Nelson, R. J. 2008 Seasonal changes in vertebrate immune activity: mediation by physiological trade-offs. *Phil. Trans. R. Soc. B* **363**, 321–339. (doi:10.1098/rstb.2007.2142)
- Matson, K. D., Tieleman, B. I. & Klasing, K. C. 2006 Capture stress and the bactericidal competence of blood and plasma in five species of tropical birds. *Physiol. Biochem. Zool.* **79**, 556–564. (doi:10.1086/501057)
- Medzhitov, R. & Janeway Jr, C. A. 2002 Decoding the patterns of self and nonself by the innate immune system. *Science* **296**, 298–300. (doi:10.1126/science.1068883)
- Muller, G., Ertl, J., Gerl, M. & Preibisch, G. 1997 Leptin impairs metabolic actions of insulin in isolated rat adipocytes. *J. Biol. Chem.* **272**, 10 585–10 593.
- Nelson, R. J. & Demas, G. E. 1996 Seasonal changes in immune function. *Q. Rev. Biol.* **71**, 511–548. (doi:10.1086/419555)
- Puchalski, W. & Lynch, G. R. 1986 Evidence for differences in the circadian organization of hamsters exposed to short day photoperiod. *J. Comp. Physiol. A* **159**, 7–11. (doi:10.1007/BF00612490)
- Saladin, R., De Vos, P., Guerre-Millo, M., Leturque, A., Girard, J., Staels, B. & Auwerx, J. 1995 Transient increase in obese gene expression after food intake or insulin administration. *Nature* **377**, 527–529. (doi:10.1038/377527a0)
- Schwartz, M. W. & Kahn, S. E. 1999 Insulin resistance and obesity. *Nature* **402**, 860–861. (doi:10.1038/47209)
- Su, A. I. et al. 2002 Large-scale analysis of the human and mouse transcriptomes. *Proc. Natl Acad. Sci. USA* **99**, 4465–4470. (doi:10.1073/pnas.012025199)
- Tups, A., Helwig, M., Stohr, S., Barrett, P., Mercer, J. G. & Klingenspor, M. 2006 Photoperiodic regulation of insulin receptor mRNA and intracellular insulin signaling in the arcuate nucleus of the Siberian hamster, *Phodopus sungorus*. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **291**, R643–R650.
- Viardot, A., Grey, S. T., Mackay, F. & Chisholm, D. 2007 Potential antiinflammatory role of insulin via the preferential polarization of effector T cells toward a T helper 2 phenotype. *Endocrinology* **148**, 346–353. (doi:10.1210/en.2006-0686)
- Warne, J. P., Akana, S. F., Ginsberg, A. B., Horneman, H. F., Pecoraro, N. C. & Dallman, M. F. 2009 Disengaging insulin from corticosterone: roles of each on energy intake and disposition. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **296**, R1366–R1375.
- Weil, Z. M., Bowers, S. L. & Nelson, R. J. 2007 Photoperiod alters affective responses in collared lemmings. *Behav. Brain Res.* **179**, 305–309. (doi:10.1016/j.bbr.2007.02.003)
- Wolowczuk, I., Verwaerde, C., Viltart, O., Delanoye, A., Delacre, M., Pot, B. & Grangette, C. 2008 Feeding our immune system: impact on metabolism. *Clin. Dev. Immunol.* **2008**, 639 803.
- Yellon, S. M., Teasley, L. A., Fagoaga, O. R., Nguyen, H. C., Truong, H. N. & Nehlsen-Cannarella, S. L. 1999 Role of photoperiod and the pineal gland in T cell-dependent humoral immune reactivity in the Siberian hamster. *J. Pineal Res.* **27**, 243–248. (doi:10.1111/j.1600-079X.1999.tb00622.x)
- Zysling, D. & Demas, G. 2007 Metabolic stress suppresses humoral immune function in long-day, but not short-day, Siberian hamsters (*Phodopus sungorus*). *J. Comp. Physiol. B* **177**, 339–347.
- Zysling, D. A., Greives, T. J., Breuner, C. W., Casto, J. M., Dernas, G. E. & Ketterson, E. D. 2006 Behavioral and physiological responses to experimentally elevated testosterone in female dark-eyed juncos (*Junco hyemalis carolinensis*). *Horm. Behav.* **50**, 200–207. (doi:10.1016/j.yhbeh.2006.03.004)