



Published in final edited form as:

Open Pain J. 2009 ; 2: 30–40. doi:10.2174/1876386300902010030.

## Dual Roles for Endothelin-B Receptors in Modulating Adjuvant-Induced Inflammatory Hyperalgesia in Rats

Alla Khodorova<sup>1</sup>, Shiping Zou<sup>2</sup>, Ke Ren<sup>2</sup>, Ronald Dubner<sup>2</sup>, Gudarz Davar<sup>1,3</sup>, and Gary Strichartz<sup>\*,1</sup>

<sup>1</sup>Pain Research Center, Department of Anesthesiology, Perioperative and Pain Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

<sup>2</sup>Department of Biomedical Sciences, Dental School, Program in Neuroscience, University of Maryland, Baltimore, MD, USA

### Abstract

Injection of endothelin-1 (ET-1) into the plantar rat hindpaw causes acute pain at high concentrations and tactile sensitization at low concentrations. The pro-nociceptive actions are driven through ET<sub>A</sub> receptors for both levels of [ET-1], but the ET<sub>B</sub> receptors are only pro-nociceptive for allodynia from low [ET-1] and anti-nociceptive for pain from high [ET-1]. The goal of the present work was to discriminate the roles of the ET receptors in the acute hyperalgesia from inflammation by complete Freund's adjuvant (CFA, 20 mg/paw) into the rat hindpaw. Selective antagonists were injected 10 min before and then together with CFA. An ET<sub>A</sub> receptor antagonist, BQ-123, reduced CFA-induced thermal hyperalgesia (by up to 50%), as did an ET<sub>B</sub> receptor antagonist, BQ-788 (by up to 66%). BQ-123 and BQ-788 also delayed the onset (by 1.5 – 2 h) but insignificantly reduced the maximum degree of CFA-induced allodynia (~10%). Surprisingly, an ET<sub>B</sub> receptor *agonist*, IRL-1620, also reduced maximum thermal hyperalgesia induced by CFA, suppressed peak allodynia and delayed its occurrence by ~ 3 h. The latter actions of IRL-1620 were reversed by co-administration of BQ-788, naloxone hydrochloride and the peripherally restricted opiate receptor antagonist naloxone methiodide, and by antiserum against  $\beta$ -endorphin. These findings demonstrate an important role for endogenous ET-1 in acute inflammatory pain and a dual action of ET<sub>B</sub> receptors, including a pro-algesic action along with the important activation of a local analgesic pathway, implying that at least two different ET<sub>B</sub> receptors contribute to modulation of inflammatory pain.

### Keywords

Inflammatory hyperalgesia; endothelin-1; pro-nociception; anti-nociception; allodynia; pain

## INTRODUCTION

Inflammation releases substances that excite or sensitize primary afferent nerve fibers and cause pain and hyperalgesia [1,2]. Endothelin-1 (ET-1) is a peptide released following tissue injury and over-secreted in inflammatory conditions [3], and is derived from various cells in skin: keratinocytes [4], vascular endothelial cells [5], immune cells [6,7] and mast cells [8]. Sensory afferents themselves [9-11] and satellite cells of DRG [12] contain ET-1. Thus, both

© Khodorova et al.; Licensee Bentham Open.

\*Address correspondence to this author at the Pain Research Center/BWH, 75 Francis Street, Boston, MA 02115-6110, USA; Tel: (617) 732-7802; Fax: (617) 730-2801; gstrichz@zeus.bwh.harvard.edu.

<sup>3</sup>Current address: BiogIdec, Inc., 14 Cambridge Center, Cambridge, MA, USA

cells of the skin and those that innervate it may release ET-1 in normal and pathological conditions, and thereby contribute to pain (see review, [13]).

ET-1 potentiates the pain from pro-inflammatory mediators, e.g., PGE<sub>2</sub> [14] as well as pain-related activities of the capsaicin-heat-proton-activated receptor TRPV1, detected at the cellular [15-18] and at the whole animal, behavioral level [19].

Endothelin-1 can simultaneously activate both nociceptive and analgesic pathways, [20-32]. Although at first these opposing effects might be explained by the different actions of the two different G protein-coupled receptors for ET-1, called ET<sub>A</sub> and ET<sub>B</sub>, the problem is more complex. Exogenous ET-1 evokes acute pain, [21,24,25,33] and similarly enhances actions of other algogens, e.g. in experimental arthritic pain [23,24,34,35], both *via* ET<sub>A</sub> receptors. In contrast, activation of ET<sub>B</sub> receptors has been shown to have both an antihyperalgesic/antinociceptive action [24,26,27] and a pro-algesic action, e.g., causing mechanical hypernociception in rats [30]. A major objective of this paper is to address the separate, opposing effects of the ET<sub>B</sub> receptor, in inflammatory hyperalgesia that involves endogenous ET-1.

Endogenously-released ET-1 mediates pain (in the inflamed knee) via both ET<sub>A</sub> and ET<sub>B</sub> receptors [35]. ET<sub>B</sub> receptors contribute positively to pain from intraperitoneal inflammation in mice [21,36]. Although both complete Freund's adjuvant (CFA) and carrageenan have been reported to provoke thermal hyperalgesia in mice solely via ET<sub>A</sub> receptors, mechanical hyperalgesia in mice is mediated by both ET<sub>A</sub> and ET<sub>B</sub> receptors [29]. Carrageenan injected into peripheral tissues is known to rapidly increase local and plasma ET-1 levels [37] and chronic constriction of the rat sciatic nerve causing thermal and mechanical hyperalgesia (due to a substantial contribution from local inflammation [38]), elevates both ET-1 and ET<sub>A</sub> receptors at the injury site [39]. The behavioral signs of this injury-induced pain are reversed by an ET<sub>A</sub> receptor antagonist.

In summary, the ET<sub>A</sub> receptor appears always to promote inflammatory pain, but the role of ET<sub>B</sub> receptors is controversial and seems to depend on many factors: the procedure, the species, and the inflammatory state. Since we have previously shown an anti-hyperalgesic action of ET<sub>B</sub> receptors in the un-inflamed rat paw, in this work we sought to determine if ET<sub>B</sub> receptors were anti- or pro-algesic on the acute inflammatory pain induced by CFA in the rat paw. A preliminary report of these findings was presented at the 2003 meeting of the Society for Neuroscience\*.

## MATERIALS AND METHODOLOGY

All procedures used in these studies adhered to guidelines approved by the *Institutional Animal Care and Use Committee* of the University of Maryland Dental School and performed according to the ethical standards prescribed by the *Committee for Research and Ethical Issues* of the International Association for the Study of Pain. Experiments were performed on 144 adult (250-300 g), male Sprague-Dawley rats (Harlan, Indianapolis, IN). Rats were housed in cages (2-3 per cage) in a viral antibody-free facility on a 12/12 h light/dark cycle with food and water *ad libitum*. Prior to beginning experiments, animals were handled for 1-3 days to acclimate them to both the experimenters and the testing environment. Before measurements the rats were placed in a clear plastic chamber on a glass surface and allowed to acclimate for 15 - 30 min.

---

\*Zou S, Ren K, Dubner R, Khodorova A, Davar G. Endothelin receptor mechanisms of adjuvant-induced hyperalgesia in rats. 2003 (Abstract No 588.15). In: Program of the Society for Neuroscience 33<sup>rd</sup> Annual Meeting, November 8-12.

## Drugs

Complete Freund's adjuvant (CFA, *Mycobacterium tuberculosis*; Sigma-Aldrich, St. Louis, MO), used as the inflammatory agent, was suspended in an oil/saline (1:1) emulsion and administered at a final concentration of 0.5 mg/ml. All drugs were diluted in phosphate buffered saline (PBS, pH=7.4, Invitrogen, Carlsbad, CA) as stock solutions and stored at +4°C. Crude  $\beta$ -endorphin antiserum (C-55, a gift of Dr. G. Mueller, Uniformed Services University of the Health Sciences, Bethesda, Maryland) was stored at -20°C. Prior to the experiment, stock aliquots were diluted with PBS or mixed with undiluted CFA (1 mg/ml) at 1:1 (v/v). During experiments, working solutions were kept on ice to minimize breakdown. Crude C-55 was centrifuged for 5 min at 14,000x g in a microcentrifuge, then the supernatant was collected and used for injections. The ET<sub>A</sub> receptor selective antagonist, BQ-123 (D-Trp-D-Asp-Pro-D-Val-Leu), the ET<sub>B</sub> receptor selective antagonist, BQ-788 (*N-cis*-2,6-Dimethyl-piperidinocarbonyl-L-gamma-methylleucyl-D-1-thoxycarbonyltryptophanyl-D-Nle); and the ET<sub>B</sub> receptor selective agonist, IRL-1620 (Suc-Asp-Glu-Glu-Ala-Val-Tyr-Phe-Ala-His-Leu-Asp-Ile-Ile-Trp); were supplied by American Peptides Co. (Sunnyvale, CA). Naloxone hydrochloride and methyl-naloxone iodide were obtained from Sigma-Aldrich Chemical. The dose of naloxone used for local injection was based on previously described reports of efficacy in rat models of cutaneous pain [26,40,41].

## Injection Procedures

Injections of 40  $\mu$ l were delivered subcutaneously through a 28 gauge needle (regular bevel, 12.7 mm length, BD Medical) into the mid-plantar hindpaw, about 1 cm distal to the heel (for thermal testing), or into the lateral edge of the hindpaw (for mechanical testing). Only one paw per rat was injected and tests were completed on that paw and on the contralateral paw. Drugs were delivered with regard to the unilateral delivery of CFA as follows: An ET-1 receptor agonist or two antagonists, or naloxone, were injected twice, first pre-emptively (10 min before), and then, a few seconds before CFA. The latter delivery ("second injection", as noted in Results), given into the same hind paw site, was followed immediately by CFA (20  $\mu$ g/paw), as the third injection. (These procedures are referred to as *CFA + agent* in the **Data Analysis** section, below). In "control" experiments, the first two injections (prior to CFA) contained vehicle only (referred to as *CFA + vehicle*). ET receptor agonist, antagonists, and opioid receptor antagonists, or antiserum to  $\beta$ -endorphin were always injected at the same concentration for both first and second injections. In several previous reports of these agents acting in the rat skin we have shown that these concentrations of agents appear to be selective and effective, although in all cases they had to be used at several orders of magnitude above their equilibrium dissociation constant values. Concerns about this large ratio are addressed in the Discussion.

## Thermal Nociceptive Testing

The thermal nocifensive response was tested using the method of Hargreaves *et al.* [42], that allows for side-by-side comparisons of drug effects on inflamed and uninflamed paws within the same animal. The paw withdrawal latency, to the nearest 0.1 s, in response to paw heating by radiant energy was determined. If a rat failed to withdraw the heated paw by 20 s (cut off value), the trial was terminated. Initially, withdrawal latencies were measured in both left and right, naïve paws (pre-CFA level). Then, 15 min after CFA administration testing re-started and continued three more times for the next 3h, and then daily for up to 3 days after injection.

## Responses to Mechanical Stimulation

Calibrated Semmes-Weinstein (S-M) monofilaments (von Frey filaments, Stoelting, Wood Dale, IL) were used to mechanically stimulate the hindpaw. The bending force of the filaments ranged from 1 to 257 g. The testing method has been described in detail previously [43,44].

Briefly, rats were habituated to stand on their hindpaws and lean against the experimenter's hand covered by a regular leather work glove (Sears Inc., Balto, MD). The testing filament was pressed in the medial direction against the lateral edge of the hindpaw. The filaments were applied in an ascending series until the rat lifted the stimulated hindpaw. A descending series of the filaments were used when the rat responded to the starting filament. Each filament was tested 5 times, separated by intervals of a few seconds. If paw-withdrawal due to stimulation was observed, it was registered as a response to a filament. The response frequencies [(number of responses/number of stimuli) × 100%] to a range of von Frey filament forces were determined and a stimulus-response frequency curve was plotted. Non-linear regression analysis allowed determination of an EF<sub>50</sub> value, defined as the von Frey filament force (g) that produces a 50% response frequency and used as the measure of mechanical sensitivity. Prior to injection of CFA, there was no significant difference between the baseline stimulus-response frequency curves among the different groups of animals.

### Data Analysis

Data are reported as means ± S.E.M. Thermal hyperalgesia from CFA (preceded by “control”, vehicle injections, see p.8) was determined at the different times from the change from the baseline, pre-CFA value of the Paw Withdrawal Latency (PWL, in secs.). The degree of inhibition of the response by different doses of the different test agents was quantitated by taking the difference in the change in PWL between the *CFA + vehicle* injection and the *CFA + agent* injection, and dividing it by the change in PWLs between *Baseline* and *CFA + vehicle*:

$$\% \text{ inhibition} = 100 \times \frac{(\text{PWL:CFA+agent}) - (\text{PWL:CFA+vehicle})}{(\text{Baseline PWL}) - (\text{PWL:CFA+vehicle})}$$

To establish significant differences between the effects of *CFA + vehicle* and *CFA + agent*, multi-group ANOVA was performed with post-hoc application of Fisher's protected least significant difference test.  $P < 0.05$  was considered significant in all cases.

## RESULTS

### General Observations of Inflammatory Pain

Injection of CFA into the rat hindpaw produced a rapid onset of both thermal and mechanical hyperalgesia, as previously described [45,46]. Within 15 min after CFA injection (20 µg/paw), the latency to paw withdrawal (PWL) in response to a noxious thermal stimulus was significantly reduced, and persisted so for at least 3 h (Fig. 1). About 25% recovery had occurred at day 1 and about 70% by day 3, although thermal hypersensitivity was still significant at that time.

Single injections of the same volume (40 µl) of phosphate buffered saline (PBS) into the paw caused no significant change in PWL. Three injections of this volume, with the same intervals between injections as those used for the delivery of antagonists before and with CFA (see next), caused ~ 20% shortening in PWL ( $P > 0.05$  compared to baseline) at 15 min after the third injection, a reduction that slowly declined to zero over the next 90 min. The fall in PWL induced by saline was unaffected by an ET<sub>A</sub> receptor antagonist, indicating that it was not due to ET-1 released by the needle puncture.

CFA-induced hyperalgesia was accompanied by erythema and swelling of the hindpaw, similar to that reported for CFA given at higher doses [46-48]. Licking and guarding behavior of the injected hindpaw were also observed, as previously described. No significant changes in

thermal or mechano-responsiveness were detected in the contralateral paw after CFA (Fig. 1 for thermal); contralateral paw data are not reported further in this paper.

### **ET<sub>A</sub> Receptor Blockade Inhibits Thermal Hyperalgesia Evoked by CFA**

To evaluate the contribution of ET<sub>A</sub> receptors to CFA-induced thermal hyperalgesia, we injected the selective ET<sub>A</sub> receptor antagonist, BQ-123 (0.008-3.28 mM; total dose 6.56-262 nmol/paw), subcutaneously into the rat plantar hindpaw 10 min prior to and then just before CFA ( $n = 23$ ). BQ-123 inhibited ipsilateral thermal hyperalgesia maximally at 45 -180 min, by 20-50% over the antagonist's concentration range (BQ-123 vs. PBS; Fig. 1A).

### **ET<sub>B</sub> Receptor Blockade Partially Inhibits Thermal Hyperalgesia Evoked by CFA**

Subcutaneous paw injection of the selective ET<sub>B</sub> receptor antagonist, BQ-788 (0.075-1.5 mM; total dose 6-120 nmol/paw), also reduced hyperalgesia, maximally at 3 h after CFA (by 21-66%;  $n=14$ , over this respective concentration range; Fig. 1B). Thermal hyperalgesia in control (CFA + vehicle treated) paws and paws treated with the lower concentrations of BQ-788 remained significant at day 3, but had reversed to baseline values for paws injected with 120 nmoles BQ-788.

### **ET<sub>B</sub> Receptor Activation Inhibits Thermal Hyperalgesia Evoked by CFA**

To assess the capacity of activated ET<sub>B</sub> receptors to affect CFA-induced hyperalgesia, the ET<sub>B</sub> receptor agonist IRL-1620 (0.0055-0.55 mM; total dose 0.11-11 nmol/paw) was injected 10 min prior to and then just before CFA ( $n=22$ ). As shown in Fig. (2), thermal hyperalgesia was strongly reduced by IRL-1620; at a total dose of 11 nmol/paw (0.55 mM) inhibition was observed from 15 min to day 1 after CFA, including 75% inhibition at 45 and 90 min,  $n=12$ , and at a total dose of 1.1 nmol/paw (0.055 mM), 48% inhibition at 15 and 45 min ( $n=6$ ). The lowest dose of IRL-1620, 0.11 nmol/paw, however, gave no significant inhibition ( $n=4$ ).

Some inhibition of CFA-induced hyperalgesia was also observed when 11 nmoles (total dose) of IRL-1620 was injected subcutaneously at the neck ( $36 \pm 4\%$  inhibition,  $n=4$ ) (data not shown). This effect was half that resulting from the same concentration/dose injected directly into the paw and was equal to the effect when 0.1 of this total dose, i.e., 1.1 nmoles, was injected in the paw. It appears that a portion of the anti-hyperalgesic action of the ET<sub>B</sub> agonist resulted from its systemic distribution.

*Concentration vs. response* curves for the inhibition of CFA-induced thermal hyperalgesia by these antagonists of ET<sub>A</sub> and ET<sub>B</sub> receptors and the ET<sub>B</sub> receptor agonist IRL-1620 are shown in Fig. (3). (Here the injected doses are expressed as injected concentrations to permit comparison with published  $K_i$  values, reported as concentrations.) The data are too sparse for serious fitting of a non-linear function, e.g. a Hill equation, but data points above and below the 50% inhibition line allow estimates of IC<sub>50</sub> values. From such interpolation, the IC<sub>50</sub>s equal about 3 mM and 0.7 mM for the respective antagonists, BQ-123 and BQ-788, and about 0.1 mM for the agonist, IRL-1620. These values are in the *same rank order* as the reported affinities of these ligands for their respective receptors, a result that is consistent with the reported selectivity for the intended targets (see Discussion).

### **IRL-1620-Induced Anti-Hyperalgesia Operates through ET<sub>B</sub> Receptors, is Naloxone-Sensitive and Mediated by $\beta$ -Endorphin**

To confirm that an ET<sub>B</sub> receptor mediates the observed anti-hyperalgesic actions of IRL-1620, we co-injected this agonist (0.55 mM; total dose 11 nmol/paw) together with the ET<sub>B</sub> receptor antagonist, BQ-788 (0.75 mM; total dose 60 nmol/paw) before CFA. The peak anti-hyperalgesic action of IRL-1620 (~75% inhibition of the shortening of PWL caused by CFA,

at 45 min,  $n=12$ ) was reduced by about 2/3 (to ~24%,  $n=6$ , inhibition) by BQ-788 (Fig. 4), close to the value from inhibition from the systemic delivery of this dose (see above). This effect on CFA-induced hyperalgesia is consistent with IRL-1620's specific binding to local  $ET_B$  receptors to effect cutaneous anti-hyperalgesia, as we have shown previously for IRL-1620's effect on ET-1-induced pain behavior [26]. The residual anti-hyperalgesia when the local  $ET_B$  antagonist was co-injected with the agonist suggests that this antagonist may not distribute systemically to the same extent as IRL-1620.

Antinociception from IRL-1620 against the pain from exogenous ET-1 in glabrous paw skin is mediated by  $\beta$ -endorphin that is locally released from keratinocytes and then bound to  $\mu$ -opioid receptors, most probably located on nociceptor terminals [27]. We hypothesized that  $\beta$ -endorphin also mediates IRL-1620's inhibitory actions on CFA-induced hyperalgesia. Indeed, antisera against  $\beta$ -endorphin (C-55, 200  $\mu$ g/10  $\mu$ l) [49], injected subcutaneously into the plantar hindpaw 15 min before IRL-1620+CFA, almost completely prevented the inhibitory actions of IRL-1620, (~75% suppression by IRL-1620 alone, compared to ~12% suppression for IRL-1620 in C-55 pre-treated paws,  $n=6$ ;  $P<0.005$ ; Fig. 4). Naive rabbit serum (NRS), lacking antibodies against  $\beta$ -endorphin, did not affect IRL-1620's anti-hyperalgesia ( $n=4$ ; Fig. 4).

To verify that IRL-1620's inhibitory actions on CFA-induced hyperalgesia are mediated by opioid receptors, we used the opioid receptor antagonist naloxone ((-)-naloxone hydrochloride, NX). Co-injection of NX (0.69 mM; total dose 55 nmol/paw) with IRL-1620 lessened the inhibition of CFA-induced hyperalgesia, (Fig. 4; ~42% inhibition at 45 min;  $n=6$ , compared to ~75% inhibition by IRL-1620 alone before CFA;  $P < 0.05$ ).

In order to separate the peripheral and central nervous system effects of NX, we co-injected a peripherally-restricted opioid receptor antagonist, methyl-naloxone iodide (mNX, 0.69 mM; total dose 55 nmol/paw). This antagonist also substantially prevented IRL-1620's anti-hyperalgesia (from ~75% inhibition of the CFA-induced shortening in PWL by IRL-1620 alone vs. ~25% inhibition with mNX + IRL-1620,  $n=6$ ,  $P < 0.005$ ; Fig. 4). To control for possible systemic actions of mNX, it was injected subcutaneously at the neck with the same total dose as injected into the paw prior to injection of IRL-1620+CFA into the hindpaw. This treatment resulted in a weak and insignificant reduction of the anti-hyperalgesic actions of local IRL-1620 (from ~75% to ~51%,  $n=4$ , change in PWL,  $P>0.05$ ) (data not shown).

### Blockade of $ET_A$ and $ET_B$ Receptors Inhibits Mechanical Hyperalgesia Evoked by CFA

Mechanical hypersensitivity following CFA administration into the lateral edge of the rat hindpaw was characterized by both an increase in responses to supra-threshold stimuli (mechanical hyperalgesia) and the appearance of responses to weak stimuli that in naive animals did not produce nocifensive behavior (tactile allodynia). These changes, together causing a drop in the  $EF_{50}$  for paw withdrawal, were apparent within 15 min after injection, continued to increase up to 3 h, were maintained to day 1, and had partially recovered by day 3 (Fig. 5).

In the inflamed paw there is at least a 10-fold increase in mechanical sensitivity, evident when  $EF_{50} = 10$  g at 15 min after CFA, and which continues to fall over 3 hrs (to ~3 g). This substantial mechanical hypersensitivity is sustained for 1 day, with partial recovery at 3 days. When the  $ET_A$  receptor is blocked by BQ-123 (0.82 mM; total dose 66 nmol/paw) injected subcutaneously into the lateral edge of the rat hindpaw before CFA, the occurrence of mechanical hyperalgesia was delayed by several hours ( $n=6$ ; Fig. 5). However, BQ-123 did not significantly elevate the maximum CFA-altered reduction in  $EF_{50}$  measured from 3 h until day 3. Neither injection of CFA alone nor of BQ-123 + CFA affected responses to mechanical stimulation of the contralateral paw (not shown).

Blockade of ET<sub>B</sub> receptors by BQ-788 (0.75 mM; total dose 60 nmol/paw), injected before CFA, had a similar effect, delaying the development of mechanical hyperalgesia, with significant differences from CFA (after vehicle) at 15 and 45 min, but not effecting the EF<sub>50</sub>s at later times ( $n=4$ ; Fig. 5). The inhibition, by BQ-788, was briefer (maximum at 45 min vs. 90 min), and significantly smaller ( $P<0.001$ ) at 45 min after CFA, than the inhibition by almost equimolar (0.82 mM, from 66 nmoles) BQ-123, showing a potency rank of BQ-123>BQ-788, in contrast to the ca. 3-fold greater molar potency of BQ-788 over BQ-123 in suppressing thermal hyperalgesia (Fig. 3, above). BQ-788 treatment did not affect the mechanical responsiveness of the contralateral paw (not shown).

### Activation of an ET<sub>B</sub> Receptor Suppresses Mechanical Hyperalgesia Evoked by CFA

To assess the ability of ET<sub>B</sub> receptor activation to inhibit CFA-evoked mechanical hyperalgesia, IRL-1620 (total dose 11 nmol/paw) was injected before CFA ( $n = 6$ ). IRL-1620 prevented any decrease in EF<sub>50</sub> from CFA for the first 90 min (Fig. 5). However, subsequent mechanical responses measured 3 h and longer after injection of IRL-1620 + CFA, were not different from *vehicle* + CFA controls.

## DISCUSSION

The results reported here show that acute thermal and mechanical hyperalgesia, from inflammation induced by subcutaneous CFA in the rat hind paw, were inhibited by blockade of both local ET<sub>A</sub> and ET<sub>B</sub> receptors. In addition, activation of an ET<sub>B</sub> receptor also strongly reduced thermal and mechanical hyperalgesia. This latter anti-hyperalgesic effect was shown to be naloxone-sensitive and is probably dependent on keratinocyte release of the endogenous opioid peptide,  $\beta$ -endorphin [27]. Although a minor portion of the anti-hyperalgesic effect of the ET<sub>B</sub> antagonist, was due to systemic effects, about 75% was attributable to local actions in the paw.

It is essential to establish the receptor specificity of the ET receptor ligands used here. In other *in vivo* studies, conducted in our laboratory, measuring behavior or electrophysiological responses, high concentrations were used with total abolition of pain responses to ET-1 by BQ-123 and total reversal of ET<sub>B</sub>-mediated analgesia by BQ-788 [25-27,50]. Despite these high concentrations, the rank order of potency, IRL-1620> BQ-788~ BQ-123, in the present study is the same as the published *in vitro* inhibitory potencies for the respective ET receptor:  $K_i = 16$  pM for IRL-1620 at ET<sub>B</sub> receptors [51],  $K_i=1-100$  nM for BQ-788 at ET<sub>B</sub> receptors [52,53] and  $K_i=3.3-22$  nM for BQ-123 at ET<sub>A</sub> receptors [54-56].

In every report of ET-1-related effects, ET<sub>A</sub> receptor activation is pro-algesic, consistent with the observed inhibitory effect of the ET<sub>A</sub> receptor antagonist BQ-123 on CFA-induced hyperalgesia [13]. Activation of an ET<sub>B</sub> receptor by subcutaneous IRL-1620 has been shown previously to suppress nociception, and with the same apparent dependence on an opioidergic pathway as shown here. There is a hypothetical possibility that IR-1620 might also act on ET<sub>A</sub> receptors ( $K_i = 1.9$   $\mu$ M for ET<sub>A</sub>; [51]), however, in the case of ET<sub>A</sub> activation such an effect would favor hyperalgesia, not inhibit it. The fact that IRL-1620's anti-hyperalgesic effect is reversed by an ET<sub>B</sub> receptor antagonist, and by naloxone and the  $\beta$ -endorphin antibody, is completely inconsistent with ET<sub>A</sub> receptor blockade. The specificity of BQ-788 for inhibition of ET<sub>B</sub> receptors is testified to by its ability to abolish the analgesia from IRL-1620, whereas if it were acting at ET<sub>A</sub> receptors its effect would be anti-hyperalgesic. The reported effects are therefore fully consistent with the proposed specificity. It seems likely that the requirement for the high concentrations of subcutaneously administered agents results from the relatively impermeant nature of the dermis to such molecules when they are delivered subcutaneously, coupled with the requirement to reach nerve endings and keratinocytes located in the epidermis in order to act. In addition, these antagonists are peptides that are proteolytically degraded *in*

*vivo*, with half-lives of one hour or so [57], a factor that will determine not only the effective concentration that can reach the epidermal compartment but also the duration profile for the agents, possibly contributing to the < 1 day period of effective inhibition (cf. Fig. 1).

### Endogenous ET-1 and ET-Receptors in Adjuvant-Induced Thermal Hyperalgesia

The ET<sub>A</sub> receptor antagonist BQ-123 significantly relieved thermal hyperalgesia in CFA-treated rats, implying that endogenously released ET-1 causes part of this elevated pain response to CFA. Administered within the range of concentrations previously shown to completely abolish pain behavior evoked by exogenous ET-1 in rats [25,50], BQ-123 nonetheless only inhibited thermal hyperalgesia from CFA by ~50%. This inhibition reached its maximum at 45-90 min after CFA, consistent with the time course of stimulation-induced ET-1 production in different tissues *in vivo* (see [3]). The results with BQ-123 in the present study are evidence of an important, but limited role of ET<sub>A</sub> receptors in endogenous ET-1's actions in CFA-induced thermal hyperalgesia in rats.

Previous work has shown physiological effects of ET<sub>A</sub> receptor activation, on the soma of sensory neurons *in vitro* [58,59] and on impulses of nociceptive axons recorded *in vivo* after delivery of ET-1 to the plantar footpad (or to the sciatic nerve) [25,50]. Physiological actions of ET-1 applied to bare nerve cells *in vitro* or ensheathed fibers *in vivo*, both purely ET<sub>A</sub> mediated effects, have been previously proposed to completely account for the generation of impulses by *endogenous* ET-1 in the skin, e.g. after an incision [22]. In agreement with this proposition, Baamonde *et al.* [29] reported that only antagonists of ET<sub>A</sub> receptors were able to attenuate thermal inflammatory hyperalgesia in mice. However, the current results, showing that a selective ET<sub>B</sub> receptor antagonist partially decreases inflammatory hyperalgesia in rats, suggests that both ET<sub>B</sub> and ET<sub>A</sub> receptors contribute to inflammatory hyperalgesia. Moreover, the role of ET<sub>B</sub> receptors in CFA-induced hyperalgesia in rats is even more complex, as shown by the anti-hyperalgesia caused by ET<sub>B</sub> receptor activation (see **Dual effects from ET<sub>B</sub> receptors**, below).

### Endogenous ET-1 and ET-Receptors in Inflammatory Mechanical Hyperalgesia

Local blockade of either ET<sub>A</sub> or ET<sub>B</sub> receptors delayed the development of acute mechanical allodynia induced by CFA (Fig. 5). In contrast to the inhibitory actions on thermal hyperalgesia, the maximum extent of mechanical hyperalgesia was not affected by these agents, only the progression was slowed. Furthermore, at a time after CFA injection (45 min) when thermal hyperalgesia was only partially suppressed by BQ-123 (Fig. 1A) or BQ-788 (Fig. 1B), the same concentrations/doses of these antagonists almost totally prevented tactile allodynia (Fig. 5). Mechanisms involving ET-1 pathways therefore may be more important for suppressing the earlier phases of tactile mechanical hyperalgesia, but play a more constant role throughout all the stages of thermal hyperalgesia. Whether this difference is due to differences in the location of CFA injections in these two sensory modes, or to a differential distribution of endothelin receptors on the respective fiber types coding these separate modalities [60-62], i.e., to a peripheral differentiation, or to different central processing by spinal units that discriminate between inputs from fibers activated by different modality sensations, i.e., to CNS differentiation [63], remains to be shown.

### Dual Effects from ET<sub>B</sub> Receptors: a Plurality of Functions

The most remarkable observation here is that both an ET<sub>B</sub> receptor agonist and an ET<sub>B</sub> receptor antagonist reduced inflammatory thermal and mechanical hyperalgesia. These results indicate that under the conditions of acute inflammation ET<sub>B</sub> receptors are able to simultaneously mediate both pro- and anti-nociceptive actions.



The anti-hyperalgesia caused here by the ET<sub>B</sub> receptor antagonist BQ-788 agrees with previous reports that implicate ET<sub>B</sub> receptors in, 1. the pathogenesis of mechano-sensitivity in inflammatory pain [29,35], and with 2. the observations that ET<sub>B</sub> receptors (together with ET<sub>A</sub> receptors) mediate mechanical hyperalgesia induced by relatively low concentrations (30 nM – micromolar) of exogenous ET-1 [19,29,30,64]. These anti-hyperalgesic effects of BQ-788, however, contrast sharply with the proalgesic actions of BQ-788 shown for responses to high exogenous ET-1, responses that include the exacerbation by ET-1 of capsaicin-stimulated paw licking in mice [24] and the hindpaw flinching induced by ET-1 in rats [26].

On the other hand, the anti-hyperalgesic actions of an agonist of ET<sub>B</sub> receptors, here seen for CFA-induced hyperalgesia, has also been reported for other tests. The agonist IRL-1620, given pre-emptively, strongly inhibited the acute nociception from ET-1 [26,27], diminished the carrageenan-evoked hyper-nociception in the rat knee joint and reversed the increase in incapacitation caused by algogens delivered there [31]. Moreover, ET<sub>B</sub> receptor blockade enhanced both spontaneous and movement-evoked pain in a model of murine osteolytic cancer pain [65].

What might explain the similar effects from antagonists and agonists of ET<sub>B</sub> receptors? One possibility is that there are different sub-types of ET<sub>B</sub> receptors, with the one isoform blocked supporting pain, and another isoform suppressing pain. The particular contribution of ET<sub>B</sub> receptors, which may include more than one functional type [3,66-68], to pain processing could depend on the conditions, e.g. whether the periphery is normal or inflamed, implying that certain inflammatory mediators can modify the expression of ET receptors, or the receptors/channels to which they couple, e.g., TRPV1 [16], to enhance both primary receptor activation and the downstream coupling pathways. Evidence in support of different functional types of the ET<sub>B</sub> receptor is found in the description of at least two types of ET<sub>B</sub> receptors, characterized by nanomolar and picomolar K<sub>D</sub> values for ET-1, that are involved in the G-protein-mediated activation of different signal transduction pathways in different tissues/cells (see [68]). An equally attractive alternative to requiring more than one ET<sub>B</sub> receptor subtype to explain these opposing effects is to have the same receptor located on different cell types, whose separate outputs have opposing effects on pain. For example, pro-nociceptive ET<sub>B</sub>s might be present on nociceptors and sensitize them to the local excitatory actions of ET-1 acting through ET<sub>A</sub> and to excitation by noxious stimulation [19,29], while anti-hyperalgesic ET<sub>B</sub> would be present on keratinocytes where their activation triggers a widespread release of potent opioid peptides from these cells, e.g., β-endorphin, acting directly on nociceptive fibers and effecting a more powerful anti-hyperalgesic action. In fact, a very recent paper reports the presence of ET<sub>B</sub> receptors on the cell bodies of sensory neurons of the rat trigeminal ganglion, along with pharmacological data that such receptors contribute to nerve injury-induced thermal hyperalgesia [69].

## CONCLUSION

In summary, we have shown that endogenous ET-1 plays an important role in thermal and mechanical hyperalgesia during acute inflammation. Both types of ET receptors mediate these hyperalgesic responses, and whereas ET<sub>A</sub> regulates only pro-nociceptive actions, activation of ET<sub>B</sub> appears to play a dual role in modulating the final magnitude of pathological hypersensitivity. Activation of ET<sub>B</sub> receptors was highly effective in suppressing thermal and delaying the development of mechanical hyperalgesia of inflammatory origin, implying that ET receptors act differently to induce these different forms of inflammatory hyperalgesia.

## Acknowledgments

Thanks to Mr. Jamie Bell for technical assistance with the figures. Supported by USPHS grants CA080153 (to GS), DE11964 (to KR) and NS41384 (to RD).

## ABBREVIATIONS

CFA	Complete Freund's adjuvant
ET-1	Endothelin-1
ET <sub>A</sub>	Endothelin receptor-A
ET <sub>B</sub>	Endothelin receptor-B
NX	Naloxone
mNX	Methyl-naloxone
PBS	Phosphate buffered saline
PWL	Paw withdrawal latency
Veh	Vehicle
NRS	Naïve rabbit serum
EF <sub>50</sub>	Force (g) that produces a 50% withdrawal response frequency


## REFERENCES


- Kidd BL, Urban LA. Mechanisms of inflammatory pain. *Br J Anesthesiol* 2001;87:3–11.
- Schaible HG, Ebersberger A, Von Banchet GS. Mechanisms of pain in arthritis. *Ann NY Acad Sci* 2002;966:343–54. [PubMed: 12114291]
- Rae, GA.; Henriques, MG. Endothelins in inflammation.. In: Said, S., editor. *Pro-inflammatory and Anti-inflammatory Peptides*. Marcel and Dekker; New York: 1998. p. 163-202.
- Tsuboi R, Sato C, Shi CM, Nakamura T, Sakurai T, Ogawa H. Endothelin-1 acts as an autocrine growth factor for normal human keratinocytes. *J Cell Physiol* 1994;159:213–20. [PubMed: 8163562]
- Yanagisawa M, Kurihara H, Kimura S, Goto K, Masaki T. A novel peptide vasoconstrictor, endothelin, is produced by vascular endothelium and modulates smooth muscle Ca<sup>2+</sup> channels. *J Hypertens Suppl* 1988;6:S188–91. [PubMed: 2853725]
- Ehrenreich H, Anderson RW, Fox CH, et al. Endothelins, peptides with potent vasoactive properties, are produced by human macrophages. *J Exp Med* 1990;172:1741–48. [PubMed: 1701822]
- Sessa WC, Kaw S, Hecker M, Vane JR. The biosynthesis of endothelin-1 by human polymorphonuclear leukocytes. *Biochem Biophys Res Commun* 1991;174:613–18. [PubMed: 1993057]
- Ehrenreich H, Burd PR, Rottem M, et al. Endothelins belong to the assortment of mast cell-derived and mast cell-bound cytokines. *New Biol* 1992;4:147–56. [PubMed: 1313283]
- Szolcsanyi J, Oroszi G, Nemeth J, Szilvassy Z, Blasig IE, Tosaki A. Functional and biochemical evidence for capsaicin-induced neural endothelin release in isolated working rat heart. *Eur J Pharmacol* 2001;419:215–21. [PubMed: 11426844]
- Giaid A, Gibson SJ, Ibrahim BN, et al. Endothelin 1, an endothelium-derived peptide, is expressed in neurons of the human spinal cord and dorsal root ganglia. *Proc Natl Acad Sci USA* 1989;86:7634–38. [PubMed: 2678110]
- Franco-Cereceda A, Rydh M, Lou YP, Dalsgaard CJ, Lundberg JM. Endothelin as a putative sensory neuropeptide in the guinea-pig: different properties in comparison with calcitonin gene-related peptide. *Regul Pept* 1991;32:253–65. [PubMed: 1866471]
- Kar S, Chabot JG, Quirion R. Quantitative autoradiographic localisation of [125I]- endothelin-1 binding sites in spinal cord and dorsal root ganglia of the rat. *Neurosci Lett* 1991;133:117–20. [PubMed: 1665213]

13. Khodorova A, Montmayeur J-P, Strichartz G. Endothelin receptors and pain. *J Pain* 2009;10:4–28. [PubMed: 19111868]
14. Ferreira SH, Romitelli M, de Nucci G. Endothelin-1 participation in overt and inflammatory pain. *J Cardiovasc Pharmacol* 1989;13(Suppl 5):S220–22. [PubMed: 2473319]
15. Dymshitz J, Vasko MR. Endothelin-1 enhances capsaicin-induced peptide release and cGMP accumulation in cultures of rat sensory neurons. *Neurosci Lett* 1994;167:128–32. [PubMed: 7513837]
16. Yamamoto H, Kawamata T, Ninomiya T, Omote K, Namiki A. Endothelin-1 enhances capsaicin-evoked intracellular Ca<sup>2+</sup> response *via* activation of endothelin receptors in apertin kines C epsilon-dependent manner in dorsal root ganglion neurons. *Neuroscience* 2006;137:949–60. [PubMed: 16298080]
17. Plant TD, Zollner C, Mousa SA, Oschke A. Endothelin-1 potentiates capsaicin-induced TRPV1 currents *via* the endothelin A receptor. *Exp Biol Med* 2006;231:1161–4.
18. Plant TD, Zollner C, Kepura F, Mousa SS, Eichhorst J, Schaefer M. Endothelin potentiates TRPV1 *via* ETA receptor-mediated activation of protein kinase C. *Mol Pain* 2006;3:35–42. [PubMed: 18001466]
19. Balonov K, Khodorova A, Strichartz G. Tactile allodynia initiated by local subcutaneous endothelin-1 is prolonged by activation of TRPV1 receptors. *Exp Biol Med (Maywood)* 2006;231:1165–70. [PubMed: 16741070]
20. Rubanyi GM, Polokoff A. Endothelins: molecular biology, biochemistry, pharmacology, physiology, and pathophysiology. *Pharmacol Rev* 1994;48:325–415. [PubMed: 7831383]
21. Raffa R, Schupsky J, Jacoby H. Endothelin-induced nociception in mice: mediation by ETA and ETB Receptors. *J Pharmacol Exp Ther* 1996;276:647–51. [PubMed: 8632332]
22. Mujenda FH, Duarte AM, Reilly EK, Strichartz GR. Cutaneous endothelin-A receptors elevate post-incisional pain. *Pain* 2007;133:161–73. [PubMed: 17467172]
23. Piovezan AP, D'Orleans-Juste P, Tonussi CR, Rae GA. Effect of endothelin-1 on capsaicin-induced nociceptive in mice. *Eur J Pharmacol* 1998;351:15–22. [PubMed: 9698200]
24. Piovezan AP, D'Orleans-Juste P, Souza GE, Rae GA. Endothelin-1-induced ET(A) receptor-mediated nociception, hyperalgesia and oedema in the mouse hind-paw: modulation by simultaneous ET(B) receptor activation. *Br J Pharmacol* 2000;129:961–68. [PubMed: 10696096]
25. Gokin AP, Fareed MU, Pan HL, Hans G, Strichartz GR, Davar G. Local injection of endothelin-1 produces pain-like behavior and excitation of nociceptors in rats. *J Neurosci* 2001;21:5358–66. [PubMed: 11438612]
26. Khodorova A, Fareed MU, Gokin A, Strichartz GR, Davar G. Local injection of a selective endothelin-B receptor agonist inhibits endothelin-1 induced pain-like behavior and excitation of nociceptors in a naloxone-sensitive manner. *J Neurosci* 2002;22:7788–96. [PubMed: 12196602]
27. Khodorova A, Navarro B, Jouaville LS, et al. Endothelin-B receptor activation triggers an endogenous analgesic cascade at sites of peripheral injury. *Nat Med* 2003;9:1055–61. [PubMed: 12847519]
28. Menéndez L, Lastra A, Hidalgo A, Baamonde A. Nociceptive reaction and thermal hyperalgesia induced by local ET-1 in mice: a behavioural and fos study. *Naunyn-Schmiedeberg Arch Pharmacol* 2003;367:28–34. [PubMed: 12616338]
29. Baamonde A, Lastra A, Villazon M, Bordallo J, Hidalgo A, Menendez L. Involvement of endogenous endothelins in thermal and mechanical inflammatory hyperalgesia in mice. *Naunyn Schmiedeberg Arch Pharmacol* 2004;369:245–51. [PubMed: 14661069]
30. Da Cunha JM, Rae GA, Ferreira SH, de Chunha Q. Endothelins induce ETB receptor-mediated mechanical hypernociception in rat hindpaw: roles of cAMP and protein kinase C. *Eur J Pharmacol* 2004;501:87–94. [PubMed: 15464066]
31. Daher JB, Souza GE, D'Orleans-Juste P, Rae GA. Endothelin ETB receptors inhibit articular nociception and priming induced by carrageenan in the rat knee-joint. *Eur J Pharmacol* 2004;496:77–85. [PubMed: 15288578]
32. Houck CS, Khodorova A, Reale AM, Strichartz GR, Davar G. Sensory fibers resistant to the actions of tetrodotoxin mediate nocifensive responses to local administration of endothelin-1 in rats. *Pain* 2004;110:719–26. [PubMed: 15288413]


33. Davar G, Hans G, Fareed MU, Sinnott C, Strichartz G. Behavioral signs of acute pain produced by application of endothelin-1 to rat sciatic nerve. *Neuroreport* 1998;9:2279–83. [PubMed: 9694215]
34. De-Melo JD, Tonussi CR, D'Orleans-Juste P, Rae GA. Effects of endothelin-1 on inflammatory incapacitation of the rat knee joint. *J Cardiovasc Pharmacol* 1998a;31(Suppl 1):S518–20. [PubMed: 9595530]
35. De-Melo JD, Tonussi CR, D'Orleans-Juste P, Rae GA. Articular nociception induced by endothelin-1, carrageenan and LPS in naive and previously inflamed knee-joints in the rat: inhibition by endothelin receptor antagonists. *Pain* 1998b;77:261–69. [PubMed: 9808351]
36. Griswold DE, Douglas SA, Martin LD, et al. Endothelin B receptor modulates inflammatory pain and cutaneous inflammation. *Mol Pharmacol* 1999;56:807–12. [PubMed: 10496965]
37. Bertelli A, Clerico A, Chicca A, Giovannini L, Gorio A, Romano MA. Role of endothelin-1 in carrageenin-induced inflammation. *Int J Tissue React* 1992;14:225–30. [PubMed: 1300308]
38. Maves TJ, Pechman PS, Gebhart GF, Meller ST. Mechanisms of neuropathic pain following chronic gut ligation of the sciatic nerve in the rat. *Pain* 1993;54:57–69. [PubMed: 8378104]
39. Klass M, Hord A, Wilcox M, Denson D, Csete M. A role for endothelin in neuropathic pain after chronic constriction injury of the sciatic nerve. *Anesth Analg* 2005;101:1757–62. [PubMed: 16301255]
40. Eisenberg E, Vos BP, Strassman AM. The peripheral antinociceptive effect of morphine in a rat model of facial pain. *Neuroscience* 1996;72:519–25. [PubMed: 8737420]
41. Stein C, Gramsch C, Herz A. Intrinsic mechanisms of antinociception in inflammation: local opioid receptors and beta-endorphin. *J Neurosci* 1990;10:1292–98. [PubMed: 2158530]
42. Hargreaves K, Dubner R, Brown F, Flores C, Joris J. A new and sensitive method for measuring thermal nociception in cutaneous hyperalgesia. *Pain* 1988;32:77–88. [PubMed: 3340425]
43. Ren K. An improved method for assessing mechanical allodynia in the rat. *Physiol Behav* 1999;67:711–16. [PubMed: 10604842]
44. Anseloni VCZ, Weng H-R, Terayama R, et al. Age- dependency of analgesia elicited by intraoral sucrose in acute and persistent pain models. *Pain* 2002;97:93–103. [PubMed: 12031783]
45. Ji R-R, Befort K, Brenner GJ, Woolf CJ. ERK MAP kinase activation in superficial spinal cord neurons induces prodynorphin and NK-1 upregulation and contributes to persistent inflammatory pain hypersensitivity. *J Neurosci* 2002;22:478–85. [PubMed: 11784793]
46. Guo W, Zou S, Guan Y, et al. Tyrosine phosphorylation of the NR2B subunit of the NMDA receptor in the spinal cord during the development and maintenance of inflammatory hyperalgesia. *J Neurosci* 2002;22:6208–17. [PubMed: 12122079]
47. Iadarola MJ, Douglass J, Civelli O, Naranjo JR. Differential activation of spinal cord dynorphin and enkephalin neurons during hyperalgesia: evidence using cDNA hybridization. *Brain Res* 1988;455:205–12. [PubMed: 2900057]
48. Ren K, Hylden J, Williams GM, Ruda MA, Dubner R. The effects of a non-competitive NMDA receptor antagonist, MK-801, on behavioral hyperalgesia and dorsal horn neuronal activity in rats with unilateral inflammation. *Pain* 1992;50:331–44. [PubMed: 1454389]
49. Hargreaves K, Flores C, Dionne R, Mueller G. The role of pituitary beta- endorphin in mediating corticotropin-releasing factor-induced antinociception. *Am J Physiol* 1990;258:E235–42. [PubMed: 2137676]
50. Fareed MU, Hans G, Atanda A, Strichartz G, Davar G. Pharmacological characterization of acute pain behavior produced by application of endothelin-1 to rat sciatic nerve. *J Pain* 2000;1:46–53.
51. Takai M, Umemura I, Yamasaki K, et al. A potent and specific agonist, Suc-[Glu9,Ala11,15]-endothelin-1(8-21), IRL 1620, for the ETB receptor. *Biochem Biophys Res Commun* 1992;184:953–59. [PubMed: 1315540]
52. Ishikawa K, Ihara M, Noguchi K, et al. Biochemical and pharmacological profile of a potent and selective endothelin B-receptor antagonist, BQ-788. *Proc Natl Acad Sci USA* 1994;91:4892–96. [PubMed: 8197152]
53. Webber KM, Pennefather JN, Head GA, van den Buuse M. Endothelin induces dopamine release from rat striatum *via* endothelin-B receptors. *Neuroscience* 1998;86:1173–80. [PubMed: 9697124]

54. Ishikawa K, Fukami T, Nagase T, et al. Cyclic pentapeptide endothelin antagonists with high ETA selectivity. Potency- and solubility-enhancing modifications. *J Med Chem* 1992;35:2139–42. [PubMed: 1317926]
55. Marsault R, Feolde E, Frelin C. Receptor externalization determines sustained contractile responses to endothelin-1 in the rat aorta. *Am J Physiol* 1993;264(3 Pt 1):C687–93. [PubMed: 8096364]
56. Ihara M, Yamanaka R, Ohwaki K, et al. [3H]BQ-123, a highly specific and reversible radio-ligand for the endothelin ETA receptor subtype. *Eur J Pharmacol* 1995;274:1–6. [PubMed: 7768260]
57. Ihara M, Ishikawa K, Fukuroda T, et al. *In vitro* biological profile of a highly potent novel endothelin (ET) antagonist BQ-123 selective for the ETA receptor. *J Cardiovasc Pharmacol* 1992;20(Suppl 12):S11–4. [PubMed: 1282942]
58. Zhou QL, Strichartz G, Davar G. Endothelin-1 activates ET(A) receptors to increase intracellular calcium in model sensory neurons. *Neuroreport* 2001;12:3853–57. [PubMed: 11726808]
59. Zhou Z, Davar G, Strichartz G. Endothelin-1 (ET-1) selectively enhances the activation gating of slowly inactivating tetrodotoxin-resistant sodium currents in rat sensory neurons: a mechanism for the pain-inducing actions of ET-1. *J Neurosci* 2002;22:6325–30. [PubMed: 12151509]
60. Ossipov MH, Bian D, Malan TP Jr, Lai J, Porreca F. Lack of involvement of capsaicin-sensitive primary afferents in nerve-ligation induced tactile allodynia in rats. *Pain* 1999;79:127–33. [PubMed: 10068158]
61. Mansikka H, Shiotani M, Winchurch R, Raja SN. Neurokinin-1 receptors are involved in behavioral responses to high-intensity heat stimuli and capsaicin-induced hyperalgesia in mice. *Anesthesiol* 1999;90:1643–49.
62. Mansikka H, Sheth RN, DeVries C, Lee H, Winchurch R, Raja SN. Nerve injury-induced mechanical but not thermal hyperalgesia is attenuated in neurokinin-1 receptor knockout mice. *Exp Neurol* 2001;162:343–49. [PubMed: 10739640]
63. Treede RD, Meyer RA, Raja SN, Campbell JN. Peripheral and central mechanisms of cutaneous hyperalgesia. *Progr Neurobiol* 1992;38:397–421.
64. Motta EM, Calixto JB, Rae GA. Mechanical hyperalgesia induced by endothelin-1 in rats is mediated via phospholipase C, protein kinase C, and MAP kinases. *Exp Biol Med* 2006;231:1141–45.
65. Peters CM, Lindsay TH, Pomonis JD, et al. Endothelin and tumorigenic component of bone cancer pain. *Neuroscience* 2004;126:1043–52. [PubMed: 15207337]
66. Bax WA, Saxena PR. The current endothelin receptor classification: time for reconsideration? *Trends Pharmacol Sci* 1994;15:379–86. [PubMed: 7809954]
67. Yoneyama T, Hori M, Makatani M, et al. Subtypes of endothelin ETA and ETB receptors mediating tracheal smooth muscle contraction. *Biochem Biophys Res Commun* 1995;207:668–74. [PubMed: 7864858]
68. Sokolovsky M. Endothelin receptor heterogeneity, G-proteins, and signaling *via* cAMP and cGMP cascades. *Cell Mol Neurobiol* 1995;15:561–71. [PubMed: 8719041]
69. Chichorro JG, Zampronio AR, Cabrini DA, Franco CR, Rae GA. Mechanisms operated by endothelin ET(A) and ET(B) receptors in the trigeminal ganglion contribute to orofacial thermal hyperalgesia induced by infraorbital nerve constriction in rats. *Neuropeptides* 2009;43:133–42. doi:10.1016/j.npep.2008.12.001. [PubMed: 19157542]






**Fig. (1).** Time course of the anti-hyperalgesic effects of BQ-123 and BQ-788. Injection of Complete Freund's Adjuvant (CFA, 20  $\mu$ g/paw, with PBS vehicle) into the plantar hindpaw shortened the latency for paw withdrawal from a noxious thermal stimulus (PWL). **(A)** Local injection into the same paw of the selective ET<sub>A</sub> antagonist, BQ-123 ( $n=23$ ), at the total doses shown, partially restored the thermal paw withdrawal latency to its baseline value. **(B)** Local injection of the selective ET<sub>B</sub> antagonist, BQ-788 ( $n=14$ ), reduced CFA-induced thermal hyperalgesia at 45 min to 3 h after adjuvant injection. \* indicates significant difference ( $P < 0.05$ ) from control, i.e., CFA + vehicle ( $n=11$ ); <sup>x</sup> significantly different from baseline (pre-CFA) values in the ipsilateral paw (assessed for day 3 only).




**Fig. (2).** Robust anti-hyperalgesic effect of IRL-1620. Inhibition of CFA-induced thermal hyperalgesia occurred when ipsilateral injection of the ET<sub>B</sub> agonist IRL-1620 preceded CFA ( $n=22$ ). \* $P < 0.05$  indicates significant difference from control, CFA + vehicle ( $n=11$ );  $^xP < 0.05$  for comparison with baseline (pre-CFA) values in the ipsilateral paw.






**Fig. (3).** Concentration vs inhibition of thermal hyperalgesia (shortening of paw withdrawal latency) by the antagonists of ET<sub>A</sub> and ET<sub>B</sub>, and the agonist of ET<sub>B</sub>, BQ-123 and BQ-788, and IRL-1620, respectively. Intercepts of the dashed line indicate the EC<sub>50</sub>s.



**Fig. (4).**

Modulation of IRL-1620's anti-hyperalgesic action. The vertical axis shows the percent change, due to local injection of IRL-1620 (total dose 11 nmol/paw), in the paw withdrawal latency caused by CFA injection, calculated as:  $[(\text{post-CFA: PWL} - \text{pre-CFA: PWL}) / \text{pre-CFA: PWL}] \times 100\%$ , where a negative value results from reduction of PWL, indicative of hyperalgesia. Data show the peak values at 45 min time point. The inhibitory effect of co-administered, local IRL-1620 on CFA-induced thermal hyperalgesia is  $\text{ET}_B$ -receptor mediated (restored by BQ-788), naloxone (NX)- and naloxone methiodide (mNX)-sensitive, is prevented by antisera against  $\beta$ -endorphin (C55, 200  $\mu\text{g}$  in 10  $\mu\text{l}$ ), but not naïve anti-serum (NRS). # $P < 0.05$  indicates significant differences from IRL-1620; the number of experiments is indicated in the corresponding column.



**Fig. (5).**

Effects of the selective ET-receptor antagonists and ET<sub>B</sub> agonist on tactile hyperesthesia. EF<sub>50</sub>s, defined as the von Frey filament force (g) that produces withdrawal response half the time, were determined from interpolation of stimulus-response functions and used as a measure of mechanical sensitivity. Local pre-treatment (10 min prior to CFA and at the time of CFA injection) of rats with BQ-123 (total dose 65.6 nmol/paw), or BQ-788 (total dose 60 nmol/paw) delayed the fall of EF<sub>50</sub> in the inflamed paw, when compared to CFA + vehicle treated rats. Pre-treatment with the ET<sub>B</sub> agonist IRL-1620 (total dose 11 nmol/paw) prevented mechano-allodynia for 1.5 h. From 3 h onward there was no difference in EF<sub>50</sub>s between any of the ET-receptor agent-treated rats and those receiving CFA alone. (\**P* < 0.05 indicates significant differences from controls (CFA injection 10 min after vehicle injection). <sup>x</sup>*P* < 0.001 for CFA + vehicle or CFA + antagonist/agonist vs. the baseline, pre-CFA, values. <sup>+</sup>*P* < 0.05 for BQ-123 + CFA or IRL-1620 + CFA vs. BQ-788 + CFA).