

Vipp1 and PspA

Related but not twins

Jelle B. Bultema,³ Eva Fuhrmann,^{1,2} Egbert J. Boekema³ and Dirk Schneider^{1,*}

¹Institut für Biochemie und Molekularbiologie; ZBMZ; ²Fakultät für Biologie; Albert-Ludwigs-Universität; Freiburg, Germany;

³Groningen Biomolecular Sciences and Biotechnology Institute; University of Groningen; The Netherlands

The *Vesicle Inducing Protein in Plastids 1* (Vipp1) was suggested to be involved in thylakoid membrane formation in both chloroplasts and cyanobacteria. The protein shows sequence homology to the *Phage Shock Protein A* (PspA) from bacteria, and both proteins have similar secondary structures. 2D-structures of PspA and of Vipp1 have been determined by electron microscopy in the recent years. Both PspA and Vipp1 form large homooligomeric rings with high molecular masses but their ring dimensions differ significantly. Furthermore, Vipp1 forms rings with different rotational symmetries whereas PspA appears to form rings with singular rotational symmetry. In this article addendum we compare the structures of PspA and Vipp1. Furthermore, we suggest a spatial structural model of the observed Vipp1 rings.

In 2001 a gene was identified in *Arabidopsis thaliana*, deletion of which resulted in thylakoid membrane perturbations as well in disturbed vesicle formation.¹ Due to this observation, the encoded protein was named *Vesicle Inducing Protein in Plastids 1* (Vipp1). Homologous *vipp1* genes were found in the genomes of cyanobacteria, and while the gene in the cyanobacterium *Synechocystis* sp. PCC 6803 could not be completely deleted, depletion of the gene product resulted in a severe reduction of internal thylakoid membranes and in decreased levels of active photosystems.²⁻⁴ While the protein appears to operate similar in cyanobacteria and chloroplasts, its exact physiological functions remain

mystic. Interestingly, the amino acid sequence of Vipp1 shows similarity to the *Phage Shock Protein A* (PspA) from bacteria.⁵ Deletion experiments indicated that PspA is not essential in bacteria whereas Vipp1 is essential for normal development of both cyanobacteria and chloroplasts.^{1-3,5,6} Although the exact physiological function of PspA is essentially elusive, recent observations indicate that the protein is involved in maintaining the integrity of the *E. coli* cytoplasmic membrane.⁷ PspA from *E. coli* binds to specific lipids of the cytoplasmic membrane,⁷ and also Vipp1 associates tightly with both the cytoplasmic or inner envelope and the thylakoid membrane in cyanobacteria and chloroplasts.^{1,8,9}

A 2D projection structure of the 1 MDa homooligomeric PspA ring from *E. coli* has been determined by electron microscopy (EM), and more recently the structure of Vipp1 rings were reported.^{8,10} Although both structures have similar arrangements, there are significant differences, which could correlate to separate physiological roles.

The amino acid sequences of PspA from *E. coli* as well as for PspA and Vipp1 from *Synechocystis* align from their N-terminus up to amino acid residue 218 without insertions or deletions (Fig. 1). The two PspA proteins share a sequence identity of 27% and a sequence similarity of 51%. In line with rather low sequence conservation, the *Synechocystis* PspA protein and *Synechocystis* Vipp1 also share just 31% sequence identity (51% similarity). While the amino acid sequence of the proteins is not strictly conserved, the proteins have a highly conserved secondary structure.

Key words: PspA, Vipp1, ring, structure, function

Submitted: 11/05/09

Accepted: 11/05/09

Previously published online:

www.landesbioscience.com/journals/cib/article/10529

*Correspondence to: Dirk Schneider;
Email: Dirk.Schneider@biochemie.uni-freiburg.de

Addendum to: Fuhrmann E, Bultema JB, Kahmann U, Rupprecht E, Boekema EJ, Schneider D. The vesicle inducing protein I from *Synechocystis* sp. PCC 6803 organizes into diverse higher ordered ring structures. *Mol Biol Cell* 2009; 20:4620-8; PMID: 19776353.




Figure 1. Amino acid sequence alignment of Vipp1 from *Synechocystis* and PspA from *Synechocystis* and *E. coli* with Clustal X.¹⁴ Identical (*), strongly similar (:) and weakly identical (.) amino acid residues are indicated. The grey regions represent α -helical domains predicted with PSIPRED, V2.6.¹⁵

Table 1. A comparison of structure and symmetry of PspA and Vipp1 rings, as determined by electron microscopy and single particle image analysis

	Width (nm)	Height (nm)	Symmetry	Copies/ring	Reference
PspA	20	8.5	9 fold	36	10
Vipp1	25–33	22	12–17 fold	48–68	8

The predicted secondary structure for all proteins is essentially purely α -helical, apart from some coil interruptions, which is in excellent agreement with experimental determination of the Vipp1 secondary structure.⁸ Both PspA and Vipp1 contain large regions likely to form coiled-coils of helices when analyzed with the program PCOILS.¹¹ In contrast to PspA, Vipp1 contains a C-terminal extension of 45 amino acids, which forms an additional α -helix. This domain, which is not involved in oligomer formation, appears to be a unique feature of Vipp1 when compared to PspA.

Both PspA and Vipp1 form large homooligomeric rings. The structural details of these rings, including dimensions and observed symmetry are compared in Table 1. While PspA forms only a single ring population with 9-fold rotational symmetry, Vipp1 rings have been observed with 12 to 17-fold rotational symmetry and increased ring diameters. The unique

PspA ring is formed by 36 PspA molecules, whereas 48 to 68 Vipp1 monomers contribute to the formation of the various Vipp1 rings. The PspA ring has a height of ~8.5 nm which is considerable smaller than the Vipp1 rings which all have a constant height of ~22 nm.

So far, no crystal structure for Vipp1 is available but a potential tertiary structure can be obtained by 3D structure prediction. For this purpose the *Synechocystis* Vipp1 sequence was submitted to the PHYRE web server. A description about and case study using PHYRE is given by Kelley and Sternberg.¹² All generated 3D models had either one long α -helix or coiled-coil structures. The model based on the 3D structure of smooth muscle α -actinin (pdb entry 1sjj) was chosen for further modelling of the oligomeric Vipp1 ring. This starting model (Fig. 2A) contains amino acids 4–253 from Vipp1 (~93% of the total Vipp1 sequence).

The Vipp1 starting model has a length of 17 nm and a diameter of ~2.5 nm. Partly, it is a three-start coiled-coil in the central part of the structure as shown in Figure 2A.

Already at this stage it becomes obvious that the tertiary structure of PspA and Vipp1 must significantly differ. The suggested Vipp1 model has a length of 17 nm which later on (compared below) translates to a ring height of >17 nm. Since the PspA ring has only a height of 8.5 nm, the generated model cannot be used to derive the 3D structure of the PspA ring. It is e.g., possible that the long α -helices in the PspA monomer are kinked to form a higher-ordered helix-bundle with a length of ~8 to 9 nm. This difference in tertiary structure may determine the specificity of PspA vs. Vipp1.

Ring structures with 15 outward and 15 inward pointing spikes (15-fold rotational symmetry) are in the middle of the




Figure 2. Model of the Vip1 structure. (A and B) Side and top view ribbon representations of four predicted Vip1 monomers arranged in a dimer of dimers configuration. (C) Projection map showing a Vip1 ring with 15-fold symmetry. (D–F) Top, tilted and side view of the modelled Vip1 ring, respectively. In (D) the model is placed on top of the projection map shown in (C). The scale bar in (C) equals 10 nm which is also valid for (D–F).

observed range of Vip1 symmetries. To each spike two densities can be attributed, both with a diameter of ~ 2.5 nm. Since the diameters of the observed 2.5 nm densities and of the predicted Vip1 monomer are roughly identical, we speculate that each density could originate from a Vip1 monomer. Thus, the spikes, as seen in the EM projection maps, are each occupied by two Vip1 proteins.

Two Vip1 monomers were arranged manually such that the total dimer length was ~ 22 nm, which corresponds to the ring height determined by single particle analysis.⁸ Furthermore, two dimers were placed next to each other based on the spikes in the EM projection map with 15-fold symmetry (Fig. 2B–D). Biochemical studies have also indicated that the smallest repeating unit within

the Vip1 rings is a dimer of dimers.⁸ The 15-fold ring model (Fig. 2D–F below) was generated from this tetramer using routines from the CCP4 package.¹³ The α -helical configurations of the model are in good agreement with the overall ring dimensions, and the presented model can give an impression of how the Vip1 ring structure could look like.

Acknowledgements

This work was supported by a grant from the Deutsche Forschungsgemeinschaft (SCHN 690/3-1) and the Council of Chemical Sciences (CW) of the Netherlands Science Foundation NWO. We thank Marcel Bokhove (Department of Biophysical Chemistry, University of Groningen) for help with modelling.

References

1. Kroll D, Meierhoff K, Bechtold N, Kinoshita M, Westphal S, Vothknecht UC, et al. VIPP1, a nuclear gene of *Arabidopsis thaliana* essential for thylakoid membrane formation. *Proc Natl Acad Sci USA* 2001; 98:4238-42.
2. Fuhrmann E, Gathmann S, Rupprecht E, Golecki J, Schneider D. Thylakoid membrane reduction affects the photosystem stoichiometry in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol* 2009; 149:735-44.
3. Westphal S, Heins L, Soll J, Vothknecht UC. Vippl deletion mutant of *Synechocystis*: a connection between bacterial phage shock and thylakoid biogenesis? *Proc Natl Acad Sci USA* 2001; 98:4243-8.
4. Gao H, Xu X. Depletion of Vippl in *Synechocystis* sp. PCC 6803 affects photosynthetic activity before the loss of thylakoid membranes. *FEMS Microbiol Lett* 2009; 292:63-70.
5. Weiner L, Model P. Role of an *Escherichia coli* stress-response operon in stationary-phase survival. *Proc Natl Acad Sci USA* 1994; 91:2191-5.
6. Aseva E, Ossenbuhl F, Sippel C, Cho WK, Stein B, Eichacker LA, et al. Vippl is required for basic thylakoid membrane formation but not for the assembly of thylakoid protein complexes. *Plant Physiol Biochem* 2007; 45:119-28.
7. Kobayashi R, Suzuki T, Yoshida M. *Escherichia coli* phage-shock protein A (PspA) binds to membrane phospholipids and repairs proton leakage of the damaged membranes. *Mol Microbiol* 2007; 66:100-9.
8. Fuhrmann E, Bultema JB, Kahmann U, Rupprecht E, Boekema EJ, Schneider D. The Vesicle Inducing Protein 1 from *Synechocystis* sp. PCC 6803 Organizes into Diverse Higher Ordered Ring Structures. *Mol Biol Cell* 2009.
9. Li HM, Kaneko Y, Keegstra K. Molecular cloning of a chloroplastic protein associated with both the envelope and thylakoid membranes. *Plant Mol Biol* 1994; 25:619-32.
10. Hankamer BD, Elderkin SL, Buck M, Nield J. Organization of the AAA(+) adaptor protein PspA is an oligomeric ring. *J Biol Chem* 2004; 279:8862-6.
11. Lupas A. Prediction and analysis of coiled-coil structures. *Methods Enzymol* 1996; 266:513-25.
12. Kelley LA, Sternberg MJ. Protein structure prediction on the Web: a case study using the Phyre server. *Nat Protoc* 2009; 4:363-71.
13. The CCP4 suite: programs for protein crystallography. *Acta Crystallogr D Biol Crystallogr* 1994; 50:760-3.
14. Thompson J, Gibson T, Plewniak F, Jeanmougin F, Higgins D. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucl Acids Res* 1997; 25:4876-82.
15. McGuffin LJ, Bryson K, Jones DT. The PSIPRED protein structure prediction server. *Bioinformatics* 2000; 16:404-5.