

# Sigma factor phosphorylation in the photosynthetic control of photosystem stoichiometry

Masanori Shimizu<sup>1</sup>, Hideki Kato, Takeshi Ogawa, Aya Kurachi, Yoshiko Nakagawa, and Hirokazu Kobayashi<sup>2</sup>

Laboratory of Plant Molecular Improvement and Global Center of Excellence (COE) Program, Graduate School of Nutritional and Environmental Sciences, University of Shizuoka, Suruga, Shizuoka 422-8526, Japan

Edited\* by Bob B. Buchanan, University of California, Berkeley, CA, and approved May 5, 2010 (received for review October 29, 2009)

An imbalance in photosynthetic electron transfer is thought to be redressed by photosynthetic control of the rate of expression of genes encoding apoproteins of photosystem (PS)-I and PS-II in response to the redox state of plastoquinone (PQ), which is a connecting electron carrier. PS stoichiometry is then adjusted to enhance photosynthetic efficiency. In prokaryotes, sigma factors are well known for their participation in the control of RNA polymerase activity in transcription, whereas there have been no reports concerning their association with redox regulation. We have found that the phosphorylation of SIG1, the major sigma factor (SIG), is regulated by redox signals and selectively inhibits the transcription of the *psaA* gene, which encodes a PS-I protein. We produced transgenic *Arabidopsis* plants with or without the putative phosphorylation sites for SIG1 and demonstrated through *in vivo* labeling that Thr-170 was involved in the phosphorylation. We analyzed the *in vivo* and *in vitro* transcriptional responses of the transgenic *Arabidopsis* plants to the redox status in regard to involvement of the phosphorylation site. We revealed an enhanced phosphorylation of SIG1 under oxidative conditions of PQ in a form associated with the molecular mass of the holoenzyme. Phosphorylation of SIG1 proved crucial through a change in the promoter specificity for sustaining balanced expression of components in PS-I and PS-II and was responsible for harmonious electron flow to maintain photosynthetic efficiency.

chloroplast | redox regulation | plastoquinone | RNA polymerase | *Arabidopsis*

Plants are required for the survival of humans as food supplies and as alternative sources of energy in the form of biofuel, and are important in conservation of the environment. These contributions of plants are based on the function of chloroplasts. Chloroplasts are thought to have originated from ancestral cyanobacteria, which became symbionts in eukaryotic host cells. Plants have evolved as sessile organisms that use their chloroplasts to exploit solar energy. They are unable to escape from undesirable environmental conditions and must adapt to survive. When plants are exposed to sunlight, their chloroplast photosystems (PSs) excite electrons donated by water, which are transferred to NADP<sup>+</sup> and other recipients. Altered light conditions influence the efficiency of electron excitation in PS-I and PS-II, leading to an imbalance in photosynthetic electron transfer (1–3) and injury caused by uncoupled excess energy. This imbalance is thought to be redressed by the photosynthetic control of PS stoichiometry (2), where the rate of expression of genes for apoproteins of PS-I and PS-II is regulated in response to the redox state of plastoquinone (PQ) (4, 5). However, it is not known how the regulation is achieved in a signal transduction pathway through redox states of PQ. We have focused on the activity of RNA polymerase as one of the critical steps. The transcription of most photosynthesis genes in the chloroplast is accomplished with bacterial-type plastid-encoded RNA polymerase (PEP), which requires multiple nucleus-encoded sigma factors (SIGs) (6–9) destined for the chloroplast. To reveal the underlying mechanisms we focused on SIGs, which are well

characterized in prokaryotes as components of RNA polymerase and engaged in transcription in chloroplasts.

## Results and Discussion

**Phosphorylation Site of the Sigma Factor.** The amino acid sequences of all of the sigma proteins, SIG1 to SIG6, were searched for sites that could be phosphorylated. Among multiple categories of predicted sites, we initiated mutagenesis in those sites that had a minimal number of candidates. SIG1 transcripts and protein are the most abundant of the six kinds of SIGs in leaves grown in light for 2–4 wk, a finding that led us to focus on SIG1 (Fig. S1). SIG1 has sites that can be phosphorylated by Ser/Thr protein kinases, and those immediately ahead of region 1.2 are conserved in *Arabidopsis* SIG3 and SIG5 (Fig. S2) and in SIG1 orthologs. We produced transgenic *Arabidopsis* derivatives carrying wild-type SIG1 (referred to as SIG1) and derivatives with individual putative phosphorylation sites that were deleted [*sig1*(S55L), *sig1*(T170V) and *sig1*(S55L/T170V); Fig. 1A]. As SIG genes were expressed in the presence of light (6), 3-wk-old *Arabidopsis* plants were maintained in the dark for 1 wk to lower endogenous SIG levels before analyzing their chloroplast gene transcripts. In each transgenic line, we investigated the expression of *rbcl*, which is known to be transcribed by PEP, to determine whether the phosphorylation of sigma factors influences transcription in chloroplasts. The transcript levels of *rbcl* were normalized with the SIG protein level by immunoblotting with antibodies against recombinant SIG1 (anti-SIG1; Fig. 1B). In lines transformed with *sig1*(T170V) and *sig1*(S55L/T170V), transcript levels were higher than in those harboring SIG1 or *sig1*(S55L) (Fig. 1B). Therefore, Ser-55 in SIG1 is not involved in transcript level regulation. This finding is consistent with it being located in the putative transit peptide that may be eliminated to generate mature SIG1, whereas Thr-170 might play a role in transcription regulation. Transcript levels were normalized with SIG1 protein content, which eliminated the possibility that the amino acid substitution interfered with the SIG1 protein degradation. Hence, we conclude that Thr-170 among the predicted phosphorylation sites is responsible for reducing the levels of transcripts of chloroplast genes in the dark. The phenotypes are reproducible in multiple independent lines [wild-type, SIG1, and *sig1*(S55L), or *sig1*(T170V) and *sig1*(S55L/T170V)] (Fig. 1B), suggesting both that the status of phosphorylation in SIG1 may affect chloroplast gene expression and that potential position effects or dominant-negative effects may not occur. This in-

Author contributions: M.S., H. Kato, and H. Kobayashi designed research; M.S., H. Kato, T.O., A.K., and Y.N. performed research; and M.S. and H. Kobayashi wrote the paper.

The authors declare no conflict of interest.


\*This Direct Submission article had a prearranged editor.

Freely available online through the PNAS open access option.

<sup>1</sup>Present address: School of Health Promotional Science, Hamamatsu University, 1230 Miyakoda, Hamamatsu 431-2102, Japan.

<sup>2</sup>To whom correspondence should be addressed. E-mail: hirokoba@u-shizuoka-ken.ac.jp.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.0911692107/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.0911692107/-DCSupplemental).



**Fig. 1.** Effects of Ser/Thr phosphorylation of SIG1 on the transcript level of a chloroplast gene. (A) Predicted Ser/Thr phosphorylation sites in *Arabidopsis* SIG1. (B) Levels of *rbcl* transcripts in wild-type plants and transgenic lines *sig1(S55L)*, *sig1(T170V)*, and *sig1(S55L/T170V)* when adapted to the dark for 1 wk. Levels determined by quantitative RT-PCR were standardized by using *ACT2* transcripts and were further normalized by the content of SIG1 protein determined by immunoblotting. Vertical bars at the top of histograms represent the SDs of three independent experiments. (C–E) Phosphorylation of SIG1 and *sig1(T170V)*. (C) Plants (3-wk-old) were incubated with [ $^{32}$ P]orthophosphate in the dark. Immunoprecipitates with anti-SIG1 were fractionated by SDS/PAGE, and radioactivity on the dried gel was detected with Storm 820 (Amersham Biosciences). (D) Immunoblotting of extracts shown in C with anti-SIG1. (E) Extracts from SIG1-transgenic plants labeled with  $^{32}$ P in the dark were prepared without phosphatase inhibitors only for this experiment and were immunoprecipitated with anti-SIG1. The immunoprecipitates were incubated with or without 25 units of calf intestinal alkaline phosphatase for 30 min, followed by SDS/PAGE and autoradiography.

ference was also supported by the analysis of the number and location of inserted T-DNA sites in the genome (SI Text).


The effect of the amino acid substitution at Thr-170 might have resulted from a change of protein conformation rather than phosphorylation status. To test this hypothesis, we dipped the roots of SIG1-transgenic plants in  $^{32}$ P in the dark and determined the amount of  $^{32}$ P in their SIG proteins. Although the amounts of SIG1 protein expressed were roughly equivalent in *Arabidopsis* transformed with SIG1 and in *sig1(T170V)* (Fig. 1D), there was a strong  $^{32}$ P signal in the SIG1-transgenic line and a faint signal in the *sig1(T170V)* transgenic line (Fig. 1C). Phosphorylated SIG1 detected by autoradiograph (Fig. 1C) was at a higher molecular weight than SIG1 detected by immunoblotting. The faint signal of the upper band in the immunoblot is due to a smaller amount of phosphorylated SIG1 than non-phosphorylated SIG1, which is consistent with the result that most of the SIG1 proteins existed without phosphorylation at Thr-170, free from the core enzyme (see below).

We determined the ratio of transcripts derived from exogenous *sig1(T170V)* to those from the endogenous wild-type SIG1 by sequencing individual cDNA clones and found that *sig1(T170V)* showed a 20-fold higher expression than endogenous SIG1. The low level of expression of endogenous SIG1 was consistent with the faint signal obtained from *sig1(T170V)* plants. The  $^{32}$ P signals diminished when extracts were treated with calf intestinal alkaline phosphatase (Fig. 1E). Hence, the Thr-170 residue of wild-type SIG1 could be phosphorylated. Together with previous experimental data, these findings show that phosphorylation of Thr-170 of SIG1 interferes with chloroplast gene transcription.

To confirm that SIG1 phosphorylation controls transcriptional activity, we performed chloroplast transcription run-on experiments (Fig. 2A). Transgenic lines for SIG1 and *sig1(T170V)*, and wild-type plants, were grown for 4 wk and transferred to the dark for 2 h. Chloroplasts were isolated and used for run-on transcription. Transcriptional activities were expressed per SIG1 molecule. In the *sig1(T170V)*-transgenic line, transcription of *psbA*, *psbD*, *psaA*, *rbcl*, and *rrn16* was higher than that for wild-type SIG1. Hence, phosphorylation of SIG1 reduces the transcription of genes encoded by the chloroplast genome to different extents. The transcript levels per SIG1 molecule in the SIG1-transgenic line were not significantly different from those

in the wild-type. Transcript levels of *psbA* and *psaA* in the SIG1-transgenic line were 0.82-fold and 0.64-fold lower, respectively, than in the *sig1(T170V)*-transgenic line. Phosphorylation of SIG1 might therefore affect *psaA* transcription more than *psbA* transcription. Pfannschmidt et al. (4) showed that the redox state of PQ controls the level of transcription of *psbA* and *psaA*. Hence, the phosphorylation of SIG1 might mediate the adjustment of stoichiometry in response to redox signals from PQ.

**Redox Regulation of SIG1 Phosphorylation.** We altered the redox state of PQ in plants by using 3-(3',4'-dichlorophenyl)-1,1'-dimethyl urea (DCMU) for the oxidative state and 2,5-dibromo-3-methyl-6-isopropyl-p-benzoquinone (DBMIB) for the reductive state. *psaA* expression in the wild-type was diminished with DCMU but not with DBMIB, whereas the reverse was observed for *psbA* expression (Fig. 2B). These agents therefore acted as expected on the basis of a previous report (4). *psaA* expression was not enhanced, but tended to diminish, under oxidative conditions with DCMU in SIG1-transgenic plants, whereas it was stimulated in *sig1(T170V)*-transgenic plants (Fig. 2B). Hence, the effect on *psaA* transcripts in *sig1(T170V)*-transgenic plants was different from that in SIG1-transgenic and wild-type plants, and the inhibitory effect was stronger to a different degree than in wild-type plants under reductive conditions with DBMIB. By contrast, *psbA* expression was similar in wild-type and SIG1-transgenic plants with DCMU, whereas it declined in *sig1(T170V)*-transgenic plants to the level found in wild-type plants under reductive conditions with DBMIB (Fig. 2B). The results for *sig1(T170V)*-transgenic plants show that SIG1 is phosphorylated by oxidative signals, and the phosphorylation of SIG1 shifts the expression pattern from the reductive type to the oxidative type. Fig. 2C shows the ratios of accumulation of the *psbA* transcript to the *psaA* transcript (*psbA/psaA*), as used to define the stoichiometry of the adjustment between the two PSs. The *psbA/psaA* ratio in wild-type and SIG1-transgenic plants varied from one to five when changing from reductive conditions to oxidative conditions, respectively. However, the *psbA/psaA* ratio under oxidative conditions was lower in *sig1(T170V)*-transgenic plants than in the other plant lines. This result has been reproduced with three additional transgenic lines for both SIG1 and *sig1(T170V)* (Fig. S3). Overall, these results indicate that the *psaA* transcript level was elevated in *sig1*




**Fig. 2.** Effects of phosphorylation of SIG1 on the run-on transcription and transcript levels of genes for PS-II (*psbA*) and PS-I (*psaA*) at different states of PQ. (A) Run-on transcription of chloroplast genes. Transcriptional activity for each gene is shown as the fraction of that of the wild-type after standardization with the SIG1 content determined by immunoblotting. Error bars represent the variations of three independent experiments. Asterisks indicate statistically significant differences ( $P < 0.05$ ). (B and C) Transcript levels in plants were determined by real-time RT-PCR. The redox status of PQ was manipulated with the inhibitors DCMU and DBMIB. (B) Ratios of transcript levels of each gene in the treated transgenic lines to those in the untreated wild-type. (C) Ratio of transcripts of *psbA* to *psaA* in each plant. "Control" means the plants were not treated. Error bars represent the variation of three independent experiments. Triple asterisks indicate statistically significant differences ( $P < 0.01$ ).

(*T170V*)-transgenic plants under oxidative conditions, whereas the *psbA* transcript level was not. Hence, the phosphorylation preferentially reduces the transcription of *psaA*. We conclude that a change in the promoter specificity of phosphorylated SIG1 leads to a difference between the transcription of *psaA* and *psbA*, and the phosphorylation of SIG1 causes the changes of stoichiometry that depend on the redox state of PQ.

Are phosphorylated SIG1 factors free or associated with the RNA polymerase core enzyme *in vivo*? Under moderate light intensity, SIG1 exists mainly in the free form in wild-type plants, based on its molecular mass in sucrose density gradients (Fig. 3A). A minor peak of SIG1 was found in fraction 2 at  $\approx 400$  kDa (Fig. 3A), where the  $\beta$ -subunit of the core was detected with antibodies against the *rpoB* product (Fig. S4). Plants transgenic for wild-type SIG1 were labeled with [ $^{32}\text{P}$ ]orthophosphate in the light and dark. After sucrose density-gradient centrifugation, the  $^{32}\text{P}$  incorporated into SIG1 was detected by immunoprecipitation with anti-SIG1 (Fig. 3B and C). The profiles of phosphorylated SIG1 molecules differed in the light where most sedimented in fraction 3 at  $\approx 200$  kDa (Fig. 3B), whereas in the dark they formed two peaks at fractions corresponding to the RNA polymerase holoenzyme (fraction 2) and free SIG1 according to molecular mass (Fig. 3C). Hence, SIG1 in the holoenzyme is phosphorylated only in the dark. The preferential inhibition of transcription of *psaA* relative to *psbA* can be ascribed to the presence of phosphorylated SIG1 in the holoenzyme. Photosynthetic control of transcription by the redox signal from the reduced and oxidized forms of PQ might be explained as follows. In the reductive state, the holoenzyme contains nonphosphorylated SIG1 and transcribes *psaA* and *psbA* equally. In the oxidative state, the holoenzyme contains phosphorylated SIG1 and transcribes *psbA* efficiently and *psaA* poorly (Fig. 4).

**Pathway of the Redox Signal Started from PQ States.** The study of the phosphorylation of sigma factors was initiated by Tiller et al. (10). Their group showed that sigma-like factors in mustard were phosphorylated and dephosphorylated in etioplasts and chloroplasts, respectively. In addition, they reported that plastid transcription kinase (PTK) affected the phosphorylation of an RNA polymerase moiety in mustard plastids through the reduction of glutathione in response to irradiance (11), and that cpCK2 was an enzyme corresponding to PTK on the criterion of their common properties (12). The group has recently found that cpCK2 phosphorylates multiple sites in SIG6, and that mutations at these sites cause the malfunction of RNA polymerase (13). We show here that the phosphorylation of sigma factors is engaged in the stoichiometry adjustment of PSs and involved in a kind of regulation different from that of cpCK2. There are two kinds of redox-signaling pathways originating from PSs, namely the "priming signal" generated from PQ and the "thiol-mediated signal" (14). The former signal transduction pathway may be involved in the phosphorylation of sigma factors observed in this investigation. This phosphorylation is promoted by the priming signal, starting from the oxidized state of PQ that results from low-intensity or higher-wavelength light. Although the components meditating the signal transduction pathway for the stoichiometry adjustment of PSs have not yet been clarified, Puthiyaveetil et al. (15) reported that chloroplast sensor kinase (CSK) is required to control the transcription of chloroplast genes after the redox state. CSK might be involved in an early step in the signal transduction pathway leading to the dephosphorylation of sigma factors under the control of the priming signal.

Various redox effects impact the transcription and transcript stability of photosynthetic genes (16, 17), transcription of nuclear genes (18, 19), and *de novo* protein synthesis (20, 21). In higher plants, Pfannschmidt et al. (4) demonstrated direct regulatory coupling between the redox state of PQ and the transcription of specific chloroplast genes. We have shown that the stoichiometry




**Fig. 3.** Effect of phosphorylation of SIG1 on association with the RNA polymerase core enzyme in vivo. (A) Leaf extracts of wild-type plants harvested in the daytime were subjected to sucrose density-gradient centrifugation, followed by immunoblotting with anti-SIG1. Molecular masses are based on those of the native proteins in each fraction, as further analyzed by nondenaturing PAGE (4–12%, wt/vol, polyacrylamide gradient). (B and C) *SIG1* transgenics (3-wk-old) were incubated with [<sup>32</sup>P]orthophosphate in light at 75 μmol·m<sup>-2</sup>·s<sup>-1</sup> (B) or in the dark (C). The extracted proteins were fractionated by sucrose density-gradient centrifugation, and each fraction was then immunoprecipitated with anti-SIG1, run on SDS/PAGE, and autoradiographed.

of the two PSs is adjusted by modification of a sigma factor in response to a redox signal. Redox regulation typically transduces signals rapidly, directly, and simply (4). Instead of the de novo synthesis of proteins and mRNA or the participation of multiple transcriptional factors, a sigma factor is modified by phosphorylation or dephosphorylation. Each bacterial sigma factor plays a specific role in gene expression under different conditions, and many cooperate to adapt to changed environments (22–24). However, we have shown that modification of a sigma factor causes a change of promoter preference in higher plants.

Nucleic acids and proteins inside prokaryotic endosymbionts or chloroplasts are exposed to oxygen-free radicals, which cause mutations and protein degradation (25). Land plants cannot escape light illumination and have developed systems for preventing

damage (1–4, 25–32). To minimize damage caused by the generation of reactive oxygen species (ROS), plants have evolved mechanisms of photoprotection (28–32). PS-I must be induced to accept electrons from PQ to prevent the generation of ROS, when PS-II is strongly active in the daytime by shorter-wavelength light. Our results elucidate a mechanism of induction of PS-I through the dephosphorylation of sigma factors by redox signals of the PQ state (Fig. 4). On the other hand, longer-wavelength light, which is enriched at sunrise and sunset, must be absorbed more efficiently by PS-I than PS-II, resulting in the excitement of electrons in PS-I and, to a lesser degree, in PS-II. Formation of PS-I may then be repressed to coordinate with the activity of PS-II, leading to a reduction of the rate of cyclic electron flow (31, 32) and the detoxification of ROS (30). Sigma factors play a role in this regulation through their phosphorylation in response to the oxidized state of PQ (Fig. 4). Consequently, the phosphorylation of sigma factors under redox control may therefore be involved in maintaining the photosynthetic efficiency.



**Fig. 4.** Model of the photosynthetic control of chloroplast gene transcription via phosphorylation of a SIG. The redox status of PQ is changed by the intensity and wavelength of light: Irradiance or shorter-wavelength light (680 nm) generates PQH<sub>2</sub>, and low light or longer-wavelength light (700 nm) at sunrise and sunset oxidizes PQ. The chemically reduced form of PQH<sub>2</sub> is produced by electron flow from PS-II and converted to PQ after electron flow to PS-I. Electron flow from PS-II to PQ can be inhibited by DCMU, and electron flow from PQ to PS-I can be inhibited by DBMIB. The expression of *psaA*, encoding the apoprotein of the PS-I reaction center, is induced by PQH<sub>2</sub> and repressed when PQH<sub>2</sub> is oxidized. The expression of *psbA* encoding the D1 protein of the PS-II reaction center is less affected by redox status. The core enzyme associated with dephosphorylated SIG has high transcriptional activity for *psaA* and *psbA*, and phosphorylation of SIG causes a greater reduction of the transcription of *psaA* than *psbA*.

## Materials and Methods

**Transformation of *Arabidopsis*.** Constructs for ectopic expression of *SIGs* were made with the 35S-GFP(S65T) vector and pSMAB701. Details of in vitro mutagenesis of *SIG1* and plasmid construction can be found in *S1 Text* and *Table S1*.

**<sup>32</sup>P-labeling and Immunoprecipitation.** Whole plants were placed in Petri dishes containing 10 mL of [<sup>32</sup>P]orthophosphate solution (H<sub>3</sub><sup>32</sup>PO<sub>4</sub>, ICN; 5.55 MBq) for 2 h. They were rinsed with distilled water to remove excess <sup>32</sup>P. Protein extracts were mixed with Protein-G-Sepharose (Amersham Biosciences) at 4 °C for 1 h. The resulting supernatants were reacted with anti-SIG1 and incubated at 4 °C for 1 h, followed by passage through Protein-G-Sepharose and incubation at 4 °C for 1 h. The immunoprecipitated complexes were washed with chilled PBS, and subjected to SDS polyacrylamide gel electrophoresis (SDS/PAGE; 10%, wt/vol, polyacrylamide).

**Analysis of Gene Expression and Protein Products.** Real-time PCR was performed with a LightCycler (Roche), and detected with LightCycler DNA Master SYBR Green I (Roche). Transcripts for cytosolic actin (33) (ACT2) were measured as an internal standard.

Chloroplasts prepared from 4-wk-old plants were incubated in the dark for 2 h and subjected to run-on assays. Values were normalized on the basis of SIG1 content determined by immunoblotting, and the means of three independent experiments are shown.

Protein extracts were subjected to SDS/PAGE (10%, wt/vol, polyacrylamide), and SIGs were detected with ECL Plus Western Blotting Detection System (Amersham Biosciences). Details of the protocols are available in *S1 Text*.

**Treatment of Plants with Electron-Transport Inhibitors.** Seedlings grown on MS solid medium in the dark for 5 d were cultured hydroponically with MS liquid medium additionally for 5 d. The medium was then replaced with that containing 0.5  $\mu$ M DCMU or 0.2  $\mu$ M BDMIB and plants were incubated in the light for 2 d.

**ACKNOWLEDGMENTS.** We thank Mamoru Isemura for guidance in the preparation of rabbit antisera, Hiroaki Ichikawa of the National Institute of Agrobiological Resources in Japan for pSMAB701, Takanori Hirano and Yasuo Niwa for intermediate constructs used in cloning in our laboratory, Csaba Koncz at Max Planck Institute for Plant Breeding Research in Cologne in Germany for *Agrobacterium tumefaciens* GV3101, Akira Ishihama for

technical comments and encouragement, Kozi Asada and Kimiyuki Satoh for discussion on photo-oxidative damage, and Philip Hawke for English editing. The research was supported by Grants-in-Aid, the 21st Century COE Program and the Global COE Program from the Ministry of Education, Culture, Sports, Science and Technology of Japan (Monbukagakusho), and by grants from the Toray Science Foundation and the Salt Science Research Foundation in Japan. This paper is dedicated to the late Lawrence Bogorad, who originally characterized chloroplast RNA polymerase, as well as the *psbA* gene for triazine receptor protein and the *psaA* gene for PSI-A1, which he described as "photogenes." H. Kobayashi and other colleagues had the honor of sharing time and common interests with him at Harvard University until he passed away in 2003.

- Fujita Y, Murakami A, Ohki K (1987) Regulation of photosystem composition in the cyanobacterial photosynthetic system: The regulation occurs in response to the redox state of the electron pool located between the two photosystems. *Plant Cell Physiol* 28:283–292.
- Chow WS, Melis A, Anderson JM (1990) Adjustments of photosystem stoichiometry in chloroplasts improve the quantum efficiency of photosynthesis. *Proc Natl Acad Sci USA* 87:7502–7506.
- Kim JH, Glick RE, Melis A (1993) Dynamics of photosystem stoichiometry adjustment by light quality in chloroplasts. *Plant Physiol* 102:181–190.
- Pfannschmidt T, Nilsson A, Allen JF (1999) Photosynthetic control of chloroplast gene expression. *Nature* 397:625–628.
- Allen JF, Pfannschmidt T (2000) Balancing the two photosystems: Photosynthetic electron transfer governs transcription of reaction centre genes in chloroplasts. *Philos Trans R Soc Lond B Biol Sci* 355:1351–1359.
- Isono K, et al. (1997) Leaf-specifically expressed genes for polypeptides destined for chloroplasts with domains of  $\sigma^{70}$  factors of bacterial RNA polymerases in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 94:14948–14953.
- Allison LA (2000) The role of sigma factors in plastid transcription. *Biochimie* 82: 537–548.
- Tsunoyama Y, et al. (2004) Blue light-induced transcription of plastid-encoded *psbD* gene is mediated by a nuclear-encoded transcription initiation factor, AtSig5. *Proc Natl Acad Sci USA* 101:3304–3309.
- Ishizaki Y, et al. (2005) A nuclear-encoded sigma factor, *Arabidopsis* SIG6, recognizes sigma-70 type chloroplast promoters and regulates early chloroplast development in cotyledons. *Plant J* 42:133–144.
- Tiller K, Link G (1993) Phosphorylation and dephosphorylation affect functional characteristics of chloroplast and etioplast transcription systems from mustard (*Sinapis alba* L.). *EMBO J* 12:1745–1753.
- Link G (2003) Redox regulation of chloroplast transcription. *Antioxid Redox Signal* 5: 79–87.
- Ogrzewalla K, Piotrowski M, Reinbothe S, Link G (2002) The plastid transcription kinase from mustard (*Sinapis alba* L.). A nuclear-encoded CK2-type chloroplast enzyme with redox-sensitive function. *Eur J Biochem* 269:3329–3337.
- Schweeer J, Turkeri H, Link B, Link G (2010) AtSIG6, a plastid sigma factor from *Arabidopsis*, reveals functional impact of cpCK2 phosphorylation. *Plant J* 62:192–202.
- Buchanan BB, Balmer Y (2005) Redox regulation: A broadening horizon. *Annu Rev Plant Biol* 56:187–220.
- Puttiyaveetil S, et al. (2008) The ancestral symbiont sensor kinase CSK links photosynthesis with gene expression in chloroplasts. *Proc Natl Acad Sci USA* 105: 10061–10066.
- Oswald O, Martin T, Dominy PJ, Graham IA (2001) Plastid redox state and sugars: Interactive regulators of nuclear-encoded photosynthetic gene expression. *Proc Natl Acad Sci USA* 98:2047–2052.
- Liere K, Link G (1997) Chloroplast endoribonuclease p54 involved in RNA 3'-end processing is regulated by phosphorylation and redox state. *Nucleic Acids Res* 25: 2403–2408.
- Escoubas JM, Lomas M, LaRoche J, Falkowski PG (1995) Light intensity regulation of *cab* gene transcription is signaled by the redox state of the plastoquinone pool. *Proc Natl Acad Sci USA* 92:10237–10241.
- Fey V, et al. (2005) Retrograde plastid redox signals in the expression of nuclear genes for chloroplast proteins of *Arabidopsis thaliana*. *J Biol Chem* 280:5318–5328.
- Danon A, Mayfield SP (1994) Light-regulated translation of chloroplast messenger RNAs through redox potential. *Science* 266:1717–1719.
- Trebitsh T, Danon A (2001) Translation of chloroplast *psbA* mRNA is regulated by signals initiated by both photosystems II and I. *Proc Natl Acad Sci USA* 98: 12289–12294.
- Kroos L, Kunkel B, Losick R (1989) Switch protein alters specificity of RNA polymerase containing a compartment-specific sigma factor. *Science* 243:526–529.
- Ishihama A (2000) Functional modulation of Escherichia coli RNA polymerase. *Annu Rev Microbiol* 54:499–518.
- Brahamsha B, Haselkorn R (1992) Identification of multiple RNA polymerase sigma factor homologs in the cyanobacterium *Anabaena* sp. strain PCC 7120: Cloning, expression, and inactivation of the *sigB* and *sigC* genes. *J Bacteriol* 174:7273–7282.
- Martin W, et al. (1998) Gene transfer to the nucleus and the evolution of chloroplasts. *Nature* 393:162–165.
- Allen JF, Raven JA (1996) Free-radical-induced mutation vs redox regulation: Costs and benefits of genes in organelles. *J Mol Evol* 42:482–492.
- Pfannschmidt T (2003) Chloroplast redox signals: How photosynthesis controls its own genes. *Trends Plant Sci* 8:33–41.
- Müller P, Li XP, Niyogi KK (2001) Non-photochemical quenching. A response to excess light energy. *Plant Physiol* 125:1558–1566.
- Allen JF (2003) Cyclic, pseudocyclic and noncyclic photophosphorylation: New links in the chain. *Trends Plant Sci* 8:15–19.
- Asada K (2006) Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol* 141:391–396.
- Munekage Y, et al. (2004) Cyclic electron flow around photosystem I is essential for photosynthesis. *Nature* 429:579–582.
- DalCorso G, et al. (2008) A complex containing PGRL1 and PGR5 is involved in the switch between linear and cyclic electron flow in *Arabidopsis*. *Cell* 132:273–285.
- An YQ, et al. (1996) Strong, constitutive expression of the *Arabidopsis* ACT2/ACT8 actin subclass in vegetative tissues. *Plant J* 10:107–121.