



Published in final edited form as:

*J Am Chem Soc.* 2008 June 18; 130(24): 7746–7755. doi:10.1021/ja7110348r.

## Insights into the Structural Specificity of the Cytotoxicity of 3-Deoxy-Phosphatidylinositols

Yanling K. Wang<sup>1</sup>, Wei Chen<sup>1</sup>, Derek Blair<sup>2</sup>, Mingming Pu<sup>1</sup>, Yingju Xu<sup>3</sup>, Scott J. Miller<sup>3</sup>, Alfred G. Redfield<sup>4</sup>, Thomas C. Chiles<sup>2</sup>, and Mary F. Roberts<sup>1</sup>

<sup>1</sup>Department of Chemistry, Boston College, Chestnut Hill, MA 02467

<sup>2</sup>Department of Biology, Boston College, Chestnut Hill, MA 02467

<sup>3</sup>Department of Chemistry, Yale University, New Haven, CT 06520

<sup>4</sup>Department of Biochemistry, Brandeis University, Waltham, MA 02454

### Abstract

*D*-3-Deoxy-phosphatidylinositol derivatives have cytotoxic activity against various human cancer cell lines. These phosphatidylinositols have a potentially wide array of targets in the phosphatidylinositol-3-kinase (PI3K)/Akt signaling network. To explore the specificity of these types of molecules, we have synthesized *D*-3-deoxy-dioctanoylphosphatidylinositol (*D*-3-deoxy-diC<sub>8</sub>PI), *D*-3,5-dideoxy-diC<sub>8</sub>PI and *D*-3-deoxy-dioctanoylphosphatidylinositol-5-phosphate and their enantiomers, characterized their aggregate formation by novel high resolution field cycling <sup>31</sup>P NMR, and examined their susceptibility to phospholipase C (PLC) and their effects on the catalytic activity of PI3K and PTEN against diC<sub>8</sub>PI and dioctanoylphosphatidylinositol-3-phosphate substrates, respectively, as well as their ability to induce the death of the U937 human leukemic monocyte lymphoma cells. Of these molecules, only *D*-3-deoxy-diC<sub>8</sub>PI was able to promote cell death; it did so with an IC<sub>50</sub> of 40 μM, well below the CMC of 0.4 mM. Under these conditions, there was little inhibition of PI3K or PTEN observed in assays of recombinant enzymes (although the complete series of deoxy-PI compounds did provide insights into ligand binding by PTEN). The *D*-3-deoxy-diC<sub>8</sub>PI was a poor substrate and not an inhibitor of the PLC enzymes. The *in vivo* results are consistent with the current thought that the PI analogue acts on Akt1 since the transcription initiation factor eIF4e, which is a downstream signaling target of the PI3K/Akt pathway, exhibited reduced phosphorylation on Ser209. Phosphorylation of Akt1 on Ser473, but not Thr308, was reduced. Since the potent cytotoxicity for U937 cells is completely lost with the *L*-3-deoxy-diC<sub>8</sub>PI as well as with modification of the hydroxyl group at the inositol C5 (either replacing the –OH with a hydrogen or phosphorylating it) in *D*-3-deoxy-diC<sub>8</sub>PI, both chirality of the phosphoinositol moiety and the hydroxyl group at C5 are major determinants of 3-deoxy-PI binding to its target in cells.

### Keywords

3-deoxyphosphatidylinositol; PTEN; field cycling NMR; short-chain phospholipids; U937 cells

### INTRODUCTION

The phosphatidylinositol 3-kinase (PI3K<sup>a</sup>) / Akt (or protein kinase B) signaling pathway is critical for cell survival and upregulated in a variety of human cancer cell lines<sup>1</sup> and solid

<sup>a</sup>To whom correspondence should be addressed: M.F.R., Merkert Chemistry Center, Boston College, 2609 Beacon Street, Chestnut Hill, MA 02467; Tel: (617)-552-3616; FAX: (617)-552-2705; mary.roberts@bc.edu.

tumors.<sup>2-4</sup> A key step in this pathway is the specific phosphorylation of the 3-hydroxyl group of the inositol ring in phosphoinositides by PI3K enzymes. These lipid products affect cell growth by binding specifically to enzymes such as Akt and recruiting them to the membrane for activation by phosphorylation.<sup>5-7</sup> The activated Akt survival signal in cells stems from its phosphorylation and inactivation of pro-apoptosis proteins. Counterbalancing PI3K is the lipid phosphatase PTEN, which specifically dephosphorylates the 3-phosphate and in so doing inhibits PI3K/Akt signaling.<sup>8</sup> Indeed, mutations of PTEN have been reported in an array of human tumors.<sup>9,10</sup>

Attempts to inhibit the PI3K/Akt pathway led to the synthesis of *D*-3-deoxy-phosphatidylinositol molecules that can no longer be phosphorylated by PI3K (for a review see Gills and Dennis (2004)).<sup>11</sup> Many of these molecules have antiproliferative properties.<sup>12-13</sup> The first of these, 3-deoxy-dipalmitoyl-PI, was shown to inhibit cancer cell (HT-29 human colon carcinoma) growth in vitro with an IC<sub>50</sub> of 35 μM.<sup>12</sup> Recent syntheses of ether linked rather than ester linked alkyl chains, e.g., *D*-3-deoxy-myoinositol 1-[(*R*)-3-(hexadecyloxy)-2-hydroxypropyl hydrogen phosphate], have generated a newer class of PI analogues that should have higher stability in vivo and may also have slightly better delivery properties since they are more like lyso-phospholipids.<sup>14</sup> 3-Deoxy-PIs have also been shown to reduce drug resistance in human leukemia cell lines.<sup>15</sup> Thus, these PI analogues may have a future in treatment of a variety of cancers.

The proposed mechanism for inhibition of cell growth by deoxy-PIs is rather curious. Previous work suggested that rather than inhibit PI3K, the 3-deoxy-PI compounds inhibit the serine/threonine protein kinase Akt by binding tightly to its PH domain, which normally binds PI(3,4)P<sub>2</sub> or PI(3,4,5), and trapping it in the cytoplasm, thereby preventing phosphorylation by effector kinases.<sup>7</sup> The spectrum of changes in cells caused by these 3-deoxy-PI molecules differs from other widely studied cell growth inhibitors.<sup>16</sup> Interestingly, their patterns of activity are most like other lipid-based compounds (e.g., miltefosine and perifosine), which do not contain inositol moieties and also appear to inhibit Akt translocation and phosphorylation,<sup>17-18</sup> and distinct from other compounds known to inhibit the PI3K/Akt pathway.

Significant synthetic effort has been used to modify the *D*-inositol ring to improve its cytotoxicity,<sup>19</sup> but little has been done to explore broader changes in the stereochemistry of the inositol ring or systematically employing further deoxygenation<sup>20, 21</sup> to assess specific interactions of 3-deoxy-PI molecules with targets. Modulating inhibitor solubility by shortening acyl chains so that it can exist as monomers may also contribute to understanding mechanisms of action of these lipids, since this modification is likely to alter uptake and localization in the cell. To this end we have synthesized a series of 3-deoxy-dioctanoyl phosphatidylinositol (3-deoxy-diC<sub>8</sub>PI) derivatives (Figure 1) with altered chirality of the inositol ring, and additional 5-deoxy-modification, or the addition of a phosphate group to the 5-hydroxyl group. The phosphate at C-5 was motivated by the possibility of producing an activator of PTEN since other groups have shown that phosphoinositides with a phosphate group at that position kinetically activate PTEN.<sup>22</sup> This series of 3-deoxy-diC<sub>8</sub>PI molecules has been examined as substrates/inhibitors of PI3K, PI-specific phospholipase Cδ1, and PTEN (enzymes that could be inhibited by or degrade deoxy-diC<sub>8</sub>PI molecules in cells), and then screened as inhibitors of the human leukemia U937 cell line. Few of the 3-deoxy-PI compounds affected the kinase or phospholipase activities at low concentrations relative to substrate. However, interesting trends were observed for PTEN inhibition by *D*- and *L*- series of lipids that help delineate how this enzyme is likely to bind substrate molecules. Of all the compounds examined, only *D*-3-deoxy-diC<sub>8</sub>PI was cytotoxic to U937 cells with an IC<sub>50</sub> of 40 μM, well below the CMC of 0.4 mM, and at a concentration where it had no effect on the recombinant enzymes examined. This very specific cytotoxicity profile for U937 cells is discussed in terms of the proposed target for this type of molecule.

## EXPERIMENTAL SECTION

### Deoxy-diC<sub>8</sub>PI lipids

Peptide-catalyzed asymmetric phosphorylation (Scheme 1) has been used previously in the total synthesis of PI(3)P analogues.<sup>23</sup> Combined with a monodebenzylation-Mitsunobu sequence, this methodology has also been applied in the syntheses of some deoxygenated PI analogues, in both enantiomeric series.<sup>24</sup> The 3-deoxygenated analogue of PI(5)P and its enantiomer, 3-deoxydiC<sub>8</sub>PI(5)P and *ent*-3-deoxy-diC<sub>8</sub>PI(5)P, were prepared following the same strategy (Scheme 1) with both protected *D*- and *L*-3-deoxy-diC<sub>8</sub>PI, steps *a* through *g* in Scheme 2, as the starting points.

**Protected *D*-3-deoxy-diC<sub>8</sub>PI(5P)**—To a stirred solution of the benzyl protected *D*-3-deoxy-diC<sub>8</sub>PI (0.020 g, 0.022 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4.0 mL) was added dibenzyl diisopropylphosphoramidite (0.14 ml, 0.43 mmol) followed by 4,5-dicyanoimidazole (0.063 g, 0.54 mmol). The reaction was stirred at room temperature for 14 h and then cooled to 0 °C. 30% H<sub>2</sub>O<sub>2</sub>/H<sub>2</sub>O (2 ml) was added, and the reaction was stirred at 0 °C for another 1 h. The reaction was then quenched with saturated Na<sub>2</sub>SO<sub>3</sub> solution (10 ml) and the mixture extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 30 ml). The organic layers were combined, dried over sodium sulfate, and then concentrated under reduced pressure to afford a clear oil. The crude product was purified using silica gel flash chromatography (using Silica Gel 60 Å (32–63 μm)) eluting with a gradient of 0–55% ethyl acetate/hexanes to afford pure product as a clear oil (0.022 g, 86% yield): <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz, referenced to tetramethylsilane at 0.00 ppm) δ 7.37–7.05 (m, 30H), 5.07 (m, 1H), 4.99–4.84 (m, 6H), 4.75 (m, 2H), 4.58–4.38 (m, 4H), 4.34–4.27 (m, 1H), 4.14–3.92 (m, 5H), 3.84 (m, 2H), 3.74 (dd, *J* = 5.6, 6.0 Hz, 1H), 2.19 (m, 5H), 1.62 (m, 1H), 1.54 (m, 4H), 1.27 (m, 16H), 0.87 (m, 6H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz, referenced to tetramethylsilane at 0.00 ppm) δ 173.1, 173.0, 172.7, 138.4, 138.3, 138.2, 136.4, 136.3, 136.2, 136.1, 135.7, 135.6, 128.7, 128.6, 128.6, 128.6, 128.5, 128.5, 128.4, 128.4, 128.4, 128.2, 128.2, 128.0, 127.9, 127.8, 127.7, 127.7, 127.6, 127.6, 127.5, 127.4, 82.4, 80.4, 78.8, 74.9, 74.4, 72.6, 72.5, 72.3, 69.8, 69.7, 69.6, 69.6, 69.5, 69.4, 69.3, 61.8, 34.4, 34.3, 32.0, 30.3, 29.4, 29.3, 29.2, 25.2, 22.9, 14.4; <sup>31</sup>P NMR (CDCl<sub>3</sub>, 121 MHz, relative to an 85% H<sub>3</sub>PO<sub>4</sub> external standard) δ -0.28, -0.75, -0.85; IR (film, cm<sup>-1</sup>) 2927, 2855, 1742, 1455, 1270, 1213, 1155, 1012, 737, 697; TLC R<sub>f</sub> 0.12 (50% ethyl acetate/hexanes); exact mass calcd. for [C<sub>67</sub>H<sub>84</sub>O<sub>15</sub>P<sub>2</sub> + H]<sup>+</sup> requires *m/z* 1191.5364, found 1191.5448 (ESI+). The optical rotation was recorded on a Rudolf Research Analytical Autopol IV Automatic polarimeter at the sodium D line (path length 50 mm): [α]<sub>D</sub> = -3.4 (4.0, CHCl<sub>3</sub>).

**Protected *L*-3-deoxy-diC<sub>8</sub>PI(5)P**—Synthesis and spectral data matched that for protected *D*-3-deoxydiC<sub>8</sub>PI(5)P above. [α]<sub>D</sub> = +3.1 (4.0, CHCl<sub>3</sub>).

***D*-3-Deoxy-diC<sub>8</sub>PI(5)P**—To a stirred solution of benzyl protected *D*-3-deoxy-diC<sub>8</sub>PI(5)P (0.020 g, 0.017 mmol) in *t*-BuOH/H<sub>2</sub>O (5:1, 3 ml) was added sodium ion-exchange resin (Chelex 100 sodium form, 50–100 dry mesh, washed with H<sub>2</sub>O) followed by Pd(OH)<sub>2</sub>/C (20 mg, washed with H<sub>2</sub>O). The reaction was then stirred at 1 atm of H<sub>2</sub> for 32 h. The reaction mixture was filtered through Celite, and the filtrate was concentrated under reduced pressure and lyophilized to afford a white solid (0.011 g, 91% yield). <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 5.12 (m, 1H), 4.26 (m, 1H), 4.10 (m, 2H), 3.90 (m, 2H), 3.82 (m, 1H), 3.74 (m, 1H), 3.63 (m, 2H), 2.22 (m, 4H), 1.93 (m, 1H), 1.42 (m, 5H), 1.11 (m, 16H), 0.68 (m, 6H); <sup>31</sup>P NMR (D<sub>2</sub>O, 121 MHz) δ 4.3, 0.5; exact mass calcd. for [C<sub>25</sub>H<sub>48</sub>O<sub>15</sub>P<sub>2</sub> - H]<sup>-</sup> requires *m/z* 649.2390, found 649.2360 (ESI-); [α]<sub>D</sub> = +3.4 (1.0, H<sub>2</sub>O at pH = 9).

***L*-3-Deoxy-diC<sub>8</sub>PI(5)P**—Synthesis and spectral data matched that for 3-deoxy-diC<sub>8</sub>PI(5)P above. [α]<sub>D</sub> = -5.2 (1.0, H<sub>2</sub>O at pH = 9).

All the purified lipids were treated with Chelex to remove contaminating paramagnetic ions (introduced via the Pd catalyst used in generating the final product). Concentrations of stock solutions were measured by  $^{31}\text{P}$  NMR (202.3 MHz) spectroscopy by comparing phosphorus peak integration (in the absence of  $^1\text{H}$  decoupling) with a standard inorganic phosphate peak.

### $^{31}\text{P}$ NMR characterization of phospholipids

Most  $^{31}\text{P}$  NMR spectra (202.3 MHz) were obtained on a Varian INOVA 500 spectrometer. Variation of the  $^{31}\text{P}$  NMR linewidth for the synthetic PI as a function of lipid concentration (0.25 to 4 mM) was used to estimate the CMC values for several of these molecules in 50%  $\text{D}_2\text{O}$  containing 100mM Tris HCl, pH 8.0, with 2 mM EDTA. In this concentration range the  $^{31}\text{P}$  linewidth of most of these compounds typically decreased 3-4 Hz to a constant value when only monomers were present. The CMC was estimated from a plot of the linewidth versus reciprocal PI concentration as the intersection of the two lines (negative slope for the micelle/monomer combination and zero slope for only monomers). Assays of recombinant enzyme activities were also done by  $^{31}\text{P}$  NMR on the Varian INOVA 500 (details below for each enzyme).

High resolution  $^{31}\text{P}$  NMR field cycling at magnetic fields from 0.004 to 11.74 Tesla (T) was carried out using a home-built shuttling system on a Varian Unity<sup>PLUS</sup> 500 spectrometer.<sup>25, 26</sup> This technique was used to characterize two deoxy-diC<sub>8</sub>PI micelles and to explore if 0.5 mM  $\text{D}$ -diC<sub>8</sub>PI(3)P had detectable micelles when dispersed in the PTEN assay buffer. The sample, sealed in a standard 10 mm tube, was pneumatically shuttled to a higher position within, or just above, the magnet, where the magnetic field is between 0.06 and 11.7 T. Lower fields (down to 0.004 T for these samples) were accessible by shuttling to a region outside the magnet in the middle of a small Helmholtz coil located just above the top of the superconducting magnet. The spin-lattice relaxation rate,  $R_1$ , at each field strength was measured using 6-8 programmed delay times.

### Recombinant enzymes

Recombinant p110 $\alpha$ /p85 $\alpha$  PI-3-kinase was purchased from Upstate. A chimeric rat PLC with the catalytic domain of phospholipase C $\delta$ 1 and the N-terminal PH domain of phospholipase C $\beta$ 1 was the gift of Dr. Suzanne Scarlata, University of New York at Stony Brook. Recombinant *Listeria monocytogenes* phosphatidylinositol-specific PLC without its signal sequence<sup>27, 28</sup> was expressed and purified using the IMPACT-CN expression system from New England Biolabs. The construction of pET28a PTEN has been described previously<sup>29</sup>; *E. coli* BL21 (DE3) cells were used as the host for protein expression. PTEN protein was purified from the bacterial extracts by using a Qiagen Ni<sup>2+</sup>-nitrilotriacetic acid agarose column according to manufacturer protocol. Fractions of pure PTEN protein as judged on SDS-PAGE were combined and dialyzed against 100 mM Tris-HCl, pH 8.0. Protein concentration was determined by Lowry assay.<sup>30</sup>

### PI3K assay

PI3K (2  $\mu\text{g}$ ) was added to a reaction mix (400  $\mu\text{l}$ ) containing 50 mM Tris-HCl, pH 7.5, 0.1 mg/ml BSA (Sigma), 50 mM NaCl, 5 mM  $\text{MgCl}_2$ , 2 mM ATP, 1 mM  $\text{D}$ -diC<sub>8</sub>PI and 0 to 3 mM of the 3-deoxy-diC<sub>8</sub>PI analogues. The reaction was incubated at 22°C for 3 h and stopped by the addition of 10 mM EDTA. The phosphorylation of PI and PI analogues was monitored by  $^{31}\text{P}$  NMR spectroscopy using the phosphodiester resonance ( $\delta_{\text{P}} = 0.23$  ppm at pH 7.5) as the standard for integration of the PI3K phosphomonoester product. The acquisition conditions followed what has been previously used for PLC assays.<sup>31, 32</sup> The amount of product produced was quantified by comparing the phosphomonoester resonance for  $\text{D}$ -diC<sub>8</sub>PI(3)P to the phosphodiester peak in the absence of  $^1\text{H}$  decoupling.

## PLC assay

PLC activities toward diC<sub>8</sub>PI and the 3-deoxy-diC<sub>8</sub>PI analogues were measured by <sup>31</sup>P NMR spectroscopy after incubation for fixed time points as described for other PI-specific PLC enzymes.<sup>31, 33</sup> The assay buffer for PLCδ1 was 50 mM Tris-HCl, 0.5 mM CaCl<sub>2</sub>, pH 7.5; incubation was at 28°C. For the recombinant *L. monocytogenes* PLC, the assay buffer was 50 mM Tris-HCl, 0.5 mg/ml BSA, pH 7.0, and incubation at 25°C. Enzyme (6.9 to 27.6 μg of PLCδ1 or 0.04 to 176 μg of *L. monocytogenes* PLC) was added to each 200 μl sample and incubation time was chosen so that less than 20% PI cleavage occurred. The reaction was stopped by the addition of 200 μl CHCl<sub>3</sub> to the sample. Both cIP and I-1-P content in the aqueous phase were quantified in the <sup>31</sup>P NMR spectrum using added glucose-6-phosphate as an internal standard.

## PTEN assay

Phosphatase assays were carried out in 50 μl assay buffer (100 mM Tris-HCl, pH 8.0, 2 mM EDTA, and 10 mM DTT). The phosphatase reaction was initiated by adding ~25 μg of purified PTEN (10 μM) and quenched by adding the malachite green reagent containing 1 M HCl after a 20 min incubation at 37°C.<sup>34</sup> A comparison of observed A<sub>660</sub> changes to those for standard Pi samples was used to calculate the reaction rate. In most assays, 0.5 mM D-diC<sub>8</sub>PI(3)P was used as substrate and various concentrations (0.05 to 2 mM) of synthetic short-chain PIs were added to test their effect on phosphatase activity. For estimation of the K<sub>m</sub> for D-diC<sub>8</sub>PI(3)P, assays were carried out in 100 μl aliquots containing six different concentrations (0.05 to 1.6 mM) of substrate. Most assays were done at least in duplicate.

## U937 cell growth and incubation with PI analogues

The U937 human leukemic monocyte lymphoma cell line was obtained from the American Type Culture Collection (Manassas, VA). U937 cells were maintained in RPMI 1640 medium supplemented with 10% fetal bovine serum (FBS), penicillin (100 U/ml), streptomycin (100 μg/ml), 2-mercaptoethanol (50 μM), and glutamine (2 mM) at a cell density of 5 × 10<sup>5</sup> to 1 × 10<sup>6</sup> cells/ml (at 37 °C in a 5% CO<sub>2</sub> atmosphere at 95% humidity). Cells were cultured in the presence or absence of the short-chain diC<sub>8</sub>PI compounds at the indicated concentration. Cells were also incubated with 10 μM LY294002 and/or 20 nM wortmannin. At the appropriate time, 2 to 5 × 10<sup>5</sup> cells were collected by centrifugation at 400 × g for 8 min, washed in FACS buffer (1 × PBS containing 0.5% BSA and 0.01% sodium azide), and resuspended in FACS buffer containing 5 μg/ml propidium iodide. Samples were incubated on ice for 10 min then analyzed by flow cytometry using a BD FACSCanto flow cytometer with BD FACS Diva software (BD Biosciences, San Jose, CA).

## Preparation of cell extracts and western blot analysis

Cells were washed twice in PBS and then incubated for 20 min at ~10<sup>8</sup>/ml in RIPA buffer containing 1 mM sodium orthovanadate, 1 mM PMSF, protein inhibitor cocktail, 1 mM okadaic acid, 1 mM sodium fluoride and 10mM β-glycerophosphate, followed by 3 freeze/thaw cycles in dry ice. Lysates were centrifuged at 14,000 × g (15 min) to remove insoluble material and then protein (20 μg/lane) was separated by polyacrylamide-SDS gel electrophoresis and transferred to a PVDF membrane. The membrane was blocked in TBS-T (20 mM Tris, pH 7.6, 137 mM NaCl, and 0.05% Tween-20) containing 5% nonfat dry milk for 1 h and then incubated overnight (4°C) with primary Ab at 1 μg/ml in TBS-T. The membrane was washed several times in TBS-T, incubated with a 1:2500 dilution of anti-rabbit IgG-coupled horseradish peroxidase Ab (60 min) and developed by enhanced chemiluminescence (ECL). Bands for the different phosphorylated proteins were quantified by densitometric analysis, using the band corresponding to an extract from cells incubated with L-3-deoxy-diC<sub>8</sub>PI as a control since this compound had no effects on the U937 cells.



## RESULTS

### Characterization of diC<sub>8</sub>PI compounds

Three target enzymes that might be inhibited by (PI3K and PTEN) or degrade (PLC) 3-deoxy-PI molecules are soluble enzymes whose natural substrates are membrane constituents. However, in detailed kinetic studies with synthetic short-chain phospholipid substrates, these types of enzymes typically display a preference for molecules aggregated into a micelle compared to monomers in solution, a phenomenon termed ‘interfacial activation.’ Both phospholipase C and PI3K exhibit this enhanced activity with interfacial substrates / inhibitors.<sup>35, 36</sup> Since the 3-deoxy-diC<sub>8</sub>PI molecules, unlike the longer acyl chain synthetic 3-deoxy-PI molecules examined previously,<sup>7, 14</sup> can exist as both monomers and micelles in solution, we require information on the physical state of the short chain PI analogues, notably their CMC as well as roughly what size micelles they form. This is also critical information for determining the distribution of the 3-deoxy-diC<sub>8</sub>PI species in cells at concentrations where they may cause cell death. The critical micelle concentration (CMC) describes the concentration above which micelles form. All the dioctanoyl-PI derivatives used in this work have CMC values in 100 mM Tris HCl, pH 8, between 0.4 and 0.7 mM as measured by <sup>31</sup>P linewidth changes as a function of lipid concentration (Table 1).

Two synthetic PIs (3 mM each), *l*-3-deoxy-diC<sub>8</sub>PI and *l*-3,5-dideoxy-diC<sub>8</sub>PI, were examined at 3 mM by high resolution <sup>31</sup>P field cycling to get a sense of the size and dynamics of these micelles. This is a novel technique that is very sensitive to the aggregation state of the micelles.<sup>25, 26</sup> The *l*-isomers of deoxydiC<sub>8</sub>PIs were chosen since they had comparable CMC values to their enantiomers and more material was available. The field dependence of the <sup>31</sup>P R<sub>1</sub> from 0.1 up to 11.74 T for phospholipid aggregates can be analyzed in terms of a contribution from three terms: (i) dipolar relaxation associated with a ‘slow’ correlation time,  $\tau_c$ , and a relaxation rate extrapolated to zero field, R<sub>c</sub>(0) that is proportional to  $\tau_c$  and inversely proportional to  $r_{PH}^6$  ( $r_{PH}$  is the effective distance of the phosphorus from the protons which relax this nucleus), (ii) chemical shift anisotropy (CSA) relaxation associated with the same slow correlation time, and (iii) CSA relaxation due to a faster motion (whose dipolar counterpart would be small and negligible in the analysis of these data). R<sub>1</sub> is then the sum of those three terms.<sup>25</sup>  $\tau_c$  is determined directly while a correlation time for a faster motion ( $\tau_{hf}$ ) can be extracted from coefficients in the fits.<sup>25</sup> In samples where phospholipid vesicles or large rod-shaped micelles form (e.g., diC<sub>7</sub>PC), there is a distinct rise in R<sub>1</sub> detected at very low fields (<0.06 T).<sup>26</sup> When well-separated from the nanosecond dispersion, a correlation time,  $\tau_v$ , that may reflect the contribution of particle tumbling to relaxation can be extracted. It has been proposed that  $\tau_v$  is equal to the rotational correlation time D<sub>r</sub>/6 where D<sub>r</sub> is the rotational diffusion constant of the individual lipid due to both diffusion of the entire aggregate, and translational diffusion of individual lipids relative to the aggregate.<sup>26</sup>

For micelles as opposed to vesicles, a detailed analysis is complicated by the fact that monomers and micelles coexist in fast exchange and that many micelles are usually rod-shaped rather than spherical. Nonetheless, the line shape of the R<sub>1</sub> versus field profile can indicate that micelles form, and whether they are large (such that a low field rise in R<sub>1</sub> is observed) or are small (in which case no low field rise is detected, but a minimum in rate at 1-2 T, suggesting a several ns correlation time). If the compound exists in solution as a monomer or very small aggregates (clusters of under 10 molecules) a single sub-nanosecond correlation time is likely to describe the behavior and R<sub>1</sub> is invariant versus field below 2 T. This type of assay for micelles works well in complex buffer where other components can complicate other micelle detection methods (for instance impurities can skew surface tension measurements and dyes that fluoresce when partitioned into micelle environments can influence the measured CMC). The control for a monomeric phospholipid was 10 mM dibutyroylphosphatidylcholine (diC<sub>4</sub>PC) whose CMC is greater than 150 mM.<sup>37</sup> The monomer lipid clearly showed a rise in

$R_1$  as the field was increased above 2 T (Figure 2A). This profile could be fit with a *single* correlation time of  $0.75 \pm 0.08$  ns. The maximum dipolar contribution,  $R_c(0)$ , extrapolated from the low field constant contribution was  $0.077 \pm 0.006$  s<sup>-1</sup>. Both deoxy-diC<sub>8</sub>PI compounds showed higher  $R_1$  values and could not be well fit with a single correlation time, behavior indicative of aggregate formation (Figure 2A). With the simplistic but useful model-free analysis described previously,<sup>25</sup> the 3-deoxy-diC<sub>8</sub>PI at this concentration was characterized by a  $1.6 \pm 0.5$  ns correlation time as well as a faster motion ( $\tau_{hf} = 290 \pm 170$  ps); the dideoxy-PI exhibited similar behavior with a  $1.7 \pm 0.5$  ns  $\tau_c$  ( $\tau_{hf} = 500 \pm 340$  ps). Although the fast motions are not well defined, they must be included for reasonable fits to the data.

The ns  $\tau_c$  clearly indicates that micelles are forming for these concentrations of the 1-3-deoxy-diC<sub>8</sub>PI and the 1-3,5-dideoxy-diC<sub>8</sub>PI. A further enhanced relaxation rate below 0.1 T can be used to provide an average particle size from the high resolution field cycling. Below 0.1 T,  $R_1$  was invariant for 3-deoxydiC<sub>8</sub>PI but increased for the dideoxy-PI analog (Figure 2B). The correlation time for the 3,5-dideoxydiC<sub>8</sub>PI low field dispersion was  $55 \pm 27$  ns (this slower correlation time is defined as  $\tau_v$ ). If the micelles were spherical this would correspond to a radius of 38 Å. For comparison, diC<sub>7</sub>PC rod-shaped micelles exhibit a 200 to 350 ns  $\tau_v$  (depending on concentration) for the low field dispersion.<sup>38</sup> Clearly, the anionic deoxy-diC<sub>8</sub>PI micelles are considerably smaller. Since 3-deoxy-diC<sub>8</sub>PI did not exhibit a low field dispersion, its micelles must be even smaller in size and likely similar to those of dihexanoylphosphatidylcholine, which are 20-25 molecules per micelle and do not show a distinct low field dispersion.<sup>38</sup> Thus, as a second hydroxyl group is removed from the inositol headgroup, the micelles formed are larger, but not as large as diC<sub>7</sub>PC micelles.

In the PI3K and PLC assays, *D*-diC<sub>8</sub>PI substrate was present at 1 or 2 mM, a concentration above the CMC. Adding more diC<sub>8</sub>PI analogs at 1 mM or higher should increase the micellar fraction in solution. Given the field cycling results on the two deoxy-PI compounds, all mixed micelles with diC<sub>8</sub>PI are likely to be relatively small (compared to short-chain PC micelles). This information on the physical state of the short chain PI analogues in assay buffers is important for understanding their interactions with enzymes that may show enhanced activity with interfacial substrates / inhibitors.

For PTEN assays, a concentration of 0.5 mM *D*-diC<sub>8</sub>PI(3)P was optimal for detection of inhibition by the different deoxy-PI compounds. This is in the vicinity of the CMC for this compound (the CMC values for the phosphorylated lipids, *D*-diC<sub>8</sub>PI(3)P which is the substrate for PTEN, and *D*-3-deoxy-diC<sub>8</sub>PI(5)P, were not very accurate since the changes in linewidth with concentration were small – less than 2 Hz). To assess whether micelles were present in assay mixtures of substrate in the absence of protein, we determined  $R_1$  for 0.5 mM *D*-diC<sub>8</sub>PI(3)P dispersed in the PTEN assay buffer (100 mM Tris-HCl, pH 8.0, 2 mM EDTA, and 10 mM DTT) at several field strengths (Figure 2C). This low concentration taxes the sensitivity of the field cycling system so that a complete field dependence profile could not be obtained. However, it does provide insight into the status of the *D*-diC<sub>8</sub>PI(3)P in solution under the conditions used in the PTEN assays. Both the phosphodiester phosphorus and the phosphate group on inositol C(3) of *D*-diC<sub>8</sub>PI(3)P exhibit significant CSA relaxation at higher fields and  $R_1$  decreases as the field decreases. The phosphodiester resonance has behavior intermediate between monomeric and aggregated phospholipids while the phosphomonoester has low field data that resembles a monomer. The higher limiting  $R_1$  value at low fields for the phosphodiester compared to the phosphomonoester could reflect the contribution of more protons to  $R_1$  of the  $-\text{CH}_2\text{O}-\text{P}(\text{O})_2-\text{OCH}-$  compared to the  $-\text{CHOPO}_3^-$  group. Of more importance, a direct comparison of the *D*-diC<sub>8</sub>PI(3)P phosphodiester <sup>31</sup>P profile to the phosphodiester in 3-deoxy-diC<sub>8</sub>PI strongly suggests that at 0.5 mM in the PTEN assay mixture there may be some small micelles as well as monomers of the substrate present. Since all diC<sub>8</sub>PI compounds have similar CMCs, the addition of inhibitors to a fixed substrate

concentration (0.5 mM) is likely to increase the proportion of micelles. However, all inhibitors should have the same effect since their CMCs are essentially the same.

### Effect of phospholipase C enzymes on 3-deoxy-diC<sub>8</sub>PI molecules

PI-specific PLC enzymes cleave PI molecules in two steps: (i) initial formation of diacylglycerol and the water-soluble product cyclic-inositol-1,2-phosphate, followed by (ii) hydrolysis of the cIP to inositol-1-phosphate.<sup>39-41</sup> The initial 3-deoxy-PI studied as a cell growth inhibitor, *D*-3-deoxy-dipalmitoyl-PI, did not appear to be a substrate for PLC. However, that phospholipid has a high gel-to-liquid-crystalline phase transition temperature and may not have been presented in structures accessible to phospholipases in the *in vitro* assays. By using *D*-diC<sub>8</sub>PI micelles as the standard substrate, we can quantify the relative cleavage of the 3-deoxy-diC<sub>8</sub>PI substrates as well as examine their ability to inhibit the enzyme, which reflects their ability to bind to PLC active sites. We examined the 3-deoxy-PI molecules as substrates and inhibitors of two different phosphatidylinositol-specific PLC enzymes – Ca<sup>2+</sup>-dependent mammalian PLC $\delta$ 1 (this is a chimera of the  $\delta$ 1 catalytic domain and the  $\beta$ 1 PH domain and was chosen because it has moderately high activity *in vitro*)<sup>42</sup> and a Ca<sup>2+</sup>-independent bacterial PLC. Mechanistically, PI cleavage occurs in a similar fashion in both types of enzymes except that an Arg replaces the active site Ca<sup>2+</sup> in the bacterial enzyme.<sup>39, 40</sup> As seen in Table 2, removal of the hydroxyl group at C3 generated a very poor substrate for the PLC $\delta$  enzyme with PI cleavage occurring at 0.5 to 1% that of diC<sub>8</sub>PI. However, in the context of a cell and over the time course of several days, these lipids are likely to be hydrolyzed by the endogenous PLC enzymes.

These compounds were also not very good inhibitors of the PLC $\delta$ . The ratio of cIP to the final water-soluble product I-1-P reflects how well the intermediate is bound to the enzyme. A tighter binding cIP generated *in situ* translates to a lower ratio of cIP/I-1-P.<sup>33, 33, 43</sup> As hydroxyl groups are removed from the inositol ring, the intermediate cIP analogue becomes more hydrophobic and its release is slow compared to attack by water and production of I-1-P (Table 2). With *D*-3,5-dideoxy-diC<sub>8</sub>PI, no cIP was observed suggesting that the enzyme must hold the cyclic intermediate sufficiently long that it is always hydrolyzed (but very slowly). Addition of a phosphate to C(5) to produce *D*-3-deoxy-diC<sub>8</sub>PI(5)P did not generate a better substrate but it did bias the enzyme so that now cIP was the dominant product (compared to hydrolysis of *D*-3-deoxy-diC<sub>8</sub>PI). This is an interesting contrast to how the enzyme hydrolyzed phosphorylated glycerol-phosphoinositols,<sup>33</sup> where a phosphorylated substrate was more efficiently hydrolyzed and I-1-P became the major product. It strongly indicates that the 3-hydroxy group of the inositol ring must make important hydrogen bond contacts with the enzyme that stabilize binding of PI analogues to the protein but not in an optimal configuration for PI cleavage.

Mammalian PLC enzymes have multiple domains that could complicate / mask effects of the deoxy-diC<sub>8</sub>PI compounds. Therefore, for comparison we also examined the effect of these compounds on the PLC from *L. monocytogenes*. This bacterial PLC is essentially the catalytic domain of the mammalian enzymes and serves to assess the effects of the deoxy-PI compounds on catalysis only. The 3-deoxy-diC<sub>8</sub>PI compounds were also poor substrates and poor inhibitors for *L. monocytogenes* PLC (Table 3). The specific activities toward *D*-3-deoxy-diC<sub>8</sub>PI and *D*-3,5-dideoxy-diC<sub>8</sub>PI were 1.3 and 0.037% that toward diC<sub>8</sub>PI, respectively. Addition of 2 to 6 mM deoxy-PI compounds had only minor effects on the activity of *L. monocytogenes* PLC toward 2 mM *D*-diC<sub>8</sub>PI. For all the deoxy- compounds, the decrease in specific activity was less than 10%, comparable to the effect of *L*-diC<sub>8</sub>PI. As with the mammalian enzyme, removal of the 3-hydroxyl group generates a compound that binds very poorly to PLC and does not compete well with substrate.



### Effect of 3-deoxy-diC<sub>8</sub>PI molecules on PI3K activity

Removal of the 3-hydroxyl group from PI generates a molecule that should inhibit PI3K phosphorylation of *D*-diC<sub>8</sub>PI. Previous work<sup>7</sup> has suggested that the 3-deoxy-PI species are not very potent inhibitors of PI3K, although this can be misleading if substrate and inhibitor have different chain lengths or solubilities. In an attempt to keep substrate and inhibitors in the same physical state we investigated the activity of the p110 $\alpha$ /p85 $\alpha$  complex towards 1 mM *D*-diC<sub>8</sub>PI in the absence and presence of 3 mM 3-deoxy-diC<sub>8</sub>PI analogs (the high deoxy-PI concentration was chosen to maximize observation of any inhibition). Since PI3K has been shown to work in a scooting mode with PI dispersed in vesicles,<sup>36</sup> the concentrations of PIs were chosen so that substrate (and in most cases inhibitor) was micellar. In fact, kinetics attempts with 0.3 or 0.5 mM substrate and comparable concentrations of the deoxy-PI molecules exhibited either no effect or an increase in activity (and a high degree of error) likely due to formation of micelles and better presentation of substrate. As shown in Table 4, both *D*- and *L*-3-deoxy-diC<sub>8</sub>PI inhibited the phosphorylation of diC<sub>8</sub>PI; removing a second hydroxyl group (data not shown) or adding a phosphate to the 5-hydroxyl group reduced the potency of these compounds as inhibitors so that very little inhibition was observed under these conditions of excess inhibitor. For comparison, we also examined the inhibitory effect of *L*-diC<sub>8</sub>PI on p110 $\alpha$ /p85 $\alpha$  activity toward *D*-diC<sub>8</sub>PI. The substrate enantiomer had no effect on *D*-diC<sub>8</sub>PI phosphorylation, indicating that the observed inhibition for the 3-deoxy-diC<sub>8</sub>PI molecules is not just surface dilution inhibition but represents a real binding effect of the enzyme with PI3K. The replacement of the 3-hydroxyl of the inositol ring of diC<sub>8</sub>PI with hydrogen (deoxygenation) not only improved the binding of 3-deoxy-diC<sub>8</sub>PI compounds to PI3K (compared to diC<sub>8</sub>PI) but also eliminated the inositol ring stereoisomer selectivity for ligand binding. A phosphorylated 3-deoxy-diC<sub>8</sub>PI (*D*-3-diC<sub>8</sub>PI(5)P) was also examined for inhibition of diC<sub>8</sub>PI phosphorylation by PI3K. Enzyme activity decreased only 20% with an excess of *D*-3-diC<sub>8</sub>PI(5)P. These *in vitro* assays suggest that the 3-deoxydiC<sub>8</sub>PI molecules, especially when monomeric, are unlikely to have a large effect on PI3K *in vivo*. They also suggest that if PI3K is involved *in vivo*, both *D*- and *L*-3-deoxy-diC<sub>8</sub>PI should have similar effects as growth inhibitors.

### Effect of 3-deoxy-diC<sub>8</sub>PI molecules on PTEN activity

Although PTEN is often mutated in some tumor cells, in others it could be a potential target for inhibition by the short-chain 3-deoxy-diC<sub>8</sub>PI molecules. This would not be productive if the aim of the treatment is to induce cell death of the tumor cells. To check this, we examined all the different 3-deoxy-PI molecules as inhibitors of PTEN (Figure 3). The substrate in this case is *D*-diC<sub>8</sub>PI(3)P at 0.5 mM, a concentration in these assay mixtures where there is a mixture of monomers and some small micelles. The dependence of specific activity on the concentration of this substrate exhibited an apparent  $K_m$  of  $0.20 \pm 0.07$  mM and a  $V_{max}$  of  $14 \pm 1$  nmol min<sup>-1</sup> mg<sup>-1</sup> at 37°C. The different 3-deoxy-PI compounds as well as diC<sub>8</sub>PI, from 0.05 up to 2.5 mM, were examined as inhibitors of PTEN (IC<sub>50</sub> values in Table 1). At inhibitor concentrations >0.5 mM (compared to 0.5 mM substrate), all the synthetic PI molecules were inhibitory, although the *L*-phosphatidylinositols (Figure 3B) were usually poorer inhibitors than the *D*-inositol lipids (Figure 3A). The compound *D*-3-deoxy-diC<sub>8</sub>PI was the most potent inhibitor towards 0.5 mM substrate with an IC<sub>50</sub> of 0.23 mM under these conditions and consistent with tighter binding than reflected in the substrate apparent  $K_m$ . Two very interesting kinetic trends were observed: (i) as the 3- and 5-hydroxy groups were replaced with hydrogen, the *L*-diC<sub>8</sub>PI derivatives became more potent inhibitors with *L*-3,5-dideoxy-diC<sub>8</sub>PI more inhibitory below 1 mM than its enantiomer, and (ii) at 0.1-0.3 mM two of the compounds, *D*-diC<sub>8</sub>PI and *D*-3-deoxy-diC<sub>8</sub>PI(5)P, activated PTEN 20-40% for *D*-diC<sub>8</sub>PI(3)P hydrolysis. The *L*-enantiomers of these compounds had no activating effect in this concentration range. Activation of PTEN by PI(4,5)P<sub>2</sub> has been reported previously.<sup>22</sup> In our system, where all PIs have similar CMC values, we can suggest that the non-substrate *D*-3-deoxy-diC<sub>8</sub>PI(5)P must be binding as a monomer in a

stereospecific manner to PTEN to enhance its hydrolysis of diC<sub>8</sub>PI(3)P, since activation is seen at low total PI concentrations. Even if some substrate aggregation occurs, the other non-activating PI molecules are likely to have the same effects on the physical distribution of substrate between monomer and very small micelles. Since they do not activate PTEN, the effect must be due to a specific PTEN/ *D*-3-deoxydiC<sub>8</sub>PI(5)P interaction. However, extrapolating these results to the *in vivo* situation, we would predict that at 3-deoxy-PI concentrations <0.05 mM, there should be little inhibition of PTEN; any activation by diC<sub>8</sub>PI compounds is also likely to be small under these conditions.

### Effect of 3-deoxy-diC<sub>8</sub>PI compounds on the growth of the U937 human lymphoma cell line

The longer chain *D*-3-deoxy-PI analogs are thought to inhibit cell growth by competing with PI(3)P for binding to the Akt1 PH domain and reducing Akt1 translocation to the plasma membrane.<sup>7</sup> The short-chain 3-deoxy-diC<sub>8</sub>PI compounds could affect cell survival in a similar fashion. The difference is that the short-chain PI analogs at low concentrations are likely to be monomeric in the cell as opposed to partitioning into membranes (expected for the long-chain PI analogs). There may also be specificity among the different deoxy-PI compounds that would shed light on key interactions of these modified PIs with diverse targets. To explore this, we incubated the human leukemic monocyte lymphoma cell line U937 with the compounds *D*-3-deoxy-diC<sub>8</sub>PI, *L*-3-deoxy-diC<sub>8</sub>PI, *D*-3,5-dideoxy-diC<sub>8</sub>PI, and *L*-3,5-dideoxy-diC<sub>8</sub>PI at various concentrations, and assessed the viability of the cells at different time points. Of all the compounds tested, only *D*-3-deoxy-diC<sub>8</sub>PI had any significant effect on cell viability, with an IC<sub>50</sub> of 40 μM (Figure 4). The other compounds had no measurable effect on viability at concentrations up to 200 μM.

To determine if the 3-deoxy-diC<sub>8</sub>PI compounds inhibited endogenous signaling via the PI3K pathway, we assessed phosphorylation of Akt1, a downstream target of PI3K, and compared the effects of the short-chain PI compounds (50 μM) to the effects of LY294002 (10 μM) and wortmannin (20 nM), known inhibitors of signaling via the PI3K/Akt pathway (Figure 5). Since *L*-3-deoxy-PI had no effect on cell growth, the intensity of the band from cells incubated with that compound was used as a phosphorylation control. Phosphorylation of the transcriptional initiation factor eIF4E was clearly reduced after incubation of the cells with *D*-3-deoxy-diC<sub>8</sub>PI for 24 h. However, reduction in Akt phosphorylation appeared to show interesting specificity. The phosphorylation of Thr308 did not appear to be affected by *D*-3-deoxy-diC<sub>8</sub>PI, while LY294002 and wortmannin did lead to some reduced phosphorylation in the same time period (and the combination of the two was extremely effective). In contrast, Ser473 phosphorylation was reduced (more than with the two other known inhibitors of the PI3K/Akt pathway at this time point) when the cells were incubated with *D*-3-deoxy-diC<sub>8</sub>PI. In previous work with *D*-3-deoxy-PI molecules,<sup>7</sup> Akt Ser473 phosphorylation was significantly reduced (to about 65% of the maximum phosphorylation for the *D*-3-deoxy-diC<sub>16</sub>PI). Here we show that the more soluble short-chain *D*-3-deoxy-diC<sub>8</sub>PI can also reduce Akt1 phosphorylation on Ser473. Whether it does so by preventing Akt translocation or by altering the interaction of the protein with the membrane remains to be seen. That this soluble, specific enantiomer, and not further deoxygenated compounds, is cytotoxic and affects Akt1 phosphorylation is an exciting observation.

## DISCUSSION

Compounds based on *D*-3-deoxy-PI have been examined as cell growth inhibitors for a variety of cancer cells.<sup>12-14</sup> The first analogue examined, *D*-3-deoxy-diC<sub>16</sub>PI, inhibited the growth of NIH3T3 mouse fibroblasts with an IC<sub>50</sub> of 17.6 μM.<sup>7</sup> Modifications of the 3-deoxy-PI to increase potency included replacing acyl chains with ether linked chains, and shortening the chain attached to the glycerol sn-2 position.<sup>14</sup> The ether bonds should make the 3-deoxy-PI

compounds stable to PLA<sub>2</sub> type activities, although most of the mammalian PLA<sub>2</sub> enzymes are not selective for PI headgroups. Work by Tabellini et al.<sup>15</sup> showed that at 5 μM the ether-linked 3-deoxy-PI analogues (2-substituted, 3-deoxy-PIs) had a small effect by themselves but enhanced induction of apoptosis when administered with either etoposide or cytarabine. At this concentration they were not toxic to human umbilical cells, although doubling the concentration of the deoxy-PI caused a pronounced increase in cell apoptosis. Shortening the sn-2 chain length may make also a more significant contribution to molecule potency. This creates a more lyso-like phospholipid that has a higher solution concentration than the diacyl long chain lipids. However, other structural aspects of 3-deoxy-PI that make it cytotoxic have not been studied.

In this work we have used synthetic diC<sub>8</sub>PI compounds, both as substrates for different target enzymes and to screen for 3-deoxy-PI inhibition *in vitro*. Knowing the CMC values for these lipids allows us to carefully interpret *in vitro* assays. In the case of PI3K and PLC enzymes, the assays use mixed micelles to assess analog inhibitory potency. For PTEN, assays have a mix of monomers and some small micelles. We have examined how inositol stereochemistry as well as specific hydroxyl groups at C(3) and C(5) on the ring contribute to the cytotoxicity of this class of phosphatidylinositols. The 3-deoxy-diC<sub>8</sub>PI molecules are poor inhibitors of PI3K and only moderate inhibitors of PTEN. They are neither good substrates nor inhibitors of PI-PLC enzymes, strongly indicating that these enzymes are not the *in vivo* target(s). The compound *D*-3-deoxy-diC<sub>8</sub>PI, and only that PI in this series, is cytotoxic to U937 cell. These leukemic cells provide a good system to evaluate effectiveness of the diverse 3-deoxy-PIs because they have an activated PI3K/Akt pathway. Since neither the enantiomer of *D*-3-deoxy-diC<sub>8</sub>PI nor *D*-3,5-dideoxy-diC<sub>8</sub>PI were cytotoxic, the *in vivo* target must bind *D*-3-deoxy-diC<sub>8</sub>PI in a very specific way.

The mechanism of action of the 3-deoxy-PIs has been proposed to be the binding of this lipid to the PH domain of Akt1.<sup>7</sup> This binding is then supposed to prevent targeting of Akt1 to the plasma membrane for phosphorylation and subsequent activation. Indeed, studies have shown that this molecule inhibits growth of cell lines where Akt1 is the major isoform of this protein kinase.<sup>12-14</sup> Akt1 is phosphorylated on Ser473 and Thr308; treatment of cells with the 3-deoxy-diC<sub>16</sub>PI and the ether-linked long chain PI analogue DPEL showed that phosphorylation of the Ser sites was reduced.<sup>7</sup> Other lipid-based antitumor compounds have been shown to alter Thr308 phosphorylation,<sup>15</sup> so it is not clear whether one or both phosphorylation sites are critical for Akt1 activation. In fact, one can ask if the reduced phosphorylation of Akt1 really represents the mechanism by which the 3-deoxy-PI molecules inhibit cell growth. All the previously examined modified phospholipids are relatively hydrophobic and could presumably be integrated into different intracellular membranes, attracting the Akt1 to a membrane devoid of its activating kinase. By using short-chain synthetic 3-deoxy-PI compounds, which should be monomeric in the cell at concentrations less than 0.1 mM, and showing that only *D*-3-deoxy-diC<sub>8</sub>PI is inhibitory *in vivo*, we provide data indicating that a very specific interaction of the 3-deoxy-PI with its target occurs. Inositol ring attachment to the glycerol backbone (only the *D*- compound is inhibitory) is critical as is the C(5) hydroxyl group. The observation that an Akt1 downstream target (eIF4E) exhibited reduced phosphorylation, strongly points to Akt1 as the likely *in vivo* target of *D*-3-deoxy-diC<sub>8</sub>PI. Furthermore, since Akt1 phosphorylation on Ser473 was reduced while that on Thr308 was unaffected, we might propose that this lipid analog has its primary effect in modulating Ser phosphorylation. Interestingly, recent work also suggests that phosphatidylinositol ether lipid analogues activate the proapoptotic stress kinase p38α that is a subgroup of the MAPK family and activated by phosphorylation by MAPKKs.<sup>16</sup> This particular interaction could also contribute to the cytotoxicity of 3-deoxy-PIs. With the *in vivo* specificity profile in hand for the soluble, synthetic 3-deoxy-diC<sub>8</sub>PI analogs, one might further explore that kinase *in vitro* to assess direct effects of the 3-deoxy-PIs.

Although there is no direct evidence for an Akt1 PH domain complex with 3-deoxy-PI, there is a crystal structure of the PH domain with a polyphosphorylated inositol bound. The crystal structure of the Akt PH domain with I(1,3,4,5)P<sub>4</sub> bound<sup>44, 45</sup> shows strong interactions of the protein with the 3- and 4-phosphate groups (not the 5-phosphate which was poised towards solvent and not tightly held). Given the importance of the phosphate monoester interactions in this structure, it might be surprising that the 3-deoxy-PIs bind at all to the PH domain. The interaction of the Akt1 PH domain with the inositol ring of the 3-deoxy compounds is likely to be quite different than its interaction with I(1,3,4,5)P<sub>4</sub>. Indeed, previous modeling studies of a 3-deoxy-PI binding to the Akt1 PH domain showed strong H-bonds of the inositol 4- and 5-hydroxyl groups with the protein.<sup>7</sup> Consistent with the modeling, we have shown that, while *D*-3-deoxy-diC<sub>8</sub>PI inhibits U937 cell growth, *D*-3,5-dideoxy-diC<sub>8</sub>PI is no longer cytotoxic. Given the very cationic nature of the Akt1 PH domain binding site, one might expect a phosphate at C(5) in *D*-3-deoxy-diC<sub>8</sub>PI to enhance analogue binding. However, we observed no growth inhibition by *D*-3-deoxy-diC<sub>8</sub>PI(5)P. It is possible that such a compound was not taken up by the U937 cells, since intracellular phosphatases would have been expected to remove the phosphate and generate *D*-3-deoxy-diC<sub>8</sub>PI, which is inhibitory. *D*-3-Deoxy-diC<sub>8</sub>PI, unlike the longer chain length analogues studied previously, will be monomeric in the tumor cells at the concentrations where it was found to be cytotoxic. This observation decouples membrane partitioning of the modified PI from specific target / 3-deoxy-PI interactions. The increased solubility without a decrease in tumor potency (compared to the ester-linked long-chain 3-deoxy-PI) may also be important for reducing toxic effect in normal cells. Tantalizing to this possibility, in preliminary experiments where 0.2 mM *D*-3-deoxy-diC<sub>8</sub>PI was incubated with primary B cells (which have all three Akt forms), there was little loss of cell viability (M. Pu, D. Blair, and T. Chiles, unpublished results). Clearly, the area is ripe for further investigation with well-defined soluble 3-deoxy-PI molecules.

## Acknowledgments


We would like to thank Dr. Suzanne Scarlata for the chimeric PLC $\delta$ 1 and Dr. Wonhwa Cho for the plasmid containing PTEN. This work was supported by NIH grants GM60418 (M.F.R.), GM68649 (S.J.M.), GM077974 (A.G.R.), and AI49994 (T.C.C.) and N.S.F. MCB-0517381 (M.F.R.).

## REFERENCES


1. Vivanco I, Sawyers CL. Nat. Rev. Cancer 2002;2:489–501. [PubMed: 12094235]
2. Seufferlein T. Int. J. Gastronint. Cancer 2002;31:15–21.
3. Ghosh PM, Malik S, Bedolla R, Kreisberg JL. Curr. Drug Metab 2003;4:487–496. [PubMed: 14683476]
4. Barnett S, Bilodeau M, Lindsley C. Curr. Top. Med. Chem 2005;5:109–125. [PubMed: 15853641]
5. Kandel ES, Hay N. Exp. Cell Res 1999;253:210–229. [PubMed: 10579924]
6. Qiao L, Nan F, Kunkel M, Gallegos A, Powis G, Kozikowski AP. J. Med. Chem 1998;41:3303–3206. [PubMed: 9719581]
7. Meuillet EJ, Mahadevan D, Vankayalapati H, Berggren M, Williams R, Coon A, Kozikowski AP, Powis G. Molec. Cancer Therap 2003;2:389–399. [PubMed: 12700283]
8. Kim D, Dan HC, Park S, Yang L, Liu Q, Kaneko S, Ning J, He L, Yang H, Sun M, Nicosia SV, Cheng JQ. Front. Biosci 2005;10:975–987. [PubMed: 15569636]
9. Li J, Yen C, Liaw D, Podsypanima K, Bose S, Wang SI, Puc J, Millaresis C, Rodgers L, McComble R. Science 1997;275:1943–1947. [PubMed: 9072974]
10. Steck PA, Pershouse MA, Jasser SA, Ung WK, Lin H, Ligon A. Nat. Genet 1997;15:356–362. [PubMed: 9090379]
11. Gills JJ, Dennis PA. Expert Opin. Investig. Drugs 2004;13:787–797.
12. Kozikowski AP, Kiddle JJ, Frew T, Berggren M, Powis G. J. Med. Chem 1995;38:1053–1056. [PubMed: 7707307]

13. Kozikowski AP, Sun H, Brognard J, Dennis PA. *J. Am. Chem. Soc* 2003;125:1144–1145. [PubMed: 12553797]
14. Andresen TL, Skytte DM, Madsen R. *Org. Biomol. Chem* 2004;2:2951–2957. [PubMed: 15480460]
15. Tabellini G, Tazzari PL, Bortul R, Billi AM, Cocco L, Martelli AM. *Br. J. Haemat* 2004;126:574–582. [PubMed: 15287952]
16. Gills JJ, Holbeck S, Hollingshead M, Hewitt SM, Kozikowski AP, Dennis PA. *Mol. Cancer Ther* 2006;2006;5:713–722. [PubMed: 16546986]
17. Ruiter GA, Zerp SF, Bartelink H, van Bitterswijk WJ, Verheij M. *Anticancer Drugs* 2003;14:167–173. [PubMed: 12569304]
18. Kondapaka SB, Singh SS, Dasmahapatra GP, Sausville EA, Roy KK. *Mol. Cancer Ther* 2003;2:1093–1103. [PubMed: 14617782]
19. Castillo SS, Brognard J, Petukhov PA, Zhang C, Tsurutani J, Granville CA, Li M, Jung M, West KA, Gills JJ, Kozikowski AP, Dennis PA. *Cancer Res* 2004;64:2782–2792. [PubMed: 15087394]
20. Lemieux RU. *Acc. Chem. Res* 1996;29:373–380.
21. Tor Y. *ChemBioChem* 2003;4:998–1007. [PubMed: 14523917]
22. Campbell RB, Liu F, Ross AH. *J. Biol. Chem* 2003;278:33617–33620. [PubMed: 12857747]
23. Sculimbrenne BR, Xu Y, Miller SJ. *J. Am. Chem. Soc* 2004;126:13182–13183. [PubMed: 15479046]
24. Xu Y, Sculimbrenne BR, Miller SJ. *J. Org. Chem* 2006;71:4919–4928. [PubMed: 16776522]
25. Roberts MF, Redfield AG. *J. Amer. Chem. Soc* 2004;126:13765–13777. [PubMed: 15493936]
26. Roberts MF, Redfield AG. *Proc. Natl. Acad. Sci. U.S.A* 2004;101:17066–17071. [PubMed: 15569928]
27. Goldfine H, Kolb C. *Infect. Immun* 1992;60:4059–4067. [PubMed: 1398918]
28. Ryan M, Zaikova TO, Keana JF, Goldfine H, Griffith OH. *Biophys. Chem* 2002;101-102:347–358. [PubMed: 12488013]
29. Das S, Dixon JE, Cho W. *Proc. Natl. Acad. Sci. U.S.A* 2003;100:7491–7496. [PubMed: 12808147]
30. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. *J. Biol. Chem* 1951;193:265–275. [PubMed: 14907713]
31. Zhou C, Wu Y, Roberts MF. *Biochemistry* 1997;36:347–355. [PubMed: 9003187]
32. Feng J, Bradley WD, Roberts MF. *J. Biol. Chem* 2003;278:24651–24657. [PubMed: 12714598]
33. Wu Y, Perisic O, Williams RL, Katan M, Roberts MF. *Biochemistry* 1997;36:11223–11233. [PubMed: 9287165]
34. Itaya K, Ui M. *Clin. Chim. Acta* 1966;14:361–366. [PubMed: 5970965]
35. Lewis KA, Garigapati VR, Zhou C, Roberts MF. *Biochemistry* 1993;32:8836–8841. [PubMed: 8395883]
36. Barnett SF, Ledder LM, Stirdivant SM, Ahern J, Conroy RR, Heimbrook DC. *Biochemistry* 1995;34:14254–14262. [PubMed: 7578025]
37. Bian J, Roberts MF. *J. Coll. & Int. Sci* 1992;153:420–428.
38. Shi, X.; Shao, C.; Zhang, X.; Zambonelli, C.; Redfield, AG.; Head, JF.; Seaton, BA.; Roberts, MF. 2007. submitted for publication
39. Griffith OH, Ryan M. *Biochim. Biophys. Acta* 1999;1441:237–254. [PubMed: 10570252]
40. Hondal RJ, Zhao Z, Kravchuk AV, Liao H, Riddle SR, Yue X, Bruzik KS, Tsai MD. *Biochemistry* 1998;37:4568–4580. [PubMed: 9521777]
41. Heinz DW, Essen LO, Williams RL. *J. Mol. Biol* 1998;275:635–650. [PubMed: 9466937]
42. Feng J, Roberts MF, Drin G, Scarlata S. *Biochemistry* 2005;44:2577–2584. [PubMed: 15709770]
43. Liu Y, Bruzik KS, Ananthanarayanan B, Cho W. *Bioorg. Med. Chem* 2003;11:2471–2475. [PubMed: 12735994]
44. Thomas CC, Deak M, Alessi DR, van Aalten DMF. *Curr. Biol* 2002;12:1256–1262. [PubMed: 12176338]
45. Kumar CC, Madison V. *Oncogene* 2005;24:7493–7501. [PubMed: 16288296]







**Figure 1.** Structures of 3-deoxy-dioctanoylphosphatidylinositol (3-deoxy-diC<sub>8</sub>PI) compounds synthesized.




**Figure 2.**  $^{31}\text{P}$  field cycling profiles for deoxy- $\text{diC}_8\text{PI}$  compounds. Dependence of spin-lattice relaxation rate,  $R_1$ , on magnetic field for 3 mM  $L$ -3-deoxy- $\text{diC}_8\text{PI}$  ( $\bullet$ ), 3 mM  $L$ -3,5-dideoxy- $\text{diC}_8\text{PI}$  ( $\circ$ ), and 10 mM  $\text{diC}_4\text{PC}$  ( $\Delta$ ) in 50 mM HEPES, pH 7.5, in the field range (A) 0.1 to 11.74 T, and (B) 0.03 to 1 T. (C)  $R_1$  for the phosphodiester  $^{31}\text{P}$  ( $\bullet$ ) and phosphate monoester ( $\circ$ ) on the inositol C(3) of 0.5 mM  $D$ - $\text{diC}_8\text{PI}(3)\text{P}$  dissolved in PTEN assay buffer. The solid line indicates the relaxation profile for micellar  $L$ -3-deoxy- $\text{diC}_8\text{PI}$  while the dashed line indicates field dependence for the  $\text{diC}_4\text{PC}$  monomer.



**Figure 3.** Effect of diC<sub>8</sub>PI derivatives on the activity of PTEN toward 0.5 mM D-diC<sub>8</sub>PI(3)P. In (A) are shown the effects of D- enantiomers and in (B) the effects of the corresponding L- enantiomers: diC<sub>8</sub>PI (●), 3-deoxy-diC<sub>8</sub>PI (○), 3,5-dideoxy-diC<sub>8</sub>PI (■), and 3-deoxy-diC<sub>8</sub>PI(5)P (□).





**Figure 4.** Viability of U937 cells after incubation with different concentrations of 3-deoxy-diC<sub>8</sub>PI compounds for 24 h: D-3-deoxy-diC<sub>8</sub>PI (●), L-3-deoxy-diC<sub>8</sub>PI (○), and D-3-diC<sub>8</sub>PI(5)P (X).




**Figure 5.** Western blots used to monitor phosphorylation of Akt1 (at both Ser473 and Thr308) as well as eIF4E (Ser209) after 24 h incubation with 50  $\mu$ M D-3-deoxy-diC<sub>8</sub>PI (D-3-d-PI), 50  $\mu$ M L-3-deoxydiC<sub>8</sub>PI (L-3-d-PI), 10  $\mu$ M LY 294002 (LY) or 20 nM wortmannin (WM) or a combination LY 294002 and wortmannin (LY+W). The numbers under each lane represent the fractional phosphorylation in the presence of the indicated compound compared to phosphorylation in the presence of L-3-deoxy-diC<sub>8</sub>PI, which has no effect on the cells.




**Scheme 1.**

General scheme for the synthesis of different enantiomers of inositol phosphates for use in generating diC<sub>8</sub>PI derivatives.



a. Phenyl chlorothionioformate, DMAP, Pyridine, CH<sub>2</sub>Cl<sub>2</sub>; 70% yield; b. TBSOTf, 2,6-lutidine, CH<sub>2</sub>Cl<sub>2</sub>; 90% yield; c. AIBN, Bu<sub>3</sub>SnH, Toluene; 70% yield; d. BnOH, NaH, THF; 91% yield; e. HF-Pyridine, THF; 75% yield; f. LiBr, acetone; g. DEAD, Ph<sub>3</sub>P, **3a**, THF; 89% yield for 2 steps; h. Dibenzyl diisopropylphosphoramidite, 4,5-dicyanoimidazole, CH<sub>2</sub>Cl<sub>2</sub>, then H<sub>2</sub>O<sub>2</sub>; 86% yield; i. Pd(OH)<sub>2</sub>/C, tBuOH, H<sub>2</sub>O, Na<sup>+</sup> ion exchange resin, H<sub>2</sub>; 91% yield.



### Scheme 2.

Synthetic scheme for generation of *D*-3-deoxy-diC<sub>8</sub>PI(5)P and *L*-3-deoxy-diC<sub>8</sub>PI(5)P.

**Table 1**

CMC values for dioctanoyl-PI compounds and IC<sub>50</sub> for recombinant PTEN hydrolysis of D-3-deoxy-diC<sub>8</sub>PI(3)P.

Phospholipids	CMC (mM) <sup>a</sup>	PTEN IC <sub>50</sub> (mM) <sup>b</sup>
D-diC <sub>8</sub> PI	0.5±0.1	0.43
L-diC <sub>8</sub> PI	0.5±0.1	> 2.5
D-diC <sub>8</sub> PI(3)P	0.7±0.2	
D-3-deoxy-diC <sub>8</sub> PI	0.42±0.08	0.23
L-3-deoxy-diC <sub>8</sub> PI	0.40±0.07	1.5
D-3,5-dideoxy-diC <sub>8</sub> PI	0.47±0.05	0.86
L-3,5-dideoxy-diC <sub>8</sub> PI	0.47±0.07	0.38
D-3-deoxy-diC <sub>8</sub> PI(5)P	0.61±0.15	0.47
L-3-deoxy-diC <sub>8</sub> PI(5)P	-	> 2.5

<sup>a</sup>Determined by analysis of <sup>31</sup>P NMR linewidth in D<sub>2</sub>O.

<sup>b</sup>Determined with 0.5 mM D-3-deoxy-diC<sub>8</sub>PI(3)P as the substrate concentration.

**Table 2**Recombinant PLC $\delta$ 1 activity toward diC<sub>8</sub>PI and deoxy-diC<sub>8</sub>PI lipids.

<u>Substrate<sup>a</sup></u>	<u>Inhibitor (mM)<sup>b</sup></u>	<u><math>\frac{\text{cIP}}{\text{I-1-P}}</math></u>	<u>Specific Activity<sup>c</sup> (<math>\frac{\mu\text{mol}}{\text{min}\cdot\text{mg}}</math>)</u>	<u>Relative Activity</u>
D-diC <sub>8</sub> PI		0.93	3.68	1.00
D-3-deoxy-diC <sub>8</sub> PI		0.22	0.039	0.011
D-3,5-dideoxy-diC <sub>8</sub> PI		0.00	0.018	0.005
D-3-deoxy-diC <sub>8</sub> PI(5)P		3.01	0.022	0.006
D-diC <sub>8</sub> PI	D-3-deoxy-diC <sub>8</sub> PI (2)	1.99	3.01	0.82
D-diC <sub>8</sub> PI	D-3-deoxy-diC <sub>8</sub> PI5P (6)	1.29	4.81	1.31
D-diC <sub>8</sub> PI	D-3,5-dideoxy-diC <sub>8</sub> PI (2)	1.77	3.25	0.88
D-diC <sub>8</sub> PI	D-3,5-dideoxy-diC <sub>8</sub> PI (6)	0.99	3.93	1.07
D-diC <sub>8</sub> PI	L-diC <sub>8</sub> PI (2)	1.02	3.08	0.84

<sup>a</sup>Substrates present at 2 mM.<sup>b</sup>The value in parentheses represents the concentration of inhibitor used in these assays.<sup>c</sup>Errors in specific activity typically <20%.

**Table 3**

Recombinant *L. monocytogenes* PI-PLC activity toward D-diC<sub>8</sub>PI molecules and inhibition by deoxy-diC<sub>8</sub>PI lipids.

<u>Substrate<sup>a</sup></u>	<u>Inhibitor (mM)<sup>b</sup></u>	<u>Specific Activity<sup>c</sup> (<math>\mu\text{mol min}^{-1} \text{mg}^{-1}</math>)</u>	<u>Relative Activity</u>
D-diC <sub>8</sub> PI		489	1.00
D-3-deoxy-diC <sub>8</sub> PI		6.44	0.013
D-3,5-dideoxy-diC <sub>8</sub> PI		0.18	$3.7 \times 10^{-4}$
D-diC <sub>8</sub> PI	D-3-deoxy-diC <sub>8</sub> PI (2)	522	1.07
D-diC <sub>8</sub> PI	D-3-deoxy-diC <sub>8</sub> PI(5)P (6)	452	0.92
D-diC <sub>8</sub> PI	D-3,5-dideoxy-diC <sub>8</sub> PI (2)	522	1.07
D-diC <sub>8</sub> PI	D-3,5-dideoxy-diC <sub>8</sub> PI (6)	539	1.10
D-diC <sub>8</sub> PI	L-diC <sub>8</sub> PI (2)	441	0.90

<sup>a</sup>Substrates present at 2 mM.

<sup>b</sup>The value in parentheses represents the concentration of inhibitor used in these PI-PLC assays.

<sup>c</sup>Errors in specific activity were  $\leq 15\%$ .



**Table 4**

Recombinant PI3K (P110 $\alpha$ /P85 $\alpha$ ) activity toward D-diC<sub>8</sub>PI (1 mM) and inhibition by L- and D- deoxy-diC<sub>8</sub>PI lipids.<sup>a</sup>

<b>Inhibitor</b>	<b>(mM)</b>	<b>Specific Activity<sup>b</sup> (<math>\mu\text{mol min}^{-1} \text{mg}^{-1}</math>)</b>	<b>Relative Activity</b>
–		0.60	1.00
D-3-deoxy-diC <sub>8</sub> PI	3	0.09	0.15
D-3-deoxy-diC <sub>8</sub> PI(5)P	3	0.49	0.81
L-diC <sub>8</sub> PI	3	0.60	1.00
L-3-deoxy-diC <sub>8</sub> PI	3	0.11	0.18

<sup>a</sup> Assay conditions included 1 mM D-diC<sub>8</sub>PI and 2 mM ATP as substrates, 5 mM Mg<sup>2+</sup>, in 50 mM Tris HCl, pH 7.5, with 3 mM of the deoxy-PI analogs.

<sup>b</sup> For several of the samples run in duplicate, the error in determining the specific activity was <10%.